MATRIX IDENTITIES CONNECTED WITH THE JACOBIAN CONJECTURE

T. Krasiński

1. Introduction

Let $f \in \mathbb{C}[x,y]$ be a polynomial in two variables with complex coefficients. f is said to be a *component of an automorphism* if there exists a polynomial $g \in \mathbb{C}[x,y]$ such that $F := (f,g) : \mathbb{C}^2 \longrightarrow \mathbb{C}^2$ is a polynomial automorphism of \mathbb{C}^2 (i.e. there exists F^{-1} and it is also polynomial).

In turn, f is said to be a Keller's component if there exists $g \in \mathbb{C}[x,y]$ such that the jacobian Jac(f,g) of the mapping $(f,g): \mathbb{C}^2 \longrightarrow \mathbb{C}^2$ is a non-zero constant. Then g is called associated with f and the pair (f,g) a Keller's mapping.

It is well-known that the famous (and unsolved so far) jacobian conjecture can be formulated in the following way:

(1) If f is a Keller's component, then f is a component of an automorphism.

It is easy to show that it sufficies to prove (1) for $f \in \mathbb{C}[x, y]$ satisfying the following three conditions which will be called in the sequel the main assumptions:

1. f is of the form

$$f(x,y) = y^m + a_1(x)y^{m-1} + \ldots + a_m(x), \ m \ge 2, \ \deg a_i \le i, \ i = 1, \ldots, m,$$

- 2. for every $\lambda \in \mathbb{C}$, $f \lambda$ is reduced in $\mathbb{C}[x, y]$,
- 3. $\deg_u f'_x > 0$.

In the paper, using one of the main results in [3], we give conditions for f (which satisfy the main assumptions) equivalent to the fact that f is a component of an automorphism and a Keller's component, respectively (in the latter case under some additional assumptions on f). These conditions are differential matrix identities connected with the matrix

$$M = \begin{bmatrix} a'_1 & a'_2 & \dots & a'_m \\ & \ddots & & & \ddots \\ & & a'_1 & a'_2 & \dots & a'_m \\ m & (m-1)a_1 & \dots & a_{m-1} \\ & \ddots & & \ddots & \\ & & m & (m-1)a_1 & \dots & a_{m-1} \end{bmatrix} \right\} (m-1) - \text{rows},$$

Typeset by AMS-TEX

which determinant is equal, up to a non-zero constant, to the resultant $R_y(f'_x, f'_y)$ of the partial derivatives f'_x, f'_y of f with respect to the variable g. Precisely, if $a'_1 = 0, \ldots, a'_k = 0$, $a'_{k+1} \neq 0$, then

$$\det M = (-1)^{k(m+1)} m^k R_y(f_x', f_y').$$

In the sequel, for a square matrix $A = [a_{ij}]_{1 \leq i,j \leq n}$, $a_{ij} \in \mathbb{C}[x]$, we shall denote by $|a_{ij}|_{x^k}$ k-th derivative of the determinant of A.

2. Component of an automorphism

Theorem A. Let $f \in \mathbb{C}[x,y]$ satisfy the main assumptions. Then f is a component of an automorphism if and only if the following identities hold (A)

$$\sum_{\substack{1,\ldots,r_i=1\\1,\ldots,s_j=1\\r_$$

for k = 1, ..., m-1, i, j = 0, 1, ..., k-1, i+j = k-1 (the above determinant is obtained from the matrix M by replacing the rows with indices $r_1, ..., r_i, (m-1)+s_1, ..., (m-1)+s_j$ by appropriate versors [0, ..., 0, 1, 0, ..., 0] with the unit at $m+r_1-1, ..., m+r_i-1, m+s_1-1, ..., m+s_j-1$ place, respectively).

Remark 1. In particular, for k = 1 we have the identity

$$\begin{vmatrix} a'_1 & a'_2 & \dots & a'_m \\ & \ddots & & & \ddots \\ & & a'_1 & a'_2 & \dots & a'_m \\ m & (m-1)a_1 & \dots & a_{m-1} \\ & \ddots & & & \ddots \\ & & m & (m-1)a_1 & \dots & a_{m-1} \end{vmatrix}_x \equiv 0,$$

and for k = 2 we have two identities

Remark 2. Obviously in (A) we may additionally assume that $r_p \neq s_q$, p = $1, \ldots, i, \quad q = 1, \ldots, j.$

Remark 3. The identities (A) give also a set of equations describing coefficients of components of polynomial automorphisms in the space of all coefficients of a polynomial in two variables of degree m.

Proof of Theorem A. Let f satisfy the main assumptions. Denote by u, vnew variables and consider the polynomials $f'_x - u, f'_y - v \in \mathbb{C}[x, y, u, v]$. Let

$$Q(x, u, v) = Q_N(u, v)x^N + Q_{N-1}(u, v)x^{N-1} + \dots + Q_0(u, v)$$

be the resultant of these polynomials with respect to the variable y. By definition (2)

$$Q(x, u, v) = R_y(f'_x - u, f'_y - v) = C \begin{vmatrix} a'_1 & a'_2 & \dots & a'_m - u \\ & \ddots & & & \ddots \\ & & a'_1 & a'_2 & \dots & a'_m - u \\ m & (m-1)a_1 & \dots & a_{m-1} - v \\ & & \ddots & & & \ddots \\ & & m & (m-1)a_1 & \dots & a_{m-1} - v \end{vmatrix},$$

where C is a non-zero constant. It is easy to check that $Q(x, u, v) \not\equiv 0$. Moreover, by main assumptions 1. and 2. we have that $R_y(f'_x, f'_y) \not\equiv 0$, i.e.

$$Q(x,0,0) \not\equiv 0.$$

In paper [3], Th.3.4 (see also [2], Th.5.2, where the inequality $\operatorname{ord}_{(0)}Q_N \neq 0$ should be replaced by the equality $\operatorname{ord}_{(0)}Q_N = 0$) the following characterization, in terms of Q, of a component of an automorphism was given:

If $f \in \mathbb{C}[x,y]$, $0 < \deg f - 1 = \deg_y f'_y$ and $0 < \deg_y f'_x$, then f is a component of an automorphism if and only if $\operatorname{ord}_{(0,0)}Q_0 = 0$ and $\operatorname{ord}_{(0,0)}Q_i \ge i$ for $i = 1, \ldots, N$.

Actually, Theorem 3.4 in [3] is slightly weaker. However, from Remark 11.2 and Theorem 9.4 of that paper we easily obtain the above form of the theorem.

From the main assumptions on f it follows that f satisfies the general assumptions of the cited theorem. So, f is a component of an automorphism if and only if $\operatorname{ord}_{(0,0)}Q_0 = 0$ and $\operatorname{ord}_{(0,0)}Q_i \geq i$ for $i = 1, \ldots, N$. These conditions are equivalent to the following

$$Q_0(0,0) \neq 0, \ \frac{\partial^{i+j}Q_k}{\partial u^i \partial v^j}(0,0) = 0, \ k = 1, \dots, N, \ i,j = 0,1,\dots,k-1, \ i+j \leq k-1.$$

Since for every k = 1, ..., N and $i, j \in \mathbb{N}$ we have

$$\frac{\partial^{i+j+k}Q}{\partial u^i\partial v^j\partial x^k}(x,0,0) = \frac{N!}{(N-k)!}\frac{\partial^{i+j}Q_N}{\partial u^i\partial v^j}(0,0)x^{N-k} + \dots + k!\frac{\partial^{i+j}Q_k}{\partial u^i\partial v^j}(0,0),$$

then (4) (taking into account (3)) is equivalent to the following:

(5)
$$\frac{\partial^{i+j+k}Q}{\partial u^i \partial v^j \partial x^k}(x,0,0) \equiv 0, \ k=1,\ldots,N, \ i,j=0,1,\ldots,k-1, \ i+j=k-1.$$

From form (2) of Q it follows that we have obtained (A) for k = 1, ..., N. Obviously, if $N \leq m-1$ then from (5) we obtain that the identities in (A) are satisfied for k = N+1, ..., (m-1). In turn, if m-1 < N then directly from identities (A) it follows that they also hold for k = m, ..., N. Hence (5) is all so satisfied for k = m, ..., N.

In consequence, (5) is equivalent to (A) in every case.

3. Keller's component

Theorem B. Let $f \in \mathbb{C}[x,y]$ satisfy the main assumptions. Then f is a Keller's component for which there exists an associated component g such

that $\deg_y g \leq m$ if and only if the following identities hold

for k = 0, ..., m - 1, where for k = 0, 1 the above identities mean

Proof. 1. Let f be a Keller's component for which there exists an associated component g such that $\deg_y g \leq m$. Then $g(x,y) = b_0(x)y^m + b_1(x)y^{m-1} + \cdots + b_m(x)$, $b_i \in \mathbb{C}[x]$, $i = 0, \ldots, m$. It is easy to show that we can assume that $b_0 \equiv 0$. In fact, it is easy to see that from the jacobian condition $\operatorname{Jac}(f,g) = \operatorname{const.} \neq 0$, it follows that $b'_0(x) \equiv 0$. Hence $b_0 \equiv 0$ or $b_0 \equiv \operatorname{const.} \neq 0$. In the first case the assertion is satisfied, whereas in the second it sufficies to replace g by $g - b_0 f$.

So, in the sequel we shall assume that

$$g(x,y) = b_1(x)y^{m-1} + \dots + b_m(x), \ b_i \in \mathbb{C}[x], \ i = 1,\dots, m.$$

Additionally, we shall assume that Jac(f,g) = 1. Then

$$1 = f'_x g'_y - f'_y g'_x = (a'_1 y^{m-1} + a'_2 y^{m-2} + \dots + a'_m)((m-1)b_1 y^{m-2} + \dots + b_{m-1})$$
$$- (my^{m-1} + (m-1)a_1 y^{m-2} + \dots + a_{m-1})(b'_1 y^{m-1} + \dots + b'_m).$$

Comparing the coefficients of the both sides of this equality we obtain the following system of polynomial equations

Considering $b_i, b'_j, i = 1, ..., m-1, j = 1, ..., m$ as unknows we have a square system of 2m-1 equations. Simple calculations give that for the determinant D of the system we have

$$D = (-1)^{\frac{m(m-1)}{2} - 1} m! \begin{vmatrix} a'_1 & a'_2 & \dots & a'_m \\ & \ddots & & & \ddots \\ & & a'_1 & a'_2 & \dots & a'_m \\ m & (m-1)a_1 & \dots & a_{m+1} \\ & \ddots & & & \ddots \\ & & m & (m-1)a_1 & \dots & a_{m+1} \end{vmatrix}$$
$$= (-1)^{\frac{m(m-1)}{2} - 1} m! CR_y(f'_x, f'_y),$$

where C is a non-zero constant. Since f is a Keller's component $R_y(f'_x, f'_y) = \text{const.} \neq 0$. In consequence, D is a non-zero constant equal to $(-1)^{\frac{m(m-1)}{2}-1}m!R$,

where $R = CR_y(f'_x, f'_y)$. So, by the Cramer's rule we get formulae for solutions of this system. Simple calculations give

$$(7)$$

$$i = 1, \dots, m - 1,$$

$$b_1' = 0,$$

$$(8)
 j = 2, \dots, m.$$

From these formulae it follows that identities (B) are fulfilled for $k = 1, \dots, m-1$ 1. Moreover, from the fact that $R_y(f'_x, f'_y) = \text{const.} \neq 0$ we obtain identity (B) for k = 0.

2. Assume that identities (B) hold for a polynomial f. From the main assumptions 1. and 2. we have that $R_y(f'_x, f'_y) \neq 0$. So, by the identity (B) for k=0 we obtain $R_y(f'_x, f'_y) = \text{const.} \neq 0$. Hence $\det M = CR_y(f'_x, f'_y)$, C=const. $\neq 0$, is also a non-zero constant. Denote it by R. Then we define polynomials b_i , i = 1, ..., m-1 by formulae (7) and polynomials b_j , j = $1, \ldots, m$ by formulae (8). Additionally, we put $b_m = \int \widetilde{b}_m dx$ (an arbitrary indefinite integral of b_m). From identities (B) we have

(9)
$$b_i' = \widetilde{b}_i, \quad i = 1, \dots, m.$$

Besides, from the formulae for b_i and b_j , i = 1, ..., m - 1, j = 1, ..., m, it follows that they are solutions of system (6). Hence and from (9) we obtain that for $g(x, y) = b_1(x)y^{m-1} + \cdots + b_m(x)$ we have Jac(f, g) = 1.

Remark 4. From the proof it follows that the assumption $\deg a_i \leq i, i = 1, \ldots, m$, in the theorem is superfluous.

Remark 5. A characterization of polynomials, associated with a given Keller's component, can be find in [1].

4 Corollaries

Corollary 1. Let $l \in \mathbb{N}$. If for every $m \leq l$ and arbitrary polynomials $a_1, \ldots, a_m \in \mathbb{C}[x]$ such that $\deg a_i \leq i$, $i = 1, \ldots, m$, identities (B) imply identities (A) then the jacobian conjecture is true for polynomials of degree $\leq l$.

Proof. Let (f,g) be an arbitrary Keller's mapping such that $m = \deg f \leq l$ and $n = \deg g \leq l$. Using a linear change of coordinates in \mathbb{C}^2 we may assume that both f and g satisfy the main assumptions. If $m \geq n$, then from Theorem B we get that the coefficients a_1, \ldots, a_m of f satisfy (B). By assumption we obtain that (A) hold. Hence, by Theorem A, f is a component of an automorphism. It easily implies that (f,g) is a polynomial automorphism.

In the case n > m considerations are similar.

From the above corollary we obtain directly

Corollary 2. If for arbitrary polynomials $a_1, \ldots, a_m \in \mathbb{C}[x]$ such that $\deg a_i \leq i$, $i = 1, \ldots, m$, identities (B) imply identities s (A), then the jacobian conjecture is true.

Before we pass to concrete cases let us notice

Remark 6. Without loss of generality we may add to the main assumptions that $a_1 \equiv 0$. It can be obtained by applying to f the known triangular automorphism $(x,y) \longmapsto (x,y-\frac{1}{n}a_1(x))$ of \mathbb{C}^2 .

Let us now consider two cases:

- 1. m = 2. It is easily seen then in this case identities (A) and (B) are equivalent.
- 2. m=3. In this case, assuming that $a_1 \equiv 0$ (see Remark 6) identities (A) are as follows

(10)
$$((a_2')^2 a_2 + 3(a_3')^2)' \equiv 0, \quad a_2'' \equiv 0, \quad a_3''' \equiv 0,$$

whereas identities (B) are

(11)
$$((a_2')^2 a_2 + 3(a_3')^2)' \equiv 0, \quad a_2'' \equiv 0, \quad a_3'' \equiv 0,$$

It is easy to show that conditions (10) and (11) are equivalent. From these cases and Corollary 1 we obtain

Corollary 3. The jacobian conjecture is true for polynomials of degree ≤ 3 .

References

- 1. Z. Charzyński, J. Chądzyński and P. Skibiński, *A contribution to Keller's jacobian conjecture II*, Deformations of Mathematical Structures, Kluwer Academic Publishers, Dordrecht–Boston –London, 1989, pp. 133–140.
- J. Chądzyński and T. Krasiński, Sur l'exposant de Lojasiewicz à l'infini pour les applications polynomiales de C² dans C² et les composantes des automorphi smes polynomiaux de C², C. R. Acad. Sci. Paris, Ser. I 315 (1992), 1399–1402.
- 3. _____, On the Lojasiewicz exponent at infinity for polynomial mappings of \mathbb{C}^2 into \mathbb{C}^2 and components of polynomial automorphisms of \mathbb{C}^2 , Ann. Polon. Math. 57 (1992), 291–302.

Łódź, December 1993

Institute of Mathematics, University of Łódź, S. Banacha 22 , 90-238 Łódź, Poland $E\text{-}mail\ address$: krasinsk@plunlo51.bitnet