A NOTE ON ELLIPTIC PLANE CURVES WITH FIXED *j*-INVARIANT

RAHUL PANDHARIPANDE

10 May 1995

1. Summary

Let N_d be the number of irreducible, reduced, nodal, degree *d* rational plane curves passing through 3d - 1 general points in the complex projective plane \mathbf{P}^2 . The numbers N_d satisfy a beautiful recursion relation ([K-M], [R-T]):

 $N_1 = 1$

$$\forall d > 1, \quad N_d = \sum_{i+j=d, \ i,j>0} N_i N_j \left(i^2 j^2 \binom{3d-4}{3i-2} - i^3 j \binom{3d-4}{3i-1} \right)$$

Let $E_{d,j}$ be the number of irreducible, reduced, nodal, degree d, elliptic plane curves with fixed *j*-invariant *j* passing through 3d - 1 general points \mathbf{P}^2 . $E_{d,j}$ is defined for $d \geq 3$ and $\infty \neq j \in \overline{M}_{1,1}$. In this note, the following relations are established:

$$\forall j \neq 0, 1728, \infty, \qquad E_{d,j} = \binom{d-1}{2} N_d, \\ j = 0, \qquad E_{d,0} = \frac{1}{3} \binom{d-1}{2} N_d, \\ j = 1728, \qquad E_{d,1728} = \frac{1}{2} \binom{d-1}{2} N_d$$

If $d \equiv 0 \mod 3$, then $3 \not\mid \binom{d-1}{2}$. Since $E_{3\hat{d},0}$ is an integer, $N_{3\hat{d}} \equiv 0 \mod 3$ for $\hat{d} \geq 1$. In fact, a check of values in [D-I] shows $N_d \equiv 0 \mod 3$ if and only if $d \equiv 0 \mod 3$ for $3 \leq d \leq 12$. P. Aluffi has calculated $E_{3,j}$ for $j < \infty$ in [A]. Aluffi's results agree with the above formulas.

Thanks are due to Y. Ruan for discussions on Gromov-Witten invariants and quantum cohomology. The question of determining the numbers $E_{d,j}$ was first considered by the author in a conversation with Y. Ruan.

2. Kontsevich's Space of Stable Maps

2.1. The Quasi-Projective Subvarieties $U_C(\Gamma, c, \overline{w})$, $U_{j=\infty}(\Lambda, \overline{w})$. Fix $d \geq 3$ for the entire paper. Let C be a nonsingular elliptic curve or an irreducible, nodal rational curve of arithmetic genus 1. Consider the coarse moduli space of 3d - 1pointed stable maps from C to \mathbf{P}^2 of degree $d \geq 3$, $\overline{M}_{C,3d-1}(\mathbf{P}^2, d)$. For convenience, the notation $\overline{M}_C(d) = \overline{M}_{C,3d-1}(\mathbf{P}^2, d)$ will be used. Let $S_d = \{1, 2, \ldots, 3d - 1\}$ be the marking set. Constructions of $\overline{M}_C(d)$ can be found in [Al], [K], [P]. Let Γ be a tree consisting of a distinguished vertex $c, k \geq 0$ other vertices v_1, \ldots, v_k , and 3d - 1 marked legs. Let $0 \leq e \leq d$. Weight the vertex c by e. Let w_1, \ldots, w_k be non-negative integral weights of the vertices v_1, \ldots, v_k satisfying

$$e + w_1 + \dots + w_k = d$$

Denote the weighting by $\overline{w} = (e, w_1, \dots, w_k)$. The marked, weighted tree with distinguished vertex $(\Gamma, c, \overline{w})$ is stable if the following implication holds for all $1 \le i \le k$:

$$w_i = 0 \Rightarrow valence(v_i) \ge 3.$$

Two marked, weighted trees with distinguished vertex $(\Gamma, c, \overline{w})$ and $(\Gamma', c', \overline{w}')$ are isomorphic if there is an isomorphism of marked trees $\Gamma \to \Gamma'$ sending c to c' and respecting the weights.

A quasi-projective subvariety $U_C(\Gamma, c, \overline{w})$ of $\overline{M}_C(d)$ is associated to each isomorphism class of stable, marked, weighted graph with distinguished vertex $(\Gamma, c, \overline{w})$. $U_C(\Gamma, c, \overline{w})$ consists of stable maps $\mu : (D, p_1, \ldots, p_{3d-1}) \to \mathbf{P}^2$ satisfying the following conditions. The domain D is equal to a union:

$$D = C \cup \mathbf{P}_1^1 \cup \cdots \cup \mathbf{P}_k^1.$$

The marked, weighted dual graph with distinguished vertex of the map μ is isomorphic to $(\Gamma, c, \overline{w})$. The distinguished vertex of the dual graph of μ corresponds to the (unique) component of D isomorphic to C. Weights of the dual graph of μ are obtained by the degree of μ on the components. Note $U_C(\Gamma, c, \overline{w}) = \emptyset$ if and only if e = 1.

Let $(\Gamma, c, \overline{w})$ be a stable, marked, weighted tree with distinguished vertex. Assume $e \neq 1$. The dimension of $U_C(\Gamma, c, \overline{w})$ is determined as follows. If $e \geq 2$, then

$$\dim U_C(\Gamma, c, \overline{w}) = 6d - 2 - k.$$

If e = 0, then

$$\dim U_C(\Gamma, c, \overline{w}) = 6d - k$$

(where k is the number of non-distinguished vertices of Γ). These calculations are straightforward.

Let C be a nonsingular elliptic curve. Every stable map in $\overline{M}_C(d)$ has domain obtained by attaching a finite number of marked trees to C. By the definition of tree and map stability:

$$\bigcup_{(\Gamma,c,\overline{w})} U_C(\Gamma,c,\overline{w}) = \overline{M}_C(d).$$

Let C be an irreducible, 1-nodal rational curve. The quasi-projective varieties $U_C(\Gamma, c, \overline{w})$ do not cover $\overline{M}_C(d)$. The curve C can degenerate into a simple circuit of \mathbf{P}^1 's. Let Λ be a graph with 1 circuit (1st Betti number equal to 1, no self edges), $k \geq 1$ vertices v_0, \ldots, v_k , and 3d - 1 marked legs. Note the different vertex numbering convention. At least 2 vertices are required to make a circuit, so $k \geq 1$. Let w_0, w_1, \ldots, w_k be non-negative, integral weights summing to d. The marked, weighted graph with 1 circuit (Λ, \overline{w}) is stable if each zero weighted vertex has valence at least 3. A quasi-projective subvariety $U_C(\Lambda, \overline{w})$ of $\overline{M}_C(d)$ is associated to each isomorphism class of stable, marked, weighted graph with 1 circuit (Λ, \overline{w}). $U_C(\Lambda, \overline{w})$ consists of stable maps with marked, weighted dual graphs isomorphic to (Λ, \overline{w}). The union

$$\bigcup_{(\Gamma,c,\overline{w})} U_C(\Gamma,c,\overline{w}) \quad \cup \quad \bigcup_{(\Lambda,\overline{w})} U_C(\Lambda,\overline{w}) = \overline{M}_C(d)$$

holds by the definition of stability.

Finally, the dimensions of the loci $U_C(\Lambda, \overline{w})$ will be required. Let (Λ, \overline{w}) be a stable, marked, weighted graph with 1 circuit. Let c_1, \ldots, c_l be the unique circuit of vertices of Λ . Let e be the sum of the weights of the circuit vertices. $U_C(\Lambda, \overline{w}) = \emptyset$ if and only if e = 1. If $e \geq 2$, then

$$\dim U_C(\Lambda, \overline{w}) = 6d - 2 - k.$$

If e = 0, then

$$\dim U_C(\Lambda, \overline{w}) = 6d - k$$

(where k + 1 is the total number of vertices of Λ). Again, these results are straightforward.

2.2. The Component $\overline{W}_1(d)$. Let $\overline{M}_1(d) = \overline{M}_{1,3d-1}(\mathbf{P}^2, d)$ be Kontsevich's space of 3d-1-pointed stable maps from genus 1 curves to \mathbf{P}^2 . There is canonical morphism

$$\pi: \overline{M}_1(d) \to \overline{M}_{1,1}$$

obtained by forgetting the map and all the markings except $1 \in S_d$ (the 3d - 1 possible choices of marking in S_d all yield the same morphism π). Let $j \in \overline{M}_{1,1}$. By the universal properties of the moduli spaces, there is a canonical bijection

$$\overline{M}_{C_j}(d) \to \pi^{-1}(j)$$

where C_j is the elliptic curve (possibly nodal rational) with *j*-invariant *j*. When $j \in \overline{M}_{1,1}$ is automorphism-free, this bijection is an isomorphism. For j = 0, 1728, the scheme theoretic fiber $\pi^{-1}(j)$ is nonreduced. Define an open locus $W_1(d) \subset \overline{M}_1(d)$ by $[\mu : (D, p_1, \ldots, p_{3d-1}) \rightarrow \mathbf{P}^2] \in W_1(d)$ if and only if *D* is irreducible. By considering the natural tautological spaces over the universal Picard variety of degree *d* line bundles over $M_{1,3d-1}$, it is easily seen that $W_1(d)$ is a reduced, irreducible open set of dimension 6d - 1. Let $\overline{W}_1(d)$ be the closure of $W_1(d)$ in $\overline{M}_1(d)$.

3. A Deformation Result

Let Φ be a stable, marked, weighted tree with distinguished vertex c determined by the data: k = 1, $(e, w_1) = (0, d)$. There are 2^{3d-1} isomorphism classes of such Φ determined by the marking distribution. Let $j \in \overline{M}_{1,1}$. The dimension of $U_j(\Phi)$ is 6d - 1. A point $[\mu] \in U_j(\Phi)$ has domain $C_j \cup \mathbf{P}^1$. There are 3d - 1 dimensions of the map $\mu|_{\mathbf{P}^1} : \mathbf{P}^1 \to \mathbf{P}^2$. The incidence point $p = C_j \cap \mathbf{P}^1$ moves in a 1-dimensional family on \mathbf{P}^1 . The remaining 3d - 1 markings move in 3d - 1 dimensions on C_j and \mathbf{P}^1 (specified by the marking distribution). 6d - 1 = 3d - 1 + 1 + 3d - 1. A technical result is needed in the computation of the numbers $E_{d,j}$.

Lemma 1. Let $I(\Phi, j) = \overline{W}_1(d) \cap U_j(\Phi) \subset \overline{M}_1(d)$. The dimension of $I(\Phi, j)$ is bounded by dim $I(\Phi, j) \leq 6d - 3$.

Proof. Let $[\mu] \in I(\Phi, j)$ be a point. Let $D = C_j \cup \mathbf{P}^1$ be the domain of μ as above. The following condition will be shown to hold: the linear series on \mathbf{P}^1 determined by $\mu|_{\mathbf{P}^1}$ has vanishing sequence $\{0, \geq 2, *\}$ at the incidence point $p = C_j \cap \mathbf{P}^1$. The existence of a point with vanishing sequence $\{0, \geq 2, *\}$ is a 1-dimensional condition on the linear series. The condition that the incidence point p has this vanishing sequence is an additional 1-dimensional constraint on p. Therefore, the dimension of $I(\Phi, j)$ is at most 6d - 1 - 1 - 1 = 6d - 3. The vanishing sequence $\{0, \geq 2, *\}$ is equivalent to $d(\mu|_{\mathbf{P}^1}) = 0$ at p.

It remains to establish the vanishing sequence $\{0, \geq 2, *\}$ at p. This result is easily seen in explicit holomorphic coordinates. Let Δ_t be a disk at the origin in \mathbb{C} with coordinate t. Let $\eta : \mathcal{E} \to \Delta_t$ be a flat family of curves of arithmetic genus 1 satisfying:

(i.) $\eta^{-1}(0) \cong C_j$. (ii.) $\eta^{-1}(t \neq 0)$ is irreducible, reduced, and (at worst) nodal. For each $1 \leq i \leq d$, let $\mathcal{G}_i = \mathcal{H}_i \subset \mathcal{E}$ be the open subset of \mathcal{E} on which the morphism η is *smooth*. Consider the fiber product:

$$X = \mathcal{G}_1 \times_{\bigtriangleup_t} \cdots \times_{\bigtriangleup_t} \mathcal{G}_d \times_{\bigtriangleup_t} \mathcal{H}_1 \times_{\bigtriangleup_t} \cdots \times_{\bigtriangleup_t} \mathcal{H}_d$$

X is a nonsingular open set of the 2*d*-fold fiber product of \mathcal{E} over Δ_t . Let $Y \subset X$ be the subset of points $y = (g_1, \ldots, g_d, h_1, \ldots, h_d)$ where the two divisors $\sum g_i$ and $\sum h_i$ are linearly equivalent on the curve $\eta^{-1}(\eta(y))$. Y is a nonsingular divisor in X.

Let $p \in C_j = \eta^{-1}(0)$ be a nonsingular point of C_j . Certainly $p \in \mathcal{G}_i, \mathcal{H}_i$ for all *i*. Let $\gamma : \Delta_t \to \mathcal{E}$ be any local holomorphic section of η such that $\gamma(0) = p$. Let *V* be a local holomorphic field of vertical tangent vectors to \mathcal{E} on an open set containing *p*. The section γ and the vertical vector field *V* together determine local holomorphic coordinates (t, v) on \mathcal{E} at *p*. Let $\phi_V : \mathcal{E} \times \mathbb{C} \to \mathcal{E}$ be the holomorphic flow of *V* defined locally near $(p, 0) \in \mathcal{E} \times \mathbb{C}$. The coordinate map

$$\psi: (t,v) \to \mathcal{E}$$

is determined by $\psi(t, v) = \phi_V(\gamma(t), v)$.

Local coordinates on X near the point $x_p = (p, \ldots, p, p, \ldots, p) \in X$ are given by

$$(t, v_1, \ldots, v_d, w_1, \ldots w_d).$$

The coordinate map is determined by:

$$\psi_X(t, v_1, \dots, v_d, w_1, \dots, w_d) = (\psi(t, v_1), \dots, \psi(t, v_d), \psi(t, w_1), \dots, \psi(t, w_d)) \in X.$$

Note $x_p \in Y$. Let $f(t, v_1, \ldots, v_d, w_1, \ldots, w_d)$ be a local equation of Y at x_p . Since f is identically 0 on the line $(t, 0, \ldots, 0, 0, \ldots, 0)$,

(1)
$$\forall k \ge 0, \quad \frac{\partial^k f}{\partial t^k}|_{x_p} = 0.$$

The tangent directions in the plane t = 0 correspond to divisors on the fixed curve C_i . Here, it is well know (up to \mathbb{C}^* - factor)

(2)
$$\frac{\partial f}{\partial v_i}|_{x_p} = +1, \quad \frac{\partial f}{\partial w_i}|_{x_p} = -1.$$

Equations (1) and (2) are the only properties of f that will be used.

Let $\hat{\eta} : \hat{\mathcal{E}} \to \triangle_t$ be the family obtained by blowing-up \mathcal{E} at p and adding 3d - 1-marking. Let $\mu : \hat{\mathcal{E}} \to \mathbf{P}^2$ be a morphism. Let $\hat{\eta}^{-1}(0) = D = C_j \cup \mathbf{P}^1$. Assume the following conditions are satisfied:

(i.) μ , $\hat{\eta}$, and the 3d-1 markings determine a family of Kontsevich stable pointed maps to \mathbf{P}^2 .

- (ii.) The markings of D are distributed according to Φ .
- (iii.) $deg(\mu|_{C_j}) = 0, deg(\mu|_{\mathbf{P}^1}) = d.$

Let L_1 , L_2 be general divisors of $\mu^*(\mathcal{O}_{\mathbf{P}^2}(1))$ that each intersect \mathbf{P}^1 transversely at d distinct points. For $1 \leq \alpha \leq 2$, L_α breaks into holomorphic sections $s_{\alpha,1} + \ldots + s_{\alpha,d}$ of $\hat{\eta}$ over a holomorphic disk at $0 \in \Delta_t$. These sections $s_{\alpha,i}$ $(1 \leq \alpha \leq 2, 1 \leq i \leq d)$ determine a map $\lambda : \Delta_t \to Y$ locally at $0 \in \Delta_t$. Let an affine coordinate on \mathbf{P}^1 be given by ξ corresponding to the normal direction

(3)
$$\frac{d\gamma}{dt}|_{t=0} + \xi \cdot V(p)$$

Let $s_{1,i}(0) = \nu_i \in \mathbb{C} \subset \mathbf{P}^1$, $s_{2,i}(0) = \omega_i \in \mathbb{C} \subset \mathbf{P}^1$ be given in terms of the affine coordinate ξ . The map λ has the form

$$\lambda(t) = (t, \nu_1 t, \dots, \nu_d t, \omega_1 t, \dots, \omega_d t)$$

to first order in t (written in the coordinates determined by ψ_X). Equations (1), (2), and the condition $f(\lambda(t)) = 0$ implies

(4)
$$\sum_{i=1}^{d} \nu_i = \sum_{1}^{d} \omega_i.$$

 $L_1 \cap \mathbf{P}^1$ is a degree *d* polynomial with roots at ν_i . Condition (4) implies that the sums of the roots (in the coordinates (3)) of general elements of the linear series $\mu|_{\mathbf{P}^1}$ are the same. Therefore, a constant *K* exists with the following property. If

$$\beta_0 + \beta_1 \xi + \ldots + \beta_{d-1} \xi^{d-1} + \beta_d \xi^d$$

is an element of the linear series $\mu|_{\mathbf{P}^1}$, then $\beta_{d-1} + K \cdot \beta_d = 0$. The vanishing sequence at $\xi = \infty$ is therefore $\{0, \ge 2, *\}$. The point $\xi = \infty$ is the intersection $C_j \cap \mathbf{P}^1$.

Suppose $\tilde{\eta} : \tilde{\mathcal{E}} \to \Delta_t$ is obtained from \mathcal{E} by a sequence of n blow-ups over p. The fiber $\tilde{\eta}^{-1}(0)$ is assumed to be C_j union a chain of \mathbf{P}^1 's of length n. Each blow-up occurs in the exceptional divisor of the previous blow-up. Let \mathbf{P} denote the extreme exceptional divisor. Let $\mu : \tilde{\mathcal{E}} \to \mathbf{P}^2$ be of degree d on \mathbf{P} and degree 0 on the other components of the special fiber $\tilde{\eta}^{-1}(0)$. Let there be 3d - 1 markings as before. It must be again concluded that the linear series on \mathbf{P} has vanishing sequence $\{0, \geq 2, *\}$ at the node.

Let γ be section of η such that the lift of γ to $\tilde{\eta}$ meets P. Let the coordinates (t, v)on \mathcal{E} be determined by this γ (and any V). An affine coordinate ξ is obtained on \mathbf{P} in the follow manner. Let γ_{ξ} be the section of η determined in (t, v) coordinates by

$$\gamma_{\xi}(t) = (t, \xi t^n).$$

Let $\tilde{\gamma}_{\xi}$ be the lift of γ_{ξ} to a section of $\tilde{\eta}$. The association

$$\mathbb{C} \ni \xi \mapsto \tilde{\eta}(0) \in \mathbf{P}$$

is an affine coordinate on \mathbf{P} . Let L_1, L_2 be divisors in the linear series μ intersecting \mathbf{P} transversely. As before, L_{α} breaks into holomorphic sections $s_{\alpha,1}$. Let $s_{1,i} = \nu_i \in \mathbb{C} \subset \mathbf{P}$, $s_{2,i} = \omega_i \in \mathbb{C} \subset \mathbf{P}$. As before, a map $\lambda : \Delta_t \to Y$ is obtained from the sections $s_{\alpha,i}$. In the coordinates determined by ψ_X ,

$$\lambda(t) = (t, \nu_1 t^n + O(t^{n+1}), \dots, \nu_d t^n + O(t^{n+1}), \omega_1 t^n + O(t^{n+1}), \dots, \omega_d t^n + O(t^{n+1})).$$

As before $f(\lambda(t)) = 0$. The term of leading order in t of $f(\lambda(t))$ is

$$\left(\sum_{i=1}^d \nu_i - \sum_{i=1}^d \omega_i\right) \cdot t^n.$$

This follows from equations (1) and (2). The vanishing sequence $\{0, \ge 2, *\}$ is obtained as before.

By definition, an element $[\mu] \in I(\Phi, j)$ can be obtained as the special fiber of family of Kontsevich stable maps where the domain is a smoothing of the node p. After resolving the singularity in the total space at the node p by blowing-up, a family $\tilde{\mathcal{E}}$ is obtained. The above results show the linear series on \mathbf{P}^1 has vanishing sequence $\{0, \geq 2, *\}$ at p. \Box

The markings play no role in the preceding proof. An identical argument establishes the following:

Lemma 2. Let Φ be a stable, marked, weighted tree with distinguished vertex satisfying e = 0 and $w_i = d$ for some i. Let k be the number of non-distinguished vertices of Φ . Let $j \in \overline{M}_{1,1}$. Let $I(\Phi, j) = \overline{W}_1(d) \cap U_j(\Phi)$. The dimension of $I(\Phi, j)$ is bounded by dim $I(\Phi, j) \leq 6d - k - 2$

Lemma 3. Let Ω be a stable, marked, weighted graph with 1 circuit. Let v_i be a non-circuit vertex with weight $w_i = d$ (this implies e = 0). Let k + 1 be the total number of vertices of Ω . Let $I(\Omega, \infty) = \overline{W}_1(d) \cap U_{\infty}(\Omega)$. The dimension of $I(\Omega, \infty)$ is bounded by dim $I(\Omega, \infty) \leq 6d - k - 2$.

The vanishing sequence $\{0, \geq 2, *\}$ condition reduces the dimensions of $U_C(\Phi), U_{\infty}(\Omega)$ by 2.

4. The NUMBERS $E_{d,i}$

The space of maps $\overline{M}_1(d)$ is equipped with 3d-1 evaluation maps corresponding to the marked points. For $i \in S_d$, let $e_i : \overline{W}_1(d) \to \mathbf{P}^2$ be the restriction of the i^{th} evaluation map to $\overline{W}_1(d)$. let $\mathcal{L}_i = e_i^*(\mathcal{O}_{\mathbf{P}^2})$ Let

$$Z = c_1(\mathcal{L}_1)^2 \cap \ldots \cap c_1(\mathcal{L}_{3d-1})^2$$

Let $\pi_{\overline{W}} : \overline{W}_1(d) \to \overline{M}_{1,1} \cong \mathbf{P}^1$ be the restriction of π to $\overline{W}_1(d)$. Let

 $T = c_1(\pi_{\overline{W}}^*(\mathcal{O}_{\mathbf{P}^1}(1))).$

Note $\overline{W}_1(d)$ is an irreducible, projective scheme of dimension 6d - 1. The top intersection of line bundles on $\overline{W}_1(d)$, $Z \cap T$, is an integer.

Lemma 4.

$$\forall j \neq 0, 1728, \infty, \qquad Z \cap T = E_{d,j} ,$$

 $j = 0, \qquad Z \cap T = 3 \cdot E_{d,0} ,$
 $j = 1728, \qquad Z \cap T = 2 \cdot E_{d,1728} .$

Proof. Via pull-back, lines in \mathbf{P}^2 yield representative classes of $c_1(\mathcal{L}_i)$. Therefore 3d-1 general points in \mathbf{P}^2 , $\overline{x} = (x_1, \ldots, x_{3d-1})$, determine a representative cycle $Z_{\overline{x}}$ of the the class Z. Let $\infty > j \in \overline{M}_{1,1}$. Let π_W be the restriction of π to $W_1(d)$. It is first established for a general representative $Z_{\overline{x}}$,

(5)
$$Z_{\overline{x}} \cap \pi_{\overline{W}}^{-1}(j) \subset \pi_{W}^{-1}(j).$$

The statement (5) is proven by considering the quasi-projective strata of $\overline{M}_{C_i}(d)$.

Note $\pi_W^{-1}(j)$ is the strata $U_{C_j}(\Gamma, c, \overline{w})$ where $(\Gamma, c, \overline{w})$ is the trivial, stable, marked, weighted tree with distinguished vertex. Assume now $(\Gamma, c, \overline{w})$ is not the trivial tree. By the equations for the dimension of $(\Gamma, c, \overline{w})$

$$dimU_{C_i}(\Gamma, c, \overline{w}) \le 6d - 3$$

unless e = 0 and k = 1, 2. Since the linear series determined by the evaluation maps are base point free, the general intersection (5) will miss all loci of dimension less than 6d - 2.

It remains to consider the trees $(\Gamma, c, \overline{w})$ where e = 0 and k = 1, 2. If k = 1, $(\Gamma, c, \overline{w}) = \Phi$ satisfies the conditions of Lemma (1). By Lemma (1),

$$\dim I(\Phi, j) \le 6d - 3$$

Hence, the general intersection (5) will miss all the loci $U_C(\Phi, c, (0, d))$.

If k = 2, there are two cases to consider. If there exists a vertex of weight d, then $(\Gamma, c, \overline{w}) = \Phi$ satisfies the conditions of Lemma (2). By Lemma (2),

$$\dim I(\Phi, j) \le 6d - 4.$$

If $w_1 + w_2 = d$ is a positive partition, then the image of every map $[\mu] \in U_C(\Gamma, c, \overline{w})$ is the union of two rational curves of degrees w_1 and w_2 . No such unions pass through 3d - 1 general points. The proof of claim (5) is complete.

For $\infty > j \neq 0,1728$, $\pi_W^{-1}(j)$ is a reduced, irreducible divisor of $W_1(d)$. Since the linear series determined by the evaluation maps are base point free, the general intersection cycle

is a reduced collection of $Z \cap T$ points. The general intersection cycle (6) also consists exactly of the reduced, nodal, degree d elliptic plane curves with j-invariant j passing through the points \overline{x} .

The argument for j = 0,1728 is identical except that $\pi_W^{-1}(0)$, and $\pi_W^{-1}(1728)$ are divisors in $W_1(d)$ with multiplicity 3, 2 respectively. These multiplicities arise from the extra automorphisms for j = 0,1728. Therefore the cycle (6) is a collection of

$$\frac{1}{3} \cdot Z \cap T,$$
$$\frac{1}{2} \cdot Z \cap T$$

triple and double points respectively. \Box

It remains to evaluate $Z \cap T$.

Lemma 5. $Z \cap T = \binom{d-1}{2} N_d$.

Proof. It is first established for a general representative $Z_{\overline{x}}$,

(7)
$$Z_{\overline{x}} \cap \pi_{\overline{W}}^{-1}(\infty) \subset \pi_{W}^{-1}(\infty)$$

The statement (7) is proven by considering the quasi-projective strata of $\overline{M}_{\infty}(d)$.

By arguments of Lemma (4), all the loci $U_{\infty}(\Gamma, c, \overline{w})$ where $(\Gamma, c, \overline{w})$ is not the trivial tree are avoided in the general intersection (7). Only the strata $U_{\infty}(\Lambda, \overline{w})$ remain to be considered. Let $k + 1 \ge 2$ be the total number of vertices of Λ . By the equations for the dimensions of $U_{\infty}(\Lambda, \overline{w})$,

$$\dim U_{\infty}(\Lambda, \overline{w}) \le 6d - 2 - k \le 6d - 3$$

unless all the circuit vertices have weight zero. If all circuit vertices have weight zero, $k + 1 \ge 3$. Now

$$\dim U_{\infty}(\Lambda, \overline{w}) \le 6d - k \le 6d - 3$$

unless k = 2.

Only one stable, marked, weighted, graph with 1-circuit (Λ, \overline{w}) need be considered. Vertices c_1, c_2 form a weightless circuit. Vertex v_3 is connected to c_2 and $w_3 = d$. $(\Lambda, \overline{w}) = \Omega$ satisfies the conditions of Lemma (3). Therefore,

$$\dim I(\Omega, \infty) \le 6d - 4.$$

Claim (7) is now proven.

The divisor $\pi_W^{-1}(\infty)$ is reduced and irreducible in $W_1(d)$. As above,

is a reduced collection of $Z \cap T$ points. The general intersection cycle (8) also consists exactly of degree d maps of the 1-nodal rational curve C_{∞} passing through \overline{x} . The image of such a map must be one of the N_d degree d, nodal, rational plane curves passing through \overline{x} . The number of distinct birational maps (up to isomorphism) from C_{∞} to a $\binom{d-1}{2}$ -nodal plane curve is exactly $\binom{d-1}{2}$. Therefore, $Z \cap T = \binom{d-1}{2}N_d$. \Box

References

- [Al] V. Alexeev, Moduli Spaces $M_{g,n}(W)$ For Surfaces, preprint 1994.
- [A] P. Aluffi, How Many Smooth Plane Cubics With Given j-Invariant Are Tangent To 8 Lines In General Position?, Contemporary Mathematics 123, 1991, p.15-29.
- [D-I] P. Di Francesco and C. Itzykson, *Quantum Intersection Rings*, preprint 1994.
- [K] M. Kontsevich, Enumeration of Rational Curves Via Torus Actions, preprint 1994.
- [K-M] M. Kontsevich and Y. Manin, Gromov-Witten Classes, Quantum Cohomology, and Enumerative Geometry, preprint 1994.
- [P] R. Pandharipande, Notes On Kontsevich's Compactification Of The Moduli Space Of Maps, Course Notes, 1995.

Department of Mathematics, University of Chicago

rahul@math.uchicago.edu