A REMARK ON VIRTUAL ORIENTATIONS FOR COMPLETE INTERSECTIONS

VADIM SCHECHTMAN

The aim of this note is to give a simple definition of genus zero *virtual orientation classes* (or *fundamental classes*) for projective complete intersections or, more generally, complete intersections in convex varieties, and to prove a *push forward formula* (see Lemmas 1 and 2 below) for them.

1. Let X be a smooth complex projective variety. Let X_{β} denote the stack of stable maps $f: C \longrightarrow X$ of genus g curves with n marked points, such that $f_*([C]) = \beta \in H_2(X)$. Here n and g will be fixed and suppressed from the notations. Let $\pi_{\beta}: \tilde{X}_{\beta} \longrightarrow X_{\beta}$ be the universal curve, and $\psi_{\beta}: \tilde{X}_{\beta} \longrightarrow X$ be the canonical map.

A vector bundle N over X is called *convex* if $R^1\pi_{\beta*}\psi_{\beta}^*N=0$ for all β . In this case we will use the notation N_{β} for the vector bundle $\pi_*\psi^*N$ over X_{β} . The variety X is called convex if the tangent bundle \mathcal{T}_X is convex.

Example. Let $X = \mathbb{P}^N$, and g = 0. Then X is convex, and any bundle of the form $N = \bigoplus \mathcal{O}(l_a)$ $(l_a \geq 0)$ is convex.

2. Let X be a convex variety and N a convex bundle over X. Let $s: X \longrightarrow N$ be a regular (i.e. transversal to the zero section) section of $N, i: Y := s^{-1}(0) \hookrightarrow X$ the subscheme of its zeros.

Set $Y_{\beta} := \coprod Y_{\gamma}$, the disjoint union over all $\gamma \in H_2(Y)$ mapping to β . The section s induces the section $s_{\beta} : Y_{\beta} \longrightarrow N_{\beta}$, and we have $Y_{\beta} = s_{\beta}^{-1}(0)$. We denote by i_{β} the embedding $Y_{\beta} \hookrightarrow X_{\beta}$.

Definition 1. The virtual dimension $\dim^{virt}(Y_{\beta})$ is the number $\dim(X_{\beta}) - \operatorname{rk}(N_{\beta})$.

The section s is not necessarily regular. If it is, then the virtual dimension of Y_{β} coincides with its usual dimension.

3. Construction. Our goal is to define certain class in the homology Chow group $A_{\dim^{virt}(Y_{\beta})}(Y_{\beta})$. We will use the key construction from the intersection theory, [F], 6.1. We have a cartesian square

$$\begin{array}{ccc} Y_{\beta} & \xrightarrow{i_{\beta}} & X_{\beta} \\ i_{\beta} \downarrow & & \downarrow s_{d} \\ X_{\beta} & \xrightarrow{s_{\beta}} & N_{\beta} \end{array}$$

where s_0 denotes the zero section. Let C be the normal cone of Y_{β} in X_{β} . We have the canonical closed embedding $C \hookrightarrow i_{\beta}^* N_{\beta}$, whence the class $[C] \in A_d(i_{\beta}^* N)$ where d =

 $\dim(C) = \dim(X_{\beta})$. We set by definition

$$[Y_{\beta}]^{virt} := (p^*)^{-1}([C]).$$

Here $p^*: A_*(Y_\beta) \xrightarrow{\sim} A_{*+\mathrm{rk}(N)}(i_\beta^*N)$ is the isomorphism induced by the projection $p: i_\beta^*N \longrightarrow Y_\beta$.

4. Push forward formula. Lemma 1. We have

$$i_{\beta*}([Y_{\beta}]^{virt}) = [X_{\beta}] \cap e(N_{\beta})$$

where e denotes the Euler (top Chern) class.

This formula was written down as a conjecture by Yu.I.Manin (talk at MPI, February 1997).

Proof. We can identify $s_0: X_{\beta} \longrightarrow N_{\beta}$ with the embedding of the scheme of zeros of the regular diagonal section of the vector bundle $N_{\beta} \times N_{\beta} \longrightarrow N_{\beta}$, and then apply [F], 6.3.4. \square

5. Iteration. Let M be a convex vector bundle over Y, and $j: Z \hookrightarrow Y$ be the subscheme of zeros of a regular section $t: Y \longrightarrow M$. We set by definition,

$$\dim^{virt}(Z_{\beta}) := \dim^{virt}(Y_{\beta}) - \operatorname{rk}(M_{\beta}).$$

We define

$$[Z_{\beta}]^{virt} \in A_{\dim^{virt}(Z_{\beta})}(Z_{\beta})$$

by the same construction as in no. 3, with $[X_{\beta}]$ replaced by $[Y_{\beta}]^{virt}$.

If $M = i^*N'$ and t is the restriction of a regular section $s': X \longrightarrow N'$ then Z may be defined as the subscheme of zeros of the regular section $s \oplus s': X \longrightarrow N \oplus N'$, and the definition of $[Z_{\beta}]^{virt}$ at one step using $s \oplus s'$ coincides with the previous one (which used first s, then t).

We have a generalization of Lemma 1:

6. Push forward formula. II. Lemma 2. We have

$$j_{\beta*}([Z_{\beta}]^{virt}) = [Y_{\beta}]^{virt} \cap e(M_{\beta}).$$

The proof is the same as in Lemma 1.

7. The above constructions have obvious equivariant versions.

I am grateful to A.Goncharov and especially to Yu.I.Manin, for very useful discussions.

References

[F] W.Fulton, Intersection theory, Springer-Verlag, Berlin et al., 1984.

 ${\it Max-Planck-Institut}$ für Mathematik, Gottfried-Claren-Strasse 26, 53225 Bonn, Germany

 $E\text{-}mail\ address{:}\ \mathtt{vadik@mpim-bonn.mpg.edu}$