AN ELLIPTIC CONIC BUNDLE IN \mathbb{P}^4 ARISING FROM A STABLE RANK-3 VECTOR BUNDLE

HIROTACHI ABO, WOLFRAM DECKER, NOBUO SASAKURA†

0. Introduction

In this paper we give a cohomological proof for the existence of a new family of smooth surfaces in \mathbb{P}^4 . In fact, we show:

Theorem 0.1. (i) Let $X \subset \mathbb{P}^4$ be an elliptic conic bundle with degree d = 8 and sectional genus $\pi = 5$. Then the ideal sheaf of X arises as a cokernel

$$0 \to 4\mathcal{O}(-1) \to \mathcal{G} \to \mathcal{J}_X(3) \to 0$$

where \mathfrak{G} is a rank-5 vector bundle on \mathbb{P}^4 with Chern-classes

$$c_1 = -1, c_2 = 2, c_3 = -2$$
 and $c_4 = -3$.

9 is isomorphic to the cohomology bundle of a monad

$$(M) \qquad 0 \to \Omega^3(3) \overset{\alpha}{\to} \Omega^2(2) \oplus \Omega^1(1) \overset{\beta}{\to} \mathcal{O} \to 0,$$

with

$$\alpha = \begin{pmatrix} e_4 \\ e_0 \wedge e_2 + e_1 \wedge e_3 \end{pmatrix}$$

and

$$\beta = (e_0 \wedge e_2 + e_1 \wedge e_3 - e_4),$$

where e_0, \ldots, e_4 is a basis of the underlying vector space V of \mathbb{P}^4 . In particular, \mathfrak{G} is uniquely determined up to isomorphisms and coordinate transformations.

(ii) Conversely, if \mathfrak{G} is the cohomology bunndle of the monad (M) as in (i), then $\mathfrak{G}(1)$ is globally generated. Therefore the dependancy locus of four general sections of $\mathfrak{G}(1)$ is a smooth surface $X \subset \mathbb{P}^4$. In fact, X is an elliptic conic bundle with invariants as above. \square

For a geometric construction of our surfaces we refer to [Ra].

The new surfaces are missing in a series of classification papers. First of all, they are falsely ruled out in the classification of smooth degree 8 surfaces in \mathbb{P}^4 by Okonek [Ok2] and, independently, Ionescu [Io]. As a consequence, they are e.g. also missing in the first version of two papers which are concerned with the classification of conic bundles in \mathbb{P}^4 [ES], [BR]. The correct result is:

Theorem 0.2. [ES], [BR] Let $X \subset \mathbb{P}^4$ be a smooth surface ruled in conics. Then X is either rational or an elliptic conic bundle with d=8 and $\pi=5$. In the first case X is either a Del Pezzo surface of degree 4 or a Castelnuovo surface. \square

This result is important in the context of adjunction theory [So], [SV], [VdV]. Recall, that for a smooth surface $X \subset \mathbb{P}^4$ the adjunction map is defined unless X is a plane or a scroll. If the adjunction map is defined, then it has a 2-dimensional image unless X is a Del Pezzo surface or a conic bundle. Therefore the classification of scrolls in \mathbb{P}^4 [La], [Au] and the above result imply:

Corollary 0.3. Let $X \subset \mathbb{P}^4$ be a smooth surface of degree $d \geq 9$. Then the adjunction map is defined and has a 2-dimensional image. \square

The new family is one of a few known families of irregular smooth surfaces in \mathbb{P}^4 . In fact, up to pullbacks via finite covers $\mathbb{P}^4 \to \mathbb{P}^4$, our surfaces are the first such surfaces which do not possess a Heisenberg symmetry (compare [ADHPR]). Moreover, they provide a counterexample to a conjecture of Ellingsrud and Peskine. According to this conjecture there should be no irregular m-ruled surface in \mathbb{P}^4 for $m \geq 2$.

We first came across the elliptic conic bundles when studying a stable rank-3 vector bundle \mathcal{E} on \mathbb{P}^4 with Chern classes

$$c_1 = 4, c_2 = 8$$
 and $c_3 = 8$.

 \mathcal{E} has been found by the stratification theoretical method of the third author (compare [Sa] for this method). The dependancy locus of two sections of \mathcal{E} is a smooth surface of the desired type. In fact, \mathcal{E} is a cokernel

$$0 \to 20 \to \mathcal{G}(1) \to \mathcal{E} \to 0$$
,

where \mathfrak{G} is the rank-5 vector bundle from Theorem 0.1.

Our paper is organized as follows: In Section 1 we review Beilinson's theorem [Bei] in the context of smooth surfaces in \mathbb{P}^4 . In Section 2 we follow the cohomological approach of Okonek and find the rank-5 vector bundle \mathcal{G} and thus the elliptic conic bundles via Beilinson's generalized monads. In Section 3 we take a quick look at the stratification theoretical method of the third author by studying some examples. In particular, we will construct the rank-3 bundle \mathcal{E} and we will explain its relation to the rank-5 bundle \mathcal{G} .

The third author, Nobuo Sasakura, died in June 1997. His death has been a great loss to us.

Acknowledgement. We thank Kristian Ranestad and Frank-Olaf Schreyer for helpful discussions. The second author is grateful to the Tokyo Metropolitan University and to M. Maruyama and the Kyoto University for their hospitality. The first author would like to thank the German research society for its financial support (DFG-Schwerpunkt "Algorithmische Zahlentheorie und Algebra") and the Universität des Saarlandes for its hospitality.

Notations 0.4. $\mathbb{P}^4 = \mathbb{P}^4(x) = \mathbb{P}(V)$ will be the projective space of lines in a 5-dimensional \mathbb{C} -vector space V and

$$R = \mathbb{C}[x_0, \dots, x_4] = \bigoplus_{m \ge 0} S^m V^*$$

its homogeneous coordinate ring. We write $\Omega^i = \Lambda^i \mathfrak{T}^*_{\mathbb{P}^4}$ for the i^{th} bundle of differentials on \mathbb{P}^4 . If X is a smooth surface in \mathbb{P}^4 , then we denote by

- H its hyperplane class
- $d = H^2$ its degree
- $K = K_X$ its canonical divisor
- $\pi = \frac{1}{2}H \cdot (H+K) + 1$ its sectional genus
- $q = p_g p_a$ its irregularity
- $\chi = \chi(\mathcal{O}_X) = 1 q + p_g$ the Euler characteristic of its structure sheaf. \square

1. Beilinson's Theorem

In this section we review Beilinson's theorem [Bei] in the context of smooth surfaces in \mathbb{P}^4 . The general theorem tells us that the derived category of coherent sheaves on \mathbb{P}^n is generated by the twisted bundles of differentials $\Omega^i(i)$, $0 \le i \le n$. More concretely, every coherent sheaf on \mathbb{P}^n is the cohomology of a certain generalized monad involving bundles of differentials. In the next section we apply Beilinson's theorem in order to study our surfaces in \mathbb{P}^4 from a cohomological point of view.

Theorem 1.1. [Bei] For any coherent sheaf S on \mathbb{P}^4 there is a complex \mathfrak{K} with

$$\mathcal{K}^i \cong \bigoplus_j H^{i+j}\left(\mathbb{P}^4, \mathcal{S}(-j)\right) \otimes \Omega^j(j),$$

such that

$$H^i(\mathcal{K}^{\cdot}) = \left\{ egin{array}{ll} \mathbb{S} & i = 0 \\ 0 & i
eq 0 \end{array} \right. \qquad \square$$

The differentials of \mathcal{K} are given by matrices with entries in the exterior algebra ΛV over the underlying vector space V of \mathbb{P}^4 :

Remark 1.2. The Koszul complex on $\mathbb{P}^4 = \mathbb{P}(V)$ is the exact sequence

$$0 \leftarrow \mathcal{O} \stackrel{\kappa_1}{\leftarrow} V^* \otimes \mathcal{O}(-1) \stackrel{\kappa_2}{\leftarrow} \Lambda^2 V^* \otimes \mathcal{O}(-2) \stackrel{\kappa_3}{\leftarrow} \dots \stackrel{\kappa_5}{\leftarrow} \Lambda^5 V^* \otimes \mathcal{O}(-5)) \leftarrow 0,$$

defined by contraction with the tautological subbundle

$$\mathcal{O}(-1) \to V \otimes \mathcal{O}$$
.

The syzygy bundles of this complex are the bundles of differentials:

$$(\ker \kappa_i) \cong \Omega^i, \quad 0 < i < 4.$$

Via the short exact sequences associated to the Koszul complex we may compute the intermediate cohomology modules of the Ω^i :

$$H^j_*\left(\mathbb{P}^4,\Omega^i\right)\cong \left\{ \begin{array}{ll} \mathbb{C} & 1\leq j=i\leq 3\\ 0 & 1\leq j\neq i\leq 3 \end{array} \right. .$$

In the same way we obtain for $0 \le i, j \le 4$:

$$\operatorname{Hom}(\Omega^{i}(i), \Omega^{j}(j)) \cong \Lambda^{i-j}V,$$

the isomorphism being defined by contraction. The composition of homomorphisms coincides with multiplication in ΛV . \square

We may therefore express conditions on a coherent sheaf on $\mathbb{P}(V)$ as conditions on certain matrices with entries in ΛV . For example we need:

Remark 1.3. Let

$$t\Omega^i(i) \stackrel{A}{\to} s\Omega^{i-1}(i-1)$$

be a vector bundle homomorphism, i.e., let A be a $s \times t$ -matrix with entries in V. Then a necessary condition for A to be surjective is: If (a_1, \ldots, a_t) is a non-trivial linear combination of the rows of A, then

$$\dim \operatorname{span}(a_1,\ldots,a_t) \ge i+1:$$

A is surjective iff its dual map is pointwise injective iff

$$s\Lambda^{i-1}V \wedge x \stackrel{A}{\to} t\Lambda^i V \wedge x$$

is injective for any $\langle x \rangle \in \mathbb{P}(V)$. \square

In order to apply Beilinson's theorem to the ideal sheaf (suitably twisted) of a smooth surface $X \subset \mathbb{P}^4$ we need information on the dimensions $h^i \mathcal{J}_X(m)$. First we recall Riemann-Roch:

Proposition 1.4. Let $X \subset \mathbb{P}^4$ be a smooth surface. Then

$$\chi(\mathcal{J}_X(m)) = \chi(\mathcal{O}_{\mathbb{P}^4}(m)) - \binom{m+1}{2}d + m(\pi-1) - 1 + q - p_g$$
.

Moreover one knows:

Proposition 1.5. [DES] Let $X \subset \mathbb{P}^4$ be a smooth, non-general type surface which is not contained in any cubic hypersurface. Then we have the following table for the $h^i \mathcal{J}_X(m)$:

	$\left i \right $					
0	0	0	0	0	0	
N+1	p_g	0	0	0	0	
0	q	$h^2 \mathcal{J}_X(1)$	$h^2 \mathcal{J}_X(2)$	$h^2 \mathcal{J}_X(3)$	$h^2 \mathcal{J}_X(4)$	
0	0	0	$h^1 \mathcal{J}_X(2)$	$h^1 \mathcal{J}_X(3)$	$h^1 \mathcal{J}_X(4)$	
0	0	0	0	0	$h^0 \mathcal{J}_X(4)$	

where

$$N = \pi - q + p_g - 1 . \qquad \Box$$

In the sequel we represent a zero in a cohomology table by an empty box.

2. A MONAD CONSTRUCTION

In this section we show, that every elliptic conic bundle in \mathbb{P}^4 with d=8 and $\pi=5$ is given as the dependancy locus of four sections in a unique rank-5 vector bundle $\mathfrak{G}(1)$. We give a monad construction for \mathfrak{G} , which conversely provides a proof for the existence of our surfaces since $\mathfrak{G}(1)$ is globally generated.

We first mention a classification result concerning the numerical invariants of those surfaces we are interested in here. This result is a consequence of generalized Serre correspondence [Ok1, Theorem 2.2] and adjunction theory [So], [SV], [VdV].

Proposition 2.1. [Ok2] Let $X \subset \mathbb{P}^4$ be a smooth surface with d=8 and $\pi=5$. Then $p_g=0$ and $q\leq 1$. Moreover:

(i) If q = 0, then X is the blow-up of \mathbb{P}^2 in eleven points,

$$X = \mathbb{P}^2(p_0, \dots, p_{10}),$$

embedded by the linear system (with obvious notations)


$$H = 7L - E_0 - \sum_{i=1}^{10} 2E_i.$$

(ii) If q = 1, then the adjunction map $\Phi_{|K+H|}$ exhibits X as a conic bundle over an elliptic normal curve C in \mathbb{P}^3 . There are precisely eight singular fibres of $\Phi_{|K+H|}$. These singular fibres are pairs of (-1)-lines E_i , \tilde{E}_i with $E_i \cdot \tilde{E}_i = 1$.

Proof of (ii). By classification (see [La] and [Au]) X is not a scroll. It follows from [So], [VdV] that $\mathcal{O}_X(K+H)$ is spanned, i.e., that the adjunction map is a well-defined morphism $X \to \mathbb{P}^N$ with $N = \pi - q + p_g - 1 = 3$. We have $H \cdot K = 0$ since $\pi = 5$. So we obtain $K^2 = -8$ and thus $(K + H)^2 = 0$ from the double point formula (cf. [Ha, Appendix A, 4.1.3])

$$d^2 - 10d - 5H \cdot K - 2K^2 + 12\chi = 0.$$

By [So] $\Phi_{|K+H|}$ exhibits X as a conic bundle over a smooth elliptic curve in \mathbb{P}^3 of degree $\pi - 1 = 4$. The singular fibres of $\Phi_{|K+H|}$ are pairs of (-1)-lines E_i , \tilde{E}_i with $E_i \cdot \tilde{E}_i = 1$. By blowing down one irreducible component for each singular fiber we obtain a ruled surface Y of irregularity q = 1 together with a commutative diagram



By comparing $K_X^2=-8$ with $K_Y^2=8(1-q)=0$ we find that Φ has precisely eight singular fibers. \square

The existence of surfaces of type (i) has been verified by Alexander [Al]. Okonek [Ok2] mistakenly claimed that surfaces X of type (ii) do not exist. Essentially, he claimed that the generalized monad of Beilinson for the twisted ideal sheaf $\mathcal{J}_X(2)$ cannot exist. He gave no argument, however, and in fact such a monad can be easily constructed. It is even more convenient to apply Beilinson's theorem to $\mathcal{J}_X(3)$.

Proposition 2.2. Let $X \subset \mathbb{P}^4$ be a smooth surface with d = 8, $\pi = 5$ and q = 1. Then: (i) \mathcal{J}_X has the following cohomology table:

	$\uparrow i$					
4						
	1	1				
			1	1		
					6	
	•	•	•	•	-	

In particular, X is cut out by 6 quartics.

(ii) There exists an exact sequence

$$0 \to 4\mathcal{O}(-1) \to \mathcal{G} \to \mathcal{J}_X(3) \to 0,$$

where \mathfrak{G} is a rank-5 vector bundle on \mathbb{P}^4 with Chern-classes

$$c_1 = -1, c_2 = 2, c_3 = -2$$
 and $c_4 = -3$.

G is isomorphic to the cohomology bundle of a monad

$$(M)$$
 $0 \to \Omega^3(3) \xrightarrow{\alpha} \Omega^2(2) \oplus \Omega^1(1) \xrightarrow{\beta} 0 \to 0,$

with

$$\alpha = \begin{pmatrix} e_4 \\ e_0 \wedge e_2 + e_1 \wedge e_3 \end{pmatrix}$$

and

$$\beta = (e_0 \wedge e_2 + e_1 \wedge e_3 \quad -e_4),$$

where e_0, \ldots, e_4 is a basis of the underlying vector space V of \mathbb{P}^4 . In particular, \mathfrak{G} is uniquely determined up to isomorphisms and coordinate transformations.

(iii) 9 has a minimal free resolution of type

So \mathcal{J}_X has syzygies of type

$$0 \leftarrow \mathcal{J}_X \leftarrow 60(-4) \xrightarrow{40(-5)} 0(-6)$$

$$\searrow \oplus \leftarrow \oplus \\
50(-6) & 40(-7) & 0(-8) \leftarrow 0.$$

Proof. (i) X is a ruled surface by Proposition 2.1, (ii). In particular, X is not of general type. Moreover, X is not contained in a cubic hypersurface by general classification results (see [Ro], [Au] and [Ko]). Therefore \mathcal{J}_X has a cohomology table of type

,	$\bigcap_{i} i$					
4						
	1	1	a	*	*	
			*	*	*	
					*	
-	•		5	5		

by Proposition 1.5. Beilinson's theorem applied to $\mathcal{J}_X(2)$ implies the existence of a surjective map $\Omega^1(1) \to a\mathbb{O}$. By Remark 1.3 this is only possible if $a = h^2 \mathcal{J}_X(2) = 0$. Then $h^2 \mathcal{J}_X(m) = 0$ for every $m \geq 2$. In particular, $h^1 \mathcal{J}_X(2) = h^1 \mathcal{J}_X(3) = 1$ by Riemann-Roch. Applying the same argument again gives $h^1 \mathcal{J}_X(m) = 0$ for every $m \geq 4$ and $h^0 \mathcal{J}_X(4) = 6$.


(ii) The adjoint linear system

$$H^0(\mathcal{O}_X(K+H)) \cong \operatorname{Ext}^1(\mathcal{J}_X(3), \mathcal{O}(-1)) =: W$$

has dimension 4 and generates $\mathcal{O}_X(K+H) \cong \omega_X(1)$ (compare the proof of Proposition 2.1). Therefore the identy in


$$W^* \otimes W \cong \operatorname{Ext}^1(\mathcal{J}_X(3), W^* \otimes \mathcal{O}(-1))$$

defines an extension as in the assertion with a locally free sheaf \mathfrak{G} . By construction and (i), $\mathfrak{G}(-3)$ has the following cohomology table:



Beilinson's theorem implies that \mathcal{G} is the cohomology bundle of a monad with bundles of differentials as in the assertion. Expressing the monad conditions in terms of the exterior algebra ΛV shows, that, up to isomorphisms, also the arrows of the monad are of the claimed type (compare Remark 1.2 and Remark 1.3).

(iii) The kernel $\mathcal{K} = \ker \beta$ is a rank-9 bundle which fits into a commutative diagram with exact rows and columns as follows:



With respect to suitably chosen bases of U and $\Lambda^1 V^* \oplus \Lambda^0 V^*$ the map ϕ is given by the matrix

Resolving ϕ shows that \mathcal{K} has syzygies of type

$$0 \leftarrow \mathcal{K} \leftarrow 150(-1) \underbrace{50(-2)0(-3)}_{} \oplus \leftarrow \oplus \underbrace{50(-3)}_{} 0(-5) \leftarrow 0.$$

Comparing with the Koszul resolution of $\Omega^3(3)$ gives the result. \square

Remark 2.3. Conversely, by starting with (M) and \mathfrak{G} , we obtain a proof for the existence of smooth surfaces in \mathbb{P}^4 with d=8, $\pi=5$ and q=1: Since $\mathfrak{G}(1)$ is globally generated, Kleiman's Bertini-type theorem [Kl] implies that the dependancy locus of 4 general sections of $\mathfrak{G}(1)$ is indeed a smooth surface (with invariants as above). \square

Remark 2.4. (i) Let $X \stackrel{\Phi}{\to} C$ be an elliptic conic bundle as above. Ionescu [Io] studied the union W of the planes of the conics which are the fibres of Φ . He showed, that W is a quartic hypersurface in \mathbb{P}^4 which is a cone with vertex a point. The base S of the cone is an

elliptic ruled quartic surface in \mathbb{P}^3 with two double lines l_1 and l_2 . The rulings of S form a divisor of class (2,2) in the $\mathbb{P}^1 \times \mathbb{P}^1 \cong l_1 \times l_2$ of lines joining l_1 and l_2 . Ionescu considered the cone over a pair of rulings through a point on l_1 but overlooked the presence of l_2 . His residual quadric at the end of [Io, 6.4] is the double plane over l_2 .

(ii) The vertex of the cone W is the distinguished point $\langle e_4 \rangle$ in our construction. The projection from $\langle e_4 \rangle$ maps X 2:1 to S. The conics on X are mapped 2:1 onto the rulings of S.


Remark 2.5. In the minimal free resolution of \mathfrak{G} (and thus also of \mathfrak{J}_X) there are two maps which are given by four linear forms. In fact, for both maps, these forms generate the ideal of the distinguished point $\langle e_4 \rangle$. \square

3. Vector bundles via stratifications


In this section we take a quick look at the approach of the third author to the construction of vector bundles on \mathbb{P}^n . In particular, we construct a stable rank-3 bundle \mathcal{E} on \mathbb{P}^4 such that the dependancy locus of two general sections of \mathcal{E} is an elliptic conic bundle. This rank-3 bundle was the starting point of our investigations. It is related to the rank-5 bundle 9 from Section 2 via an exact sequence

$$0 \to 20 \to \mathfrak{G}(1) \to \mathcal{E} \to 0.$$

Let \mathcal{E} be a rank-r vector bundle on \mathbb{P}^n with first Chern class c_1 . Minimal systems of generators s_1, \ldots, s_l of $H^0_*\mathcal{E}$ and $\sigma_1, \ldots, \sigma_k$ of $H^0_*\mathcal{E}^{\vee}$ give rise to a commutative diagram



where \mathcal{L} and \mathcal{K} are direct sums of line bundles of ranks l and k resp. S is a $k \times l$ -matrix with polynomial entries. The entries in the j-th column of S define the zero-scheme $X_{s_j} = \{s_j = 0\}$. The entries in the i-th row of S define the zero-scheme $X_{\sigma_i} = \{\sigma_i = 0\}$. Suppose that \mathcal{K} has a direct summand of type $r\mathcal{O}(m)$, and let $\sigma_{i_1}, \ldots, \sigma_{i_r}$ be the corresponding elements of $H^0_*\mathcal{E}^\vee$. By projecting onto this direct summand we may represent \mathcal{E} as a subsheaf of $r\mathcal{O}(m)$:



In fact, θ is a monomorphism of vector bundles outside the divisor defined by the form $f = \sigma_{i_1} \wedge \cdots \wedge \sigma_{i_r} \in H^0(\mathbb{P}^n, \mathcal{O}((r \cdot m - c_1))) \cong H^0(\mathbb{P}^n, \Lambda^r \mathcal{E}^{\vee}(m))$. The *j*-th column of

T represents the section $t_j := H^0(\theta)(s_j)$ of $r\mathcal{O}(m)$, and we have the relations of forms $t_{j_1} \wedge \cdots \wedge t_{j_r} = f \cdot (s_{j_1} \wedge \cdots \wedge s_{j_r})$. In order to detect properties of \mathcal{E} it is often enough to study the matrix T. For example, one may verify that \mathcal{E} is indeed a vector bundle by looking at the ideal I generated by the maximal minors of T and by checking that the ideal quotient (I:f) defines the empty subset of \mathbb{P}^n .

Conversely, one can construct vector bundles \mathcal{E} on \mathbb{P}^n by starting with convenient matrices T. In the examples below f is of the form $(f_{\text{red}})^{(r-1)}$, where $f_{\text{red}} = x_0 \cdot \ldots \cdot x_{c_1}$ is a product of coordinates, and T is of the form $T = (T' T'') : \mathcal{L} \to r\mathcal{O}(c_1)$, where T' is the appropriate identy matrix multiplied by f_{red} . In each example the minimal free resolution of \mathcal{E} is obtained by resolving T (use e.g. Macaulay [BS]).

Example 3.1. The Nullcorrelation bundle \mathcal{E} on \mathbb{P}^3 (compare e.g. [Ba]). In this case $r=2, c_1=2$, and we may choose $f=f_{\rm red}=x_0x_1$ and

$$T = \begin{pmatrix} x_0 x_1 & 0 & x_0 x_3 & x_1 x_2 & x_2 x_3 \\ 0 & x_0 x_1 & x_0^2 & x_1^2 & x_0 x_2 + x_1 x_3 \end{pmatrix}.$$

Note that the sections of $\mathcal{E}^{\vee}(2) \cong \mathcal{E}$ corresponding to the rows of T vanish along the union of two skew lines and a double line resp. The minimal free resolution of \mathcal{E} is of type

$$0 \leftarrow \mathcal{E} \leftarrow 50 \leftarrow 40(-1) \leftarrow 0(-2) \leftarrow 0$$
.

Example 3.2. The Horrocks-Mumford bundle \mathcal{E} on \mathbb{P}^4 [HM]. In this case $r=2, c_1=5$, and we may choose $f=f_{\rm red}=x_0x_1x_2x_3x_4$ and

$$T = (T_0 \quad T_1 \quad T_2 \quad T_3),$$

with T_0, \ldots, T_3 as follows:

$$T_0 = \begin{pmatrix} \gamma_0 & 0 & \gamma_3 & \gamma_4 \\ 0 & \gamma_0 & \gamma_1 & \gamma_2 \end{pmatrix},$$

where $\gamma_0, \ldots, \gamma_4$ are the five Horrocks-Mumford quintics

$$\begin{array}{ll} \gamma_0 = y_0 y_1 y_2 y_3 y_4 & \gamma_1 = \sum_{i \in \mathbb{Z}_5} y_i y_{i+2}^2 y_{i+3}^2 & \gamma_2 = \sum_{i \in \mathbb{Z}_5} y_i^3 y_{i+2} y_{i+3} \\ \gamma_3 = \sum_{i \in \mathbb{Z}_5} y_i^3 y_{i+1} y_{i+4} & \gamma_4 = \sum_{i \in \mathbb{Z}_5} y_i y_{i+1}^2 y_{i+4}^2. \end{array}$$

The entries of T_1, \ldots, T_3 are sextics:

$$T_{1} = (x_{i+1}x_{i+2}x_{i+3} t(x_{i+2}x_{i+3}x_{i+4} x_{i+1}x_{i+4}^{2}))_{i \in \mathbb{Z}_{5}},$$

$$T_{2} = (x_{i+2}x_{i+3}x_{i+4} t(x_{i+2}x_{i+3}x_{i+4} x_{i+1}x_{i+4}^{2} + x_{i}x_{i+2}^{2}))_{i \in \mathbb{Z}_{5}},$$

$$T_{3} = (x_{i+1}x_{i+3}x_{i+4} t(x_{i+2}x_{i+3}^{2} + x_{i}x_{i+4}^{2} x_{i+1}x_{i+3}x_{i+4}))_{i \in \mathbb{Z}_{5}}.$$

The zero-schemes of the sections of \mathcal{E} are called *Horrocks-Mumford surfaces*. They are the minimal abelian surfaces in \mathbb{P}^4 and the degenerations of these smooth surfaces [HM] (compare [BHM]). Those sections of $\mathcal{E}^{\vee}(5) \cong \mathcal{E}$ corresponding to the rows of T vanish along the unions of five double planes, namely $\{x_i^2 = x_{i+2}^2 = x_i x_{i+2} = x_{i+2} x_{i+3}^2 + x_i x_{i+4}^2 = 0\}$,

 $i \in \mathbb{Z}_5$, and $\{x_i^2 = x_{i+1}^2 = x_i x_{i+1} = x_{i+1} x_{i+4}^2 + x_i x_{i+2}^2 = 0\}$, $i \in \mathbb{Z}_5$, resp. (compare e.g. [ADHPR, Section 9] for these degenerations). The minimal free resolution of \mathcal{E} is of type

$$0 \leftarrow \mathcal{E} \leftarrow \bigoplus_{15\mathcal{O}(-1)}^{4\mathcal{O}} \times 35\mathcal{O}(-2) \leftarrow 20\mathcal{O}(-3) \times 2\mathcal{O}(-5) \leftarrow 0$$

(compare e.g. [De]). \square

Example 3.3. A stable rank-2 reflexive sheaf \mathcal{E} on \mathbb{P}^4 such that $\mathcal{E}(-3)$ has Chern classes $c_1 = -1$, $c_2 = 9$, $c_3 = 25$ and $c_4 = 50$ [ADHPR, Section 7].

In this case r = 2, $c_1 = 5$, and we may choose $f = f_{red} = x_0 x_1 x_2 x_3 x_4$ and

$$T = (T_0, T_1),$$

with T_0 and T_1 as follows:

$$T_0 = \begin{pmatrix} \gamma_0 & 0 & \gamma_2 & \gamma_4 \\ 0 & \gamma_0 & \gamma_1 & \gamma_3 \end{pmatrix},$$

where $\gamma_0, \ldots, \gamma_4$ are the Horrocks-Mumford quintics from 3.2. The entries of T_1 are septics:

$$T_1 = (x_{i+1}x_{i+2}x_{i+3}x_{i+4} t(x_{i+2}x_{i+4}^2 + x_{i+1}^2x_{i+3} x_{i+3}^2x_{i+4} + x_{i+1}x_{i+2}^2))_{i \in \mathbb{Z}_5}.$$

This time \mathcal{E} is locally free only outside the union of the 25 Horrocks-Mumford lines (compare [HM] for these lines). The zero-schemes of the sections of \mathcal{E} are non-minimal abelian surfaces in \mathbb{P}^4 and the degenerations of these smooth surfaces. They are (5,5)-linked to Horrocks-Mumford surfaces. The zero-schemes of those sections of $\mathcal{E}^{\vee}(5) \cong \mathcal{E}$ corresponding to the rows of T consist of five components, namely $\{x_i = x_{i+1}^2 x_{i+3} = x_{i+2} x_{i+4}^2 = 0\}, i \in \mathbb{Z}_5$, and $\{x_i = x_{i+1} x_{i+2}^2 + x_{i+3}^2 x_{i+4} = 0\}, i \in \mathbb{Z}_5$, resp. The minimal free resolution of \mathcal{E} is of type

$$0 \leftarrow \mathcal{E} \leftarrow \bigoplus_{50(-2)}^{40} \nwarrow 150(-3) \leftarrow 100(-4) \leftarrow 20(-5) \leftarrow 0.$$

Compare [ADHPR, Section 7 and 9]. \square

Example 3.4. A stable rank-3 vector bundle \mathcal{E} on \mathbb{P}^4 with Chern classes $c_1 = 4$, $c_2 = 8$ and $c_3 = 8$.

We choose $f = f_{red} = x_0 x_1 x_2 x_3$ and

$$T = (T' \quad t_4 \quad t_5 \quad t_6 \quad t_7 \quad t_8),$$

with

$$T' = \begin{pmatrix} x_0 x_1 x_2 x_3 & 0 & 0 \\ 0 & x_0 x_1 x_2 x_3 & 0 \\ 0 & 0 & x_0 x_1 x_2 x_3 \end{pmatrix},$$

and

$$t_4 = x_0 x_3^{t} (x_4^2 + x_1 x_3 - x_0 x_2 \quad x_2 x_3 - x_0 x_1 \quad -x_0 x_3),$$

$$t_{5} = x_{0}x_{1} \ ^{t}(x_{4}^{2} + x_{1}x_{3} + x_{0}x_{2} - x_{0}x_{1} \ x_{1}x_{2} - x_{0}x_{3}),$$

$$t_{6} = x_{1}x_{2} \ ^{t}(-x_{4}^{2} + x_{1}x_{3} - x_{0}x_{2} \ x_{0}x_{1} - x_{2}x_{3} - x_{1}x_{2}),$$

$$t_{7} = x_{2}x_{3} \ ^{t}(-x_{4}^{2} + x_{1}x_{3} + x_{0}x_{2} - x_{2}x_{3} \ x_{0}x_{3} - x_{1}x_{2}),$$

$$t_{8} = \begin{pmatrix} -x_{4}^{4} + x_{1}^{2}x_{3}^{2} + x_{0}^{2}x_{2}^{2} \\ x_{4}^{2}(x_{0}x_{1} - x_{2}x_{3}) - (x_{0}x_{1} + x_{2}x_{3})(x_{1}x_{3} + x_{0}x_{2}) \\ (x_{0}x_{2} - x_{1}x_{3})(x_{1}x_{2} + x_{0}x_{3}) + x_{4}^{2}(x_{0}x_{3} - x_{1}x_{2}) \end{pmatrix}.$$

A check on the maximal minors of T shows that \mathcal{E} is locally free (use e.g. Macaulay [BS]). Let X_1, X_2 and X_3 be the zero-schemes of the sections of $\mathcal{E}^{\vee}(4)$ corresponding to the rows of T. Then X_1 is the union of four plane quadrics, whereas X_2 and X_3 are the unions of four double planes (use e.g. the primary decomposition package of Singular [GPS] to obtain the explicit equations). The minimal free resolution of \mathcal{E} is of type

The minimal free resolution of \mathcal{E}^{\vee} is of type

$$0(-2)$$

$$0 \leftarrow \mathcal{E}^{\vee} \leftarrow 40(-3) \qquad 20(-4)$$

$$0 \leftarrow \mathcal{E}^{\vee} \leftarrow 40(-3) \qquad 20(-4)$$

$$0 \leftarrow \mathcal{E}^{\vee} \leftarrow 40(-3) \qquad 20(-4)$$

$$0 \leftarrow \mathcal{E}^{\vee} \leftarrow 40(-6)$$

$$0 \leftarrow 40(-6)$$

$$0 \leftarrow$$

In particular, \mathcal{E} is stable since the normalized bundle $\mathcal{E}(-2)$ and its dual twisted by -1 have no non-zero sections.

Let (M) be the monad of the rank-5 vector bundle \mathcal{G} as in Proposition 2.2. By taking sections in the display of (M) we identify $H^0\mathcal{G}(1)$ in terms of the exterior algebra ΛV^* as the kernel

$$0 \to H^0 \mathfrak{G}(1) \to \Lambda^1 V^* \oplus \Lambda^0 V^* \stackrel{\beta}{\to} V^* \to 0$$

Two general elements of $H^0\mathfrak{G}(1)$ are nowhere dependant. In fact,

$$\sigma_1 = \left(\begin{array}{c} e_0 \wedge e_1 + e_2 \wedge e_3 \\ 0 \end{array}\right)$$

and

$$\sigma_2 = \begin{pmatrix} e_1 \wedge e_2 - e_0 \wedge e_3 \\ e_1 \wedge e_2 \wedge e_4 - e_0 \wedge e_3 \wedge e_4 \end{pmatrix}$$

give rise to an exact sequence

$$0 \to 20 \xrightarrow{\sigma} \mathcal{G}(1) \to \mathcal{E} \to 0.$$

This can be easily seen by comparing the syzygies of the cokernel of σ with those of \mathcal{E} (use e.g. Macaulay [BS]). It follows that $\mathcal{E}(-1)$ is isomorphic to the cohomology bundle of the monad

$$0 \to 2\mathfrak{O}(-1) \oplus \Omega^3(3) \xrightarrow{(\sigma,\alpha)} \Omega^2(2) \oplus \Omega^1(1) \xrightarrow{\beta} \mathfrak{O} \to 0. \qquad \Box$$

In each of these examples we have a stratification

$$L^{m_0} \subset \cdots \subset L^1 \subset \mathbb{P}^n, \quad m_0 = \min\{n, c_1\},$$

where we use the following notations: For any subset I of \mathbb{Z}_{m_0} we set $L_I = \bigcap_{i \in I} L_i$, where $L_i = \{x_i = 0\}$, and for any $m \in \mathbb{Z}_{m_0}$ we write

$$L^m = \bigcup \{L_I \mid I \in \mathcal{P}_m(\mathbb{Z}_{m_0})\},\,$$

where $\mathcal{P}_m(\mathbb{Z}_{m_0})$ is the collection of subsets $I \subset \mathbb{Z}_{m_0}$ whith cardinality |I| = m. One may use this filtration to find local frames for \mathcal{E} in a more systematic way, and to construct a filtration of sheaves

$$\mathcal{E}^0 \cong r\mathcal{O} \subset \mathcal{E}^1 \cdots \subset \mathcal{E}^{m_0} \cong \mathcal{E}$$
,

such that

$$\mathcal{E}^m/\mathcal{E}^{m-1} \cong \bigoplus_{I \in \mathcal{P}_m(\mathbb{Z}_{m_0})} \mathcal{O}_{L_I}(d_{L_I}),$$

with convenient twists d_{L_I} . Via this filtration one can e.g. compute the cohomology of \mathcal{E} (use Čech cohomology). We refer to [Sa] and [SEKS] for more details.

References

- [ADHPR] Aure, A.B., Decker, W., Hulek, K., Popescu, S., Ranestad, K., Syzygies of abelian and bielliptic surfaces in \mathbb{P}^4 , Int. J. of Math. (to appear).
- [Al] Alexander, J., Surfaces rationelles non-speciales dans \mathbb{P}^4 , Math. Z. **200** (1988), 87–110.
- [Au] Aure, A.B., On surfaces in projective 4-space, Thesis, Oslo 1987.
- [Ba] Barth, W., Some properties of stable rank-2 vector bundles on \mathbb{P}^n , Math. Ann. **226** (1977), 125–150.
- [Bei] Beilinson, A., Coherent sheaves on \mathbb{P}^N and problems of linear algebra, Funct. Anal. Appl. 12 (1978), 214–216.
- [BHM] Barth, W., Hulek, K., Moore, R., Degenerations of Horrocks-Mumford surfaces, Math. Ann. 277 (1987), 735–755.
- [BR] Braun, R., Ranestad, K., Conic bundles in projective fourspace, to appear.
- [BS] Bayer, D., Stillman, M., Macaulay: A system for computation in algebraic geometry and commutative algebra, Source and object code available for Unix and Macintosh computers. Contact the authors, or download from math.harvard.edu via anonymous ftp.
- [De] Decker, W., Monads and cohomology modules of rank 2 vector bundles, Compositio Math. **76** (1990), 7–17.
- [ES] Ellia, P.R., Sacchiero, G., Smooth surfaces of \mathbb{P}^4 ruled in conics, to appear.
- [GPS] Greuel, G.-M., Pfister, G., Schönemann, H., Singular Reference Manual, in: Reports On Computer Algebra, number 12. Centre for Computer Algebra, University of Kaiserslautern, May 1997. http://www.mathematik.uni-kl.de/~zca/Singular.
- [Ha] Hartshorne, R., Algebraic geometry, Springer, Berlin, Heidelberg, New York, Tokyo, 1977.
- [HM] Horrocks, G., Mumford, D., A rank 2 vector bundle on \mathbb{P}^4 with 15,000 symmetries, Topology 12 (1973), 63–81.

- [Io] Ionescu, P., Embedded projective varieties of small invariants III, in: Algebraic Geometry (L'Aquila 1988), LNM **1417**, Springer, Berlin, Heidelberg, New York, Tokyo, 1990, 138–154.
- [Kl] Kleiman, S., Geometry on grassmanians and applications to splitting bundles and smoothing cycles, Publ. Math. I.H.E.S. **36** (1969), 281–297.
- [Ko] Koelblen, L., Surfaces de \mathbb{P}_4 tracées sur une hypersurface cubique, J. reine und angew. Math. 433 (1992), 113–141.
- [La] Lanteri, A., On the existence of scrolls in \mathbb{P}^4 , Lincei-Rend. Sc. fis.mat.e.nat **LXIX** (1980), 223–227.
- [Ok1] Okonek, C., Reflexive Garben auf \mathbb{P}^4 , Math. Ann. **269** (1982), 211–237.
- [Ok2] _____, Flächen vom Grad 8 im \mathbb{P}^4 , Math. Z. **191** (1986), 207–223.
- [Ra] Ranestad, K., A geometric construction of elliptic conic bundles in \mathbb{P}^4 , to appear.
- [Ro] Roth, L., On the projective classification of surfaces, Proc. of London Math. Soc. 42 (1937), 142–170.
- [Sa] Sasakura, N., A stratification theoretical method of construction of holomorphic vector bundles, in: Complex analytic singularities, Advanced studies in Pure Mathematics 8 (1987), 527–581.
- [SEKS] Sasakura, N., Enta, Y., Kagesawa, M., Sakurai, T., Rank two reflexive sheaves formed from quadratic residue graph, Tokyo metropolitan university mathematical preprint series 1 (1996).
- [So] Sommese, A.J., Hyperplane sections of projective surfaces I. The adjunction mapping, Duke Math. J. 46 (1979), 377–401.
- [SV] Sommese, A.J., Van de Ven, A., On the adjunction mapping, Math. Ann. 278 (1987), 593–603.
- [VdV] Van de Ven, A., On the 2-connectedness of very ample divisors on a surface, Duke Math. J. 46 (1979), 403–407.

Нікотасні Аво

DEPARTMENT OF MATHEMATICS, TOKYO METROPOLITAN UNIVERSITY, MINAMI-OHSAWA 1-1, HACHIOJI-SHI TOKYO, 192-03 JAPAN

E-mail address: abo@math.metro-u.ac.jp

Wolfram Decker

FACHBEREICH MATHEMATIK, UNIVERSITÄT DES SAARLANDES, D 66041 SAARBRÜCKEN, GERMANY E-mail address: decker@math.uni-sb.de