# A Sparse Effective Nullstellensatz<sup>1</sup>

Martín Sombra  $^2$ 

Departamento de Matemática, Universidad Nacional de La Plata, Calle 50 y 115, 1900 La Plata, Argentina. E-mail: sombra@mate.unlp.edu.ar

**Abstract.** We present bounds for the sparseness and for the degrees of the polynomials in the Nullstellensatz. Our bounds depend mainly on the unmixed volume of the input polynomial system. The degree bounds can substantially improve the known ones when this polynomial system is sparse, and they are, in the worst case, simply exponential in terms of the number of variables and the maximum degree of the input polynomials.

**Keywords.** Cohen–Macaulay ring, Effective Nullstellensatz, Newton polytope, Degree of a polynomial system of equations.

AMS Subject Classification. 13P10.

## Introduction

We consider the problem of bounding the sparseness of the polynomials in the Nullstellensatz in the case when the input is a sparse polynomial system.

Let us denote by k a field and by  $\overline{k}$  its algebraic closure. As usual we denote by  $\mathbb{A}^n$  the affine space of n dimensions over  $\overline{k}$ . Let be given polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  without common zeros in  $\mathbb{A}^n$ . Classical Hilbert's Nullstellensatz states then that there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  satisfying the Bézout equation

$$1 = g_1 f_1 + \dots + g_s f_s. \tag{1}$$

Let d denote the maximum degree of the polynomials  $f_1, \ldots, f_s$  and assume that  $n \geq 2$ . In the previous situation we have indeed that there exist polynomials  $g_1, \ldots, g_s$  verifying the degree bound

 $\deg g_i f_i \le \max\{3, d\}^n.$ 

This result is due to Kollár, and it is optimal provided that  $d \ge 3$  [19]. In the case when d = 2 and the characteristic of k is different from two, Sabia and Solernó proved that the sharper bound deg  $g_i f_i \le n 2^{n+2}$  holds true [27]. In fact more precise estimations are valid, we refer to the original papers for the exact statements.

<sup>&</sup>lt;sup>1</sup>Date of this version: 6.8.97

<sup>&</sup>lt;sup>2</sup>Partially supported by CONICET PID 3949/92, UBA CyT EX. 001 and Fundación Antorchas.

We note that such a degree bound allows us, given polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$ , to determine whether the equation (1) is solvable or not, and in the case it is, to actually find a solution, as it reduces the original problem to the solving of a k-linear system of equations.

The study of this Bézout identity is the object of much research, due to both its theoretical and practical importance, mainly in the context of computational algebraic geometry and diophantine approximation. Thus it has been approached from many points of view and with different objectives. In this respect we refer to the research papers [2], [4], [5], [7], [11], [13], [15], [20], [24], [27], [28], [29]. We also refer to the surveys [3], [23], [34] for a broad introduction to the history of this problem, main results and open questions.

Let be given a Laurent polynomial  $f = \sum_{i \in \mathbb{Z}^n} a_i x^i \in k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$ . The *support* of f is defined as the set  $\{i : a_i \neq 0\}$ , that is, the set of exponents of the nonzero monomials of f. More generally, the support of a family of Laurent polynomials  $f_1, \ldots, f_s$  is defined as the set of exponents of the nonzero monomials of any of the  $f_i$  for  $i = 1, \ldots, s$ .

The Newton polytope  $\mathcal{N}(f_1, \ldots, f_s)$  is defined as the convex hull of the support of  $f_1, \ldots, f_s$ . Let  $\rho$  denote the dimension of this polytope. Then the unmixed volume  $\mathcal{U}(f_1, \ldots, f_s)$  of the family of Laurent polynomials  $f_1, \ldots, f_s$  is defined as  $\rho$ ! times the volume of the polytope  $\mathcal{N}(f_1, \ldots, f_s)$ .

The degree of a polynomial is bounded by a nonnegative integer d if and only if its Newton polytope is contained in d times the standard simplex of  $\mathbb{R}^n$ . Thus the notion of Newton polytope gives a sharper characterization of the monomial structure of a polynomial system than just degree. This concept is in the basis of sparse elimination theory. Within this theory, algorithms for elimination problems are designed trying to exploit the sparseness of the involved polynomials, and sparseness is then usually measured in terms of the Newton polytope of these polynomials. This is the point of view introduced by Sturmfels in his foundational work [31] and followed, for instance, in [8], [18], [25], [26], [35].

The sparse aspect in the Nullstellensatz has also been considered by Canny and Emiris, who obtained a sparse effective Nullstellensatz but only for the case of n + 1 generic n-variate Laurent polynomials [8, Theorem 7.2].

We obtain the following result, which in this context can be seen as a bound for the sparseness of the output polynomials in terms of the sparseness of the input system.

**Theorem 1.** Let be given polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  without common zeros in  $\mathbb{A}^n$ . Let  $\mathcal{N}$  denote the Newton polytope of the polynomials  $x_1, \ldots, x_n, f_1, \ldots, f_s$ , and let  $\mathcal{U}$  denote the unmixed volume of this polytope. Then there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  satisfying

$$1 = g_1 f_1 + \dots + g_s f_s,$$

with  $\mathcal{N}(g_i f_i) \subseteq n^{n+3} \mathcal{U} \mathcal{N}$  for  $i = 1, \dots, s$ .

Let us keep the notations of Theorem 1 and let d denote the maximum degree

of  $f_i$  for i = 1, ..., s. We readily derive from the previous result the degree bound

$$\deg g_i f_i \le n^{n+3} d \mathcal{U}.$$

We obtain from this the worst-case bound  $\deg g_i f_i \leq n^{n+2} d^{n+1}$ , as the unmixed volume of the polynomials  $x_1, \ldots, x_n, f_1, \ldots, f_n$  is bounded by  $d^n$ . We show however that our degree bound can considerably improve the usual one in the case when the input polynomial system is sparse and  $d \geq n$  (Example 2.14).

We note that the naive notion of sparseness, based just on the counting of the number of nonzero monomials in each polynomial, does not yield better bounds for the degrees than the usual ones, in view of the well known example of Mora, Lazard, Masser and Philippon [5].

We obtain an analogous result for the case of Laurent polynomials.

**Theorem 2.** Let be given Laurent polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$ without common zeros in  $(\overline{k}^*)^n$ . Let  $\mathcal{N}$  denote the Newton polytope of  $f_1, \ldots, f_s$ , and let  $\mathcal{U}$  denote the unmixed volume of this polytope. Then there exist  $a \in \mathbb{Z}^n$  and  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$  satisfying

$$1 = g_1 f_1 + \dots + g_s f_s,$$
  
with  $a \in n^{2n+3} \mathcal{U}^2 \mathcal{N}$  and  $\mathcal{N}(g_i f_i) \subseteq n^{2n+3} \mathcal{U}^2 \mathcal{N} - a$  for  $i = 1, \dots, s$ .

The proof of both results is similar. It takes as its first step the translation of the original system of equations over the affine space or the torus into a system of linear equations over an appropriate toric variety. The resulting system is then solved by appealing to an effective Nullstellensatz for linear forms in a Cohen-Macaulay graded ring. This key lemma is proved following for the most part the lines of a previous paper [30] which in turn is based on previous work of Dubé [9] and Almeida [1], although we introduce at this time some simplifications into the arguments. In particular we eliminate the use of estimations for the Hilbert function.

As a by-product, we obtain an effective Nullstellensatz which holds not only for linear forms, but for arbitrary homogeneous elements in a Cohen–Macaulay graded ring (Theorem 1.9).

Besides we apply these arguments in two other situations. First we consider the usual effective Nullstellensatz and we obtain the following degree bound.

**Theorem 3.** Let be given polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  without common zeros in  $\mathbb{A}^n$ . Let  $d_i$  denote the degree of  $f_i$  for  $i = 1, \ldots, s$  and assume that  $d_1 \geq \cdots \geq d_s$  holds. Then there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  satisfying

$$1 = g_1 f_1 + \dots + g_s f_s,$$

with  $\deg g_i f_i \leq 2 d_s \prod_{j=1}^{\min\{n,s\}-1} d_j$  for  $i = 1, \dots, s$ .

In the case when the polynomials  $f_1, \ldots, f_s$  are quadratic this estimation improves the bound  $\deg g_i f_i \leq n 2^{n+2}$  to the bound  $\deg g_i f_i \leq 2^{n+1}$ , which is very close to the expected  $2^n$ .

Finally we obtain another bound for the degrees in the Nullstellensatz. This bound depends on the number of variables and the maximum degree of the input polynomials and also on an additional parameter, called the *algebraic degree* of the input polynomial system, which we are going to introduce.

Let  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  be polynomials without common zeros in  $\mathbb{A}^n$ . Let  $\lambda = (\lambda_{ij})_{ij} \in \overline{k}^{s \times s}$  be an arbitrary  $s \times s$  matrix with entries in  $\overline{k}$ . We note by  $h_i(\lambda)$  the linear combinations  $\sum_j \lambda_{ij} f_j$  induced by the matrix  $\lambda$  for  $i = 1, \ldots, s$ .

Consider the set  $\Gamma$  of  $s \times s$  matrices such that for any  $\lambda$  in  $\Gamma$  the polynomials  $h_1(\lambda), \ldots, h_{t-1}(\lambda)$  form a regular sequence in  $\overline{k}[x_1, \ldots, x_n]$  and  $1 \in (h_1(\lambda), \ldots, h_t(\lambda))$  for some  $t = t(\lambda) \leq \min\{n, s\}$ . This set is nonempty, and indeed it contains a nonempty open set of  $\overline{k}^{s \times s}$ .

For each  $\lambda \in \Gamma$  and  $i = 1, \ldots, t - 1$  we denote by  $J_i(\lambda) \subseteq k[x_0, \ldots, x_n]$  the homogenization of the ideal  $(h_1(\lambda), \ldots, h_i(\lambda))$ . Then let  $\delta(\lambda)$  denote the maximum degree of the homogeneous ideal  $J_i(\lambda)$  for  $i = 1, \ldots, t - 1$ .

The algebraic degree of the polynomial system  $f_1, \ldots, f_s$  is defined as

$$\delta(f_1,\ldots,f_s) := \min \{\delta(\lambda) : \lambda \in \Gamma\}.$$

This is the algebraic analogue of the notion of geometric degree of a system of equations of Giusti et al. [14], Krick, Sabia and Solernó [21] and Sombra [30]. We refer to Section 3 for a comparison between both notions.

There have been obtained degree bounds for the polynomials in the Nullstellensatz which mainly depend on the geometric degree [13], [21], [30]. We show that a similar bound holds by replacing the geometric degree of the input polynomial system by the algebraic degree.

**Theorem 4.** Let be given polynomials  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  without common zeros in  $\mathbb{A}^n$ . Let d denote the maximum degree of  $f_i$  for  $i = 1, \ldots, s$  and let  $\delta$ denote the algebraic degree of this polynomial system. Then there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  satisfying

 $1 = g_1 f_1 + \dots + g_s f_s,$ 

with deg  $g_i f_i \leq \min\{n, s\}^2 d\delta$  for  $i = 1, \ldots, s$ .

Let us denote by  $d_i$  the degree of  $f_i$  for  $i = 1, \ldots, s$  and assume that  $d_1 \geq \cdots \geq d_s$ holds. Then the Bézout bound  $\delta(f_1, \ldots, f_s) \leq d_s \prod_{i=1}^{\min\{n,s\}-2} d_i$  holds true, and therefore we essentially recover from the previous result the known bounds for the degrees in the Nullstellensatz. The algebraic degree is bounded by the geometric degree, and so we also recover the known degree bounds in the Nullstellensatz which depend on the geometric degree. We show however that the algebraic degree is much smaller than the geometric degree in some particular instances, and by force, than the Bézout bound  $d^{n-1}$  (Example 3.22). We conclude that the obtained degree bound is much sharper in these cases than the known ones.

The outline of the paper is as follows. In Section 1 we obtain the effective Nullstellensatz for homogenous elements in a Cohen–Macaulay graded ring. In Section 2 we prove both Theorems 1 and 2 and we derive some of their consequences. In Section 3 we devote to the degree bounds in the usual Nullstellensatz.

## 0. Notations and Conventions

We denote by k an infinite field with an algebraic closure denoted by  $\overline{k}$ . All the rings to be considered in the sequel are Noetherian commutative, and more precisely finitely generated k-algebras. The polynomial ring  $k[x_0, \ldots, x_n]$  is alternatively denoted by S.

As usual  $\mathbb{P}^n$  and  $\mathbb{A}^n$  denote the projective and the affine spaces of n dimensions over  $\overline{k}$ , respectively. A variety is not necessarily irreducible.

Let J be a homogeneous ideal in the ring S/I. The dimension of J is defined as the Krull dimension of the quotient ring (S/I)/J and it is denoted by dim J. The degree of J is defined as  $(\dim J - 1)!$  times the principal coefficient of the Hilbert polynomial of the graded k-algebra (S/I)/J.

Let  $\varphi : A \to B$  be a morphism between two rings, and let I and J be ideals of A and B respectively. Then  $I^e$  denotes the extension of the ideal I to B and  $J^c$  denotes the contraction of the ideal J to A. Given an element  $\alpha \in A$ , we denote by  $\overline{\alpha}$  the element  $\varphi(\alpha) \in B$ . We shall make use of this notation when it is clear from the context the morphism to which we refer.

#### 1. An Effective Nullstellensatz over Cohen–Macaulay Graded Rings

Let be given a Cohen-Macaulay homogeneous ideal I in the ring  $k[x_0, \ldots, x_n]$ . By this we mean that the quotient ring  $k[x_0, \ldots, x_n]/I$  is a Cohen-Macaulay ring. Let r denote the dimension of I. We also suppose that there is given a homogeneous element p in S/I which is not a zero-divisor. Let  $\eta_1, \ldots, \eta_s \in S/I$  be homogeneous elements of degree one —or for short, linear forms— which define the empty variety in the open set  $\{p \neq 0\}$  of V(I). In this situation, Hilbert's Nullstellensatz implies that p belongs to the radical of the ideal  $(\eta_1, \ldots, \eta_s)$ , that is,  $p \in \sqrt{(\eta_1, \ldots, \eta_s)}$ . Equivalently we have that 1 lies in the ideal  $(\overline{\eta}_1, \ldots, \overline{\eta}_s)$  spanned by  $\overline{\eta}_1, \ldots, \overline{\eta}_s$  in the ring  $(S/I)_p$ .

We are going to give a bound for the minimal  $D \in \mathbb{N}$  such that  $p^D$  falls into the ideal  $(\eta_1, \ldots, \eta_s)$  (Main Lemma 1.7). This bound depends on the number of linear forms, and on the dimension and the degree of the ideal I. As a consequence of this result we derive an effective Nullstellensatz for Cohen-Macaulay graded rings (Theorem 1.9 and Corollary 1.10).

Let A be a ring and let  $\alpha_1, \ldots, \alpha_t$  be elements of A. Then  $\alpha_1, \ldots, \alpha_t$  is called a *weak regular sequence* if  $\overline{\alpha}_i$  is not a zero-divisor in the ring  $A/(\alpha_1, \ldots, \alpha_{i-1})$  for  $i = 1, \ldots, t$ . We note that this definition differs from usual notion of regular sequence only in one point, namely that it allows  $\overline{\alpha}_t$  to be a unit in  $A/(\alpha_1, \ldots, \alpha_{t-1})$ .

**Lemma 1.1** Let notations be as before. Then there exist linear forms  $\zeta_1, \ldots, \zeta_t \in (\eta_1, \ldots, \eta_s)$  for some  $t \leq \min\{r, s\}$  such that  $\overline{\zeta}_1, \ldots, \overline{\zeta}_t$  is a weak regular sequence in  $(S/I)_p$  and  $1 \in (\overline{\zeta}_1, \ldots, \overline{\zeta}_t)$ 

*Proof.* Let  $z_i := \sum_{j=1}^s \lambda_{ij} \overline{\eta}_j$  be a generic k-linear combination of  $\overline{\eta}_1, \ldots, \overline{\eta}_s$  for  $i = 1, \ldots, s$ . We obtain a maximal secant sequence  $z_1, \ldots, z_t$  for some  $t \leq s$ , as the field k is infinite. By this we mean that  $\dim(z_1, \ldots, z_i) = r - i - 1$  for  $i = 1, \ldots, t - 1$  and that  $1 \in (z_1, \ldots, z_t)$ . In particular  $t \leq \min\{r, s\}$  holds true.

The ring  $(S/I)_p$  is Cohen-Macaulay as it is a localization of a Cohen-Macaulay ring. Then  $z_1, \ldots, z_t$  is a regular sequence in  $(S/I)_p$ .

We take  $\zeta_i := \sum_{j=1}^s \lambda_{ij} \eta_j$  for i = 1, ..., t. Thus  $\zeta_i \in (\eta_1, ..., \eta_s)$  and  $\overline{\zeta}_i = z_i$  for i = 1, ..., s as required.

Thus we can suppose without loss of generality that  $\overline{\eta}_1, \ldots, \overline{\eta}_s$  is a weak regular sequence in  $(S/I)_p$  and that  $s \leq r$  holds. We assume this from now on. Next we are going to show that  $\eta_1, \ldots, \eta_s$  can be replaced by polynomials of controlled degree which form a regular sequence in S/I (Corollary 1.3).

The following lemma is a generalization of [17, Remark 4].

**Lemma 1.2** Let be given a homogeneous unmixed ideal  $K \subseteq k[x_0, \ldots, x_n]$  and points  $\xi_1, \ldots, \xi_m \in \mathbb{P}^n$  not lying in V(K). Then there exists a homogeneous polynomial g in K such that deg  $g \leq \deg K$  and that  $g(\xi_i) \neq 0$  hold.

*Proof.* The case when K is a homogeneous prime ideal follows easily from [17, Remark 4].

Let us consider the general case. For each associated prime ideal P of K we take a homogeneous polynomial  $g_P$  such that  $\deg g_P \leq \deg P$  and  $g_P(\xi_i) \neq 0$  for  $i = 1, \ldots, m$ . Let  $Q_P$  be the corresponding P-primary ideal in the decomposition of K. Let  $l(Q_P)$  denote the length of  $Q_P$ , that is, the length of  $(S/Q_P)_P$  as a S/P-module. Then let

$$g := \prod_P g_P^{l(Q_P)},$$

where the product is taken over all the associated prime ideals of K. Then  $g(\xi_i) \neq 0$ for  $i = 1, \ldots, m$ , and we have also that the polynomial g lies in the ideal K by [6, Lemma 1] and the degree bound  $\deg g \leq \sum_P l(Q_P) \deg P = \deg K$  holds true by [36, Proposition 1.49]. It follows that g satisfies the stated conditions.  $\Box$ 

In the sequel we shall denote by  $J_i$  the contraction to the ring S/I of the ideal  $(\overline{\eta}_1, \ldots, \overline{\eta}_i) \subseteq (S/I)_p$  and by  $\delta_i$  the degree of the homogeneous ideal  $J_i$  for  $i = 1, \ldots, s$ .

**Corollary 1.3** Let notations be as before. Then there exist homogeneous elements  $h_1, \ldots, h_s \in S/I$  satisfying the following conditions:

- i)  $h_i \equiv p^{c_i} \eta_i \mod J_{i-1}$  for some  $c_i \ge 0$ ,
- ii)  $h_1, \ldots, h_s$  is a regular sequence,
- *iii)*  $\deg h_i \leq \deg J_{i-1} + \deg p 1$ ,

for i = 1, ..., s.

*Proof.* We proceed by induction on *i*. By assumption *p* is not a zero-divisor in S/I so that the canonical morphism  $S/I \to (S/I)_p$  is injective. The fact that  $\overline{\eta}_1$  is not a zero-divisor in  $(S/I)_p$  implies then that  $\eta_1$  is not a zero-divisor in S/I.

Now let  $i \geq 2$  and assume that the elements  $h_1, \ldots, h_{i-1}$  are already constructed. Let  $H_{i-1}$  denote the ideal spanned by  $h_1, \ldots, h_{i-1}$  in S/I. Let  $H_{i-1} = (\bigcap_j Q_j) \cap (\bigcap_l R_l)$  be the primary decomposition of  $H_{i-1}$ , with  $p \notin \sqrt{Q_j}$  and  $p \in \sqrt{R_l}$ . Our objective is to find a homogeneous element  $h_i$  in S/I not lying in any of the associated primary ideals of  $H_{i-1}$ .

We recall that the ideal  $H_{i-1}$  has no imbedded component as it is spanned by a regular sequence in a Cohen–Macaulay ring. On the other hand the ideal  $J_{i-1}$  has the primary decomposition  $\cap_j Q_j$  and so it follows that  $V(R_l) \not\subseteq V(J_{i-1})$  holds for each l. We choose a point  $\xi_l$  lying in  $V(R_l) - V(J_{i-1})$  and a homogeneous element  $g \in J_{i-1}$  such that  $\deg g \leq \deg J_{i-1}$  and  $g(\xi_l) \neq 0$  for each l. The existence of g is guaranteed by the previous lemma. By eventually multiplying g with linear forms we can suppose without loss of generality that  $\deg g = c_i \deg p + 1$  holds for some  $c_i \geq 0$ . In particular we can assume that  $\deg g \leq \deg J_{i-1} + \deg p - 1$  holds. Finally we set

$$h_i := ag + p^{c_i}\eta_i$$

for some indeterminate scalar  $a \in k$ . Then  $h_i$  is homogeneous and  $h_i \equiv p^{c_i}\eta_i \mod J_{i-1}$  holds true. Therefore  $h_i$  does not belong to  $\sqrt{Q_j}$ , as both p and  $\eta_i$  are not zero-divisors modulo  $J_{i-1}$ . We have also that  $h_i(\xi_l) = a g(\xi_l) + (p^{c_i}\eta_i)(\xi_l) \neq 0$  for a generic choice of a, which forces  $h_i \notin \sqrt{R_l}$ .

We fix the following notation. Let  $h_1, \ldots, h_s \in S/I$  be the homogeneous polynomials introduced in Corollary 1.3, and let  $H_i := (h_1, \ldots, h_i)$  and  $L_i := (\eta_1, \ldots, \eta_i)$  denote the homogeneous ideals successively generated by  $h_1, \ldots, h_s$  and  $\eta_1, \ldots, \eta_s$  respectively.

Let us write  $h_i = l_i + p^{c_i} \eta_i$  for some  $l_i \in J_{i-1}$  and  $c_i \ge 0$ . Then set  $\gamma_i := \delta_{i-1} - \delta_i$ , and let  $\lambda_i := \sum_{j=1}^i (\gamma_j + c_j)$  and  $\mu_i := \sum_{j=1}^i ((i-j+1)\gamma_j + (i-j)c_j)$  for  $i = 1, \ldots, s$ .

Given an ideal K in S/I we denote by  $K^u$  the unmixed part of K, that is, the unmixed ideal given as the intersection of the primary components of K of maximal dimension.

**Lemma 1.4** Let be given an element  $q \in J_i$  for some  $1 \leq i \leq s$ . Then  $p^{\gamma_i}q \in (J_{i-1}, \eta_i)^u$ .

Proof. Let  $(\bigcap_j Q_j) \cap (\bigcap_l R_l)$  be the primary decomposition of the ideal  $(J_{i-1}, \eta_i)^u$ , with  $p \notin \sqrt{Q_j}$  and  $p \in \sqrt{R_l}$ . Then the ideal  $J_i$  has the primary decomposition  $\bigcap_j Q_j$ . Let  $K_i := \bigcap_l R_l$  be the intersection of the other primary components. Then  $K_i$  is an unmixed ideal which lies in the hypersurface  $\{p = 0\}$ .

The ideals  $(J_{i-1}, \eta_i)^u$  and  $(J_{i-1}, \eta_i)$  have the same degree because they only differ in an ideal of codimension at least i + 1. The degree of  $(J_{i-1}, \eta_i)$  is  $\delta_{i-1}$ , as  $\eta_i$  is not a zero-divisor mod  $J_{i-1}$ , and so the degree of  $K_i$  equals  $\gamma_i = \delta_{i-1} - \delta_i$ . Therefore  $p^{\gamma_i}$ lies in the ideal  $K_i$  [6, Lemma 1] and we conclude that  $p^{\gamma_i}q \in (\bigcap_j Q_j) \cap (\bigcap_l R_l) =$  $(J_{i-1}, \eta_i)^u$  as stated.  $\Box$ 

The following two statements (Lemmas 1.5 and 1.6) are plain extensions of [9, Lemmas 6.1 and 6.2].

**Lemma 1.5** Let be given an element  $q \in J_i$  for some  $1 \le i \le s$ . Then  $p^{\lambda_i}q \in H_i$ .

*Proof.* We proceed by induction on *i*. First  $p^{\gamma_1}q \in (\eta_1)^u$  by Lemma 1.4. We have also that  $(\eta_1)^u = (\eta_1)$  and so the assertion is true for i = 1.

Let  $i \geq 2$  and assume that the statement holds for i - 1. By Lemma 1.4,  $p^{\gamma_i}q \in (J_{i-1}, \eta_i)^u$ , that is,  $p^{\gamma_i}q$  belongs to the intersection of the primary components of dimension r - i of the ideal  $(J_{i-1}, \eta_i)$ . The intersection of the other primary components is an ideal of codimension at least i + 1. Then there exists a regular sequence  $w_1, \ldots, w_{i+1}$  in this ideal, as S/I is a Cohen–Macaulay ring. We have that  $w_j p^{\gamma_i}q \in (J_{i-1}, \eta_i)$  and so there exist  $u_j \in J_{i-1}$  and  $v_j \in S/I$  such that  $w_j p^{\gamma_i}q = u_j + v_j\eta_i$  for  $j = 1, \ldots, i + 1$ . Then

$$w_j p^{\gamma_i + c_i} q = p^{c_i} u_j + p^{c_i} v_j \eta_i = p^{c_i} u_j + v_j (h_i - l_i) = (p^{c_i} u_j - v_j l_i) + v_j h_i.$$

Therefore  $p^{\gamma_i+c_i}u_j - v_jl_i \in J_{i-1}$  and by the inductive hypothesis  $p^{\lambda_{i-1}}(p^{\gamma_i+c_i}u_j - v_jl_i)$  lies in the ideal  $H_{i-1}$ . Then  $w_j p^{\lambda_i} q \in H_i$  holds for  $j = 1, \ldots, i+1$ , as  $\lambda_i = \lambda_{i-1} + \gamma_i - c_i$ .

The ideal  $H_i$  is spanned by a regular sequence  $h_1, \ldots, h_i$  and so it is an unmixed ideal of dimension r - i. Thus for each associated prime ideal P of  $H_i$  there exists some j such that  $w_j \notin P$ . We conclude that  $p^{\lambda_i}q \in H_i$ .

**Lemma 1.6** Let be given an element  $q \in J_i$  for some  $1 \le i \le s$ . Then  $p^{\mu_i}q \in L_i$ .

*Proof.* We shall proceed by induction on *i*. The case i = 1 follows in the same way as in the preceding lemma because  $L_1 = H_1$  and  $\mu_1 = \lambda_1$ .

Let  $i \geq 2$ . Then  $p^{\lambda_i}q$  lies in  $H_i$  by Lemma 1.5. Let us write  $p^{\lambda_i}q = u + v h_i$  for some  $u \in H_{i-1}$  and  $v \in S/I$ . Therefore  $p^{\lambda_i}q - v h_i \in H_{i-1}$  and thus  $p^{\lambda_i}q - p^{c_i}v \eta_i$ lies in the ideal  $J_{i-1}$  because  $H_{i-1} \subseteq J_{i-1}$  and  $h_i \equiv p^{c_i} \eta_i \mod J_{i-1}$ . This implies in turn that  $p^{\lambda_i - c_i}q - v \eta_i \in J_{i-1}$ .

From the inductive hypothesis we get that  $p^{\mu_{i-1}}(p^{\lambda_i-c_i}q-v\eta_i)$  lies in  $L_{i-1}$  and so  $p^{\mu_{i-1}+\lambda_i-c_i}q \in L_i$ . The statement follows from the observation that  $\mu_i = \mu_{i-1}+\lambda_i-c_i$ .  $\Box$ 

**Main Lemma 1.7** Let  $I \subseteq k[x_0, \ldots, x_n]$  be a homogeneous Cohen–Macaulay ideal of dimension r. Let be given in addition a homogeneous element  $p \in k[x_0, \ldots, x_n]/I$ which is not a zero–divisor and linear forms  $\eta_1, \ldots, \eta_s \in k[x_0, \ldots, x_n]/I$  such that p lies in the radical of the ideal  $(\eta_1, \ldots, \eta_s)$ . Then

$$p^D \in (\eta_1, \ldots, \eta_s)$$

holds, with  $D := \min\{r, s\}^2 \deg I$ .

*Proof.* By Lemma 1.1 we can suppose without loss of generality that  $\overline{\eta}_1, \ldots, \overline{\eta}_s$  is a weak regular sequence in  $(S/I)_p$  and that  $s \leq r$ . After Lemma 1.6 it only remains to bound  $\mu_s$ . We make use of the estimations  $\gamma_i, c_i \leq \delta_{i-1}$  and we get the bound

$$\mu_{s} = \sum_{j=1}^{s} ((s-j+1)\gamma_{j} + (s-j)c_{j})$$
  

$$\leq \sum_{j=1}^{s} ((s-j+1)\delta_{j-1} + (s-j)\delta_{j-1}) \leq s^{2} \deg I.$$

Compare with Caniglia, Galligo and Heintz [7, Proposition 10], Smietanski [29, Lemma 1.44], and Sturmfels, Trung and Vogel [33, Theorems 2.1 and 5.3]

The rest of the section is devoted to the extension of the previous result to the case when we are given homogeneous elements of arbitrary degree instead of linear forms. First we establish some generalities about the Veronese imbedding.

Let us denote by N the integer  $\binom{n+d}{d} - 1$  and let  $a_0, \ldots, a_N$  denote the exponents of the different monomials of degree d in S. Let

$$v_d: \mathbb{P}^n \to \mathbb{P}^N, \qquad \qquad x := (x_0: \dots: x_n) \mapsto (x^{a_0}: \dots: x^{a_N})$$

be the Veronese map. This is a regular morphism of projective varieties and so its image is a closed subvariety of  $\mathbb{P}^N$ . This variety is called the Veronese variety and it is denoted by  $v_{n,d}$ . Let  $I(v_{n,d})$  be its defining ideal and let us denote by  $S(d) := k[y_0, \ldots, y_N]/I(v_{n,d})$  its homogeneous coordinate ring. The Veronese map induces then an inclusion of k-algebras  $i_d : S(d) \hookrightarrow S$  defined by  $y_j \mapsto x^{a_j}$  for  $j = 0, \ldots, N$ .

Let be given an ideal J in S and let J(d) denote its contraction to the ring S(d). Identifying the quotient ring S(d)/J(d) with its image in S/J through the inclusion  $i_d: S(d)/J(d) \hookrightarrow S/J$  we obtain the decomposition in graded parts

$$S(d)/J(d) = \bigoplus_j (S/J)_{dj}.$$

Let  $h_{J(d)}$  and  $h_J$  denote the Hilbert functions of J(d) and J respectively. Then  $h_{J(d)}(m) = h_J(dm)$  for  $m \in \mathbb{N}$ . It follows that the ideals J(d) and J have the same dimension and that their degrees are related by the formula  $\deg J(d) = d^{\dim J-1} \deg J$ .

**Lemma 1.8** Let J be a homogeneous Cohen–Macaulay ideal in S and let J(d) denote its contraction to the ring S(d). Then J(d) is a Cohen–Macaulay ideal.

*Proof.* Let us denote by A and B the quotient rings S(d)/J(d) and S/J respectively. We identify A with its image in B through the inclusion  $i_d$ .

We are going to prove that A is a Cohen–Macaulay ring. As A is a graded ring it suffices to exhibit a regular sequence of homogeneous elements of length equal to the dimension of A.

Let e denote the dimension of the ring B, which is also the dimension of A. Let  $\beta_1, \ldots, \beta_e$  be a regular sequence in B of homogeneous elements. Let  $\alpha_i := \beta_i^d$  for  $i = 1, \ldots, e$ . Then  $\alpha_1, \ldots, \alpha_e$  are elements of A which form a regular sequence in B, by [22, Theorem. 16.1]. We affirm that they also form a maximal regular sequence in A. We only have to prove that  $\alpha_i$  is not a zero-divisor in  $A/(\alpha_1, \ldots, \alpha_{i-1})$  for  $i = 1, \ldots, e$ . Let  $\zeta \in A$  be an element such that  $\zeta \alpha_i \in (\alpha_1, \ldots, \alpha_{i-1})$ . Then there exist homogeneous elements  $\zeta_1, \ldots, \zeta_{i-1} \in B$  such that  $\zeta = \zeta_1 \alpha_1 + \cdots + \zeta_{i-1} \alpha_{i-1}$  because  $\alpha_1, \ldots, \alpha_{i-1}$  is a regular sequence in B. An easy verification shows that  $\zeta_1, \ldots, \zeta_{i-1}$  can be chosen to lie in A, from where it follows that  $\zeta \in (\alpha_1, \ldots, \alpha_{i-1})$ .  $\Box$ 

**Theorem 1.9** Let  $I \subseteq k[x_0, \ldots, x_n]$  be a homogeneous Cohen–Macaulay ideal. Let be given in addition an homogeneous element  $p \in k[x_1, \ldots, x_n]/I$  which is not a zero–divisor and homogeneous elements  $f_1, \ldots, f_s \in k[x_0, \ldots, x_n]/I$  such that p lies in the radical of the ideal  $(f_1, \ldots, f_s)$ . Let r denote the dimension of I and let d the maximum degree of  $f_i$  for  $i = 1, \ldots, s$ . Then

$$p^D \in (f_1, \ldots, f_s)$$

holds, with  $D := r^2 d^r \deg I$ .

*Proof.* First we note that the zero locus in V(I) of the polynomials  $\{f_i\}_i$  equals the zero locus in V(I) of the polynomials  $\{x_j^{d-\deg f_i}f_i\}_{ij}$ . We have also that  $x_j^{d-\deg f_i}f_i$  lies in the ideal  $(f_1, \ldots, f_s)$  for all i and j. Therefore we can suppose without loss of generality that  $f_i$  is a homogeneous polynomial of degree d for  $i = 1, \ldots, s$ . We note however that the number of input polynomials have been enlarged in this preparative step.

Let  $i_d : S(d) \hookrightarrow S$  be the inclusion of k-algebras induced by the Veronese map and let I(d) denote the contraction of the ideal I to the ring S(d). Then we have the inclusion  $i_d : S(d)/I(d) \hookrightarrow S/I$  and the decomposition in graded parts  $i_d(S(d)/I(d)) = \bigoplus_j (S/I)_{dj}$ . We take a linear form  $\eta_i \in S(d)/I(d)$  such that  $i_d(\eta_i) =$  $f_i$  for  $i = 1, \ldots, s$ , which exists as the inclusion  $i_d$  is a bijection in degree one. We take also a homogeneous element  $q \in S(d)/I(d)$  such that  $i_d(q) = p^d$ .

The map  $v_d: V(I) \to V(I(d))$  is a dominant regular map of projective varieties and so it is surjective. Therefore the zero locus of the linear forms  $\eta_1, \ldots, \eta_s$  lies in the image of the zero locus of the polynomials  $f_1, \ldots, f_s$ . The common zeros of  $f_1, \ldots, f_s$  lie in the hypersurface  $\{p^d = 0\}$  of V(I) and we have in addition that  $v_d(\{p^d = 0\}) = \{q = 0\}$ . Then the subvariety of V(I(d)) defined by  $\eta_1, \ldots, \eta_s$  lies in the hypersurface  $\{q = 0\}$ . By Lemma 1.8 the ideal I(d) is Cohen-Macaulay, and we have also that q is not a zero-divisor modulo I(d). Then we are in the hypothesis of the Main Lemma 1.7. As a consequence we obtain that

$$q^{r^2 \deg I(d)} \in (\eta_1, \dots, \eta_s)$$

holds. Finally we apply the morphism  $i_d$  to the previous expression and we get that

$$p^{dr^2(d^{r-1}\deg I)} \in (f_1,\ldots,f_s)$$

holds, as deg  $I(d) = d^{r-1} \deg I$ .

**Corollary 1.10** Let be given an ideal  $I \subseteq k[x_1, \ldots, x_n]$ . Assume furthermore that the homogenization  $I^h$  of the ideal I in the ring  $k[x_0, \ldots, x_n]$  is a Cohen-Macaulay ideal. Let  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  be polynomials without common zeros in the affine variety V(I). Then there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  such that

$$1 \equiv g_1 f_1 + \dots + g_s f_s \tag{mod } I)$$

holds, with  $\deg g_i f_i \leq (r+1)^2 d^{r+1} \deg I^h$  for i = 1, ..., s.

*Proof.* By assumption the ideal  $I^h$  is a Cohen–Macaulay homogeneous ideal of dimension r + 1. We have also that  $x_0$  is not a zero–divisor modulo  $I^h$ .

Let  $f_i^h$  denote the homogenization of  $f_i$  for i = 1, ..., s. The homogeneous polynomials  $f_1^h, \ldots, \tilde{f}_s^h$  have no common zero in  $V(\tilde{I})$  outside the hyperplane  $\{x_0 = 0\}$ . By Theorem 1.9 there exist homogeneous polynomials  $v_1, \ldots, v_s \in S$  such that

$$x_0^{(r+1)^2 d^{r+1}} = v_1 f_1^h + \ldots + v_s f_s^h \qquad (\text{mod } I^h)$$

holds, with  $\deg v_i f_i^h = (r+1)^2 d^{r+1}$ . The corollary then follows by evaluating  $x_0 := 1$ .

Let notations be as in Corollary 1.10. In the case when I is the zero ideal, that is, in the setting of the classic effective Nullstellensatz, we get the degree bound

$$\deg g_i f_i \le (r+1)^2 \, d^{r+1}.$$

#### 2. The Sparse Effective Nullstellensatz

In this section we shall devote to the sparse effective Nullstellensätze (Theorems 1 and 2) and to the derivation of some of their consequences.

First we introduce notation and state some basic facts from polyhedral geometry and toric varieties. We refer to the books [12] and [32] for the proofs of these facts and for a more general background on these topics.

Let us be given a finite set of integer vectors  $\mathcal{A} \subseteq \mathbb{Z}^n$ . The convex hull of  $\mathcal{A}$  as a subset of  $\mathbb{R}^n$  is denoted by  $\operatorname{conv}(\mathcal{A})$ . The cone over  $\operatorname{conv}(\mathcal{A})$  is denoted by  $\operatorname{pos}(\mathcal{A})$ .

The set  $\mathcal{A}$  is graded if there exists an integer vector  $\omega \in \mathbb{Z}^n$  such that  $\langle a, \omega \rangle = 1$  holds for every  $a \in \mathcal{A}$ , that is, when the set  $\mathcal{A}$  lies in an affine hyperplane which does not contain the origin.

Let  $\mathbb{Z}\mathcal{A}$  denote the  $\mathbb{Z}$ -module generated by  $\mathcal{A}$ . Let  $\mathbb{R}\mathcal{A}$  denote the linear space spanned by  $\mathcal{A}$ , so that  $\mathbb{Z}\mathcal{A}$  is a lattice in  $\mathbb{R}\mathcal{A}$ . Let  $\rho$  denote the dimension of this linear space. Then we consider the euclidian volume form in  $\mathbb{R}\mathcal{A}$ , normalized in such a way that the lattice  $\mathbb{Z}\mathcal{A}$  has covolume  $\rho$ ! or equivalently, such that each primitive lattice simplex has unit volume. The normalized *volume* Vol( $\mathcal{A}$ ) of the set  $\mathcal{A}$  is defined as the volume of its convex hull with respect to this volume form.

We get readily from the definition the bound

$$\operatorname{Vol}(\mathcal{A}) \le \rho! \operatorname{vol}(\operatorname{conv}(\mathcal{A})),$$

where vol  $(\operatorname{conv}(\mathcal{A}))$  denotes the volume of the convex hull of  $\mathcal{A}$  with respect to the usual non-normalized volume form of  $\mathbb{R}^n$ .

Let  $\mathbb{N}\mathcal{A}$  denote the semigroup spanned by  $\mathcal{A}$ . This semigroup is always contained in the semigroup  $\operatorname{pos}(\mathcal{A}) \cap \mathbb{Z}\mathcal{A}$ . The set  $\mathcal{A}$  is said to be *normal* or *saturated* if the equality  $\mathbb{N}\mathcal{A} = \operatorname{pos}(\mathcal{A}) \cap \mathbb{Z}\mathcal{A}$  holds.

A polytope  $\mathcal{P}$  is said to be *integral* if it is the convex hull of a finite set of integer vectors.

Let S be an integral simplex. Then S is said to be *unimodular* if its interior contains no integral vector. Let  $\{s_1, \ldots, s_k\}$  be the set of vertices of S. Then we have that S is unimodular if and only if the set of integer vectors  $\{s_2-s_1, \ldots, s_k-s_1\}$  is normal.

Let  $\mathcal{P}$  be an integral polytope. A subdivision of  $\mathcal{P}$  is said to be *unimodular* if it is formed by unimodular integral simplices.

Given integral polytope  $\mathcal{P}$  in  $\mathbb{R}^n$ , we denote by  $\mathcal{A}(\mathcal{P})$  the set  $\{1\} \times (\mathcal{P} \cap \mathbb{Z}^n)$ , which is a graded set of integral vectors in  $\mathbb{Z}^{n+1}$ . We note that the set  $\mathcal{A}(\mathcal{P})$  is normal in the case when  $\mathcal{P}$  admits a unimodular subdivision.

With respect to toric geometry, we shall follow the lines of [32]. This point of view differs from the usual one in algebraic geometry. It is more combinatorial and suits better for our purposes.

Let us be given again a finite set of integer vectors  $\mathcal{A} = \{a_1, \ldots, a_N\}$  in  $\mathbb{Z}^n$ . We associate to the set  $\mathcal{A}$  the morphism

$$\varphi_{\mathcal{A}}: k[y_1, \dots, y_N] \to k[x_1, \dots, x_n, x_1^{-1}, \dots, x_n^{-1}], \qquad y_i \to x^{a_i}.$$

The kernel of this map is a prime ideal  $I_{\mathcal{A}}$  of  $k[y_1, \ldots, y_N]$ , called the *toric ideal* associated to the set  $\mathcal{A}$ . This ideal defines an *affine toric variety*  $X_{\mathcal{A}}$  as its zero locus in  $\mathbb{A}^N$ . This variety is irreducible and its dimension equals the rank of the  $\mathbb{Z}$ -module  $\mathbb{Z}\mathcal{A}$ .

The *k*-algebra  $k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$  is the coordinate ring of the torus  $(\overline{k}^*)^n$ . Thus the map  $\varphi_{\mathcal{A}}$  induces a dominant map  $(\overline{k}^*)^n \to X_{\mathcal{A}}$ . The image of this map is called the *torus*  $T_{\mathcal{A}}$  of the affine toric variety  $X_{\mathcal{A}}$ . This torus equals the open set  $\{y_1 \cdots y_N \neq 0\}$  of  $X_{\mathcal{A}}$ .

The ideal  $I_{\mathcal{A}}$  is homogeneous if and only if the set  $\mathcal{A}$  is graded. In this case the set  $\mathcal{A}$  defines a *projective toric variety*  $Y_{\mathcal{A}}$  as the zero locus of the ideal  $I_{\mathcal{A}}$  in the projective space  $\mathbb{P}^{N-1}$ . The dimension of  $Y_{\mathcal{A}}$  equals then the rank of  $\mathbb{Z}\mathcal{A}$  minus one, and its degree equals the normalized volume of the set  $\mathcal{A}$ .

Let  $\mathcal{A} = \{a_1, \ldots, a_N\} \subseteq \mathbb{Z}^n$  be a graded set. The intersection of the projective variety  $Y_{\mathcal{A}}$  with the affine chart  $\{y_i \neq 0\} \cong \mathbb{A}^{N-1}$  equals the affine toric variety associated to the set

$$\mathcal{A} - a_i := \{a_1 - a_i, \dots, a_{i-1} - a_i, a_{i+1} - a_i, \dots, a_N - a_i\}.$$

In fact  $Y_{\mathcal{A}}$  is irredundantly covered by the affine varieties  $X_{\mathcal{A}-a_i}$ , where  $a_i$  runs over the vertices of the polytope conv $(\mathcal{A})$ .

The k-algebra  $k[y_1, \ldots, y_N]/I_A$  is canonically isomorphic to the semigroup algebra  $k[\mathbb{N}A]$ . This algebra is normal if and only if the set A is normal. We recall Hochster's theorem that the k-algebra  $k[\mathbb{N}A]$  is a Cohen-Macaulay domain in the case when the set A is normal.

Let be given an integral polytope  $\mathcal{P}$  in  $\mathbb{R}^n$ . This polytope determines a fan  $\Delta_{\mathcal{P}}$ and an *abstract* complete toric variety  $X_{\mathcal{P}} = X(\Delta_{\mathcal{P}})$ . This variety comes equipped with an ample Cartier divisor  $D_{\mathcal{P}}$ . This Cartier divisor defines then a map  $\varphi_{\mathcal{P}}$ :  $X_{\mathcal{P}} \to \mathbb{P}^{N-1}$ , where N denotes the cardinality of the set  $\{\mathcal{P} \cap \mathbb{Z}^n\}$ . The image of this map is the projective variety  $Y_{\mathcal{A}(\mathcal{P})}$ , where the set  $\mathcal{A}(\mathcal{P})$  is defined as before as  $\{1\} \times (\mathcal{P} \cap \mathbb{Z}^n)$  [12, Section 3.4]. The divisor  $(n-1)D_{\mathcal{P}}$  is very ample [10], and so the graded set  $\mathcal{A}((n-1)\mathcal{P})$  is normal.

**Theorem 2.11** Let be given polynomials  $p, f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  such that p lies in the radical of the ideal  $(f_1, \ldots, f_s)$ . Let  $\mathcal{P}$  be an integral polytope which contains the Newton polytope of the polynomials  $1, x_1, \ldots, x_n, f_1, \ldots, f_s$ . Assume furthermore that  $\mathcal{A}(\mathcal{P})$  is a normal set of integer vectors in  $\mathbb{Z}^{n+1}$ . Then there exist  $D \in \mathbb{N}$  and  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  such that

$$p^D = g_1 f_1 + \dots + g_s f_s$$

holds, with  $D \leq n! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P})$  and  $\mathcal{N}(g_i f_i) \subseteq (1 + \deg p) n! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P}) \mathcal{P}$  for  $i = 1, \ldots, s$ .

*Proof.* Let  $\mathcal{B} = \{b_0, \ldots, b_N\}$  denote the set of integer vectors  $\mathcal{P} \cap \mathbb{Z}^n$ , so that  $\mathcal{A}(\mathcal{P}) = \{1\} \times \mathcal{B}$ . Assume that  $b_0 = (0, \ldots, 0)$ . We consider the morphism of k-algebras

$$\psi: k[y_1, \dots, y_N] \to k[x_1, \dots, x_n], \qquad \qquad y_i \mapsto x^{b_i}$$

The kernel of this morphism is the defining ideal  $I_{\mathcal{B}-b_0}$  of the affine toric variety  $X_{\mathcal{B}-b_0}$ . This affine variety is the intersection of the projective toric variety  $Y_{\mathcal{A}(\mathcal{P})}$  with the affine cart  $\{y_0 \neq 0\}$  of  $\mathbb{P}^N$ . In addition the map  $\psi$  induces an isomorphism  $\mathbb{A}^n \to X_{\mathcal{B}-b_0}$ .

Let  $\zeta_i$  be a polynomial of degree one in  $k[y_1, \ldots, y_N]$  such that  $\psi(\zeta_i) = f_i$  for  $i = 1, \ldots, s$ . We take also a polynomial q in  $k[y_1, \ldots, y_N]$  of degree less or equal to the degree of p such that  $\psi(q) = p$ . Then  $\zeta_1, \ldots, \zeta_s$  have no common zero in  $X_{\mathcal{B}-b_0}$  outside the hypersurface  $\{q = 0\}$ .

Let  $\eta_1, \ldots, \eta_s, u$  denote the homogenization of  $\zeta_1, \ldots, \zeta_s, q$  in  $k[y_0, \ldots, y_N]$  respectively. Then the linear forms  $\eta_1, \ldots, \eta_s$  have no common zero in  $Y_{\mathcal{A}(\mathcal{P})}$  outside the hypersurface  $\{y_0 \ u = 0\}$ .

By assumption the set  $\mathcal{A}(\mathcal{P})$  is normal, and so  $I_{\mathcal{A}(\mathcal{P})}$  is a Cohen–Macaulay prime homogeneous ideal of  $k[y_0, \ldots, y_N]$  of dimension less or equal that n + 1. We have also that  $y_0 u$  is not a zero–divisor modulo  $I_{\mathcal{A}(\mathcal{P})}$ . Then we are in the hypothesis of the Main Lemma 1.7. Let D denote the integer  $\min\{n+1,s\}^2 \deg Y_{\mathcal{A}(\mathcal{P})}$ . We obtain that there exist homogeneous elements  $\alpha_1, \ldots, \alpha_s \in k[y_0, \ldots, y_N]/I_{\mathcal{A}(\mathcal{P})}$  of degree  $(1 + \deg u) D - 1$  satisfying

$$(y_0 u)^D = \alpha_1 \eta_1 + \dots + \alpha_s \eta_s.$$

Finally we evaluate  $y_0 := 1$  and we apply the map  $\psi$  to the preceding identity. We get

$$p^D = g_1 f_1 + \dots + g_s f_s,$$

where we have set  $g_i(x) := \alpha_i(1, x^{b_1}, \ldots, x^{b_N})$  for  $i = 1, \ldots, s$ . We have the estimations deg  $u \leq \deg p$  and deg  $Y_{\mathcal{A}(\mathcal{P})} \leq n! \operatorname{vol}(\mathcal{P})$ . We conclude that  $D \leq n! \min\{n+1,s\} \operatorname{vol}(\mathcal{P})$  and that the polytope  $\mathcal{N}(f_i g_i)$  is contained in  $((1 + \deg p) n! \min\{n+1,s\}^2 \operatorname{vol}(\mathcal{P})) \mathcal{P}$  for  $i = 1, \ldots, s$ .

We derive from the previous theorem the following degree bound.

**Corollary 2.12** Let notations be as in Theorem 2.11. Let d denote the maximum degree of the polynomials  $f_i$  for i = 1, ..., s. Then there exist  $D \in \mathbb{N}$  and  $g_1, ..., g_s \in k[x_1, ..., x_n]$  such that

 $p^D = g_1 f_1 + \dots + g_s f_s$ 

holds, with  $D \le n! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P})$  and  $\deg g_i f_i \le d(1 + \deg p) n! \min\{n+1, s, \}^2 \operatorname{vol}(\mathcal{P})$  for  $i = 1, \ldots, s$ .

We are going to show with an example that this degree bound can be much more precise than the usual one in the case when the input system is sparse. First we need the following auxiliary result.

**Lemma 2.13** Let  $\mathcal{P}_d$  denote the integral polytope  $\{x = (x_1, \ldots, x_n) \in \mathbb{R}^n : x_1, \ldots, x_n \geq 0, x_1 + \cdots + x_{n-1} \leq 1, x_n \leq d\}$  for some  $d \in \mathbb{N}$ . Then  $\mathcal{P}_d$  admits a unimodular subdivision.

*Proof.* Let us denote by  $e_1, \ldots, e_{n-1}$  the standard unit vectors of  $\mathbb{R}^{n-1}$ . Let  $\mathcal{S}$  denote the simplex of  $\mathbb{R}^{n-1}$  with vertices  $(0, \ldots, 0), e_1, \ldots, e_{n-1}$ , so that  $\mathcal{P}_d = \mathcal{S} \times [0, d]$ .

We have the subdivision  $\mathcal{P}_d = \bigcup_{i=0}^{d-1} (\mathcal{P}_1 + ie)$ , where  $e \in \mathbb{R}^n$  denotes the vector  $(0, \ldots, 0, 1)$ . Thus it suffices to show that  $\mathcal{P}_1$  admits a unimodular subdivision.

Let  $\alpha_i, \beta_i \in \mathbb{R}^n$  denote the vectors  $e_i \times \{0\}, e_i \times \{1\}$  respectively, for  $i = 1, \ldots, n-1$ , and  $(0, \ldots, 0, 0), (0, \ldots, 0, 1)$  for i = n. Let  $\mathcal{S}_i$  be the integral simplex determined by the vectors  $\alpha_1, \ldots, \alpha_i, \beta_i, \ldots, \beta_n$  for  $i = 1, \ldots, n$ .

It is easily checked that the simplices  $S_i$  and  $S_j$  intersect in a proper face for  $i \neq j$ , so that  $S_1, \ldots, S_n$  are essentially disjoint. We have also that  $\operatorname{vol}(S) = 1/(n-1)!$ and that  $\operatorname{vol}(S_i) = 1/n!$  for  $i = 1, \ldots, n$ , and so  $S_1, \ldots, S_n$  form a unimodular subdivision of  $\mathcal{P}_1$  as desired.  $\Box$ 

**Example 2.14** Let  $f_1, \ldots, f_s \in k[x_1, \ldots, x_n]$  be polynomials without common zeros in  $\mathbb{A}^n$ . Assume in addition that  $\deg_{x_1,\ldots,x_{n-1}} f_i \leq 1$  and that  $\deg_{x_n} f_i \leq d$  holds for  $i = 1, \ldots, s$ . Let  $\mathcal{P}_d$  denote as before the polytope  $\{x \in \mathbb{R}^n : x_1, \ldots, x_n \geq 0, x_1 + \cdots + x_{n-1} \leq 1, x_n \leq d\}$ . Then  $\mathcal{P}_d$  contains the Newton polytope of the polynomials  $1, x_1, \ldots, x_n, f_1, \ldots, f_s$  and the set  $\mathcal{A}(\mathcal{P})$  is normal by the preceding lemma. Thus we are in the hypothesis of Corollary 2.12 and we can conclude that there exist  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  such that

$$1 = g_1 f_1 + \dots + g_s f_s$$

holds, with  $\mathcal{N}(g_i f_i) \subseteq d n \min\{n+1, s\}^2 \mathcal{P}_d$  for  $i = 1, \ldots, s$ , as the volume of  $\mathcal{P}_d$  equals d/(n-1)!. In particular we get the degree bound  $\deg g_i f_i \leq (n+1)^3 (d+1)^2$ , which is much sharper than the estimation  $\deg g_i f_i \leq (d+1)^n$  which follows from direct application of the usual degree bound.

Let notations be again as in Theorem 2.11. Let  $\mathcal{N}$  denote the Newton polytope of the polynomials  $1, x_1, \ldots, x_n, f_1, \ldots, f_s$  and let  $\mathcal{U}$  denote the unmixed volume of this polytope. Assume that  $n \geq 2$ . In this situation we can then take the polytope  $\mathcal{P}$  to be  $(n-1)\mathcal{N}$ . Then we get the bounds

$$D \le n^{n+2}\mathcal{U}, \qquad \qquad \mathcal{N}(g_i f_i) \subseteq \left((1 + \deg p) n^{n+3}\mathcal{U}\right) \mathcal{N}.$$

It is easy to check that these bounds hold also when n = 1. Thus Theorem 1 follows from this observation in the particular case p = 1. We observe that in this case the condition  $0 \in \mathcal{P}$  is redundant.

We obtain a similar result in the case of Laurent polynomials.

**Theorem 2.15** Let be given Laurent polynomials  $p, f_1, \ldots, f_s \in k[x_1^{-1}, \ldots, x_n^{-1}, x_1, \ldots, x_n]$ such that p lies in the radical of the ideal  $(f_1, \ldots, f_s)$ . Let  $\mathcal{P}$  be an integral polytope which contains the Newton polytope of  $p, f_1, \ldots, f_s$ . Let  $\rho$  denote its dimension. Assume furthermore that  $\mathcal{A}(\mathcal{P})$  is a normal set of integer vectors in  $\mathbb{Z}^{n+1}$ . Then there exist  $D \in \mathbb{N}$ ,  $a \in \mathbb{Z}^n$  and  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$  such that

$$p^D = g_1 f_1 + \dots + g_s f_s$$

holds, with  $D \leq \rho! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P})$ ,  $a \in (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 \mathcal{P}$  and  $\mathcal{N}(g_i f_i) \subseteq (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 \mathcal{P} - a$  for  $i = 1, \ldots, s$ .

*Proof.* As before, we denote by  $\mathcal{B} = \{b_0, \ldots, b_N\}$  the set of integer vectors  $\mathcal{P} \cap \mathbb{Z}^n$ . Assume for the moment that  $b_0 = (0, \ldots, 0)$ . We consider the morphism

$$\psi: k[y_1, \dots, y_N] \to k[x_1, \dots, x_n, x_1^{-1}, \dots, x_n^{-1}], \qquad \qquad y_i \mapsto x^{b_i}$$

The kernel of this morphism is the defining ideal  $I_{\mathcal{B}-b_0}$  of the affine toric variety  $X_{\mathcal{B}-b_0}$ . Let T denote the torus of this toric variety. Then we have that  $X_{\mathcal{B}-b_0}$  equals the intersection of the projective variety  $Y_{\mathcal{A}(\mathcal{P})}$  with the affine cart  $\{y_0 \neq 0\}$  of  $\mathbb{P}^N$ , and that T is also the torus of  $Y_{\mathcal{A}(\mathcal{P})}$ . We recall that this torus equals the open set  $\{y_0 \cdots y_N = 0\}$  of  $Y_{\mathcal{A}(\mathcal{P})}$ .

The map  $\psi$  induces a surjection  $(\overline{k}^*)^n \to T$ . Let  $\zeta_1, \ldots, \zeta_s, q$  be elements of degree one in  $k[y_1, \ldots, y_N]$  such that  $\psi(\zeta_i) = f_i$  for  $i = 1, \ldots, s$  and  $\psi(q) = p$ . Then  $\zeta_1, \ldots, \zeta_s$  have no common zero in T outside the hyperplane  $\{q = 0\}$ .

Let  $\eta_1, \ldots, \eta_s$ , *u* denote the homogenization of  $\zeta_1, \ldots, \zeta_s$ , *q* in  $k[y_0, \ldots, y_N]$  respectively. Then the linear forms  $\eta_1, \ldots, \eta_s$  have no common zero in  $Y_{\mathcal{A}(\mathcal{P})}$  outside the hypersurface  $\{y_0 \cdots y_N u = 0\}$ .

Let  $V(\eta_1, \ldots, \eta_s)$  denote the subvariety of  $Y_{\mathcal{A}(\mathcal{P})}$  defined by the linear forms  $\eta_1, \ldots, \eta_s$ . By Bézout inequality, the number of irreducible components of  $V(\eta_1, \ldots, \eta_s)$  does not exceed the degree of  $Y_{\mathcal{A}(\mathcal{P})}$ . Let us denote by  $\delta$  the degree of  $Y_{\mathcal{A}(\mathcal{P})}$ , so that  $\delta \leq \rho! \operatorname{vol}(\mathcal{P})$  holds. In our situation this implies that  $V(\eta_1, \ldots, \eta_s)$  lies in the union of at most  $\delta$  hyperplanes. These hyperplanes are defined by variables  $y_{i_1}, \ldots, y_{i_l}$ , and eventually also by the linear form u, depending on whether  $\eta_1, \ldots, \eta_s$  have a common zero in T in the hyperplane  $\{u = 0\}$  or not. Let  $\Pi$  denote the product of these equations, which is a polynomial of degree less or equal that  $\delta$ .

By assumption the set  $\mathcal{A}(\mathcal{P})$  is normal and so  $I_{\mathcal{A}(\mathcal{P})}$  is a Cohen–Macaulay prime homogeneous ideal of  $k[y_0, \ldots, y_N]$ . We have also that  $\Pi$  is not a zero–divisor modulo this ideal. Thus we are again in the hypothesis of the Main Lemma 1.7. Let E denote the integer  $\min\{n+1,s\}^2 \deg Y_{\mathcal{A}(\mathcal{P})}$ . Then there exist homogeneous elements  $\alpha_1, \ldots, \alpha_s \in k[y_0, \ldots, y_N]/I_{\mathcal{A}(\mathcal{P})}$  of degree  $\deg \Pi E - 1$  such that

$$\Pi^E = \alpha_1 \eta_1 + \dots + \alpha_s \eta_s$$

holds. We evaluate  $y_0 := 1$  and we apply the map  $\psi$  to the preceding identity. We get

$$p^D = g_1 f_1 + \dots + g_s f_s,$$

where we have set  $g_i(x) := (x^{b_{i_1}} \cdots x^{b_{i_l}})^{-1} \alpha_i(1, x^{b_1}, \dots, x^{b_N})$  for  $i = 1, \dots, s$  and D := E in the case when u appears as a factor of  $\Pi$  and D := 1 in the other case. Then  $D \leq \rho! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P})$  holds and the polytope  $\mathcal{N}(g_i f_i)$  is contained in  $(\rho! \operatorname{vol}(\mathcal{P}) E - 1) \mathcal{P} - (b_{i_1} + \cdots + b_{i_l})$  for  $i = 1, \dots, s$ . We have that  $\deg \Pi \leq \deg Y_{\mathcal{A}(\mathcal{P})} \leq \rho! \operatorname{vol}(\mathcal{P})$  and that  $i_1 + \ldots + i_k \in \deg Y_{\mathcal{A}(\mathcal{P})} \mathcal{P}$ .

Now we consider the general case. Let  $b_0$  be any integer vector in  $\mathcal{P}$ , and let  $\mathcal{Q}$  denote the polytope  $\mathcal{P} - b_0$ . By the previous considerations there exist  $D \in \mathbb{N}$ ,  $a_0 \in \mathbb{Z}^n$  and  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n, x_1^{-1}, \ldots, x_n^{-1}]$  such that

$$p^D = g_1 f_1 + \dots + g_s f_s$$

holds, with  $D \leq \rho! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{Q})$ ,  $a_0 \in \rho! \operatorname{vol}(\mathcal{Q}) \mathcal{Q}$  and  $\mathcal{N}(g_i f_i) \subseteq (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{Q}))^2 \mathcal{Q} - a_0$  for  $i = 1, \ldots, s$ .

Let *a* be the integer vector  $a_0 + (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 b_0$ . Then *a* lies in the polytope  $(\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 \mathcal{P}$  and we have also that  $\mathcal{N}(g_i f_i) \subseteq (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 \mathcal{P} - a$  holds for  $i = 1, \ldots, s$  as stated.  $\Box$ 

Let notations be as in Theorem 2.15. Let  $\mathcal{N}$  denote the Newton polytope of  $p, f_1, \ldots, f_s$  and let  $\mathcal{U}$  denote the unmixed volume of this polytope. Assume in addition that  $n \geq 2$ . In this situation we can then take the polytope  $\mathcal{P}$  to be  $(n-1)\mathcal{N}$ . We get the bounds

$$D \le n^{n+2} \mathcal{U}, \qquad \qquad \mathcal{N}(g_i f_i) \subseteq (n^{2n+3} \mathcal{U}) \mathcal{N} - a.$$

for some  $a \in (n^{2n+3}\mathcal{U})\mathcal{N}$ . As before, it is easy to verify that the same bounds hold also when n = 1. Thus Theorem 2 follows from this observation in the particular case h = 1.

Let be given a rational function  $q \in k(x_1, \ldots, x_n)$  and let q = f/g be a representation of q as the quotient of two polynomials without common factors. Then the *degree* of q is defined as  $\deg q := \max\{\deg f, \deg g\}$ .

We derive from Theorem 2.15 the following degree bound.

**Corollary 2.16** Let notations be as in Theorem 2.15. Let d denote the maximum degree of the Laurent polynomials  $f_i$  for i = 1, ..., s. Then there exist  $D \in \mathbb{N}$  and  $g_1, ..., g_s \in k[x_1, ..., x_n, x_1^{-1}, ..., x_n^{-1}]$  such that

$$p^D = g_1 f_1 + \dots + g_s f_s$$

holds, with  $D \leq \rho! \min\{n+1, s\}^2 \operatorname{vol}(\mathcal{P})$ ,  $a \in (\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2 \mathcal{P}$  and  $\deg(g_i f_i) \leq d((1 + \deg p)\rho! \min\{n+1, s\} \operatorname{vol}(\mathcal{P}))^2$  for  $i = 1, \ldots, s$ .

#### 3. Improved Bounds for the Degrees in the Nullstellensatz

In this section we consider the degree bounds in the Nullstellensatz. We shall apply the methods used in Section 1 in a direct way —without any reference to the Veronese map— in the setting of the classic effective Nullstellensatz. The proof follows closely the same lines and so we shall skip some verifications in order to avoid unnecessary repetitions.

Assume that we are given homogeneous polynomials  $f_1, \ldots, f_s$  in  $k[x_0, \ldots, x_n]$  without common zeros at finite distance. In this situation we are going to give a bound for the minimal  $D \in \mathbb{N}$  such that  $x_0^D \in (f_1, \ldots, f_s)$ .

We shall assume without loss of generality that  $s \leq n+1$  and that  $\overline{f}_1, \ldots, \overline{f}_s$ is a weak regular sequence in  $k[x_0, \ldots, x_n]_{x_0}$ . Let  $d_i$  denote the degree of  $f_i$  for  $i = 1, \ldots, s$ . We shall also suppose that  $d_2 \ge \cdots \ge d_s$  and that  $d_s \ge d_1$  hold. As before these polynomials can be obtained as linear combinations of the original polynomials, eventually multiplied by powers of  $x_0$ .

Let us denote by  $J_i$  the contraction to the ring S of the ideal  $(\overline{f}_1, \ldots, \overline{f}_i) \subseteq S_{x_0}$ for i = 1, ..., s. We make the convention  $J_0 := (0)$ .

**Lemma 3.17** Let notations be as before. Then there exist homogeneous polynomials  $h_1, \ldots, h_s \in k[x_0, \ldots, x_n]$  satisfying the following conditions:

- i)  $h_i \equiv x_0^{c_i} f_i \mod J_{i-1}$  for some  $c_i \in \mathbb{N}$ ,
- ii)  $h_1, \ldots, h_s$  is a regular sequence,
- *iii*) deg  $h_i < \max \{ \deg J_{i-1}, \deg f_i \},$

for i = 1, ..., s.

We introduce the following notation. Let  $\delta_i$  denote the degree of the homogeneous ideal  $J_i$  for  $i = 0, \ldots, s$ . We recall the Bézout bound  $\delta_i \leq \prod_{j=1}^i d_j$ . Then we denote by  $\gamma_i$  the integer  $d_i \delta_{i-1} - \delta_i$  for  $i = 1, \dots, \min\{n, s\}$  and  $\gamma_{n+1} := \delta_n + d_{n+1} - 1$ . We also let  $\delta := \max\{\delta_i : i = 1, ..., s - 1\}$  and  $d := \max\{d_i : i = 1, ..., s - 1\}$ .

We recall that given an ideal I in S we denote by  $I^u$  its unmixed part.

**Lemma 3.18** Let be given a polynomial  $q \in J_i$  for some  $1 \leq i \leq s$ . Then  $x_0^{\gamma_i} q \in$  $(J_{i-1},\eta_i)^u$ .

*Proof.* The case  $i \leq n$  is exactly as in Lemma 1.4. Thus we only consider the case i = n + 1.

The ideal  $J_n$  is has dimension one and its degree is  $\delta_n$ . Then  $(J_n, f_{n+1})_m = S_m$ for  $m \ge \delta_n + d_{n+1} - 1$  as  $f_{n+1}$  is not a zero-divisor modulo  $J_n$  [30, Theorem. 2.23]. It follows that  $x_0^{\gamma_{n+1}} \in (J_n, f_{n+1})$  and in particular  $x_0^{\gamma_{n+1}} q \in (J_n, f_{n+1})^u$ . 

Now let  $h_1, \ldots, h_s$  be the homogeneous polynomials introduced in Lemma 3.17. We set  $\mu_i := \sum_{j=2}^{i} ((i-j+1)\gamma_j + (i-j)c_j)$  for  $i = 1, \dots, \min\{n, s\}$  and  $\mu_{n+1} :=$  $\mu_n + \gamma_{n+1}$ , where  $c_i$  denotes the integer deg  $h_i - \text{deg } f_i$ .

We denote by  $L_i$  the homogeneous ideal  $(f_1, \ldots, f_i)$  for  $i = 1, \ldots, s$ .

**Lemma 3.19** Let be given a polynomial  $q \in J_i$  for some  $1 \leq i \leq s$ . Then  $x_0^{\mu_i} q \in L_i$ .

*Proof.* The case  $i \leq n$  is exactly as in Lemma 1.6. Thus we only consider the case i = n + 1.

By the previous lemma  $x_0^{\gamma_{n+1}} q \in (J_n, f_{n+1})^u = (J_n, f_{n+1})$  and so  $x_0^{\gamma_{n+1}} q - u f_{n+1} \in$  $J_n$  for some polynomial  $u \in S$ . We apply then the inductive hypothesis and we obtain that  $x_0^{\mu_n}(x_0^{\gamma_{n+1}}q - uf_{n+1}) \in L_n$  from where it follows that  $x_0^{\mu_{n+1}}q \in L_{n+1}$ .  $\Box$ 

Thus it only remains to bound  $\mu_s$ . We shall be concerned with two different types of bounds. One depends as usual on the number of variables and on the degrees of the input polynomials, and the other depends also on the degree of some ideals associated to these polynomials.

**Lemma 3.20** Let notations be as before. Then  $\mu_s \leq \min\{n,s\}^2 d\delta$ . In the case when deg  $f_i \geq 2$  for  $i = 1, \ldots, s$  we have that  $\mu_s \leq 2 \prod_{j=1}^{\min\{n,s\}} d_j$ .

*Proof.* We decompose the integer  $\mu_s$  in two terms and we estimate them separately. First we consider the term  $\sum_{j=2}^{s} (s-j) c_j$ . We have that  $c_i \leq \max\{\delta_{i-1} - d_i, 0\}$ . In particular  $c_2 = 0$  as  $\delta_1 = d_1$  and  $d_1 \leq d_2$ . Then

$$\sum_{j=2}^{s} (s-j) c_j \leq \sum_{j=3}^{s-1} (s-j) (d_1 \cdots d_{j-1} - d_j)$$
  
$$\leq (\sum_{j=3}^{s-1} (s-j)/d_j \cdots d_{s-2}) d_1 \cdots d_{s-2} - \sum_{j=2}^{s-1} (s-j) d_j$$
  
$$\leq 4 d_1 \cdots d_{s-2} - d_{s-1},$$

under the assumption  $d_i \ge 2$  for i = 1, ..., s. We have also  $\sum_{j=2}^{s-1} (s-j) c_j \le \sum_{j=2}^{s-1} (s-j) \delta = \frac{1}{2} (s-2)(s-1) \delta$ .

Now we estimate the other term. We consider first the case  $s \leq n$ . Then

$$\begin{split} \sum_{j=2}^{s} \left( s - j + 1 \right) \gamma_{j} &= \sum_{j=2}^{s} \left( s - j + 1 \right) \left( d_{j} \, \delta_{j-1} - \delta_{j} \right) \\ &= \left( s - 1 \right) d_{2} \, \delta_{1} + \sum_{j=3}^{s} \left( \left( s - j + 1 \right) d_{j} - \left( s - j \right) \right) \delta_{j-1} - \delta_{n} \\ &\leq d_{1} \cdots d_{s} - \delta_{s}, \end{split}$$

from where we obtain the bound  $\mu_s = \sum_{j=2}^s (s-j+1) \gamma_j + \sum_{j=2}^{s-1} (s-j) c_j \le (d_1 \cdots d_s - \delta_s) + (4 d_1 \cdots d_{s-2} - d_{s-1}) \le 2 d_1 \cdots d_s.$ 

In the case s = n + 1 we have that  $\mu_{n+1} = \mu_n + \gamma_{n+1}$  from where it follows that  $\mu_{n+1} \leq (2 d_1 \cdots d_n - \delta_n - d_{n-1}) + (\delta_n + d_{n+1} - 1) \leq 2 d_1 \cdots d_n$ .

On the other hand we have also the estimation  $\sum_{j=2}^{s} (s-j+1) \gamma_j \leq \frac{1}{2} (s-1) s d \delta$ from where we conclude that  $\mu_s \leq \frac{1}{2} (s-1) s d \delta + \frac{1}{2} (s-2) (s-1) \delta \leq (s-1)^2 d \delta$ holds, as stated.  $\Box$ 

**Theorem 3.21** Let be given homogeneous polynomials  $f_1, \ldots, f_s \in k[x_0, \ldots, x_n]$ such that  $x_0$  lies in the radical of the ideal  $(f_1, \ldots, f_s)$ . Let  $d_i$  denote the degree of  $f_i$  for  $i = 1, \ldots, s$  and assume that  $d_1 \geq \cdots \geq d_s$  holds. Then

$$x_0^D \in (f_1, \dots, f_s)$$

holds, with  $D := 2 d_s \prod_{i=1}^{\min\{n,s\}-1} d_i$ .

*Proof.* After Lemmas 3.19 and 3.20 it only remains to consider the case when some  $f_i$  has degree one.

By assumption  $f_1, \ldots, f_s$  are ordered in such a way that  $d_1 \ge \cdots \ge d_s$  holds. Let r be maximum such that  $d_r \ge 2$ , so that the polynomials  $f_{r+1}, \ldots, f_s$  have all degree one. We can assume without loss of generality that they are k-linearly independent. We can also suppose that neither 1 nor  $x_0$  lie in the k-linear space spanned by  $f_{r+1}, \ldots, f_s$  as if this is the case the statement is trivial.

Let  $y_0, \ldots, y_{n+r-s-1} \in S$  be polynomials of degree one which complete  $f_{r+1}, \ldots, f_s$ to a linear change of variables. We suppose in addition that  $y_0 = x_0$ . Then the natural inclusion  $k[y_0, \ldots, y_{n+r-s-1}] \hookrightarrow k[x_0, \ldots, x_n]/(f_{r+1}, \ldots, f_s)$  is an isomorphism. Let  $v_i$  be an homogeneous polynomial in  $k[y_0, \ldots, y_{n+r-s-1}]$  such that  $v_i \equiv f_i \mod (f_{r+1}, \ldots, f_s)$  for  $i = 1, \ldots, r$ . Then  $x_0$  lies in the radical of the ideal  $(v_1, \ldots, v_r)$  of  $k[y_0, \ldots, y_{n+r-s-1}]$  and deg  $v_i \leq d_i$  holds for  $i = 1, \ldots, r$ .

Let E denote the integer  $2 \prod_{i=1}^{r} \deg v_i$  so that  $E \leq D := 2 d_s \prod_{i=1}^{\min\{n,s\}-1} d_i$ . Then  $x_0^D \in (v_1, \ldots, v_r)$  from where it follows that  $x_0^D \in (f_1, \ldots, f_s)$  as stated.  $\Box$ 

Then Theorem 3 follows from this result by homogenizating the input polynomials and by considering the degree of the polynomials in a representation of  $x_0^D$ .

Now we are going to prove Theorem 4. First we recall the definition of algebraic degree of a polynomial system.

Let  $g_1, \ldots, g_s \in k[x_1, \ldots, x_n]$  be polynomials without common zeros in  $\mathbb{A}^n$ . Given a  $s \times s$  matrix  $\lambda = (\lambda_{ij})_{ij}$  with entries in  $\overline{k}$  we denote by  $h_i(\lambda)$  the linear combinations  $\sum_j \lambda_{ij} g_j$  induced by  $\lambda$  for  $i = 1, \ldots, s$ . We consider the set of  $s \times s$  matrices  $\Gamma$  such that for any  $\lambda$  in  $\Gamma$  the polynomials  $h_1(\lambda), \ldots, h_{t-1}(\lambda)$  form a regular sequence in  $\overline{k}[x_1, \ldots, x_n]$  and  $1 \in (h_1(\lambda), \ldots, h_t(\lambda))$  for some  $t = t(\lambda) \leq \min\{n, s\}$ .

For each  $\lambda \in \Gamma$  and  $i = 1, \ldots, t - 1$  we denote by  $J_i(\lambda) \subseteq k[x_0, \ldots, x_n]$  the homogenization of the ideal  $(h_1(\lambda), \ldots, h_i(\lambda))$ . Then let  $\delta(\lambda)$  denote the maximum degree of the homogeneous ideals  $J_i(\lambda)$  for  $i = 1, \ldots, t - 1$ .

The algebraic degree of the polynomial system  $g_1, \ldots, g_s$  is defined as the minimum of  $\delta(\lambda)$  over all matrices  $\lambda \in \Gamma$ .

The notion of geometric degree of [21] and [30] is defined in an analogous way as the minimum of  $\delta(\lambda)$  for  $\lambda \in \Gamma$ , with the additional hypothesis that the ideals  $J_i(\lambda)$  are radical for  $i = 1, \ldots, t-1$ . Another difference is that in the case when the characteristic of k is positive the polynomials  $h_j(\lambda)$  are taken as linear combinations of the polynomials  $\{x_j, f_i\}_{ij}$ .

The notion of geometric degree of [14] is similar to that of [21], [30], the only difference is that it is not defined as a minimum but as the value of  $\delta(\lambda)$  for a generic choice of  $\lambda$ .

Thus the algebraic degree is bounded by the geometric degree, whichever version of the later one we consider. The following example shows that in fact it can be much smaller. It is a variant of [21, Example].

**Example 3.22** Let us consider the polynomial system

$$f_1 := 1 - x_1 x_2^d, \ f_2 := x_2 - x_3^d, \ \dots, \ f_{n-1} := x_{n-1} - x_n^d, \ f_n := x_n^2$$

for some  $d \ge 3$ . It is easy to check that these polynomials have no common zero in  $\mathbb{A}^n$ . Let us denote by  $\delta_q$  and by  $\delta_a$  the geometric degree —in the sense of [21], [30]—

and the algebraic degree of this polynomial system, respectively. We are going to compute both integers for this particular example.

Let  $\lambda$  be a  $l \times n$  matrix with entries in  $\overline{k}$  for some  $l \leq n$ , and let  $h_i := \sum_j \lambda_{ij} f_j$  be the induced linear combinations for  $i = 1, \ldots, n$ .

We apply to this matrix the Gauss elimination method by rows, by using as pivots the columns of  $\lambda$  by ascending order. Invertible elementary operations by rows produces polynomials  $q_1, \ldots, q_l$  which generate the same ideal than  $h_1, \ldots, h_l$ . For our particular polynomial system this corresponds to the successive elimination of the variables  $x_1, \ldots, x_n$  in the equations  $h_1, \ldots, h_l$ . Thus in the case when  $l \leq n-1$ the affine variety defined by  $h_1, \ldots, h_l$  can be parametrized by expressing the pivot variables as rational functions of the free ones. It follows that in this case the ideal  $(h_1, \ldots, h_l)$  has dimension at least n - l.

First we consider the geometric degree of this system. The polynomials  $f_1, \ldots, f_n$  form a weak regular sequence,  $1 \in (f_1, \ldots, f_n)$  and the ideal  $(f_1, \ldots, f_i)$  is radical for  $i = 1, \ldots, n-1$ . We have that  $\deg V(f_1, \ldots, f_i) = d^i$  for  $i = 1, \ldots, n-1$  from where it follows that  $\delta_g \leq d^{n-1}$ .

On the other hand, let  $\lambda$  be a  $l \times n$  matrix with entries in  $\overline{k}$  and let  $h_1, \ldots, h_l$ be the corresponding linear combinations. Assume that  $h_1, \ldots, h_l$  is a weak regular sequence,  $1 \in (h_1, \ldots, h_l)$  and that  $(h_1, \ldots, h_i)$  is a radical ideal for  $i = 1, \ldots, l$ . In particular we get that l equals n and that  $(h_1, \ldots, h_{n-1})$  is a one dimensional radical ideal.

We apply the elimination method described above to the matrix formed by the first n-1 rows of  $\lambda$  and we denote by  $q_1, \ldots, q_{n-1}$  the obtained polynomials. We affirm that no columns *i* fails to be a pivot for  $i = 1, \ldots, n-1$ . Suppose that this is not the case. Then  $q_1, \ldots, q_{n-2}$  are polynomials which do not depend on  $x_n$  and which span a one dimensional ideal in  $\overline{k}[x_1, \ldots, x_{n-1}]$ . In addition  $q_{n-1} = x_n^2$  and so the ideal  $(q_1, \ldots, q_{n-1}) \subseteq \overline{k}[x_1, \ldots, x_n]$  is not radical, contradicting our assumption.

Thus there exist scalars  $a_1, \ldots, a_{n-1} \in \overline{k}$  such that  $q_i = f_i + a_i f_n$  for  $i = 1, \ldots, n-1$ . We deduce that the variety  $V(h_1, \ldots, h_{n-1})$  can be parametrized by a map  $\varphi : \mathbb{A}^1 \to \mathbb{A}^n$  defined by  $t \mapsto \varphi(t) = (\varphi_1(t), \ldots, \varphi_n(t))$ , where  $\varphi_i \in \overline{k}(t)$  is a rational function of degree  $d^{n-i}$  for  $i = 1, \ldots, n$ . We get that deg  $V(h_1, \ldots, h_{n-1}) = d^{n-1}$  from where it follows the lower bound  $\delta_g \geq d^{n-1}$ . Combining this with the previous estimation we conclude that  $\delta_g = d^{n-1}$ .

Now we consider the algebraic degree of the system. The polynomials  $f_n, \ldots, f_1$  form a weak regular sequence and  $1 \in (f_n, \ldots, f_1)$ . We have that  $(f_n, \ldots, f_{n-i+1}) = (x_n^2, x_{n-1}, \ldots, x_{n-i+1})$  for  $i = 1, \ldots, n$  from where it follows that  $\delta_a \leq 2$ . In addition, any nontrivial linear combination of  $f_1, \ldots, f_n$  has degree at least two and so  $\delta_a \geq 2$ . We conclude that  $\delta_a = 2$ .

We obtain the following degree bound by direct application of Lemmas 3.19 and 3.20.

**Theorem 3.23** Let be given homogeneous polynomials  $f_1, \ldots, f_s \in k[x_0, \ldots, x_n]$ such that  $x_0$  lies in the radical of the ideal  $(f_1, \ldots, f_s)$ . Let  $f_i^a$  denote the affinization of  $f_i$  for  $i = 1, \ldots, s$ . Let d denote the maximum degree of  $f_i$  for  $i = 1, \ldots, s$  and let  $\delta$  denote the degree of the polynomial system  $f_1^a, \ldots, f_s^a$ . Then

$$x_0^D \in (f_1, \dots, f_s)$$

holds, with  $D := \min\{n, s\}^2 d\delta$ .

Then Theorem 4 follows from this result in the same way that Theorem 3 was derived from Theorem 3.21.

If we apply this degree bound to the previous example we obtain that there exist  $g_1, \ldots, g_n \in k[x_1, \ldots, x_n]$  satisfying

$$1 = g_1 f_1 + \dots + g_n f_n$$

with  $\deg g_i f_i \leq 2 n^2 d$  for  $i = 1, \ldots, s$ . In fact we have the identity

$$1 = f_1 + x_1 x_2^{d-1} f_2 + x_1 x_2^{d-1} x_3^{d-1} f_3 + \dots + x_1 x_2^{d-1} \dots x_{n-1}^{d-1} x_n^{d-2} f_n$$

Acknowledgements. This work originated in several conversations with Bernd Sturmfels. He motivated me to think about the sparse Nullstellensatz and suggested me several lines to approach it. Special thanks are due to him. I am also grateful to Alicia Dickenstein and Joos Heintz for helpful discussions and suggestions, and to Pablo Solernó for providing me a counterexample to a conjecture in an early version of this paper.

I also thank the Departments of Mathematics of the Universities of Alcalá and of Cantabria, Spain, where part of this paper was written during a stay in the spring of 1997.

# References

- M. S. ALMEIDA, Función de Hilbert de álgebras graduadas y Nullstellensatz afín efectivo, Tesis de Licenciatura, Univ. Buenos Aires, 1995.
- [2] F. AMOROSO, On a conjecture of C. Berenstein and A. Yger, in L. González-Vega and T. Recio, eds., Algorithms in algebraic geometry and applications, Proceedings MEGA'94, Birkäuser Progress in Math. 143, Bikhäuser, Basel, 1996, pp. 17–28.
- [3] C. A. BERENSTEIN, D. C. STRUPPA, Recent improvements in the complexity of the effective Nullstellensatz, Linear Alg. Appl. 157 (1991), pp. 203–215.
- [4] C. A. BERENSTEIN, A. YGER, Effective Bézout identities in  $\mathbb{Q}[x_1, \ldots, x_n]$ , Acta Math. 166 (1991), pp. 69–120.
- [5] W. D. BROWNAWELL, Bounds for the degrees in the Nullstellensatz, Ann. of Math. 126 (1987), pp. 577–591.

- [6] W. D. BROWNAWELL, D. W. MASSER, Multiplicity estimates for analytic functions II, Duke J. Math. 47 (1980), pp. 273–295.
- [7] L. CANIGLIA, A. GALLIGO, J. HEINTZ, Some new effectivity bounds in computational geometry, in T. Mora, ed, Proceedings AAECC-6, Lect. Notes in Comput. Sci. 357, Springer, Berlin, 1989, pp. 131–151.
- [8] J. CANNY, I. EMIRIS, A subdivision-based algorithm for the sparse resultant, Preprint, 1996.
- [9] T. W. DUBÉ, A combinatorial proof of the effective Nullestellensatz, J. Symb. Comp. 15 (1993), pp. 277–296.
- [10] G. EWALD, U. WESSELS, On the ampleness of invertible sheaves in complete toric varieties, Results Math. 25 (1991), pp. 275–278.
- [11] N. FITCHAS, A. GALLIGO, Nullstellensatz effectif et conjecture de Sèrre (théorème de Quillen-Suslin) pour le Calcul Formel, Math. Nachr. 149 (1990), pp. 231–253.
- [12] W. FULTON, Introduction to toric varieties, Ann. Math. Studies 131, Princeton Univ. Press, 1993.
- [13] M. GIUSTI, K. HÄGELE, J. HEINTZ, J. L. MONTAÑA, J. E. MORAIS, L. M. PARDO, Lower bounds for diophantine approximation, J. Pure Appl. Algebra 117 & 118 (1997), pp. 277–317.
- [14] M. GIUSTI, J. HEINTZ, J. E. MORAIS, J. MORGENSTERN, L. M. PARDO, Straight-line programs in geometric elimination theory, to appear in J. Pure Appl. Algebra.
- [15] K. HÄGELE, J. E. MORAIS, L. M. PARDO, M. SOMBRA, On the intrinsic complexity of the arithmetic Nullstellensatz, Preprint, 1997.
- [16] J. HARRIS, Algebraic geometry: a first course, Graduate Texts in Math. 133, Springer, Berlin, 1992.
- [17] J. HEINTZ, Definability and fast quantifier elimination in algebraically closed fields, Theoret. Comput. Sci. 24 (1983), pp. 239-277.
- [18] B. HUBER, B. STURMFELS, A polyhedral method for solving sparse polynomial systems, Math. Comp. 64 (1995), pp. 1541–1555.
- [19] J. KOLLÁR, Sharp effective Nullstellensatz, J. Amer. Math. Soc. 1 (1988), pp. 963–975.
- [20] T. KRICK, L. M. PARDO, A computational method for diophantine approximation, in L. González–Vega and T. Recio, eds., Algorithms in algebraic geometry and applications, Proceedings MEGA'94, Birkäuser Progress in Math. 143, Bikhäuser, Basel, 1996, pp. 193–253.
- [21] T. KRICK, J. SABIA, P. SOLERNÓ, On intrinsic bounds in the Nullstellensatz, to appear in AAECC Journal.
- [22] H. MATSUMURA, Commutative ring theory, Cambridge Univ. Press, 1986.
- [23] L. M. PARDO, How upper and lower bounds meet in elimination theory, in G. Cohen, M. Giusti and T. Mora, eds., Proceedings AAECC-11, Lect. Notes. Comput. Sci. 948, pp. 33-69, Springer, Berlin, 1995.
- [24] P. PHILIPPON, Dénominateurs dans le théorème des zeros de Hilbert, Acta Arith. 58 (1990), pp. 1–25.

- [25] J. M. ROJAS, Toric generalized characteristic polynomials, Preprint, 1997.
- [26] J. M. ROJAS, Toric laminations, sparse generalized characteristic polynomials, and a refinement of Hilbert's tenth problem, Preprint, 1997.
- [27] J. SABIA, P. SOLERNÓ, Bounds for traces in complete intersections and degrees in the Nullstellensatz, AAECC Journal 6 (1995), pp. 353–376.
- B. SHIFFMAN, Degree bounds for the division problem in polynomial ideals, Michigan Math. J. 36 (1989), pp. 163–171.
- [29] F. SMIETANSKI, Quelques bornes effectives pour le théoreme des zeros avec parametres, These, Univ. Nice–Sophia Antipolis, 1994.
- [30] M. SOMBRA, Bounds for the Hilbert function of polynomial ideals and for the degrees in the Nullstellensatz, J. Pure Appl. Algebra 117 & 118 (1997), pp. 565–599.
- [31] B. STURMFELS, Sparse elimination theory, in D. Eisenbud and L. Robbiano, eds., Proceedings of the Cortona conference on computational algebraic geometry and commutative algebra, Symposia Matematica XXXIV, Ist. Naz. di Alta Matematica, Cambridge Univ. Press, 1993, pp. 377–396.
- [32] B. STURMFELS, Gröbner bases and convex polytopes, Univ. Lect. Series 8, Amer. Math. Soc., 1996.
- [33] B. STURMFELS, N. V. TRUNG, W. VOGEL, Bounds on degrees of projective schemes, Math. Ann. 302 (1995), pp. 417–432.
- [34] B. TEISSIER, Résultats récents d'algèbre commutative effective, Sém. Bourbaki 718, Astérisque 189–190, pp. 107–131, Soc. Math. France, 1991.
- [35] J. VERSCHELDE, P. VERLINDEN, R. COOLS, Homotopies exploiting Newton polytopes for solving sparse polynomial systems, SIAM J. Numer. Anal. 31 (1994), pp. 915–930.
- [36] W. VOGEL, Lectures on results on Bézout theorem, Tata Lect. Notes 74, Springer, Berlin, 1984.
- [37] O. ZARISKI, P. SAMUEL, Commutative algebra, 2 vols., Van Nostrand, New York, 1958, 1960.