

CHVCs: Galactic Building Blocks at $z = 0$

Robert Braun

*Netherlands Foundation for Research in Astronomy, P.O. Box 2, 7990
AA Dwingeloo, The Netherlands*

W. Butler Burton

Sterrewacht Leiden, P.O. Box 9513, 2300 RA Leiden, The Netherlands

Abstract. A distinct sub-class of anomalous velocity HI emission features has emerged from recent high quality surveys of the Local Group environment, namely the compact high velocity clouds (CHVCs). A program of high-resolution imaging with the Westerbork array and the Arecibo telescope has begun to provide many insights into the nature of these objects. Elongated core components with a velocity gradient consistent with rotation ($V_{Rot} \sim 15 \text{ km s}^{-1}$) are seen in many objects. Comparison of volume and column densities has allowed the first distance estimates to be made ($600 \pm 300 \text{ kpc}$). The objects appear to be strongly dark-matter dominated with dark-to-gas mass ratios of 30–50 implied if the typical distance is 700 kpc.

1. Introduction

The nature of anomalous velocity HI discovered during $\lambda 21 \text{ cm}$ surveys of the Galaxy has been the subject of much debate in the past three decades. There appear to be at least four reasonably distinct classes of this gas, namely: (1) Localized outflows and subsequent inflows associated with massive star formation which are commonly referred to as a “galactic fountain” (Shapiro & Field, 1976; Bregman, 1980). (2) The stream of tidal debris from the interaction of the Galaxy with the Magellanic Clouds is another major source of anomalous velocity HI. Recent imaging (Putman & Gibson, 1999) has begun to reveal the true extent of this debris system, and allow it to be distinguished from other components. (3) The extended HVC complexes (named A, C, H, M, ...) which span ten’s of degrees on the sky and have recently been determined to have both nearby distances of about 10 kpc (Van Woerden et al. 1999) and rather low (~ 0.1 solar) metal abundance (Wakker et al. 1999). These diffuse HI structures appear to be currently merging with the Galaxy and each account for some $10^7 M_{\odot}$ of fresh gas. And finally (4) the system of compact high velocity clouds (CHVCs) cataloged by Braun & Burton (1999).

The CHVCs form a distinct class of compact, high-contrast HI emission features at anomalous velocity, which can be readily distinguished from the more diffuse components in the recent high quality surveys; the Leiden/Dwingeloo Survey in the North (Hartmann & Burton 1997) and the Parkes Multibeam

Survey in the South (see Putman & Gibson 1999). Their average angular size is only 1 degree FWHM and the total velocity width 30 km s^{-1} FWHM. The 65 confirmed members of the CHVC class share a well-defined kinematic pattern. A global search for the reference frame that minimizes the line-of-sight velocity dispersion of the system, returns the Local Group Standard of Rest with high confidence. In this system the velocity dispersion of the population is only 70 km s^{-1} , although it is in-falling toward the Local Group barycenter at about 100 km s^{-1} . These properties suggest that the objects are: (1) associated with the Local Group rather than the Galaxy as such, (2) that they are likely to reside at quite substantial distances and (3) have as yet had little tidal interaction with the more massive Local Group members.

2. High Resolution Imaging

A program of high resolution imaging of the CHVCs has begun to provide further insights into the nature of these sources. An initial sub-sample of six CHVCs was imaged with the upgraded Westerbork array (WSRT) in December 1998. A complete description of those results can be found in Braun & Burton (2000). The sub-sample was chosen to span a wide range of positions on the sky and in line-of-sight velocity, as well as gas content and HI linewidth. An additional eight sources have been imaged with the WSRT during the summer of 1999, specifically targeting objects suspected of having very bright and narrow HI emission features. A sample of ten CHVCs have also been imaged with the upgraded Arecibo telescope in November of 1999. In addition to fully-sampled images of the central square degree, a very deep cross-cut of two degree length was obtained for each source reaching an rms sensitivity of about $\Delta N_{\text{HI}} = 10^{17} \text{ cm}^{-2}$ at 10 km s^{-1} resolution. Full accounts of these recent results will be given elsewhere.

2.1. Core/Halo Morphology and Physics

All of the imaged sources share a number of common properties. Each source contains between one and ten compact cores (of 1 to 10 arcmin extent) embedded in a diffuse halo of about 1 degree diameter. An average of 40% of the HI line flux arises in the compact cores (although the range varies between 1 and 60%). The peak column density in these cores varies between 10^{19} and 10^{21} cm^{-2} . The fractional surface area of each source which exceeds a column density of 5×10^{18} is only 15% on average, although this also varies greatly from source to source. The remainder of the HI emission arises in a diffuse halo which reaches peak column densities of $10^{18.5} - 10^{19.5} \text{ cm}^{-2}$ but declines smoothly with an exponential profile to values of $10^{17.5} \text{ cm}^{-2}$ or less (as illustrated in Fig.2).

The resolved HI linewidths toward the core components (with 1 arcmin resolution) are typically only 5 km s^{-1} FWHM (but range from as little as 2 to as much as 15 km s^{-1}). All of the core linewidths are sufficiently narrow that it is possible to unambiguously identify these with the Cool Neutral Medium phase of HI, with typical kinetic temperature of 100 K. The diffuse halos, on the other hand, all have resolved HI linewidths (with 3 arcmin resolution) of $21 - 25 \text{ km s}^{-1}$ FWHM, as seen in Fig.2d. The one-to-one correspondence of halo emission with linewidths equal to or exceeding the thermal linewidth




Figure 1. **A:** Westerbork image of CHVC 204+30+075 showing N_{HI} at 1 arcmin resolution; contours are drawn at levels of 20, 50, 100, 200, and $300 \times 10^{18} \text{ cm}^{-2}$. **B:** Intensity-weighted line-of-sight velocity, with contours of v_{LSR} drawn in steps of 5 km s^{-1} from 40 to 85 km s^{-1} . **C:** Rotation velocities fit to the two principal components of CHVC 204+30+075. The solid lines show the rotation curves of Navarro, Frenk, & White (1997) cold-dark-matter halos of the indicated mass and 0.7 Mpc distance. **D: top** Equilibrium temperature curves for HI in an intergalactic environment characterized by a metallicity of 10% of the solar value and a dust-to-gas ratio of 10% of that in the solar neighborhood, calculated for two values of the neutral shielding column depth, namely 10^{19} cm^{-2} (solid line) and 10^{20} cm^{-2} (dashed line). **D: bottom** Mass volume density, in units of hydrogen nuclei per cubic centimeter, as a function of radius for an NFW cold dark matter halo.

of 8000 K gas (21 km s^{-1} FWHM) provides an unambiguous identification of these halos with the Warm Neutral Medium phase of H I. A shielding column of WNM is one prerequisite for the condensation of CNM cores in calculations of H I thermodynamics (eg. Wolfire et al. 1995). The second prerequisite for CNM condensation is a sufficiently high thermal pressure, in excess of about $P/k = 100 \text{ cm}^{-3}\text{K}$. Both of these aspects are illustrated in the H I phase diagram shown in Fig.1d. An adequate shielding column of WNM is clearly detected in the CHVCs, as just discussed above. We will return to the question of an adequate thermal pressure below.

Another general result is that the CNM core components are dynamically decoupled from their WNM halos. This is illustrated in Fig.2, where a substantial velocity gradient of some 30 km s^{-1} is seen within the major core component of CHVC158–39–285. Such large gradients are not observed in the halo gas, which instead is always centered near the systemic velocity of the system. This dichotomy suggests that while the cores may represent flattened rotating disks, the halos are more nearly spherical distributions.

2.2. Distance, Metallicity and Dark Matter Content

One of the objects imaged with the WSRT, CHVC125+41–207, has proven particularly interesting for several reasons. This object has a number of opaque core components of only $\theta = 90''$ angular size, a brightness temperature in H I emission of 75 K, and linewidths so narrow (less than 2 km s^{-1} FWHM) that the kinetic temperature ($T_k = 85 \text{ K}$), H I opacity ($\tau = 2$) and column density ($N_{\text{HI}} = 10^{21} \text{ cm}^{-2}$), can be accurately derived. A good estimate of the volume density ($n_H = 2 \pm 1 \text{ cm}^{-3}$) for this object has been made by modeling the thermodynamics (Wolfire et al., priv. comm.). With these quantities in hand and only the assumption of crude spherical symmetry for the cores, it is possible to estimate the source distance from $D = N_{\text{HI}} / (n_H \theta) = 600 \pm 300 \text{ kpc}$.

The same object has also allowed the first measurement of CHVC metallicity, since the bright UV source Mrk 205 is located behind its diffuse H I halo. Bowen, Blades and Pettini (1995) detect unsaturated MgII absorption, with 0.15 \AA EW at the CHVC velocity, while we detect a neutral column $N_{\text{HI}} = 5 \times 10^{18} \text{ cm}^{-2}$ at the same position. The implied metal abundance is about 0.05 solar.

Although many of the CHVCs have only one or two core components, a handful of these objects might best be called CHVC clusters. As many as ten cores are seen in close proximity (within 30–50 arcmin), each with its own systemic velocity and internal kinematics (which we address below) but sharing the same diffuse halo of shielding gas. The most extreme case is that of CHVC115+13–275, where the cores have relative velocities as high as 70 km s^{-1} and angular separations as large as 30 arcmin. If these are gravitationally bound systems (and their cool gas content and isolation on the sky make this seem plausible) then they have a dynamical mass, $M_{\text{dyn}} = Rv^2/G = 2.3 \times 10^5 R_{\text{kpc}} v_{\text{km/s}}^2$, and gas mass $M_{\text{gas}} = 1.4 M_{\text{HI}} = 3.2 \times 10^5 S D_{\text{Mpc}}^2$. If for example the distance were 0.7 Mpc, then $M_{\text{dyn}} = 10^{8.93} M_{\odot}$ and $M_{\text{gas}} = 10^{7.22} M_{\odot}$. The dark-to-gas mass ratio in this case has the rather substantial value of $\Gamma = 51$, which scales with $1/D$ for other distance estimates.

As already noted above, many of the CHVC cores give an indication for organized internal kinematics. Many of the cores have a systematic velocity




Figure 2. **A:** Arecibo image of CHVC 158–39–285 showing N_{HI} at 3 arcmin resolution; contours are drawn at levels of 5, 10, 20, and $30 \times 10^{18} \text{ cm}^{-2}$. **B:** Intensity-weighted line-of-sight velocity, with contours of v_{LSR} drawn at the indicated velocities. **C:** Position, velocity cut at a fixed declination of +16:17:30. Contours are in units of brightness temperature. **D:** Variation of line-of-sight velocity, velocity FWHM and N_{HI} with position.

gradient along the long axis of their elliptical distribution which is very suggestive of rotation in a flattened disk system. Several examples are shown in Figs.1 and 2. The best-resolved examples of this pattern have been subjected to the standard tilted ring fitting algorithms used in deriving galactic rotation curves. Good solutions for rotation are found (Fig.1c) which slowly rise over some 6 arcmin to constant values of 15 to 20 km s⁻¹. The shapes of these rotation curves are in very good agreement with those predicted by Navarro, Frenk & White (1997) for a CDM halo at 0.7 Mpc distance (as shown in Fig.1c). The dark-to-visible mass ratios in these two cases are $\Gamma = 36$ and 29.

The presence of a significant dark matter component is important not only for understanding the kinematics of the individual cores and the CHVC clusters, but also for understanding the HI thermodynamics. As noted earlier, a significant thermal pressure in excess of about $P/k = 100 \text{ cm}^{-3}\text{K}$ is required to allow condensation of the cool cores observed in the CHVCs (see Fig.1d, upper). The mass density provided within the central few kpc of an NFW halo in this mass range (Fig.1d, lower) is sufficient to provide the required hydrostatic pressure in combination with the observed warm halo column densities of about 10^{19}cm^{-2} (see also Braun & Burton 2000).

Acknowledgments. We are grateful to M.G. Wolfire, A. Sternberg, D. Hollenbach, and C.F. McKee for providing the equilibrium temperature curves shown in Fig.1d. The Westerbork Synthesis Radio Telescope is operated by the Netherlands Foundation for Research in Astronomy, under contract with the Netherlands Organization for Scientific Research. The Arecibo Observatory is part of the National Astronomy and Ionosphere Center, which is operated by Cornell University under a cooperative agreement with the National Science Foundation.

References

- Bowen, D.V., Blades, J.C., & Pettini, M. 1995, ApJ, 448, 662
 Braun R., & Burton W.B. 1999, A&A, 341, 437
 Braun R., & Burton W.B. 2000, A&A, in press, astro-ph/9912417
 Bregman J. 1980, ApJ, 236, 577
 Hartmann D., & Burton W.B. 1997, "Atlas of Galactic Neutral Hydrogen",
 Cambridge: Cambridge University Press
 Navarro J.F., Frenk C.S., & White S.D.M., 1997, ApJ, 490, 493
 Putman M.E., & Gibson B.K. 1999, PASA, 16, 70
 Shapiro P.R., & Field G. 1976, ApJ, 205, 762
 Wakker B.P., Howk J.C., Savage B.D., van Woerden H., Tufte S.L., Schwarz
 U.J., Benjamin R., Reynolds R.J., Peletier R.F. & Kalberla P.M.W. 1999,
 Nature, 402, 388
 van Woerden H., Schwarz U.J., Peletier R.F., Wakker B.P., & Kalberla P.M.W.
 1999, Nature, 400, 138
 Wolfire M.G., Hollenbach D., McKee C.F., Tielens A.G.G.M., & Bakes E.L.O.
 1995, ApJ, 443, 152