# New Velocity Distribution for Cold Dark Matter in the Context of the Eddington Theory

J.D. Vergados<sup>1</sup>

Department of Physics, Unisa, Pretoria, South Africa.

vergados@cc.uoi.gr

and

D. Owen <sup>2</sup>

Department of Physics, Ben Gurion University, Israel.

owen@bgumail.bgu.ac.il

October 24, 2018

### ABSTRACT

Exotic dark matter together with the vacuum energy (associated with the cosmological constant) seem to dominate the Universe. Thus its direct detection is central to particle physics and cosmology. Supersymmetry provides a natural dark matter candidate, the lightest supersymmetric particle (LSP). One essential ingredient in obtaining the direct detection rates is the density and velocity distribution of the LSP. The detection rate is proportional to this density in our vicinity. Furthermore, since this rate is expected to be very low, one should explore the two characteristic signatures of the process, namely the modulation effect, i.e. the dependence of the event rate on the Earth's motion and the correlation of the directional rate with the motion of the sun. Both of these crucially depend on the LSP velocity distribution. In the present paper we study simultaneously density profiles and velocity distributions based on the Eddington theory.

Subject headings: Cosmology: Eddington theory, velocity profiles, rotational curves-Cold Dark Matter: velocity distribution, direct detection rates.

<sup>&</sup>lt;sup>1</sup>Permanent address:Theoretical Physics Division, University of Ioannina, Ioannina, Gr 451 10, Greece.

<sup>&</sup>lt;sup>2</sup>Visiting the Theoretical Physics Division, University of Ioannina, Ioannina, Gr 451 10, Greece.

#### 1. Introduction

In recent years the consideration of exotic dark matter has become necessary in order to close the Universe (Jungman et al. 1996). The COBE data (Smoot et al. 1992) suggest that CDM (Cold Dark Matter) component is at least is at least 60% (Gawser et al. 1988; Gross et al. 1998) of the total mass. On the other hand evidence from two different teams, the High-z Supernova Search Team (Riess et al. 1998) and the Supernova Cosmology Project (Somerville et al. 2002), (Perlmutter et al 1999; Perlmutter et al. 1997) suggests that the Universe may be dominated by the cosmological constant  $\Lambda$ . Thus the situation can be adequately described by a baryonic component  $\Omega_B = 0.1$  along with the exotic components  $\Omega_{CDM} = 0.3$  and  $\Omega_{\Lambda} = 0.6$ . In a more detailed  $\Lambda CDM$  analysis (Primack 2002) one finds:

$$\Omega_b = 0.040 \pm 0.002, \ \Omega_m = \Omega_{CDM} = 0.33 \pm 0.035, \ \Omega_{HDM} \le 0.05, \ \Omega_{\Lambda} = 0.73 \pm 0.08$$

(see also (Turner 1990) and (Einasto 2001)). Since the non exotic component cannot exceed 40% of the CDM (Jungman et al. 1996), (Alcock et al. 1995), there is room for the exotic WIMP's (Weakly Interacting Massive Particles). In fact the DAMA experiment (Bernabei et al. 1996) has claimed the observation of one signal in direct detection of a WIMP, which with better statistics has subsequently been interpreted as a modulation signal (Bernabei et al. 1998; Bernabei et al. 1999).

In the most favored scenario of supersymmetry the LSP can be simply described as a Majorana fermion, a linear combination of the neutral components of the gauginos and Higgsinos (Jungman et al. 1996; Vergados 1990; Gomez and Vergados 2001; Gomez et al. 2000a,b; Gomez and Vergados 2001; Anrowit and Nath 1995, 1996; Bottino et al. 1997; Bednyakov et al. 1994).

Since this particle is expected to be very massive,  $m_{\chi} \geq 30 GeV$ , and extremely non relativistic with average kinetic energy  $T \leq 100 KeV$ , it can be directly detected (Vergados 1996; Spira et al. 1995; Kosmas and Vergados 1997) mainly via the recoiling of a nucleus (A,Z) in elastic scattering.

In order to compute the event rate one needs the following ingredients:

- 1) An effective Lagrangian at the elementary particle (quark) level obtained in the framework of supersymmetry as described, e.g., in (Jungman et al. 1996; Anrowit and Nath 1995, 1996; Bottino et al. 1997; Bednyakov et al. 1994).
- 2) A procedure in going from the quark to the nucleon level, i.e. a quark model for the nucleon. The results depend crucially on the content of the nucleon in quarks other than u and d. This is particularly true for the scalar couplings as well as the isoscalar axial coupling (Drees and Noijiri 1993b,c; Cheng 1988, 1989).

- 3) Compute the relevant nuclear matrix elements (Ressel et al. 1993; Divari et al. 2000) using as reliable as possible many body nuclear wave functions. The situation is a bit simpler in the case of the scalar coupling, in which case one only needs the nuclear form factor.
- 4) The LSP density and velocity distribution. Among other other things, since the detection rates are expected to be very small, the velocity distribution is crucial in exploiting the characteristic experimental signatures provided by the reaction, namely: a) the modulation of the event rates due to the earth's revolution around the sun (Vergados 1998, 1999)—(Vergados 2000) and b) the correlation of the rates with the Sun's direction of motion in directional experiments, i.e. experiments in which the direction of the recoiling nucleus is observed (Vergados 1990; Buckland et al. 2000). To obtain the right density and velocity distributions is the purpose of the present paper.

In the past various velocity distributions have been considered. The most popular one is the isothermal Maxwell-Boltzmann velocity distribution with  $\langle v^2 \rangle = (3/2)v_0^2$  where  $v_0$  is the velocity of the sun around the galaxy, i.e.  $220 \ km/s$ . Extensions of this M-B distribution were also considered, in particular those that were axially symmetric with enhanced dispersion in the galactocentric direction (Drucker et al. 1986; Vergados 2000). In such distributions an upper cutoff  $v_{esc} = 2.84v_0$  was introduced by hand.

Non isothermal models have also been considered. Among those one should mention the late infall of dark matter into the galaxy, i.e caustic rings (Sikivie 1999(, 1998; Vergados 2001a; Green 2001; Gelmini and Gondolo 2001), and dark matter orbiting the Sun (Copi et al. 1999).

The correct approach in our view is to consider the Eddington approach (Eddington 1916), i.e. to obtain both the density and the velocity distribution from a mass distribution, which depends both on the velocity and the gravitational potential. This approach has been extensively studied by Merritt (Merritt 1985) and recently applied to dark matter by Ullio and Kamionkowski (Ullio and Kamiokowski 2001)

#### 2. Density Profiles

As we have seen in the introduction the matter distribution can be given as follows

$$dM = 2\pi f(\Phi(r), \nu_r, \nu_t) dx dy dz \nu_t d\nu_t d\nu_r$$
 (1)

where the function f the distribution function, which depends on r through the potential  $\Phi(r)$  and the tangential and radial velocities  $v_t$  and  $v_r$  (we assume axial symmetry in velocity

space). Thus the density of matter  $\rho$  satisfies the equation

$$d\rho = 2\pi \ f(\Phi(r), \nu_r, \nu_t) \ \nu_t \ d\nu_t \ d\nu_r \tag{2}$$

It is more convenient instead of the velocities to use the total energy E and the angular momentum J via the equations

$$J = v_t r$$
 ,  $2E = v_r^2 + \frac{J^2}{r^2} + 2 \Phi(r)$  (3)

The use of these variables, which are constants of motion, is very useful, when one wants to study equilibrium states. We thus find

$$\rho = \frac{2\pi}{r^2} \int \int \frac{f(E,J) J}{\sqrt{2(E - \Phi(r)) - J^2/r^2}} dJ dE$$
 (4)

The limits of integration for E are from  $\Phi$  to 0 and for J from 0 to  $[2r^2(E-\Phi(r))]^{1/2}$ .

Following Eddington we will choose a distribution function of the form

$$f(E,J) = K_{\lambda}(-2E)^{\lambda} \tag{5}$$

(E is negative for a bound system), where  $\lambda$  is a parameter, which will depend on the type of matter, and  $K_{\lambda}$  is a normalization constant, which will be related to the density at some point. With this choice of the distribution function it is quite straightforward to find the relationship between the density  $\rho$  and the potential. The result is

$$\rho = K_{\lambda} 2^{\lambda + 3/2} \pi |\Phi(r)|^{\lambda + 3/2} \beta(\lambda + 1, 3/2) \tag{6}$$

with

$$\beta(a,b) = \frac{\Gamma(a) \ \Gamma(b)}{\Gamma(a+b)} \tag{7}$$

Eq. (6) can be cast in the simple form:

$$\rho = \rho_{\lambda}(0)(\chi(x))^{\lambda + 3/2} \tag{8}$$

with

$$\chi(x) = \frac{\Phi(x)}{\Phi_0}, \quad x = r/r_s \tag{9}$$

with  $\Phi_0 = \Phi(0)$  and  $r_s$  the galactic radius (position of the sun). The constant  $K_{\lambda}$  is related to the density at the origin via the relation:

$$K_{\lambda} = \frac{\rho_{\lambda}(0)}{\pi(2|\Phi_{0}|)^{\lambda+3/2}\beta(\lambda+1,3/2)}$$
(10)

The above distribution function is isotropic. Following Merritt (Merritt 1985) we can introduce an anisotropy by modifying the distribution function as follows:

$$f(E,J) = K_{\lambda}(-2E)^{\lambda}(1 \pm \frac{J^2}{r_a^2})$$
 (11)

Instead of the parameter  $r_a$  we find it convenient for our applications later on (see below) to adopt the more recent conventions and write the above equation as follows:

$$f(E,J) = K_{\lambda}(-2E)^{\lambda} [1 + \alpha_s \frac{J^2}{(r_s \nu_m)^2}]$$
 (12)

where  $v_m$  is the maximum velocity allowed by the potential, to be specified below, and  $\alpha_s$  the asymmetry parameter. Proceeding as above we find that this induces a correction to the density of the form:

$$\Delta \rho = K_{\lambda} \frac{4}{3} 2^{\lambda + 3/2} \pi |\Phi(r)|^{\lambda + 5/2} \beta(\lambda + 1, 5/2) \frac{\alpha_s}{v_m^2} x^2$$
 (13)

Combining Eqs (6) and (13) we get

$$\rho_{\lambda}(r) = \rho_{\lambda}(0)\psi_{\lambda}(x) \tag{14}$$

$$\psi_{\lambda}(x) = (\chi(x))^{\lambda + 3/2} \left[1 + \frac{4}{3} a \frac{\beta(\lambda + 1, 5/2)}{\beta(\lambda + 1, 3/2)} x^2 \chi(x)\right]$$
(15)

with  $a = \alpha_s |\Phi_0| / v_m^2$ .

Since the scale of the potential appears only via the parameter a one, in principle, could have two a parameters, one for Matter  $(a_m)$  and one for dark matter  $(a_{dm})$ . In the present work we will assume that they are equal. We remind the reader that  $\alpha_s$  is the asymmetry parameter to be treated phenomenologically.

#### 3. Allowed Density Functions

The central question is to specify density  $\psi_{\lambda}(x)$ , the potential  $\chi(x)$  and the mass density distribution entering Eq. (1). One can adopt one of two procedures:

1) Start out with a given density, obtained, e.g., phenomenologically, and find the potential by solving Poisson's equation. This way one obtains, at least parametrically, e.g. with the radial coordinate as a parameter, the proper relation between the density and the potential. This approach has been adopted by a lot of researchers, see e.g. (Widrow

2000), (Evans 1994), (Henrriksen and Widrow 1994), (Ullio and Kamiokowski 2001). From this one can obtain the mass density function  $f(\Phi, v)$ . This approach leads, in general, to non analytic, i.e. complicated, velocity distribution, hard to implement in dark matter calculations (Ullio and Kamiokowski 2001), especially for not spherically symmetric mass distributions.

2) Use the above simple relation between the density and the potential and solve the differential equation resulting from Poisson's equation and thus obtain each one of them. One hopes that this way one will get a density distribution, which describes adequately dark matter distribution globally, via fitting the rotational curves, and in our vicinity. This approach can easily deal with asymmetric density distributions. The simplicity of the relation between the density and the potential thus yields the bonus that the obtained velocity distribution is analytic and can easily implemented in obtaining the event rates for direct dark matter detections. It can also be extended to include more than one power in  $\lambda$ . In this paper we will follow this approach, and leave it for the future to use semi-analytic mass distributions obtained in the approach outlined in the previous paragraph.

For a spherically symmetric potential Poisson's equation leads to a differential equation of the type:

$$x\chi''(x) + 2\chi'(x) = -\Lambda \ x \ (\chi(x))^n \left[1 + \frac{2a}{n+1}x^2\chi(x)\right]$$
 (16)

with  $n = \lambda + 1/2$  and

$$\Lambda = -\frac{4\pi G_N r_s^2 \rho_\lambda(0)}{\Phi_0} \tag{17}$$

with  $G_N$  Newton's gravitational constant. The dimensionless quantity  $\Lambda$  is assumed to be positive (attractive potential). Since the asymmetry parameter is assumed to be small, the first term in the right hand side dominates. Introducing the variable  $\xi = \sqrt{\Lambda}x$  and  $\chi(x) = u(x\sqrt{\Lambda})$  we arrive at:

$$\xi u''(\xi) + 2u'(\xi) = -\xi \left(u(\xi)\right)^n \left[1 + \frac{2b}{n+1}\xi^2 u(\xi)\right] , \quad b = \frac{a}{\Lambda}$$
 (18)

The last equation is a non linear differential equation, which must be solved for  $\xi > 0$ , with the conditions u(0) = 1 and u'(0) = 0. We also demand that the solution remains positive, i.e. the solution drops from unity to zero. For  $\alpha_s = 0$  the density is a rapidly decreasing function of  $\xi$ .

The above equation can be solved analytically in the special case a=0 and only when  $\lambda = -1/2$  (n=1) or  $\lambda = 7/2$  (n=5). In these cases the solutions are:

$$\lambda = -1/2 \to u(\xi) = \frac{\sin \xi}{\xi} \tag{19}$$

$$\lambda = 7/2 \to u(\xi) = \frac{1}{(1+\xi^2/3)^{1/2}}$$
 (20)

The case  $\lambda = -1/2$  may not have physical meaning, but it will serve as an illustrative example for the realistic cases to be discussed below.

We should stress that the last solution associated with  $\lambda=7/2$ , admissible over all space, can be considered as providing an adequate description of ordinary matter<sup>3</sup>. In the absence of asymmetry no constraint is imposed on the parameter  $\Lambda$  by the positivity and finiteness condition of the solution. The solution given by Eq. (19) is constrained in the range  $0 \le \xi \le \pi$ . If we demand that the range of the potential for dark matter extends far out in the halo, e.g. up to  $x_r = r/r_s = 20$ , we must have  $\Lambda = [\pi/x_r]^2 = 0.025$ . This gives a useful constraint between  $\Phi_0$  and  $\rho_{\lambda}(0)$ .

The differential equation (18) can, in general, be solved only numerically. For  $\lambda = 1/2$  we find that there is a singularity at  $\xi = 12.43$ , while the first zero of the potential occurs at  $\xi = 4.35$ . This leads to the constraint  $\Lambda = (4.35/20)^2 = 0.047$ . For  $\lambda = 1$  the corresponding value is  $\Lambda = 0.065$ .

The above results are significantly modified when the asymmetry parameter is turned on. One now can see that, depending on the parameters  $\Lambda$  and  $\alpha_s$ , the shape of the density is significantly modified.

We will examine the special case of  $\lambda = 7/2$ , which corresponds to the Eddington solution for ordinary matter. As we have seen in the absence of asymmetry the solution can be found exactly. In the presence of asymmetry we distinguish two cases:

## 1) Positive values of $\alpha_s$ .

For small values of b the solution has no roots. Beyond a critical value of b the solution attains the value zero. Then further out for still larger  $\xi$  it becomes negative. At some point it becomes singular. The extracted values of  $\Lambda$  range between 1.5 and 0.12.

## 2) Negative values of $\alpha_s$ .

In this case when the absolute value of b becomes sufficiently large the solution goes through zero. Again a value of  $\Lambda$  can be extracted. The situation is quite unstable. Generally

<sup>&</sup>lt;sup>3</sup>The designation of  $\lambda = 7/2$  as representing ordinary matter is due to the work of Eddington (Eddington 1916), where he points out that this value leads to the Plummer distribution. Furthermore this distribution provides an adequate description of the galaxy (of ordinary matter) he was concerned with. A discussion of some of the thermodynamical aspects of the  $\lambda = 7/2$  distribution is also given in this paper.

speaking for arbitrary values of b the solution blows up to infinity without going through zero. One cannot extract values of  $\Lambda$  in this case.

In any case, these observations are intended as illustrations of what is expected for dark matter where the exact solution cannot be obtained. In the case of ordinary matter we will not consider  $\alpha_s$  different from zero, since the range of the solution is small ( the density is assumed to vanish for  $r \geq r_s$ ).

Let us return to Eq. (12) and express it in terms of the velocity in our vicinity. We obtain:

$$f(v) = N[-2\Phi(r_s) - v^2]^{\lambda} (1 + \alpha_s \frac{v_t^2}{v_m^2})$$
 (21)

where N is a normalization constant, which depends on  $\lambda$ ,  $\alpha_s$  and  $\nu_m$ . From this we see that the maximum velocity in our vicinity,  $\nu_m$ , depends on the value of the potential, i.e.

$$v_m^2 = -2 \ \Phi(r_s) = -2 \ \Phi_0 \ u(\sqrt{\Lambda}) \tag{22}$$

This means that

$$b = \frac{a}{\Lambda} = \frac{\alpha_s}{2 \Lambda u(\sqrt{\Lambda})} \tag{23}$$

Our strategy is clear:

- 1) Vary b so that the solution has its first zero at some value  $\xi_0$  and is monotonically decreasing up to that point.
  - 2) Determine  $\Lambda$  so that  $\xi_0$  corresponds to a range  $x_r$  of the potential  $\chi(x)$ .
  - 3) From Eq. (23) obtain an acceptable (self-consistent) value of  $\alpha_s$ ,  $\alpha_s = 2\Lambda b \ u(\sqrt{\Lambda})$ .
  - 4) The parameter  $v_m$  is obtained via the relation:

$$v_m^2 = -2 \Phi_0 u(\sqrt{\lambda}) = 8\pi G_N r_s^2 \frac{u(\sqrt{\Lambda}) \rho_{\lambda}(0)}{\Lambda}$$
 (24)

or equivalently

$$v_{m} = v_{0} \sqrt{\eta_{\chi}} h_{v}(n, a, \Lambda) , h_{v}(n, a, \Lambda) = \left[\frac{u(\sqrt{\Lambda})}{\Lambda((u(\sqrt{\Lambda}))^{n} \left[1 + \frac{2a}{n+1} u(\sqrt{\Lambda})\right]}\right]^{1/2} , n = \lambda + 3/2$$
(25)

where  $v_0$  is our velocity of rotation around the center of our galaxy (220 km/s). In obtaining the last equation we used our solution to relate the density  $\rho_{\lambda}(0)$  to the density of dark matter in our vicinity,  $\rho_0$ , via the equation:

$$\rho_{\lambda}(0) = \frac{\rho_0}{((u(\sqrt{\Lambda}))^n [1 + \frac{2a}{n+1} u(\sqrt{\Lambda})]} , \quad n = \lambda + 3/2$$
 (26)

The parameter  $\eta_{\chi}$  is defined by  $\eta_{\chi} = \rho_0/(0.3 \text{ GeV/cm}^3)$ .  $\eta_{\chi}$  is normally taken to be unity (Jungman et al. 1996).

It is clear that in our treatment for a given  $\lambda$  we have three independent parameters, b,  $x_r$  and  $\rho_0$ . The obtained results for some interesting typical cases of  $\alpha_s$  are shown in Table 1. The value of the parameter  $h_v$  has the meaning of  $y_{esc}$  (see section 6) and it is rewarding that it is not far from the value of 2.84 deduced phenomenologically (Drucker et al. 1986). The potentials obtained for the various such parameters are shown in Figs 1 - 4. It is clear that the shapes of these potentials are similar.

The densities obtained with the same set of parameters are shown in Figs 5 - 8. We observe that, even though the potentials are similar, the densities differ substantially. The asymmetry parameter has a big effect on the density, especially for  $\alpha_s$  greater than zero, when it changes substantially around x = 8. For  $\alpha_s$  negative there is small reduction of the density in the region of interest, but, as we have already mentioned, only special values of b are acceptable. A detailed rigorous mathematical study of the behavior of the solutions for b negative is currently under study and it will appear elsewhere.

#### 4. Rotational Velocities

We are now in position to obtain the rotational velocity curves. The rotational velocity is given by

$$v_{\lambda}(x) = \frac{v_0}{\sqrt{2}} \sqrt{g_{\lambda}(x)} \tag{27}$$

with

$$g_{\lambda}(x) = \frac{1}{x} \int_{0}^{x} (x')^{2} dx' \psi_{\lambda}(x')$$
 (28)

We see that the scale of the rotational velocity is set by  $\frac{v_0}{\sqrt{2}}$  in agreement with the data, see, e.g., the recent review (Jungman et al. 1996). For dark matter the above integrals can only be performed numerically. The obtained rotational velocities are shown in Figs 9 - 11. For ordinary matter ( $\lambda = 7/2$ ) we make the choice of  $\Lambda = 3$  to get the familiar solution

$$\chi(x) = \frac{1}{(1+x^2)^{1/2}}, \quad \psi_{7/2} = \psi(x) = \frac{1}{(1+x^2)^{5/2}}$$
(29)

(We will drop the index 7/2 and use the index m, for matter, when necessary). Thus the rotational velocity due to ordinary matter is now given by

$$v_m(x) = \frac{v_0}{\sqrt{2}} \sqrt{g_m(x)} \tag{30}$$




Fig. 1.— The potential function in units of  $\Phi_0$  for  $\lambda=1/2$ . The notation of the curves is: Thick solid line, intermediate thickness solid line, dotted line, fine dashing, long-fine dashing, intermediate-fine dashing and long-long dashing in the order of the asymmetry of Table 1 (here  $\alpha_s = 0.000, -0.021, -0.038, -0.080, 0.062, 0.100, 0.150, 0.200$  and similarly for the other values of  $\lambda$ ).




Fig. 2.— The potential function in units of  $\Phi_0$  for  $\lambda = 1$ .




Fig. 3.— The potential function in units of  $\Phi_0$  for  $\lambda = -1/2$ .




Fig. 4.— The potential function in units of  $\Phi_0$  for ordinary matter.




Fig. 5.— The density in units of  $\rho(0)$  for  $\lambda=1/2$ .




Fig. 6.— The density in units of  $\rho(0)$  for  $\lambda = 1$ .




Fig. 7.— The density in units of  $\rho(0)$  for  $\lambda=-1/2.$ 




Fig. 8.— The density in units of  $\rho(0)$  for ordinary matter,  $\lambda = 7/2$ .




Fig. 9.— The rotational velocities in units of  $\frac{v_0}{\sqrt{2}}$  for  $\lambda = 1/2$ .




Fig. 10.— The rotational velocities in units of  $\frac{v_0}{\sqrt{2}}$  for  $\lambda = 1$ .




Fig. 11.— The rotational velocities in units of  $\frac{v_0}{\sqrt{2}}$  for  $\lambda = -1/2$ .

For x < 1 we write

$$g_m(x) = g(x) (31)$$

while for x > 1 we have:

$$g_m(x) = \frac{1}{x} g(1)$$
 (32)

We now distinguish two cases:

i) Spherical galaxy.

Using the above density we find:

$$g(x) = \frac{1}{3} \frac{x^2}{(1+x^2)^{3/2}} \tag{33}$$

ii) Spiral galaxy (disk).

In this case we will use the same density as above. In the Eddington theory the equation, which relates the potential and the density is no longer of the above simple form. One finds:

$$\frac{\Phi}{\Phi_0} = \frac{\Lambda}{12} \left[ 2 \left( \frac{\rho}{\rho(0)} \right)^5 - \ln \left[ \frac{1 - (\rho/\rho(0))^5}{1 + (\rho/\rho_0)^5} \right] \right]$$
 (34)

This complexity is of no concern to us since we do not need to use this complicated equation. In the case of ordinary matter we do not need to obtain the velocity distribution. Thus with the assumed density we find:

$$g(x) = \frac{1}{3x} \left[1 - \frac{1}{(1+x^2)^{3/2}}\right]$$
 (35)

The above functions are plotted in Fig. 12. In the graphs we show not only the physically interesting case, in which the density vanishes outside the radius of the galaxy, but, for illustrative purposes, the case of infinite extent of the same density function. We see that, since the density falls very fast, it makes very little difference, which form we use.

By comparing the above graphs of dark matter with those of ordinary matter we see that in the Eddington theory ordinary matter can make a significant contribution to the rotational velocities, only if the density at the origin is much bigger compared to that of dark matter. As we have mentioned above for ordinary matter there is no constraint between  $\Phi_0$  and  $\rho_0$ . So we simply rescale ordinary matter by a factor  $C_{mdm}$  and write:

$$g_{matter}(x) = C_{mdm}g_m(x) \tag{36}$$

the scaling factor can be determined by a comparison to the experimental rotational curves.




Fig. 12.— The rotational velocities in units of  $\frac{v_0}{\sqrt{2}}$  ordinary matter. Thick lines are associated with a spherical galaxy, while the thin lines are associated with a spiral galaxy. The finite extent of the matter density gives essentially the same results with that of infinite range. This happens, because the density falls relatively fast.

Thus the rotational velocity due to both matter and dark matter is given by

$$v_{\lambda}(x) = \frac{v_0}{\sqrt{2}} \sqrt{g_{\lambda}(x) + g_{matter}(x)}$$
(37)

The obtained rotational velocities for two values of  $C_{mdm}$  are shown in Figs 13 - 16 The larger value of  $C_{mdm}$  seems to be in better agreement with the data, see, e.g., the recent review (Jungman et al. 1996).

It is clear from these curves that the presence of asymmetry has a dramatic effect on the rotational velocities. Thus, in the context of the Eddington theory as discussed here, a large asymmetry is excluded by the data on rotational velocities. A detailed fitting of our input parameters to the rotational curves will not be attempted here.

## 5. Velocity distribution with respect to the galactic center

The above density via the Eddington formula leads to a velocity distribution of the form:

$$F(v,r) \propto [-2\Phi(r) - v^2]^{\lambda} (1 + \alpha_s \frac{v_t^2}{v_m^2})$$
 (38)

The above velocities and the distance r are defined with respect to the center of the matter distribution, i.e to the center of our galaxy. We note in the context of the Eddington theory the velocity distribution cannot be Maxwellian. For a given distance it goes to zero at the boundaries of the corresponding ellipsoid. It is customary to consider the value of the above distribution in our vicinity,  $r = r_s$ . This way it reduces to the product of the local density and the velocity distribution. The latter is given by Eq. (21) where N is a normalization constant, which depends on  $\lambda$ ,  $\alpha_s$  and  $v_m$ . The above notation was introduced to make the last equation coincide with the standard expression when the function f is chosen to be Maxwellian, i.e.

$$Exp(-\frac{v^2-\alpha_s v_t^2}{v_0^2}) \to (1+\alpha_s \frac{v_t^2}{v_0^2}) \ Exp(-\frac{v^2}{v_0^2})$$

for sufficiently small  $\alpha_s$ . In this limit we see that  $\alpha_s$  coincides with the parameter  $-\lambda$  of Vergados (Vergados 2000) et al Drucker et al. (1986) (in the present work  $\lambda$  is used for another purpose).

It is straightforward to find that the normalization factor  $\tilde{N}$  is given by

$$\tilde{N}^{-1}(\lambda, \alpha_s, \nu_m) = 2\pi \nu_m^{2\lambda + 3} \beta(\lambda + 1, 3/2) \left[ 1 + \frac{4}{3} \alpha_s \frac{\beta(\lambda + 1, 5/2)}{\beta(\lambda + 1, 3/2)} \right]$$
(39)




Fig. 13.— The rotational velocities in units of  $\frac{v_0}{\sqrt{2}}$  for  $\lambda=1/2$  in the case of spiral galaxy with  $_{Cmdm}=30$ .




Fig. 14.— The same as in Fig. 13 for  $C_{mdm}=2.5.$ 




Fig. 15.— The same as in Fig. 13 for a spherical galaxy.




Fig. 16.— The same as in Fig. 14 for a spherical galaxy.

with  $v_m$  given by Eq. 22. In the special case of dark matter  $(\lambda = 1/2)$  it becomes

$$\tilde{N}^{-1}(1/2, \alpha_s, \nu_m) = \frac{\pi^2}{4} \nu_m^4 \left[1 + \frac{1}{3} \alpha_s\right]$$
(40)

From the above formulas we see that the velocity of dark matter with respect to the galactic center ranges from 0 to a maximum speed  $v_m = |2\Phi(r_s)|^{1/2}$ . Since the escape velocity is determined by the potential  $\Phi(r_s)$  due to all kinds of matter, the velocity  $v_m$  is not simply related to  $v_{esc}$ . Thus, since the distribution function must remain positive, if  $\alpha_s < 0$  its absolute value is bounded. This imposes a constraint, since in the traditional analysis with only axially symmetric Gaussian distribution it leads to negative  $\alpha_s$ , i.e. enhanced dispersion in the galactocentric direction, a phenomenologically preferred result (Drucker et al. 1986). The data on rotational curves may provide an additional constraint on the negative values of  $\alpha_s$ . For positive values of  $\alpha_s$  the constraint coming from the rotational curves is, as we have seen, more stringent.

From then on one proceeds in the usual way to obtain the velocity distribution with respect to the laboratory.

#### 6. Velocity distribution with respect to the laboratory

For this transformation one needs the velocity of the sun around the galaxy  $v_0 = 220Km/s$ , a fraction of the escape velocity, which is  $v_{esc} = 625Km/s = 2.84 v_0$  (Drucker et al. 1986).

It is convenient to choose coordinate system with its polar z-axis in the direction of the disc's rotation, i.e. in the direction of the motion of the sun, the x-axis is in the outward radial direction and the y-axis is perpendicular to the plane of the galaxy. Since the axis of the ecliptic (Kosmas and Vergados 1997). lies very close to the y, z plane the velocity of the earth around the the center of the galaxy is given by

$$\boldsymbol{v}_E = v_0 \hat{\mathbf{z}} + \boldsymbol{v}_1 = v_0 \hat{\mathbf{z}} + v_1 (\sin \alpha \, \hat{\mathbf{x}} - \cos \alpha \, \cos \gamma \, \hat{\mathbf{y}} + \cos \alpha \, \sin \gamma \, \hat{\mathbf{z}})$$
(41)

where  $\alpha$  is the phase of the earth's orbital motion,  $\alpha = 2\pi(t - t_1)/T_E$ , where t is the time of observation and  $t_1$  is around second of June and  $T_E = 1$  year. The magnitude of the Earth's velocity is much smaller than that of the sun, i.e.  $\delta_1 = 2v_1/v_0 = 0.27$  The velocity of the earth around its own axis is even smaller and it is usually neglected.

One can now express the above distribution in the laboratory frame by writing  $\boldsymbol{v}' = \boldsymbol{v} + \boldsymbol{v}_E$ , where the prime indicates the velocity with respect to the center of the galaxy.

Indicating by y the velocity of the LSP (Lightest Supersymmetric Particle) in units of  $v_0$ , i.e. by defining  $y = v/v_0$ , we find

$$f(y,\theta,\phi) = N(\lambda,\alpha_s,y_m) \left[ y_m^2 - Y(y,\theta,\phi) \right]^{\lambda} \left[ 1 + \alpha_s (Y(y,\theta) - (y\sin\theta\cos\phi - \frac{\delta_1}{2}\sin\alpha)^2) \right]$$
(42)

with

$$N(\lambda, \alpha_s, y_m) = v_0^{2\lambda + 3} \tilde{N}(\lambda, \alpha_s, v_m)$$
(43)

where  $\tilde{N}(\lambda, \alpha_s, \nu_m)$  is given by Eq. 39,  $y_m = \nu_m/\nu_0$  and

$$Y(y,\theta,\phi) = 1 + \frac{\delta_1^2}{4} + y^2 + 2y\cos\theta + \delta_1[y\cos\theta\cos\alpha\sin\gamma - y\sin\theta\sin\phi\cos\alpha\cos\gamma + y\sin\theta\cos\phi\sin\alpha]$$
(44)

with In the conventional axially symmetric Gaussian velocity distribution this function is given by

$$f(y,\theta,\phi) = \frac{N(\alpha_s, y_{esc})}{\pi\sqrt{\pi}} Exp[-(\alpha_s + 1)Y(y,\theta,\phi) + \alpha_s(y \sin\theta \cos\phi - \frac{\delta_1}{2} \sin\alpha)^2]$$
(45)

In this case  $0 \le y \le y_{esc}$ , but the upper cutoff is introduced here artificially. The normalization here is defined so that  $N(\alpha_s = 0, y_{esc} \to \infty) = 1$  (Vergados 2000)

The detection rate in direct dark matter experiments is obtained by convoluting the the relevant cross section with the above velocity distribution. If the dark matter candidate is the LSP, the  $\alpha$ -dependence of the above distribution, present only when  $\delta_1 \neq 0$ , gives rise to the modulation effect, i.e. the dependence of the rate on the Earth's motion. This signal can be used to discriminate against background.

#### 6.1. The non directional rate

The non directional differential event rate is given by:

$$\frac{dR}{du} = \bar{R}\sqrt{\frac{2}{3}}T(u) , \ T(u) = a^2|F(u)|^2\Psi(a\sqrt{u})$$
 (46)

for the coherent mode and

$$\frac{dR}{du} = \bar{R}\sqrt{\frac{2}{3}}T_{spin}(u) , T_{spin} = a^2|F_{11}u|\Psi(a\sqrt{u})$$
 (47)

where  $\bar{R}$  and  $\bar{R}_{spin}$  are the total rates for the coherent and the spin contributions associated with an average LSP velocity,  $\sqrt{\langle v^2 \rangle} = \sqrt{(3/2)}v_0$ . These parameters, which carry the dependence on the SUSY parameters, are the most important ones, but they are not of interest in our present calculation. F(u) is the form factor, entering the coherent scattering and  $F_{11}(u)$  is the spin response function entering via the axial current (Vergados 2001b). The function  $\Psi$  depends on the LSP distribution velocity employed and is a function of the energy Q transferred to the nucleus

$$u = \frac{Q}{Q_0}$$
 ,  $Q_0 = 4.1 \times 10^4 A^{-4/3} KeV$  (48)

where A is the nuclear mass number and the parameter a is given by:

$$a = \left[\sqrt{2}\mu_r b v_0\right]^{-1} \tag{49}$$

where  $\mu_r$  is the reduced mass of the LSP-nucleus system and b is the (harmonic oscillator) size parameter.

The function, which is basic to us,  $\Psi$ , is given by

$$\Psi(x) = \int_0^x dy \int_0^{\pi} \sin\theta d\theta \int_0^{2\pi} d\phi \ y \ f(y,\theta,\phi)$$
 (50)

with  $0 \le x \le y_m - 1 + (\delta_1/2) \cos \alpha \sin \beta$ .

The total rate is given by:

$$R = \int_{u_{min}}^{u_{max}} \frac{dR}{du} du \tag{51}$$

where  $u_{min}$  is determined from the cutoff energy of the detector and  $u_{max} = (y_m/a)^2$ 

From now on we will specialize our results to the case  $\lambda = 1/2$ , which yields an adequate description of the rotation curves in the case of dark matter. In the case of this velocity distribution, unlike the Gaussian one, one cannot approximate the distribution by a power series in  $\delta_1$ . The reason is that there may be threshold problems, when the argument of the square root approaches zero. To simplify matters we will still make use of the fact that the velocity of the Earth around the Sun is much smaller than the velocity of the Sun around the galaxy,  $\delta_1 << 1$ . So, if we expand the previous expression into a Fourier series with respect to the phase of the Earth,  $\alpha$ , only the lowest terms will become important. In other words to leading order in  $\delta_1$  it can be put in the form:

$$R = \bar{R}[R_0 + (R_1 \cos \alpha \sin \gamma - R_2 \cos \alpha \cos \gamma + R_3 \sin \alpha)\delta_1/2]$$
 (52)

It turns out that the expansion coefficients  $R_2$  and  $R_3$  are zero. We can thus conveniently fit the rate with the formula:

$$R = \bar{R} \ t \ [1 + h \ \cos \alpha] \tag{53}$$

where h is the modulation amplitude (the difference between the maximum and the minimum is equal to 2|h|).

In the case of no modulation,  $\delta_1 = 0$ , the angular integrals can be done analytically to yield:

$$\Psi(x) = 2\pi N(1/2, \alpha_s, y_m) \left[ \frac{(1 + \alpha_s y_m^2)}{3} J_3(x, y_m) - \frac{\alpha_s}{20} J_5(x, y_m) \right]$$
 (54)

with  $0 \le x \le y_m - 1$  and

$$J_n(x, y_m) = J_{int}(n, y_m, x - 1) - J_{int}(n, y_m, x + 1) + 2 J_{int}(n, y_m, 1)$$
(55)

$$J_{int}(n, y_m, y) = \int_0^y [y_m^2 - z^2]^{n/2} dz$$
 (56)

The above integral can be done analytically to yield:

$$J_{int}(n, y_m, y) = y(y_m)^n \, _2F_1(\frac{1}{2}, \frac{-n}{2}, \frac{3}{2}, \frac{y^2}{y_m^2})$$
(57)

where  ${}_{2}F_{1}$  is the usual hypergeometric function. For the cases of interest to us here the hypergeometric function can be simplified to yield:

$$J_{int}(3, y_m, y) = \frac{1}{8} \left[ 2y(y_m^2 - y^2)^{3/2} + 3y_m^2 y(y_m^2 y^2 - y^2)^{1/2} + 3y_m^4 \sin^{-1}(y/y_m) \right]$$
 (58)

$$J_{int}(5, y_m, y) = \frac{1}{24} [4y(y_m^2 - y^2)^{5/2} + 5y_m^2 y(y_m^2 - y^2)^{3/2} + 15y_m^4 y(y_m^2 - y^2)^{1/2} + 15y_m^6 \sin^{-1}(y/y_m)]$$
(59)

$$J_{int}(7, y_m, y) = \frac{1}{192} [24y(y_m^2 - y^2)^{7/2} + 28y_m^2 y(y_m^2 - y^2)^{5/2} + 35y_m^4 y(y_m^2)^{1/2}$$

$$+ 105y_m^6 y(y_m^2)^{1/2} + 105y_m^8 \sin^{-1}(y/y_m)]$$
(60)

The corresponding expressions for the Gaussian expressions for  $\alpha_s$  cannot be done analytically. For the symmetric case,  $\alpha_s = 0$ , one finds that:

$$\Psi(x) = \frac{1}{2} [erf(x-1) - erf(x+1)] + 2 \ erf(1)$$
 (61)

with erf(x) the error function:

$$erf(y) = \frac{2}{\pi} \int_0^y e^{-t^2} dt$$
 (62)




Fig. 17.— The function  $\Psi(x)$  for dark matter in the case of the symmetric Gaussian distribution.




Fig. 18.— The function  $\Psi(x)$  in the case of dark matter for the choice  $\lambda=1/2$  of the Eddington theory. The graphs have been labeled as in Fig. 9.

The above functions  $\Psi$  are plotted in Figs 17 and 18.

From Fig. 18 we see that both the value of the function  $\Psi(x)$  as well as its range depend crucially on the parameters of the model. In the absence of asymmetry the Gaussian model, with an appropriate escape velocity put in by hand, yields results, which are almost four times larger than those of the Eddington theory. In the presence of asymmetry our results in the Eddington theory are substantially larger. This is due to the larger peak value attained as well as the larger allowed range of x. On the other hand we know that in the Gaussian model the total rate is not significantly affected by the asymmetry (Vergados 1999). The reason for the strong dependence of the results in the present model on the asymmetry is not, of course, the asymmetry per se, but the fact that all parameters change with it. In particular the value of  $y_m$  (see table 1). In the Gaussian model the introduction of asymmetry did not affect the velocity distribution in any other way, e.g. it did not affect the cutoff value of the velocity distribution. The large upper values of x in the function  $\Psi(x)$ , allowed in the present model for large  $y_m$ , are, of course, somewhat controlled by the nuclear form factor. In order to get a better feeling for such an effect on the rate, we also plot the function T(u), which is proportional to the differential non directional rate. For illustration purposes we have chosen to present results for the popular target  $^{127}I$  and a typical LSP mass of 100GeV(see Fig. 19). The results presented are for the coherent mechanism, but we expect very small changes when the spin contribution is considered. It is clear that the introduction of asymmetry has a profound effect on the rate. We have seen, however, that the large positive values of the  $\alpha_s$  can, in principle, be eliminated from the data on the rotational curves.

Effects like those discussed above may be more pronounced in the case of modulation, not studied in this work.

#### 6.2. The directional rate

The directional differential rate  $(dR/du)_{dir}$  is proportional to

$$T_{dir}(u) = \frac{1}{2\pi} \bar{R} a^2 |F(u)|^2 \Psi_{dir}(a\sqrt{u})$$

$$T_{dir}^{spin} = \frac{1}{2\pi} \bar{R}_{spin} a^2 |F_{11}u| |\Psi_{dir}(a\sqrt{u})$$
(63)

with

$$\Psi_{dir}(x) = \int_0^x dy \int_0^{\pi} \sin\theta d\theta \int_0^{2\pi} d\phi \ y \ f(y,\theta,\phi) \ X \ H(X)$$
 (64)

with H(X) the well known Heaviside function and X is given by

$$X = \cos\Theta \cos\theta + \sin\Theta \sin\theta [\sin\Phi \sin\phi + \cos\Phi \cos\phi] \tag{65}$$

where  $\Theta$  and  $\Phi$  describe the direction of observation  $\hat{e}$ 

$$\hat{e} = \sin\Theta(\cos\Phi\hat{e}_x + \sin\Phi\hat{e}_y) + \cos\Theta\hat{e}_z$$

(There should be no confusion of the angle  $\Phi$  used here with the potential  $\Phi(r)$  used earlier). Note the presence of the factor of  $1/(2\pi)$ , since the azimuthal integration of the recoiling nucleus is not present and we intend to use the same nucleon cross-section both in the directional and the non directional case.

The total rate is proportional to:

$$R_{dir} = \int_{u_{min}}^{u_{max}} \left(\frac{dR}{du}\right)_{dir} du \tag{66}$$

Taking again the lowest Fourier components of the obtained rate as a function of the phase of the Earth we get an expression similar to Eq. 52. Thus, to leading order in  $\delta_1$ , we can fit the total rate by an expression of the form:

$$R_{dir} = \frac{1}{2\pi} \bar{R} \ t_{dir} \left[ 1 + (h_1 - h_2) \cos \alpha + h_3 \sin \alpha \right]$$
 (67)

The parameters  $t_{dir}$  and  $h_i$  i = 1, 2, 3 are obviously functions of the direction of observation, i.e.  $\Theta$  and  $\Phi$ . If one observes in the direction of the Sun's velocity  $h_2 = h_3 = 0$ . Similarly if one observes in a plane perpendicular to the Sun's velocity  $h_1 = 0$ . Instead of  $t_{dir}$  it is best to use the ratio:

$$\kappa = 2\pi \frac{R_{dir}}{R} = \frac{t_{dir}}{t} \tag{68}$$

The parameter  $\kappa$  is essentially independent of the LSP mass, the nuclear parameters and the asymmetry parameter  $\alpha_s$ . But it depends strongly on the direction of observation and is expected to correlate strongly with the angle between  $\hat{e}$  and the Sun's direction of motion. This correlation provides a an experimental signature perhaps better the modulation with the Earth's motion in non directional experiments.

For  $\delta_1 \neq 0$  the above integrals over  $y, \theta, \phi$ , especially in the directional case, can only be done numerically. Such results will appear elsewhere (Braun et al. 2002).

#### 7. Conclusions

In the present paper we studied the density and velocity distributions of cold dark matter in the context of the Eddington theory, considering not only symmetric but axially symmetric distributions as well. In our approach we used standard simple distribution functions of the energy and angular momentum. This lead us to simple relations between the density  $\rho$ 

and the potential  $\Phi$ .  $\rho$  and  $\Phi$  were obtained by solving numerically poisson's equation in a suitable region of space. This procedure allowed us to determine the maximum permitted dark matter velocity. Then we were able to study both the rotational curves as well as the velocity distribution of dark matter in our vicinity. We should mention that this distribution is not Maxwellian and has an upper velocity cutoff built into into it and not put in by hand as in the traditional treatment with Gaussian distribution.

Our results depend on a minimum set of parameters, which were treated as free (see table 1).

We saw that in the context of this theory the predicted rotational velocities, see Figs 13 - 16, depending on the input parameters, can vary significantly in the presence of the asymmetry. This comes mainly from the factor  $x^2$  in the second term of Eq. 15, when the parameter a is reasonably large. By comparing our results to the observed rotational curves (Jungman et al. 1996), one may constrain the parameters of the model. The best choice seems to be the case with a small asymmetry parameter  $\alpha_s$ .

We have also made a preliminary study of the effect of the new velocity distribution on the direct detection rates for cold dark matter. For illustration purposes we have selected the case  $\lambda = 1/2$ . We have seen that, in the context of the Eddington approach, the total rates, unlike the case of the Gaussian distribution (Vergados 1999, 2000), sensitively depend on the asymmetry parameter  $\alpha_s$ . This is due to the fact that, when the asymmetry changes, the upper value of the velocity distribution also changes.

It is thus not surprising that, in the Eddington theory, the total (non directional and non modulated) event rates for direct LSP detection maybe substantially different from those of the phenomenological Gaussian distributions (compare Figs 17 and 18). The strong dependence of the rate on the asymmetry parameter remains even after the nuclear form factor has been incorporated (see Fig. 19). Results of more detailed calculations of the event rates will appear elsewhere.

The dependence of the directional and/or modulated rates on the velocity distribution is currently under study. We expect this dependence to be more pronounced than on the total rates.

### 8. acknowledgments

J.D.V. would like to thank the Physics Department of UNISA and Professor S. Sofianos for their hospitality. D.O appreciates the hospitality provided by the University of Ioannina.

#### REFERENCES

Alcock, C. et al. 1995, Phys. Rev. Lett. 74, 2967.

Arnowitt, R. and Nath, P. 1995, Phys. Rev. Lett. 74, 4952.

Arnowitt, R. and Nath, P. 1996, Phys. Rev. D 54, 2394; hep-ph/9902237.

Bednyakov, V.A., Klapdor-Kleingrothaus, H. V. and Kovalenko, S.G. 1994 Phys. Lett. B 329, 5.

Bernabei, R. et al. 1996, Phys. Lett. B 389, 757.

Bernabei, R. et al. 1998, Phys. Lett. B 424, 195.

Bernabei, R. et al. 1999, Phys. Lett B 450, 448.

Bottino, A. et al. 1997, Phys. Lett B 402, 113.

Braun, M., Owen, D., Sofianos, S. and Vergados, J.D. press, to be published.

Buckland, K.N., Lehner, M.J. and Masek, G.E. 2000, in Proc. 3nd Int. Conf. on Dark Matter in Astro- and part. Phys., Dark2000, Ed. . Klapdor-Kleingrothaus, H.V., Springer Verlag.

Cheng, T.P. 1988, Phys. Rev. D 38, 2869.

Cheng, H-Y. 1989, Phys. Lett. B 219, 347.

Copi, C., Heo, J. and Krauss, L. 1999, Phys. Lett B 461, 43.

Dawson, S. 1991, Nucl. Phys. B359, 283.

Divari, P.C., Kosmas, T.S., Vergados, J.D. and Skouras, L.D. 2000, Phys. Rev. C 61, 044612-1.

Djouadi, A. and Drees, M. 2000, Phys. Lett. B bf 484, 183.

Drees, M. and Nojiri, M.M. 1993, Phys. Rev. D 47, 376.

Drees, M. and Nojiri, M.M. 1993, Phys. Rev. D 48, 3843.

Drees, M. and Nojiri, M.M. 1993, Phys. Rev. D 47, 4226.

Drucker, A., Freeze, A., Spergel, D. 1986, Phys. Rev. D 33, 3495.

Eddington, A. S. 1916, NRAS 76, 572.

Jaan Einasto, in Dark Matter in Astro and Particle Physics, p.3, Ed. H.V. Klapdor- Kleingrothaus, Springer-Verlag, Berlin-Heidelberg, 2001.

Evans, N.W., MNRAS, 267, 333.

Gawiser, E. and Silk, J. 1988, Science 280, 1405.

Gelmini, G. and Gondolo P. 2001, Phys. Rev. D 64, 623504.

Gómez, M.E., Vergados, J.D. 2001, Phys. Lett. B 512, 252; hep-ph/0012020.

Gómez, M.E., Lazarides, G. and Pallis, C. 2000, Phys. Rev. D 61, 123512.

Gómez, M.E., Lazarides, G. and Pallis, C. 2000, Phys. Lett. B 487, 313.

Gómez, M.E. and Vergados, J.D. press, hep-ph/0105115.

Gondolo P. 2001, Phys. Rev. D 64, 023504.

Green, A 2001, Phys. Rev. D 63, 103003.

Gross, M.A.K. Somerville, R.S., Primack, J.R., Holtzman, J. and Klypin, A.A. 1998, Mon. Not. R. Astron. Soc. 301, 81.

Henriksen, R.N. and Widrow L.M., MNRAS, 276, 679.

Jungman, G. et al. (1996), Phys. Rep. 267, 195.

Kosmas, T.S. and Vergados, J.D. 1997, Phys. Rev. D 55, 1752; from Yadernaya Fisika, Vol. 61, No 7, 1166 1998.

Merritt, D. 1985, A J 90, 1027.

Perlmutter, S. et al 1999, ApJ 517, 565; astro-ph/9812133.

Perlmutter, S. et al 1997, ApJ 1997 483, 565; astro-ph/9608192.

Perlmutter, S., Turner, M.S. and White, M. 1999, Phys. Rev. Let. 83, 670.

J.R. Primack, astro-ph/0205391.

Ressell, M.T. 1993, et al., Phys. Rev. D 48, 5519.

Riess, A.G. et al 1998, AJ 116, 1009.

Sikivie, P. 1999, Phys. Rev. D 60, 063501.

Sikivie, P. 1998, Phys. Lett. 432, 139; astro-ph/9705038.

Smoot, G.F. et al. 1992, COBE data, ApJ 396, L1.

Somerville, R.S., Primack, J. R. and Fabers, S.M. press, Mon. Not. R. Astron. Soc. in press; astro-ph/9806228.

Spira, M. et al 1995, Nucl. Phys. B453, 17.

Turner, M.S. 1990, Phys. Rep. 333-334, 619; astro-ph/9904051.

Ullio, P., Kamionkowski, M. 2001, JHEP 0103, 049.

Vergados, J.D.(1990), Supersymmetric Dark Matter Detection- The Directional Rate and the Modulation Effect, hep-ph/0010151;

Vergados, J.D. 1996, J. of Phys. G 22, 253.

Vergados, J.D. 1998, Phys. Rev. D 58, 103001-1.

Vergados, J.D. 1999, Phys. Rev. Lett 83, 3597.

Vergados, J.D. 2000, Phys. Rev. D 62, 023519; astro-ph/0109296.

Vergados, J.D. 2001, Phys. Rev. D 63, 06351.

Vergados, J.D. 2001, SUSY Dark Matter in Universe- Theoretical Direct Detection Rates, Proc. NANP-01, International Conference on Non Accelerator New Physics, Dubna, Russia, June 19-23, 2001, Editors V. Bednyakov and S. Kovalenko; hep-ph/0201014.

Vergados, J.D. 2002, to be published; astro-ph/0201269.

Widrow, L.M, astro-ph/0003302.

This preprint was prepared with the AAS IATEX macros v5.0.

Table 1: The various parameters describing the dark matter distribution for  $x_r = 20$  (for the definitions see text).

	b	$\Lambda$	a	$h_v$	$\alpha_s$
	0.00	0.047	0.000	4.61	0.000
	-0.16	0.064	-0.010	3.97	-0.021
	-0.24	0.084	-0.020	3.50	-0.038
	-0.30	0.134	-0.040	2.79	-0.080
$\lambda = 1/2$	1.45	0.021	0.031	6.78	0.062
	3.34	0.015	0.050	8.02	0.100
	6.95	0.011	0.075	9.40	0.150
	12.0	0.008	0.100	10.61	0.200
	0.00	0.062	0.000	4.05	0.000
	-0.13	0.088	-0.011	3.41	-0.022
	-0.20	0.102	-0.020	3.18	-0.040
	-0.27	0.171	-0.046	2.50	-0.089
$\lambda = 1$	0.70	0.037	0.026	5.21	0.051
	2.00	0.026	0.052	6.13	0.103
	4.20	0.018	0.078	7.30	0.155
	7.50	0.013	0.099	8.50	0.201
	0.00	0.025	0.000	6.37	0.000
	-0.15	0.032	-0.005	5.57	-0.010
	-0.25	0.038	-0.008	5.15	-0.015
	-0.40	0.050	-0.012	4.51	-0.025
$\lambda = -1/2$	2.70	0.010	0.025	10.26	0.050
	9.20	0.005	0.050	13.22	0.100
	19.4	0.004	0.075	15.53	0.150
	34.0	0.003	0.101	17.53	0.200




Fig. 19.— The function T(u) for dark matter in the Eddington theory for the coherent mechanism in the case of the target  $^{127}I$ . The graphs have been labeled as in Fig. 18. For illustration purposes an LSP mass of 100GeV has been chosen.