A spectral survey of the Orion Nebula from 455 – 507 GHz

Glenn J. White^{1,2,3}, M. Araki⁴, J.S. Greaves⁵, M. Ohishi⁶, N.S. Higginbottom⁷

¹ Centre for Astrophysics & Planetary Science, University of Kent, Canterbury, Kent CT2 7NR, England

² Stockholm Observatory, SE–133 36 Saltsjöbaden, Sweden

- ³ Astrophysics Group, The Cavendish Laboratory, University of Cambridge, Madingley Road, Cambridge CB3 OHE, England
- ⁴ Institute for Physical Chemistry, University of Basel, Klingelbergstrasse 80, CH-4056 Basel, Switzerland
- ⁵ Royal Observatory, Blackford Hill, Edinburgh EH9 3HJ, United Kingdom
- ⁶ National Astronomical Observatory of Japan, 2-21-1, Osawa, Mitaka, Tokyo, 181-8588 Japan
- ⁷ Department of Physics, Queen Mary & Westfield College, University of London, Mile End Road, London E1 4NS, England

Received 11th May 2002 / Accepted 22nd May 2003

Abstract. The results of a submillimetre wavelength spectral line survey between 455.1 - 507.4 GHz of the Orion-KL hot cloud core are reported. A total of 254 lines were detected to a main beam brightness temperature sensitivity $T_{\rm mb} \sim 1$ - 3 K. The detected lines are identified as being associated with 30 different molecular species or their isotopomeric variants. The strongest line detected was the J = 4-3 transition of the CO molecule. Apart from abundant diatomic rotors such as CO and CS, the spectrum is dominated by SO, SO₂ and CH₃OH and large organic molecules such as (CH₃)₂O, CH₃CN, C₂H₃CN, C₂H₅CN and HCOOCH₃ which make up ~ 72% of the total number of lines; unidentified lines ~ 13%; and other lines the remaining ~ 15% of the total. Rotational temperatures and column densities derived using standard rotation diagram analysis techniques were found to range from 70 - 300 K, and $10^{14} - 10^{17}$ cm² respectively.

Key words. Molecules - star formation - molecular cloud

1. Introduction

The chemistry of the Orion-KL molecular cloud core has been better studied than that of any other massive star formation region in the Galaxy high spectral resolution spectroscopic surveys have been carried out by a number of authors including: 72 - 91 GHz Johansson et al. (1984), 70 - 115 GHz Turner (1989), 138 - 151 GHz Lee, Cho, & Lee (2001), 150 - 160 GHz Ziurys & McGonagle (1993), 215 - 247 GHz Sutton et al. (1985), 216 - 242 GHz Blake et al. (1986), 247 - 263 GHz Blake et al. (1987), 257 -273 GHz Greaves & White (1991), 330 - 360 GHz Jewell et al. (1989), 325 - 360 GHz Schilke et al. (1997), 342 - 359 GHz White et al. (1986), 334 - 343 GHz Sutton et al. (1995), 607 - 725 GHz Schilke et al. (2001), 780 -900 GHz Comito et al. (2003), 190 - 900 GHz Serabyn & Weisstein (1995)]. Spectral line surveys can provide an unbiased view of the molecular constituents of the gas in star forming regions, and may be used to estimate the physical and chemical environment. We report here the first high spectral resolution survey in the 600 and 650

 $\mu \mathrm{m}$ atmospheric windows between frequencies of 455 and 507 GHz.

2. The Data

The spectral line survey was made using the James Clerk Maxwell telescope in Hawaii during October 1993 over the frequency range 455.1 – 507.4 GHz. This survey extended across the most of the two atmospheric transmission windows near 650 and 600 μ m. These windows are bracketed by strong telluric H₂O absorption lines, and their transparency is highly dependent on the line of sight water vapour column. The data were collected using the JCMT facility receiver, RxC, operated in double-sideband mode. The adopted 'on-source' position was that of the 'hot core' close to IRc2 (α, δ)₁₉₅₀ = 5^h 32^m 46^s.9, -5° 24' 23''. The pointing accuracy was measured to be good to better than 2'' rms, from observations of planets and compact calibrator sources used as standards at the telescope.

The half power beam width and main beam efficiencies of the telescope were measured from observations of Mars, Jupiter and Uranus. These ranged from 11'' and 0.53 at the low frequency end of the spectral region to 10'' and 0.49 at the upper end of the band. The receiver

Send offprint requests to: g.j.white@kent.ac.uk

double sideband system noise temperatures were typically 1000 - 3000 K. The IF frequency was 3.94 GHz, and the spectra were processed using the JCMT facility 512 MHz bandwidth acousto-optical spectrometer, giving an effective spectral resolution of $\sim 0.6 \text{ km s}^{-1}$. The spectral region was covered by stepping the local oscillator in 100 MHz steps across the whole spectral region - so that any part of the spectrum was redundantly observed at least four times (i.e. least twice in each sideband - and in many cases more times). Each observation consisted of 4 - 10 minutes integration (total on and off source), which was carried out in a 'position-switched' mode, where the telescope was alternated between the on-source position, and a 'reference position' located 2100'' to the north. Previous observations of this 'reference position' have shown it to be free of significant molecular emission intense enough to affect the accuracy of the survey. The reference position was checked by position switching the telescope against several other positions that were located more than 10 degrees away from the Galactic Plane - and not known to be associated with the locations of any molecular clouds or enhanced interstellar extinction. The spectra were calibrated channel by channel using the standard JCMT three temperature chopper calibration scheme (hot and cold loads and the atmosphere). Observations of the sideband gains were measured by observations of spectral lines that were present in both sidebands. The main beam brightness temperature noise levels varied from 1 - 4 K in a 2 MHz channel (~ 1.3 km s^{-1}).

During the data reduction, we attempted to recover an estimate of the single-sideband spectrum, using data collected with a double-sideband receiver. A difficulty common to sideband-deconvolution techniques is the uniqueness of the deconvolution given the observational factors such as pointing reproducibility, sideband gain imbalances, variable calibration solutions as sky conditions change and contamination or blending with strong lines (of both terrestrial and extra-terrestrial origin) in the opposite sideband. A deconvolution technique was used that separated out the emission into the individual sidebands. The basis of this technique (which has been widely used by many observers at the JCMT and forms part of the facility software - although this paper gives the first description of the algorithm that is used) is to set up a series of linked equations for each channel in the DSB spectrum. The first equation simply describes that the DSB line temperature is the sum of two intensities, one from the upper and one from the lower sideband $(T_{11} \text{ and } T_{1}, \text{say})$. The second equation refers to the same spectral channel but with a shift in the local oscillator setting by $\Delta \nu$: the DSB signal is then the sum of T_u and $T_{(l+2\Delta\nu)}$, if we consider the upper sideband to be the frame of reference and the shift to be positive (increased frequency). Similarly we can consider the lower sideband to be the frame of reference, and obtain a third DSB signal that is the sum of $T_{(u+2\Delta\nu)} + T_{l}$. This equation set can be extended as far as desired by taking any of the line frequencies offset by $2\Delta\nu$ and establishing which upper and lower sideband frequencies contribute to the observed DSB signal. The result is always to establish a set of n equations with (n + 1) variables that are linked to a particular channel of the DSB spectrum. A solution can only be found where one of the T(DSB) is consistent with zero within the noise level of the observations. Since this implies (if there are no absorption lines) that both $T_{\rm u}$ and $T_{\rm l}$ are also zero within the noise, the equation set reduces to (n - 1) unknowns and two known values hence all n equations can be solved. In the present survey, the local oscillator was stepped in 100 MHz intervals, and the number of equations used per spectral channel was typically 4 - 5. For several spectrally crowded regions, additional spectra were taken with different local oscillator offsets to improve the reliability of the deconvolution.

Multiple coverage of individual parts of the spectrum provided sufficient redundancy to allow single sideband spectra to be reconstructed. achieving an acceptable solution at most frequencies. The veracity of the technique could also be checked as lines from the lower and upper sidebands move in opposite directions in the DSB spectrum as the local oscillator frequency is stepped (in 100 MHz intervals). The deconvolution technique worked acceptably for more than 98% of the whole spectral range covered, but the remaining $\sim 2\%$ could not be solved because there was no signal consistent with zero in the DSB signal-set. A greater number of solutions could be found by extending the number of linked equations, but since the solutions are already of the form of sums and differences of DSB signals, the noise level will be increased if more differencing is involved. It is also inherent in the technique that there are a choice of solutions (for n equations and (n-1) unknowns), so we have adopted the minimum SSB results and an initial DSB signal $\leq 2\sigma$ for the 'consistent' with zero' DSB criterion. This minimises the level of spikes and should ensure that temperature solutions are underestimated by less than $\approx 3\sigma$. The locations of the parts of the spectrum for which we did not achieve a good deconvolution, are shown as horizontal bars under the spectrum in Figs 3 - 6. We also visually inspected at the locations in a spectrum where bright lines from the opposite sideband might have left small residual artifacts (sometimes known as 'ghost features' - Schilke et al. 1997), as well as inspecting the emission in the opposite sideband to the locations of all of the 'U-lines'.

3. Data Analysis

The intention of this paper is to present the data and some basic results. The spectrum shown in Fig. 1 is crowded with many blended lines, and in many places is confused - reflecting the rich and complex chemistry. A total of 254 lines were identified, although there may be more lines than this blended together. The lines were clearly identified using the JPL Sub-millimetre spectral line catalogues of Pickett et al. (1995), Pearson et al. (1996, 2000), Müller etal. (2000) and other lists of line frequencies referred to later in this paper. A total of 98 lines (32% of the total) could not be associated with known molecular transitions



Fig. 1. The complete spectrum binned in 2 MHz channels in main beam brightness temperature units. A low order polynomial baseline was removed some of the individual spectra. The atmospheric emission spectrum that is overlaid above the Orion spectrum is purely illustrative, and just shows the main features of the atmospheric emission. Recent observational and modeling studies of the terrestrial atmospheric emission (see for example Naylor et al. 2000, Pardo, Serabyn & Cernicharo 2001a) using collisional parameters extracted from the HITRAN database with an independent radiative transfer model (ATM) and different assumptions about line shapes produce broadly similar atmospheric emission spectra to the one shown here.

and have been designated as U-lines. Although it is possible that some fraction of these may be artifacts of the deconvolution process - the fact that they are distributed throughout the survey, and do not tend to congregate in areas where the deconvolution process was not successful, it is likely that many of these are real. Table 1 gives a breakdown of the number of transitions observed from each of the known species.

These data were used to estimate the rotational temperature, $T_{\rm rot}$, and column densities, $N_{\rm col}$, of the various species using the relationship given in Eq. 1.

$$ln\left(\frac{3kc\int T_{mb}dV}{8\pi^{3}\mu^{2}\nu^{2}S}\right) = ln\left(\frac{N_{T}}{Q\left(T_{rot}\right)}\right) - \frac{E_{u}}{kT_{rot}}$$
(1)

where $\int T_{\rm mb} dV$ is the integrated intensity of the line, S is the intrinsic strength of the transition, Q is the partition function of the molecule and μ is the dipole moment. The value of $\mu^2 S$ was calculated from the value of the transition intensity listed by Pickett et al (1995), using Eq. 2.

$$I_{ba}\left(T\right) = \left(\frac{8\pi^{3}}{3hc}\right)\nu_{ba}{}^{x}S_{ba}\mu_{x}^{2}\left[\frac{e^{\frac{E_{l}}{kT}} - e^{\frac{E_{u}}{kT}}}{Q_{rs}}\right]$$
(2)

The values for $T_{\rm rot}$ and column densities are summarised in Table 1.

One of the objectives in analysing spectral line survey data is to determine molecular parameters, such as rotation temperatures, column densities etc. The approach commonly adopted has been to use a 'rotation diagram' to estimate these parameters. In this, an optically thin transition produces an antenna temperature that is proportional to the column density in the upper level of the transition being observed. If all transitions are thermalised and the kinetic temperature is known, then a single integrated line intensity and be used to estimate the total column density of the species in question. The rotational temperature diagram is a plot showing the column density per statistical weight of a number of molecular energy levels, as a function of energy above the ground state - in local thermodynamic equilibrium (LTE), this is equivalent to a Boltzmann distribution. A plot of the natural logarithm of the column density, N, divided by the degeneracy g, versus the Energy E of the final state of the level (expressed in units of degrees K) E/k, will lead to a straight line fit with a slope of 1/T, where q is the statistical weight of level u lying at E energy above the ground state, and T is the rotational temperature. This is equivalent to the kinetic temperature in the limit where all of the levels are thermalised.

One problem for the rotation diagram method is that it may underestimate the total column density if some of the lines fitted are optically thick, or LTE conditions do not hold, or if the background radiation is non-negligible (Turner 1991, Goldsmith & Langer 1999, Nummelin et al. 2000, Schilke et al. 2001). This can be however be addressed by using less abundant isotopomeric variants that allow estimates to be made of the optical depths, and then



Fig. 2. Cumulative spectrum showing the number of lines detected that exceeded some value.

using these to correct the column density estimates. It has however been widely used in past molecular line survey studies, because of its computational simplicity, and the absence of a need to have observations of an isotopomer. In this paper we give results from the traditional rotationdiagram technique (Table 1), except in cases pointed out in the Table and text where the results may be affected by high optical depth effects. We have examined a number of cases (CH₃CN, SO₂, OCS, SO, CH₃OH and ³⁴SO in some detail - see for example Sections 4.3 and 4.2 - calculating the optically thick and thin total column densities - and found the use of the rotational temperature technique to provide similar column density estimates are quite similar to the rotational diagram values (see for example discussion of opacities in Section 4.2). Another indicator as to whether opacity corrections are important, is to look at the rotational temperature diagrams (see Fig 7), to see whether there are data points that deviate noticeably from a straight line fit - and which may indicate that the line has saturated. Lee, Cho and Lee (2001) applied this kind of constraint to seven molecules (CH₃OH, HCOOCH₃, CH₃OCH₃, C₂H₅CN, SO₂, CH₃CN, and H₂CO), finding that the majority could to first order be treated using the rotational technique. The issue of opacity will be further discussed in the relevant sections dealing with molecules that may be opaque - where we also conclude that the effects of opacity are minor in the analysis of data from this survey, and that our justification of using the optically thin rotational temperature technique is adequate although there clear examples (Schilke et al. 1997, 2001) where opacity corrections are required for CH₃OH.

Full details of the detected lines are given in Tables 2 -6.

4. Discussion of Individual Species

Individual spectra are shown in more detail in Figs. 3, 4, 5 and 6, along with identifications of the prominent lines. Previous studies (Blake et al. 1987, Schilke et al. 1997,

 Table 1. JCMT Orion spectral line survey

Molecule	$N_{\rm col}$	Error	$T_{\rm rot}$	Error	Number	Note
	cm^{-3}	cm^{-3}	Κ	Κ	of lines	
$(CH_3)_2O$	$1.4 \ 10^{16}$	$1.8 10^{15}$	157	30	27(26) lines	
C_2H_3CN	$3.0 \ 10^{17}$	$3.2 10^{17}$	180	47	13(6) lines	[1]
C_2H_5CN	$2.4 10^{16}$	$8.4 10^{15}$	150	12	27(23) lines	T_{rot} from Sut85
	$8.3 \ 10^{15}$	$1.2 10^{15}$	239	12	27(23) lines	T_{rot} from Sch01
C_2H_5OH	$5.6 \ 10^{16}$	$4.0 10^{16}$	70	-	8(6) lines	T_{rot} from Ohi95
	$4.1 10^{16}$	$2.3 10^{16}$	264	196	8(6) lines	
CH_2NH	$2.4 10^{15}$	$5.9 10^{14}$	150	-	3(2) lines	T_{rot} from HNCO
CH_3CN	$3.6 10^{15}$	$4.2 10^{14}$	227	21	19 lines	
$CH_3^{13}CN$	$6.0 10^{14}$	$2.2 10^{14}$	227	-	2 lines	T_{rot} From CH_3CN
	$2.9 10^{15}$	-	74	-	2 lines	
CH_3OH	$9.3 \ 10^{16}$	$4.8 10^{16}$	599	295	24(23) lines	
H_2CO	$1.6 \ 10^{16}$	-	166	-	2(1) lines	T_{rot} from Bla87
$H_2^{13}CO$	$1.0 10^{15}$	$4.0 10^{14}$	166	-	2 lines	T_{rot} from Bla87
HC_3N	$1.5 10^{15}$	-	164	-	2 lines	
HCOOCH ₃	$5.1 \ 10^{16}$	$9.5 10^{15}$	301	95	26(24) lines	
HNCO	$4.9 10^{15}$	$4.0 10^{14}$	150	14	4(3) lines	
NH_2CN	$3.3 \ 10^{15}$	$8.5 10^{14}$	200	-	3 lines	$T_{rot} = 200$ (K) Fix
	$1.1 10^{16}$	$4.8 10^{15}$	100	-	3 lines	$T_{rot} = 100$ (K) Fix
OCS	$9.0 10^{16}$	-	106	-	2 lines	
SO	$3.3 \ 10^{17}$	-	72	-	2(1) lines	T_{rot} from Sut95
^{34}SO	$1.1 10^{16}$	$3.5 10^{15}$	89	43	5(4) lines	
SO_2	$1.2 \ 10^{17}$	$1.0 10^{16}$	136	9	35(28) lines	
$^{34}SO_2$	$8.5 \ 10^{15}$	-	156	-	3(2) lines	
^{13}CS	$2.3 \ 10^{14}$	-	120	-	1 line	T_{rot} from Zen95
30 SiO	$3.4 10^{14}$	-	50	-	1 line	T_{rot} from Sut95
CH_3CHO	$1.1 10^{16}$	-	81	-	1 line	T_{rot} from Sch97
CI	$1.2 10^{18}$	-	30	-	1 line	T_{ex} from Whi95
CO	$3.5 10^{18}$	-	200	-	4 lines (1 transition)	T_{rot} from Sut95
DCN	$1.1 \ 10^{14}$	-	200	-	1 line	T_{rot} from Bla87
HCOOH	$2.2 10^{15}$	-	100	-	1 line	T_{rot} from Sut95
HDO	$3.2 10^{16}$	-	164	-	1 line	T_{rot} from Bla87
N_2O	$4.6 10^{16}$	-	230	-	1 line	T_{rot} from Wri83
NH_2CHO	$7.5 10^{15}$	-	81	-	1 lines	T_{rot} from Sch97
$\rm NH_2D$	$8.7 10^{15}$	-	160	-	1 line	T_{rot} from Her88

LTE rotation temperatures and beam averaged $N_{\rm col}$, estimated using a Boltzmann plot. The $N_{\rm col}$ were determined using the main beam brightness temperature scale, and for species where only one line was measured, we have assumed $T_{\rm rot}$ as described in Column 7. Notes: Column 6: A(B) lines represent A: number of assigned lines, B: number of lines included in the fitting. For A lines only, all lines were included in the fitting. [1]: $v_2 = 1$ state lines were not included in the fitting, [2]: Energy levels were considered in the v = 1 vibrational state. Errors quoted are all 1σ .

2001) have shown that there are four characteristic velocity components in Orion spectra: the extended ridge which is ambient gas in the Orion Molecular cloud ($v_{\rm lsr} \sim$ 9 km s⁻¹ $\Delta\nu \sim$ 4 km s⁻¹), the compact ridge - which is a compact clump lying about 10" south-west of the hot core ($v_{\rm lsr} \sim 8 \,\rm km \, s^{-1} \Delta\nu \sim 3 \,\rm km \, s^{-1}$), the plateau - which which has been identified as the outflow, and is associated with the broadest spectral lines ($v_{\rm lsr} \sim 6$ -10 km s⁻¹ $\Delta\nu \sim$ 20 km s⁻¹), and the hot core that lies close to the infrared source IRc2 ($v_{\rm lsr} \sim 3$ -6 km s⁻¹ $\Delta\nu \sim 5$ -10 km s⁻¹). Most of the observed lines show velocities within these ranges, except a few lines which may suffer from blending (e.g. lines at 455.7798, 459.7069, 466.7817, 495.0007, 496.44095, 503.8517, 504.7281 GHz). The superposed dotted line shows the normalised atmospheric absorption (see Fig 1 caption) under conditions typical for the survey.

4.1. CO

In this survey the CO J = 4-3 transition is the most intense single line, with a full width at half maximum of 40 km s⁻¹ and full width at zero intensity of at least 120 km s⁻¹. The peak main beam brightness temperature $T_{\rm mb} = 182$ K was similar to values that we have previously measured using the JCMT (see for example White & Sandell 1995). The line profile clearly shows a small dip close to its peak, which is probably a self-absorption dip. This has also been reported by Schilke et al. (2001)

Iddie 2. Contra Chon Spectral Integration	Table	2.	JCMT	Orion	spectral	linesurvey
--	-------	----	------	-------	----------	------------

Frequency	Species	Transition	Peak	Width	Vel	Notes
MHz	1		$T_{mh} K$	$\rm km~s^{-1}$	$\rm km~s^{-1}$	
WIIIZ			1 mb 11	KIII 5	KIII 5	
455.160796	SO_2	$21_{5,17}$ - $21_{4,18}$	23.2	33.3	8.2	
455.250265	SO_2	$29_{5,25}$ - $29_{4,26}$	14.8	19.6	6.9	
455.353721	SO_2	$18_{5,13}$ - $18_{4,14}$	17.6	8.0	5.4	
455.450207	U-line		4.3	9.4		
455.617930	CH_3OH	61-50	36.1	10.6	9.1	
455.779855	SO_2	$11_{3,9}$ -10 _{2,8}	60.2	13.8	1.6	Blend with C_2H_3CN
456.271440	CH_3OH	$18_1 - 17_2$	20.2	6.2	8.4	
456.359253	SO_2	$19_{5,15}$ - $19_{4,16}$	29.5	11.6	4.4	
456.513228	U-line		18.8	4.3		
456.553270	U-line		4.7	16.6		$C_3H_2 \ 22_{0,22}-21_{0,21}$?
456.800826	CH_3OH	18_{-2} - 18_{-1}	12.7	7.9	5.3	Blend with C_2H_5CN
456.936702	$(CH_3)_2O$	$15_{3,12}$ - $14_{2,13}$	5.0	9.3	6.9	4 lines blended
457.006904	CH_3OH	$11_2 - 11_1$	23.7	7.6	8.2	
457.248931	CH_3OH	19_{-1} -18_2	9.3	15.6	9.9	
457.325926	SO_2	$16_{5,11}$ - $16_{4,12}$	30.4	11.3	2.2	Blend with C_2H_3CN
457.381156	SO_2	$15_{8,8}$ - $16_{7,9}$ + $15_{8,7}$ - $16_{7,10}$	7.5	14.6	7.8	
457.472642	SO_2	$17_{5,13}$ - $17_{4,14}$	50.1	11.4	5.2	
457.584559	U-line		7.0	3.5		
457.637840	U-line		3.5	7.3		
457.723839	C_2H_5CN	$51_{8,44}$ - $50_{8,43}$	5.1	5.2	11.5	
457.736126	$HCOOCH_3$	$40_{3,37}$ - $39_{3,36}$	9.2	6.0	7.4	
457.783725	U-line		13.3	9.4		
457.822031	U-line		9.1	28.5		
457.839618	U-line		4.7	4.3		$SO_2 v_2 = 1$?
457.878236	C_2H_5CN	$53_{2,52}$ - $52_{2,51}$	5.4	4.3	5.2	
457.890562	C_2H_5CN	$28_{5,23}$ - $27_{4,24}$	5.9	6.0	9.3	
457.947140	HCOOCH ₃	$38_{6,32}$ - $37_{5,32}$	3.4	6.0	10.5	
457.985674	HCOOH	$20_{3,17}$ -19 _{3,16}	3.8	2.8	9.6	
458.022659	C_2H_5CN	$48_{4,45}$ - $47_{3,44}$	4.9	10.3	7.2	
458.148139	CH_3OH	101-92	15.3	7.0	7.0	
458.276916	$(CH_3)_2O$	$25_{1,24}$ - $24_{2,23}$	11.6	3.9	8.8	4 lines blended
458.389953	SO_2	$15_{5,11}$ - $15_{4,12}$	40.8	25.2	6.9	
458.433457	C_2H_5OH	$28_{1,28} - 27_{1,27}$	8.2	5.3	6.1	
458.512326	HCOOCH ₃	412/3/2/2,39-403/3/2/2,38	20.7	12.2	8.5	4 lines blended
458.536360	SO_2	$14_{5,9}$ - $14_{4,10}$	34.8	16.7	5.9	
458.601426	HCOOCH ₃	$36_{11,25}$ - $35_{11,24}$	7.1	8.0	9.9	
458.619675	SO_2	$31_{5,27}$ - $31_{4,28}$	8.6	9.8	3.8	
458.998198	$\rm CH_3CN$	$J=25-24 \ K=9$	3.9	5.7	7.3	
459.054419	U-line		20.5	22.1		
459.076694	SO_2	$13_{5,9}$ - $13_{4,10}$	36.0	14.1	5.0	
459.150830	$\rm CH_3CN$	$J = 25-24 \ K = 8$	4.3	11.2	4.9	
459.257959	U-line		23.3	24.2		
459.378018	U-line		8.3	6.1		
459.395715	$\rm CH_3CN$	$J=25-24 \ K=6$	17.7	10.4	5.4	Blend with $CH_3^{13}CN J=25-24 K=3$
459.443401	${\rm CH_3}^{13}{\rm CN}$	$J=25-24 \ K=2$	4.7	13.9	9.6	
459.492742	$\rm CH_3CN$	$J=25-24 \ K = 5$	20.4	16.7	5.3	Blend with $CH_3^{13}CN J=25-24 K = 0,1$
459.511802	$\rm NH_2CN$	234,20/19-224.19/18	16.0	9.3	9.0	,
459.534263	SO_2	$11_{5,7}$ - $11_{4,8}$	39.4	19.7	5.2	
459.569872	$\rm CH_3CN$	J=25-24 K = 4	17.3	13.6	6.6	
459.631405	$\rm CH_3CN$	J = 25 - 24 K = 3	30.9	13.0	6.6	
459.678274	$\rm CH_3CN$	J=25-24 K = 2	37.8	10.8	4.8	
459.706888	$\rm CH_3CN$	J=25-24 K = 0,1	35.4	15.5	3.3	2 lines blended
459.760261	HNCO	211,21-201,20	14.8	7.2	6.1	
459.801217	SO_2	$9_{5,4}$ - $9_{4,5}$ + $9_{5,5}$ - $9_{4.6}$	35.0	22.8	8.1	
459.881164	SO_2	85,3-84,4+85,4-84,5	49.9	32.7	7.6	
450 049599	<i></i>	– –	57.0	26.2	48	
409.942020	SO_2	(5,3-(4,4	51.9	20.2	1.0	

Table	3.	JCMT	Orion	spectral	linesurvey
-------	----	------	-------	----------	------------

Frequency	Species	Transition	Peak	Width	Vel	Notes
MHz	*		$T_{mh} K$	$\mathrm{km} \mathrm{s}^{-1}$	$\mathrm{km} \mathrm{s}^{-1}$	
WIIIZ			1 110 11	KIII 5	KIII 5	
460.069658	C_2H_5CN	$31_{4,27}$ - $30_{3,28}$	2.6	23.5	7.3	
460.214973	$HCOOCH_3$	$37_{6,31}$ - $36_{6,30}$	6.5	3.6	8.8	
460.272941	$^{34}SO_2$	$6_{4,2}$ - $5_{3,3}$	7.5	31.1	7.8	
460.281266	$HCOOCH_3$	$43_{1,43}$ - $42_{1,42}$	8.8	6.9	9.6	
460.306531	C_2H_5CN	$22_{15,7/8}$ - $23_{14,10/9}$	4.0	2.3	7.1	
460.387575	HCOOCH ₃	$37_{12,26}$ - $36_{12,25}$	3.1	27.3	8.0	2 lines blended
460.466108	U-line		3.0	13.7		
460.545686	U-line		6.2	6.2		
460.591671	U-line		5.0	3.9		
460.878573	CH_3OH	92-81	24.4	6.6	7.3	
460.913504	U-line		3.8	5.0		
461.040768	CO	J = 4-3	182.0	51.0	9.0	
461.342825	U-line		5.0	7.5		
461.373909	HNCO	$21_{2,19}$ - $20_{2,18}$	7.4	6.1	5.7	
461.709888	$HCOOCH_3$	$30_{7,24}$ - $29_{6,23}$	5.5	8.4	9.6	
461.756285	CH_3OH	$15_0 - 14_1$	43.8	11.8	7.8	
461.880198	SO_2	$20_{9,11}$ - $21_{8,14}$ + $20_{9,12}$ - $21_{8,13}$	7.9	12.6	6.2	
461.910483	OCS	J=38-37	18.1	11.8	7.2	
462.035190	$CH_3OH v_t = 1$	10_{6} -11 ₅ +	8.1	13.0	9.8	
462.036358	C_2H_5OH	$13_{5,9}$ -12 _{4,8}	8.1	13.0	12.5	Blended with CH ₃ NC
462.138660	HCOOCH ₃	$37_{11,27}$ - $36_{11,26}$	3.5	7.0	8.4	
462.236037	^{34}SO		42.0	19.3		
462.334032	^{13}CS	J = 10-9	14.6	16.4	6.8	Blended with C ₂ H ₅ CN
463.017334	SO_2	$12_{2,10}$ - $11_{1,11}$	21.3	23.0	5.2	
463.122260	HNCO	$21_{1,20}$ - $20_{1,19}$	8.5	6.8	7.2	
463.329490	SO_2	$35_{4,32}$ - $35_{3,33}$	5.6	14.3	6.9	
463.720861	HC_3N	J = 51 - 50	8.2	13.2	5.4	
463.836119	$^{34}\mathrm{SO}_2$	$26_{0,26}$ - $25_{1,25}$	8.2	10.4	9.1	
464.026804	U-line		6.8	25.2		
464.091929	^{34}SO	$N_J = 11_{11} - 10_{10}$	20.8	25.8	11.6	
464.200231	CH_3CHO	$7_{4,4}$ - $6_{3,3}$	4.2	5.8	7.9	Blend with HCOOCH ₃
464.200231	HCOOCH ₃	$37_{10,28}$ - $36_{10,27}$	4.2	5.8	9.5	Blend with CH ₃ CHO
464.293855	SO_2	$33_{5,29}$ - $33_{4,30}$	4.1	10.8	5.1	
464.570359	CH_2NH	$7_{1.6}$ - $6_{1.5}$	4.4	16.8	8.5	
464.730426	U-line		12.9	6.6		
464.837513	CH_3OH	92-91	34.0	11.3	7.2	Blend with C_2H_5CN 9 ₂ -9 ₁
464.928173	HDO	$1_{0,1}$ - $0_{0,0}$	33.5	16.2	6.6	
465.061222	C_2H_3CN	$52_{2,49} - 49_{2,48} v_2 = 1$	1.9	11.1	12.3	
465.078910	U-line		3.6	4.1		
465.352350	^{34}SO	$N_J = 11_{12} - 10_{11}$	30.5	23.0	7.2	
465.512820	U-line		2.4	5.4		
465.543972	$(CH_3)_2O$	$21_{3,19}$ - $20_{2,18}$	3.6	11.2	5.9	4 lines blended
465.565977	$HCOOCH_3$	$36_{1,35}$ - $35_{1,34}$	2.4	14.7	9.0	
465.605514	C_2H_5CN	$52_{12,41/40}$ - $51_{12,40/39}$	4.0	11.2	4.4	
465.628191	SO_2	$29_{2,28}$ - $29_{1,29}$	13.0	8.4	5.1	
465.713762	C_2H_5CN	$51_{5,46}$ - $50_{5,45}$	7.4	13.2	5.9	
465.732611	$\rm NH_2CHO$	$22_{1,21}$ - $21_{1,20}$	16.1	9.7	6.7	
465.758340	SO_2	$26_{0,26}$ - $25_{1,25}$	45.3	15.8	4.4	
465.882288	$(CH_3)_2O$	$14_{4,11}$ - $13_{3,10}$	4.0	9.9	6.1	Blend with SO_2
465.882288	SO_2	$25_{10,16}$ - $26_{9,17}$ + $25_{10,15}$ - $26_{9,18}$	4.0	9.9	8.7	Blended with $(CH_3)_2O$
465.922779	$SO_2 v_2 = 1$	$18_{5,13}$ - $18_{4,14}$	2.2	9.5	7.9	
465.956339	30 SiO	J = 11 - 10	7.7	15.1	8.8	
466.781659	C_2H_3CN	$49_{13,36/37} - 48_{13,35/36} v_1 = 1$	13.8	30.7	11.9	Blended with C_2H_5CN
466.781659	C_2H_5CN	528,44-518,43	13.8	30.7	3.8	Blend with C_2H_3CN
466.892775	SO_2	406,34-405,35	5.5	5.5	5.7	
466.999547	U-line		3.1	5.6		
467.048535	HCOOCH_3	$38_{23,16/15}$ - $37_{23,15/14}$	5.6	14.7	8.9	

Table 4. JCMT Orion spectral linesurvey

Frequency	Species	Transition	Peak	Width	Vel	Notes
MHz	-		$T_{mb} K$	${\rm km}~{\rm s}^{-1}$	$\rm km~s^{-1}$	
467.212513	$HCOOCH_3$	$38_{22,17} - 37_{22,16}$	4.6	5.5	8.6	
467.282605	C_2H_3CN	$49_{16,33/34} - 48_{16,32/33} v_1 = 1$	6.9	7.6	8.2	
467.318607	HCOOCH ₃	$39_{6,34}$ - $38_{6,33}$	3.9	19.2	9.4	2 lines blended
467.475284	U-line	24 25	10.5	8.3		
467.532613	$(CH_3)_2O$	$26_{0,26}$ - $25_{1,25}$	6.2	5.4	7.7	4 lines blended
467.586402	HCOOCH ₃	$39_{5,34}$ - $38_{5,33}$	3.7	10.5	8.3	
467.627790	HCOOCH ₃	$19_{9,10}$ -18 _{8,10}	2.8	10.2	8.5	
467.658900	U-line	15 14	0.U	17.2	0.4	
467.670464	C_2H_5OH	$15_{3,12}$ $14_{2,12}$	10.4	0.4 12.0	9.4 F.C	4 lines blended
407.713023	$(C\Pi_3)_2 O$	$20_{1,26} - 20_{1,25}$	14.4	13.0 6 4	5.0	4 lines blended
401.100133	C I CN	$10_1 - 9_1 v_t = 0,1$	7.1 2.0	0.4	C F	
408.048223	$C_2 \Pi_5 CN$	000,55-040,54	2.9	12.0	0.5	Dland with U line
408.300479	Ulina	82-81	20.3 7.6	8.4 6.0	4.7	Blend with U-line
408.343184	U-line	7 6	1.0	0.0	26	
491.943519	SO_2		49.4	10.3 12.6	3.0 9.6	
491.978910	H_2CO	(1,7-0,6)	45.0 17.0	13.0	2.0	
492.100020		$P_1 - P_1$	17.9	5.U 10.4	9.0 7.0	
492.281601	CH ₃ OH	4_1 - 3_0	36.9	10.4	1.2	
492.470010	U-line		5.2	8.5	F 0	
492.544658	C_2H_5CN	$55_{11,45/44}$ - $54_{11,44/43}$	3.5	4.3	5.3 0.1	
492.695825	(CII)	0 9	9.9	4.1	8.1 9.C	o l'a co blando d
492.784191	$(CH_3)_2O$	9 _{6,4} -8 _{5,4}	20.7	8.4	8.0	8 lines blended
493.260821	C_2H_5CN	$58_{1/0,58}$ -57 $_{1/0,57}$	4.4	(.4 11.4	0.2	
493.370372	U-line	F 4	9.6	11.4	70	
493.701318	CH_3OH	53-42	23.7	11.4	7.6	
493.735068	CH ₃ OH	53-42 -	29.0	9.3	8.1	
494.460056	NH_2D	$1_{1,0} - 0_{0,0} - $	10.5	8.6	5.8	
494.484560	CH ₃ OH	72 - 71 - 72 - 72 - 72 - 72 - 72 - 72 -	50.8	10.3	7.2	
494.555486	SO_2	1327-1228	2.5	4.2	7.6	
494.557464	C_2H_5CN	$55_{7,49}$ - $54_{7,48}$	2.5	4.2	5.4	Blend with C_2H_5OH
494.557464	C_2H_5OH	$10_{9,1} - 9_{8,1}$	2.5	4.2	8.7	Blend with C_2H_5CN
494.575046	C_2H_5CN	325,27-314,28	2.6	6.4	9.5	
494.596465	$(CH_3)_2O$	$28_{9,19} - 28_{8,20} + 28_{9,20} - 28_{8,21}$	5.7	5.3	7.7	8 lines blended
494.756403	C_2H_3CN	$52_{17,35/36} - 51_{17,34/35} v = 0$	14.1	17.1	5.2	
494.781464	SO_2	$12_{3,9}$ -11 _{2,10}	45.0	26.6	7.9	
494.899521	C_2H_5OH	283,25273,24-	1.8	10.5	3.1	
495.000693	$(CH_3)_2O$	$27_{9,18} - 27_{8,19} + 27_{9,19} - 27_{8,20}$	1.6	15.1	10.9	8 lines blended
495.222806	$(CH_3)_2O$	$27_{1,26}$ - $26_{2,25}$	7.8	14.4	7.5	4 lines blended
495.486885	HCOOCH ₃	39 _{7,32} -38 _{7,31}	12.5	4.8	7.4	
495.821871	C_2H_5OH	$26_{4,22}$ - $25_{3,22}$	2.1	2.8	3.5	
495.842389	SO_2	$13_{8,6}$ - $14_{7,7}$ + $13_{8,5}$ - $14_{7,8}$	9.8	20.5	8.5	
495.979430	$(CH_3)_2O$	$24_{9,15}$ - $24_{8,16}$ + $24_{9,16}$ - $24_{8,17}$	8.7	5.8	7.1	8 lines blended with CH_3CN
495.979430	CH ₃ CN	$J=27-26\ K=7$	8.7	5.8	7.0	Blend with $(CH_3)_2O$
496.107704	CH ₃ CN	J=27-26 K=6	11.5	11.6	4.6	
496.207453	CH ₃ CN	J=27-26 K=5	8.8	13.3	6.6	
496.295793	CH ₃ CN	J=27-26 K=4	13.3	11.1	4.8	
496.361787	CH ₃ CN	J=27-26 K=3	19.2	11.6	5.1	
496.409463	CH ₃ CN	J=27-26 K=2	20.7	12.5	5.0	
496.440950	$(CH_3)_2O$	$24_{9,15}$ - $24_{8,16}$ + $24_{9,16}$ - $24_{8,17}$	26.6	17.0	10.9	8 lines blended with CH_3CN
495.175350	CH ₃ OH	7_0-6_{-1}	39.7	10.2	7.7	
496.440950	CH_3CN	J = 27 - 26 K = 1	26.6	17.0	3.2	2 lines blended, Blend with $(CH_3)_2O$
496.518825	$H_2^{13}CO$	$7_{2,6}$ - $6_{2,5}$	3.1	7.2	6.3	
496.633861	$(CH_3)_2O$	$21_{9,12}$ - $21_{8,13}$ + $21_{9,13}$ - $21_{8,14}$	7.0	5.2	8.4	8 lines blended
496.764492	C_2H_3CN	$52_{5,47}$ - $51_{5,46}$ $v_2 = 1$	2.2	14.4	14.1	
496.796267	$(CH_3)_2O$	$20_{9,11}$ - $20_{8,12}$ + $20_{9,12}$ - $20_{8,13}$	5.8	3.8	8.4	8 lines blended
496.928524	CH_3OH	$14_0 - 13_1$	16.7	9.2	6.5	
497.051216	$(CH_3)_2O$	$18_{9,9}$ - $18_{8,10}$ + $18_{9,10}$ - $18_{8,11}$	7.2	14.8	8.6	8 lines blended

Table 5. JCMT Orion spectral linesurvey

Frequency	Species	Transition	Peak	Width	Vel	Notes
MU ₂	Species		T . K	Irm a ⁻¹	$lrm c^{-1}$	
MIIZ			$1 mb \Lambda$	KIII S	KIII S	
497.085708	C_2H_5OH	97.3/2-86.2/3	3.1	10.4	10.3	
497.119025	$H_2^{13}CO$	$7_{4,4/3} - 6_{4,3/2}$	4.8	6.9	6.6	
497.149132	$(CH_3)_2O$	$17_{9,8}$ - $17_{8,9}$ + $17_{9,9}$ - $17_{8,10}$	6.4	7.3	8.5	8 lines blended
497.226494	$(CH_3)_2O$	$16_{9.7}$ - $16_{8.8}$ + $16_{9.8}$ - $16_{8.9}$	5.5	4.8	10.4	8 lines blended
497.295820	$(CH_3)_2O$	$15_{9.6}$ - $15_{8.7}$ + $15_{9.7}$ - $15_{8.8}$	5.2	3.2	8.1	4 lines blended
497.364091	U-line		3.7	7.0		
497.388286	$(CH_3)_2O$	$13_{9,4}$ - $13_{8,5}$ + $13_{9,5}$ - $13_{8,6}$	8.5	10.8	8.4	8 lines blended
497.404849	U-line	-, -,,,-	4.4	1.5		
497.417697	$(CH_3)_2O$	$12_{9,3}$ - $12_{8,4}$ + $12_{9,4}$ - $12_{8,5}$	9.4	8.4	9.4	8 lines blended
497.442262	$(CH_3)_2O$	$11_{9,2}$ - $11_{8,3}$ + $11_{9,3}$ - $11_{8,4}$	4.2	9.4	8.3	8 lines blended with $HCOOCH_3$
497.442262	HCOOCH ₃	$40_{13,28}$ - $39_{13,27}$ + $40_{15,26}$ - $39_{13,27}$	4.2	9.4	9.0	2 lines blended with $(CH_3)_2O$
497.516476	C_2H_3CN	$60_{1,59}$ - $60_{1,60}$ $v = 0$	4.8	6.1	6.8	
497.556527	U-line	, ,	1.5	9.4		
497.606988	$(CH_3)_2O$	$24_{3,22}$ - $23_{2,21}$	4.3	2.7	8.5	4 lines blended
497.661575	C_2H_5CN	$56_{3,53}$ - $55_{3,52}$	4.9	6.0	3.5	Blend with $CH_3CN v = 8$
497.661575	$CH_3CN v_8=1$	$J=27-26 \text{ K}=4_{-l}$	4.9	6.0	4.9	Blend with C_2H_5CN
497.734975	$CH_3CN v_8=1$	$J=27-26 \text{ K}=3_{-l}$	3.5	4.5	8.8	
497.828685	CH ₃ OH	82-81 -	14.0	9.4	8.7	
497.905738	$CH_3CN v_8=1$	$J=27-26 \text{ K}=4_{+l}$	3.6	6.4	5.9	
498.288253	C_2H_5CN	$57_{2.55}$ - $56_{2.54}$	2.2	6.6	5.1	
498.327509	OCS	J=41-40	12.6	10.2	6.6	
498.412959	$CH_3CN v_8=1$	$J=27-26 \text{ K}=1_{+l}$	6.4	6.8	6.9	
498.431868	$^{34}\mathrm{SO}_2$	84.4-73.5	5.9	12.2	10.0	
498.596075	HCOOCH ₃	417 35-407 34	2.2	8.1	9.2	
498.677131	$^{13}\mathrm{CH}_{3}\mathrm{OH}$	82-81	8.5	7.3	4.8	
498.785157	C ₂ H ₅ CN	554 51-544 50	2.0	9.8	4.5	
498.849325	$(CH_3)_2O$	164 13-153 12	7.7	7.0	8.5	4 lines blended with HCOOCH ₃
498.849325	HCOOCH ₃	425 37 - 415 36	7.7	7.0	6.8	Blend with $(CH_3)_2O$
498.984855	C ₂ H ₅ CN	556 49-546 48	5.4	23.7	12.0	(00)20
499.198900	HCOOCH ₃	435 39-425 38	3.0	6.5	8.1	
499.283980	C ₂ H ₂ CN	$20_{4,17}$ -193 16 $v_1 = 1$	3.4	13.9	7.3	Blend with C ₂ H ₅ CN
499.283980	C ₂ H ₅ CN	178 10/0-167 0/10	3.4	13.9	3.4	Blend with C ₂ H ₃ CN
499.367702	$(CH_3)_2O$	$27_{2,26}$ -26 _{1,25}	3.9	6.1	7.1	4 lines blended
499.892902	C ₂ H ₅ CN	$58_{1/2}$ 571/2 56	9.1	8.0	5.4	i milo stonada
499.917085	NH ₂ CN	252 23-242 22	8.1	8.1	8.2	$^{13}C^{34}S$ J=11–10 ?
499.944145	U-line	= 52,25 = 12,22	10.2	9.6	0.2	0 . 0 11 10 .
500.038140	HC ₃ N	J = 55 - 54	9.0	7.9	6.8	
500.052075	NH ₂ CN	253 22-243 21	8.0	11.8	4.9	
500.075688	$SO_2 v_2 = 1$	74 4-63 3	3.8	11.5	8.6	
500.364813	C ₂ H ₃ CN	41_{6} 35- 41_{5} 36 $v_2 = 1$	4.9	5.7	6.8	
500.432652	SO_2	18_{9} 9-198 12+189 10-198 11	20.1	6.0	8.3	
500.457324	C ₂ H ₅ CN	$12_{0.3/4} - 11_{8.4/3}$	22.3	8.2	2.9	
500.475691	CH_2NH	$2_{2,1}$ - $2_{1,2}$	18.0	7.5	6.3	
500.637890	$C_{2}H_{5}CN$	355 31-344 30	5.9	5.1	5.5	
500.661562	SO_2	$31_{2,30}$ - $31_{1,31}$	9.0	13.3	5.3	
501.087663	C_2H_3CN	$55_{1.55}$ - $54_{1.54}$ $v_1 = 1$	11.7	12.8	10.4	
501.116812	SO_2	$28_{0.28}$ - $27_{1.27}$	21.5	19.1	3.5	
501.592173	CH ₃ OH	92-91 -	32.9	11.7	7.1	
501.656920	C_2H_5CN	$59_{1/0.59}$ - $58_{1/0.58}$	5.6	8.1	8.2	
501.679997	CH_2NH	80,8-70,7	4.4	11.5	9.0	
501.891221	$(CH_3)_2O$	$30_{3,27}$ -29 _{4.26}	2.8	12.0	9.1	4 lines blended
502.181627	HCOOCH ₃	$15_{12,3/4} - 14_{11,4/3}$	4.6	7.0	8.2	
502.303684	N ₂ O	J=20-19	3.4	17.4	4.7	
503.018652	$(CH_3)_2O$	$28_{0,28}$ - $27_{1,27}$	10.4	10.7	7.4	4 lines blended
503.107069	$(CH_3)_2O$	$28_{1,28}$ - $27_{0,27}$	5.1	5.2	9.2	4 lines blended
503.210868	HCOOCH ₃	$40_{8,32}$ -39 _{8,31}	2.7	7.6	8.5	
503.518307	HCOOCH ₃	$22_{9,13}$ - $21_{8,14}$	6.2	6.1	7.6	



Fig. 3. Spectra and lines detected in the survey. The locations of parts of the spectra that were poorly deconvolved are indicated by horizontal lines placed underneath the spectrum.



Fig. 4. Spectra and lines detected in the survey. The locations of parts of the spectra that were poorly deconvolved are indicated by horizontal lines placed underneath the spectrum.



Fig. 5. Spectra and lines detected in the survey. The locations of parts of the spectra that were poorly deconvolved are indicated by horizontal lines placed underneath the spectrum.



Fig. 6. Spectra and lines detected in the survey. The locations of parts of the spectra that were poorly deconvolved are indicated by horizontal lines placed underneath the spectrum.

Frequency	Species	Transition	Peak	Width	Vel	Notes
MHz			$\mathbf{T}_{mb}~\mathbf{K}$	$\rm km~s^{-1}$	$\rm km \ s^{-1}$	
503.574370	C_2H_3CN	$53_{14,39/40} - 52_{14,38/39} v = 0$	9.5	5.6	7.9	Blend with CH ₃ OH
503.574370	CH_3OH	76-85	9.5	5.6	7.5	Blend with C ₂ H ₃ CN
503.707705	HCOOCH ₃	$41_{26,16}$ - $40_{26,15}$	3.3	6.8	8.4	
503.836578	C_2H_5CN	$28_{6,23}$ - $27_{5,22}$	3.4	2.1	6.8	
503.851690	SO	$N_J = 15_{14} - 14_{14}$	8.3	10.9	2.1	Blend with C ₂ H ₃ CN
503.921239	C_2H_3CN	$29_{6,24}$ - $29_{5,25}$ $v_2 = 1$	2.0	7.9	12.5	
504.297599	CH_3OH	71-60	34.3	8.9	6.6	
504.413582	C_2H_3CN	$23_{6,17/18} - 23_{5,18/19} v_2 = 1$	2.4	16.2	7.6	
504.510971	SO_2	$23_{10,14} - 24_{9,15} + 23_{10,13} - 24_{9,16}$	2.4	25.3	9.5	
504.680057	SO	$N_J = 4_3 - 1_2$	9.3	26.7	6.8	
504.728059	^{34}SO	$N_J = 12_{11} - 11_{10}$	20.5	22.4	6.6	Blend with C_2H_5CN
504.728059	C_2H_5CN	$57_{4,54}$ - $56_{4,53}$	20.5	22.4	-0.6	Blend with ³⁴ SO
505.431404	U-line		2.7	7.7		
505.499853	C_2H_5OH	$29_{6,23} + 28_{6,22}$	3.5	7.9	9.3	
505.765740	CH_3OH	102-101	31.4	10.3	6.8	
505.838107	H_2CO	$7_{0,7}$ - $6_{0,6}$	47.9	17.5	6.4	
506.157737	CH_3OH	$11_1 - 10_2$	26.0	8.3	6.3	
506.240528	$HCOOCH_3$	$41_{17,25/24} - 40_{17,24/23} + 41_{17,25/24} - 40_{17,23/24}$	13.7	25.5	6.5	4 lines
506.253692	^{34}SO	$N_J = 12_{12} - 11_{11}$	11.0	16.9	7.6	
506.773898	U-line		6.3	10.0		
506.831598	DCN	J = 7-6	8.7	11.8	6.6	
507.041856	U-line		4.5	5.9		
507.200247	HNCO	$23_{1,22}$ - $22_{1,21}$	9.6	10.4	6.1	
507.301540	^{34}SO	$N_J = 12_{13} - 11_{12}$	24.9	27.8	9.0	
507.337692	SO_2	$36_{6,30}$ - $35_{5,31}$	4.3	6.9	4.8	
507.383231	C_2H_3CN	$27_{3,24} - 26_{2,25} v_1 = 1$	8.7	6.6	6.1	

 Table 6. JCMT Orion spectral linesurvey

in the J = 6-5 transition, and was seen in all of our earlier unpublished JCMT spectra in this transition using both the present SIS receiver, as well as the original JCMT 460 - 490 GHz Indium Antimonide receiver (White & Padman 1991, Padman & White 1992). We have made careful checks on the many occasions that we have observed this line with the JCMT. confirming that this dip is not a result of the subtraction of emission located at the off position. Unpublished maps (in preparation) of the spatial distribution of this absorption feature show it to be spatially localised on the hot core, and that larger scale CO emission is spatially extended around the core meaning that the $T_{\rm mb}$ may overestimate the true kinetic temperature.

4.2. Sulphur Monoxide (SO) and Sulphur Dioxide (SO₂)

The high abundances of sulphur based molecules in the interstellar medium are believed to be in part due to the presence of shocks, that favour the endothermic reactions required to form these molecules. The rotational temperature diagram for SO₂ (see Fig 7) is consistent with previous estimates of the temperature and $N_{\rm col}$. The temperature is 136 K, typical of the cooler conditions found in the plateau. The value of $N_{\rm col}$ is estimated to be 9.7

 10^{16} cm⁻² in the optically thin limit, increasing to $1.2 \ 10^{17}$ $\rm cm^{-2}$ after making a correction for optical depth. Since our initial assumptions might be that this molecule should almost certainly be moderately optically thick, we ran a model, based on the Sutton et al. (1995) temperature and column density estimate, calculating the expected optical depths for the 35 SO_2 lines observed in this survey. One transition has a strong opacity $\tau = 1.46$. The optical depth of all other lines are less than 0.9 and for half of all the detected lines $\tau \leq 0.2$. Therefore the results of the optically thick and optically thin estimates of column density are understandably close to each other. This result is in reasonable agreement with studies using similar beam sizes, by Schilke et al. (2001) $(N_{\rm col} = 9.7 \ 10^{16} \ {\rm cm}^{-2}, T \sim 187$ K for a 12 - 14" beam) and Sutton et al. (1995) ($N_{\rm col} =$ 9.4 10^{16} cm⁻², $T \sim 99$ K for a 12 - 14" beam).

We further estimate that the isotopic ratio $[SO_2]/[{}^{34}SO_2] = 14.1$, which is in agreement with previous estimates by Blake et al. (1987) (14 - 16) and Schloerb et al. 1983 (11).

4.3. Methyl Cyanide (CH₃CN), Ethyl Cyanide (propionitrile) (C₂H₅CN), Vinyl Cyanide (acrylonitrile) C₂H₃CN)

The J = 25-24 and J = 27-26 lines of Methyl Cyanide were observed at 459 and 495 GHz respectively. The average LSR velocity of the lines were ~ 5.6 km s⁻¹and the line widths ~11 km s⁻¹. It is therefore likely that this species traces the hot core. The J = 25-24 k = 7 line (459.276 GHz) could not be assigned due to a strong Uline (459.267 GHz). Lines from the J = 27-26 k = 2 (-1) and k = 3 (+1) transitions were observed at 497.790 and 497.971 GHz respectively, although they were not used in the analysis due to their low intensities. The $N_{\rm col}$ and $T_{\rm rot}$ values for this molecule were 3.5×10^{15} cm⁻² and ~ 230 K - similar to those estimated by Wilner et al. 1994, Sutton et al. 1995, Lee, Cho & Lee (2001). Two lines of the isotopomeric variant CH₃¹³CN were also detected. No lines from the $v_8 = 1$ family were identified.

For this molecule, a) $T_{\rm rot}$ was fixed at 227 K, obtained from analysis of the CH_3CN line, and b) T_{rot} and N_{col} were calculated from the two observed lines. In this latter case, the value of $T_{\rm rot}$ is reduced to about one third that of CH_3CN . We consider that case a) provides a better solution, because of the difficulty of estimating $T_{\rm rot}$ from two lines whose upper state energy levels are very similar to each other. It did not prove possible to identify the J = $25-24 \ k = 0, 1, 3$ lines due to blending with CH₃CN. In view of the problems that have previously been encountered in the interpretation of CH₃CN lines using an optically thin assumption (Schilke et al. 1999, Comito et al. 2003), we ran a model to calculate the opacity of the various lines based on the above excitation conditions estimated by Wilner et al. 1994 (obtained with a similar beam size to our study). The largest value of τ amongst the present lines is estimated to be 0.2, and all other values were ≤ 0.1 . Making a correction for the optical depth only increases the value of column density by a few percent - from $N_{\rm col} = 3.5 \ 10^{15} \ {\rm cm}^{-2}$ to $3.6 \ 10^{16} \ {\rm cm}^{-2}$. We therefore conclude that with few exceptions, the optically thin assumption provides a valid and useful estimate of column density for the transitions observed in this survey.

Thirteen lines of Vinyl Cyanide and its isotopomeric variant C₂H₃CN were detected at LSR velocities consistent with the molecules being concentrated in the compact ridge. Line frequencies have been reported by Demaison etal. (1994). 5 lines were tentatively associated with the v_{15} (out-of-plane bend) = 1 vibrational excited state. However, since the frequencies of the v_{15} lines have a large error, they were not included in the fitting. We also searched at the expected frequency of the $v_{11} = 2$ state, but could not find convincing match. Grain surface reactions are thought to be the main process by which complex nitrogen bearing species are formed, due the high level of hydrogenation. It is not likely that a molecule such as CCCN would pick up the required number of hydrogen atoms to become ethyl cyanide without the intermediate step of adsorption on to a grain, allowing the hydrogenation process to occur. The high temperatures found in the hot core are sufficient to evaporate the molecules from the surface of any grains that drift into the region.

Twenty seven lines of Ethyl Cyanide were detected, based on line frequencies taken from Pearson et al. (1997) and Pearson (2000). It is expected that many observed lines of this molecule are blended with U-lines, and it was found that the simultaneous estimation of the column density and rotational temperature were difficult, since many of the lines are located very close together on the rotational diagram. To avoid this problem, we have calculated the column densities of C₂H₅CN for two fixed temperatures, deriving values of $N_{\rm col} = 2.4 \ 10^{16} \ {\rm cm}^{-2}$ (assuming a temperature of 150K based on the work of Sutton et al. (1985), or $N_{\rm col} = 8.310^{15} \text{ cm}^{-2}$ (assuming a temperature of 239K based on the work of Schilke et al. (2001). As a result, the best estimates of the column density of the C_2H_5CN from the present data are in the range $N_{col} =$ 2.4–0.83 $10^{16}~{\rm cm}^{-2}$. This value is close to that estimated by Schilke et al. (2001) $(N_{\rm col}=~3.1~10^{16}~{\rm cm}^{-2}\,)$ with a beam size of 10-12'' - which is very similar to that of the present survey. By comparison, Sutton et al. (1985) reported $N_{\rm col} = 2 \ 10^{15} \ {\rm cm}^{-2}$ with a 30" beam. Assuming that the emitting region is a core with 10'' diameter, $N_{\rm col} = \, 1.5 \ 10^{16} \ {\rm cm}^{-2} \, ,$ which is in better agreement with the estimate in the present survey value. Similar arguments applied to the Methyl Cyanide line, to resolve the differences between the column densities from this survey, and the work of Blake et al. (1985) and Schilke et al. (2001), again suggest that the size of the emitting region is $\sim 10^{\prime\prime}$.

4.4. Cyanamide (NH_2CN)

Three lines of NH₂CN were detected. However, it did not prove possible to obtain a reliable $T_{\rm rot}$ from the observed data. Consequently we assumed values of $T_{\rm rot}$ of 100 and 200K, to make a first estimate of the column densities.

4.5. Carbonyl sulphide (OCS)

Two lines of OCS were detected. The fit for $T_{\rm rot}$ and $N_{\rm col}$ are not very accurate, since the upper state energy levels of the two lines are very similar.

4.6. Methylenimeme (CH_2NH)

Methylenimine is a prolate asymmetric rotor with components of the electric dipole moment along both the a and bmolecular axes, with magnitudes 1.325 and 1.53 D, respectively. The nitrogen nucleus produces electric quadrupole hyperfine structure in low-lying transitions. CH₂NH is a likely product following the UV irradiation of icy interstellar grain mantles and may be a precursor of other complex organics which may be present in cometary ices (Bernstein et al. 1995), and may be a precursor to glycinenitrile and glycine (e.g., Dickerson 1978). Three lines of CH₂NH were used in the fitting. It was considered likely that the $2_{2,1}$ – $2_{1,2}$ line was probably blended with a U-line, since the intensity was strong compared with the other two lines.

Using only two lines, it did not prove possible to get a reasonable value for $T_{\rm rot}$. Hence, in fitting the value of $T_{\rm rot}$ was fixed at 150 K, from observations of HNCO, which has a similar dipole moment (CH₂NH $\mu({\rm a})=1.352$ D, $\mu({\rm b})=1.530$ D, HNCO $\mu({\rm a})=1.602$ D, $\mu({\rm b})=1.35$ D - from Kirchoff, Johnson & Lovas 1973, Hocking et al. 1975). The derived column density, $N_{\rm col}=2.4\ 10^{15}\ \pm 1.8\ 10^{14}\ {\rm cm}^{-2}$, is in reasonable agreement with the column density reported in the detection paper by Dickens et al. (1977) of $N_{\rm col}=6.2\pm 1.8\ 10^{14}\ {\rm cm}^{-2}$, based on observations of 5 transitions (three of which were blended with other lines) between 172 and 256 GHz obtained with a larger beam of 23 - 34'' .

4.7. Methanol (CH₃OH) and Ethanol (C_2H_5OH)

Methanol, CH₃OH, is one of the most widely observed molecules in star forming regions. Line frequencies are reported by Xu & Lovas (1997). Twenty four CH₃OH lines were detected with an average velocity of 7.8 km s⁻¹ and line width of 9.9 km s⁻¹- suggesting that it is likely to be excited in the compact ridge. The high observed abundances of this molecule imply a high abundance of the precursor ion CH_3^+ which would react with water in a ionmolecule process to form methanol. Two lines of ¹³CH₃OH (which were apparently detected at 492.695 and 498.677 GHz) would indicate that the main methanol line may have an opacity of at least 0.7. Schilke et al. (2001) have pointed out the difficulty of applying simple rotational analysis techniques to methanol due to opacity problems. The column densities reported in Table 1 have been should therefore be treated as lower limits. If the 498.677 GHz line is assigned to ${}^{13}CH_3OH-A(8_2-8_1)$, the column density of this molecule would be $8.7 \ 10^{15} \ \mathrm{cm}^{-2}$ assuming the rotational temperature is the same as that of ${}^{12}CH_3OH$ (600 K - but note that the error on this is 50%). This would indicate that the $[^{12}C]/[^{13}C]$ ratio ~ 10 - although this would also decrease if the temperature were substantially less than 600K). Neither the ${}^{13}CH_3OH-E(23_2-23_1)$ line at 503.216 GHz, nor any lines of torsionally excited methanol were identified.

Eight lines of C₂H₅OH were detected, at a velocity of ~ 8 km s⁻¹, although only 6 were used in the rotational temperature fitting. Line frequencies are given by Pearson etal. (1996). The lines at 494.899 and 495.821 had line widths that were too broad and narrow respectively, and it is likely that these are blended, or due to other species. For this molecule, we made two fits; a) simultaneously fitting both $N_{\rm col}$ and $T_{\rm rot}$, and b) $T_{\rm rot}$ was fixed at 70 K (following Ohishi et al. 1995), since simultaneous fitting of both $T_{\rm rot}$ and $N_{\rm col}$ led to large uncertainties, and a large $T_{\rm rot}$.

4.8. Methyl formate (HCOOCH₃)

The line widths and velocities of the HCOOCH₃ lines suggest an origin in the compact ridge. Line frequencies are given by Osterling etal. (1999). Lines are seen from both the E- and the A-symmetry states. The high abundance follows directly from the high abundance of methanol whose precursor ion $CH_3OH_2^+$ reacts with H₂CO. The column density we derive is similar to that estimated by Schilke et al. (1997), who have discussed difficulties in assigning an unique rotation temperature for this line.

4.9. Dimethyl ether $((CH_3)_2O)$

Dimethyl Ether, $(CH_3)_2O$, is one of the few interstellar molecules whose emission lines are affected by the presence of two internal rotors. Line frequencies are reported by Groner etal. (1998). Twenty seven lines were detected in this survey, which were best described by a fit of $N_{\rm col} =$ 1.4 10^{16} cm⁻² and $T_{\rm rot} = 157$ K for the column density and rotational temperature respectively. These values are close to those found by Sutton et al (1985) toward the compact ridge. The line widths and velocities are consistent with an origin in the compact ridge. The fact that this molecule appears to be far more abundant (from the large number of strong lines) than ethanol leads to the conclusion that the Compact ridge, where these molecules are formed, is rich in hydrogen. The molecule is formed by a similar method to methyl formate except that $CH_3OH_2^+$ reacts with methanol to form it.

4.10. Formaldehyde (H_2CO)

 H_2CO is a highly prolate asymmetric top molecule. The rotational temperature of H_2CO was fixed at 166 K following Blake et al. (1987). Two lines were associated with this molecule, however an additional line at 491.979 GHz was not included in the fitting, since its velocity appeared to be low (2.6 km s⁻¹). The derived column density is very similar to that estimated by Blake et al. (1987).

4.11. Deuterated water (HDO)

A single transition of HDO was detected at 464.925 GHz. It has a clear hot core line shape with a line width of 16.2 km s⁻¹ and a velocity of 6.6 km s⁻¹. We have attempted to model this line along with the two other lines reported and analysed by Schulz et al. (1991) and Sutton et al. (1985). Pardo et al. (2001b) have recently reported detections of the $J_{\rm Ka,Kb} = 2_{1,2} - 1_{1,1}$ and $1_{1,1} - 0_{0,0}$ lines in the 850 - 900 GHz region, which appear trace the plateau gas, rather than the hot core material which is contributes rather more to the other HDO lines observed to date. We therefore estimate values of 3.2×10^{16} cm⁻² and 164 K for $N_{\rm col}$ and $T_{\rm rot}$ respectively, in broad agreement with values reported by Plambeck & Wright (1987) and Jacq et al. (1990). This is as expected from the high abundances observed for molecules such as methanol and methyl formate

13.5

130

which require water to be present for their formation. It is thought that both water and deuterated water are formed elsewhere on grains and evaporated from the surfaces in the higher temperature conditions found in the hot core (Beckman et al. 1981, Pardo et al. 2001b).

4.12. Atomic Carbon (CI)

A lower limit on the column density of the CI ${}^{3}P_{1} - {}^{3}P_{0}$ transition was determined with the treatment of the optically thin line emission given by White & Sandell (1995). The column density, $N_{\rm col} = 1.2 \ 10^{18} \ {\rm cm}^{-2}$ is consistent with that of White & Sandell (1995). Keene et al. (1998) detected the 13 CI line toward a position 4' south of our pointing position, showing that the main isotopomeric line was optically thin. To date, this remains the best indicator that it is useful to use the optically thin column density estimate.

4.13. Other Molecules

Other detections that are of interest include HCCCN and HNCO (Isocyanic Acid), both with line shapes are typical of the hot core, and with estimated abundance similar to those reported by Schilke et al. (1997).

4.14. Unidentified Lines

At total of 33 lines were detected that could not be associated with known spectral lines. We searched carefully in the Cologne Database for Molecular Spectroscopy (Müller et al. 2001) and the JPL Molecular Spectroscopy Database, as well as other tables of isotopomeric variants of lines that were present in the survey (e.g. CH_3OH and ¹³CH₃OH from the laboratory measurements of Anderson, Herbst & de Lucia 1987, 1990, 1992). Although there were inevitably a number of lines that lay close to some of the U-line frequencies, we also used secondary criteria (line strength predictions from the Cologne database, upper energy state levels, and the presence or absence of other lines from similar levels) to make judgments as to whether lines could be associated with particular species. Six lines that were originally designated as U-lines were associated in this way, however a substantial number of intense lines remain. Assignment of lines to these frequencies is beyond the scope of this paper, and will require a sophisticated modeling effort, combined with U-lines from other published surveys. We note that the number of U-lines inferred from the present survey (33 - or 13% of the total) is similar reported from some other surveys (325 - 360 GHz $\sim 8\%$ Sutton et al. 1995), 607 - 725 GHz $\sim 14\%$ (Comito et al. 2003).

4.15. Notes on $T_{\rm rot}$ diagrams

Plots used to fit the estimates of $T_{\rm rot}$ or $N_{\rm col}$ are shown in Fig. 7.

12.0 12.4 11.5 12.2 11.0 12.0 600 200 200 400 800 100 300 400 89 K N_{col} = 1.1 10¹⁶ cm⁻² $136 \text{ K N}_{col} = 1.2 \ 10^{17} \text{ cm}^{-2}$ 15.0 15.0 14.5 14.5 14. 13. 14.0 13.0 12.5 13.5 12.0 11.5 13.0 50 100 150 400 600 200 0 200 800 1000 Fig. 7. ¡Rotational temperature diagrams. The fitted lines

were derived from a noise-weighted least squares fitting procedure.

The values for the intrinsic line strengths, level degeneracies and partition functions are derived from the literature, including the spectral line catalogues cited previously (see also Eq 1 and 2).

4.16. Line-to-continuum ratio

The data may also be used to estimate the ratio of lineto-continuum emission from the source. This is an important value to characterise for a range of molecular clouds - since a considerable fraction of the fluxes measured with



submm continuum cameras, such as the JCMT's SCUBA, and the CSO's SHARC, may be due to the integrated line emission, rather than thermal dust. Early attempts to measure, or characterise this ratio toward Orion have included the work of Groesbeck (1995), Greaves & White (1991), White & Sandell (1995), Schilke et al. 1997, 2001). The integrated line emission of the spectrum shown in Fig 1 is $\sim 1.1 \times 10^5$ K MHz. The *average* rms noise level over the whole survey range was 1.5 K per 2 MHz channel, and therefore line emission would not be detected under this level. If lines of this strength were uniformly spread across the 30 GHz spectral range, then this would contribute a further $\sim 4.5 \times 10^4$ K MHz to the integrated line emission. The total estimated integrated emission corresponds to an integrated main beam temperature of 3.7 K MHz^{-1} . Greaves & White (1991) reported an integrated emission of $\sim 1.0 \times 10^4$ K MHz over a total range of 16 GHz, which corresponded to an average temperature of 0.6 K MHz^{-1} - or about one sixth of that estimated here. This discrepancy can partially be explained by the fact that the hot core is smaller in angular extent that the beam size used in both this and the survey of Greaves & White (1991). The coupling efficiency η_c to a Gaussian source of radius σ_s is given by Eq. 3:

$$\eta_c = \frac{\sigma_s^2}{(\sigma_s^2 + \sigma^2)} \tag{3}$$

where σ is the Gaussian radius of the telescope beam. Taking $\sigma_s = 4''$ and $\sigma = 4.2''$ (for a 10" FWHM beam), $\eta_c = 0.48$, and $\eta_c = 0.25$ for the survey of Greaves & White (1991) would lead to an expectation that the current value should be double that previously estimated for this reason. The lower beam dilution in this survey could explain the rest of the discrepancy. No measurements of the continuum emission exist for the exact frequency that this survey was conducted at, but by using the measurements made by White & Sandell (1995) at 790 μ m it is possible to extrapolate and make an estimate of the emission at 600 μ m (500 GHz). The spectral index was estimated by White & Sandell (1995) to be 1.5 and the emission was 134.9 Jy in a 10" beam with an aperture efficiency of 0.3. The antenna temperature (T_A) is given by Eq. 4:

$$kT_A = \frac{1}{2}\eta_R\eta_A S_\nu A \tag{4}$$

where k is the Boltzmann constant, η_A is the aperture efficiency (0.3 for these observations), η_R is the resolution correction and A is the effective area of the antenna. This equation yields a value $T_A = 6.8$ K. Correcting this value for sideband gains, atmospheric emission and main beam efficiency gives $T_{\rm mb} = 14.4$ K. This means that the line to continuum ratio at 500 GHz is $3.7 \div 14.4 = 0.25$. This lies between the value of 10% found by Greaves & White (1991) and 30-40% found by Sutton et al. (1985) at lower frequencies. We stress that this value is the best estimate lower limit to the line-to-continuum ratio that can be made with the present data - a further measurement that simultaneously measures the continuum and line flux is desirable to obtain a more accurate value - and would for example be valuable when trying to understand the relative contributions of thermal and line emission observed at far infrared wavelengths by bolometer detectors, and far-IR fourier transform spectrometers. The value we have estimated is consistent with the prediction given by Groesbeck (1995 - also reported in Schilke et al. 2001) that the contribution of line emission to the total flux of Orion should drop from $\sim 50\%$ at 800 μm to 10% at 450 μm .

5. Conclusions

A spectral line survey of the hot core region of the OMC-1 cloud core was obtained over the frequency intervals 455 - 468 GHz and from 491.8 - 507.4 GHz.

- In this spectral line survey we find a total of 254 lines toward the hot core position at the centre of the Orion Nebula.
- Spectral lines of SO, SO₂ and CH₃OH and from large organic molecules such as (CH₃)₂O, CH₃CN, C₂H₅CN, HCOOCH₃ and C₂H₃CN make up ~ 72% of the total number of lines
- A total of 33 lines (13% of the total lines identified) could not be associated with known molecular transitions and are designated U-lines - this is comparable to the percentages of U-lines seen in other spectral lines surveys of Orion.
- The rotational temperatures and column densities derived using standard rotation diagram analysis techniques, showed ranges from 70 300 K, and $10^{14} 10^{17}$ cm² respectively (we exclude the CH₃OH line from this, as the error is large).
- The diameter of the region responsible for emitting the C_2H_5CN and C_2H_3CN lines is estimated to be $\sim 10''$, based on a comparison with earlier data for the same lines.
- The lower limit to the line-to-continuum ratio from the hot core region is 0.25, which appears to be consistent with predictions based on modeling the expected intensities of lines from the JPL Molecular Spectroscopy database.

6. Acknowledgments

We acknowledge discussions with Prof S. Saito at Fukui University, Dr Y. Fukuyama at the Institute of Physical and Chemical Research, Prof John Pearson, for discussions about frequency assignments of molecules, and the referee for helpful comments.

References

Anderson, T., Herbst, E., & de Lucia, F. 1987, ApJS, 64, 703
Anderson, T., Herbst, E., & de Lucia, F. 1990, ApJS, 74, 647
Anderson, T., Herbst, E., & de Lucia, F. 1992, ApJS, 82, 405
Beckman, J.E., Watt, G.D., White., G.J., et al. 1981, MNRAS, 201, 357.

Blake, G. A., Masson, C. R., Phillips, T. G., et al. 1986, ApJS, 60, 357

- Bernstein, M. P., Sandford, S. A., Allamandola, L. J., Chang, S., & Scharberg, M. A. 1995, ApJ, 454, 327
- Comito, C., Schilke, P., Lis, D. C., Motte, F., Phillips, T. G., & Mehringer, D. M. 2002, SFChem 2002: Chemistry as a Diagnostic of Star Formation, proceedings of a conference held August 21-23, 2002 at University of Waterloo, Waterloo, Ontario, Canada N2L 3G1. Edited by Charles L. Curry and Michel Fich. To be published by NRC Press, Ottawa, Canada, p. 18, 18
- Dickens, J. E., Irvine, W. M., De Vries, C. H., Ohishi, M. 1997, ApJ, 479, 307
- Demaison, J., Cosleou, J., Bocquet, R., & Lesarri, A.G., 1994, J. Mol. Spect., 167, 400
- Blake, G., Sutton, E., Masson, C., et al. 1987, ApJ, 315, 621 (Bla87)
- Goldsmith, P.F., Langer, W.D. 1999, ApJ, 517, 209-225
- Greaves J.S., 1990, PhD Thesis, University of London
- Greaves, J.S., White G.J. 1991, A&AS, 91, 237
- Greoesbeck. T.D., 1995, PhD Thesis, Caltech
- Groner, P., Albert, S., Herbst, E., & De Lucia, Frank C., 1998, ApJ, 500, 1059.
- Hasegawa, T., 1986, Ap&SS., 118, 421
- Hermsen, W.; Wilson, T. L.; Walmsley, C. M.; Henkel, C., 1988, A&A, 201, 285 (Her88)
- Hocking, W. H., Gerry, M. C. L. and Winnewisser, G. 1975, Can. J. Phys., 53, 1869
- Jacq, T., Walmsley, C.M., Mauersberger, R. et al. 1990, A&A, 228, 447
- Jewell, P. R., Hollis, J. M., Lovas, F., et al. 1989, ApJS, 70, 833
- Johansson, L. E. B., Andersson, C., Ellder, J., et al. 1984, A&A, 130, 227
- Keene, Jocelyn; Schilke, Peter; Kooi, J.; Lis, D. C.; Mehringer, David M.; Phillips, T. G. 1998, ApJ, 494. L107
- Kirchhoff, W. H., Johnson, D. R., Lovas, F. J. 1973, J. Phys. Chem. Ref. Data, 2, 1
- Lee, C.W., Cho, S-H., Lee, S-M. 2001, ApJ, 551, 333
- Lepp, S., Delgarno, A. 1988, ApJ, 324, 553
- Müller, H.S.P., Thorwirth, S., Rothe, D.A., et al. 2001, A&A Lett., 370, L49 (and a searchable internet form at the URL: http://www.ph1.uni-koeln.de/vorhersagen/)
- Naylor, D. A., Davis, G. R., Gom, B. G., et al. 2000, MNRAS, 315, 622
- Nummelin, A., Bergman, P., Hjalmarson, A., et al. 2000, ApJL, 128, 213
- Ohishi, M., Ishikawa, S., Yamamoto, S., et al. 1995, ApJL, 446, L43 (Ohi95)
- Oesterling, L. C., Albert, S., De Lucia, Frank C., Sastry, K. V. L. N., Herbst, E., 1999, ApJ, 521, 255
- Padman, R., White, G.J., et al. 1992, Int. Journ. IR and Mm Waves, 13, 10
- Pardo J.R., Serabyn, E., Cernicharo. 2001, JQSRT, 68, 419
- Pardo, J. R., Cernicharo, J, Herpin, F. et al. 2001, ApJ, 562, 799
- Pearson, J.C., Sastry, K.V.L.N., Herbst, E., De Lucia, F.J. 1996, J. Mol. Spect., 175, 246-261
- Pearson, J.C., Sastry, K.V.L.N., Herbst, E., De Lucia, F.J. 1997, ApJ, 480, 420
- Pearson, J. C., 2000. Private communication
- Pickett, H.M., Poyner R.L., Cohen E.A., 1995, Submillimeter, Millimeter, and Microwave Spectral line Catalogue, on the World Wide Web (http://spec.jpl.nasa.gov)

Plambeck, R.L. and Wright, M.C.H. 1987, ApJ, 317, L101

- Prased, S.S., Huntress, W.T. 1982, ApJ, 260,590
- Schilke, P., Groesbeck, T. D., Blake, G. A., Phillips, T. G. 1997, ApJS, 108, 301 (Sch97)
- Schilke, P., Phillips, T. G., Mehringer, D. M. 1999, in The Physics and Chemistry of Interstellar Medium, Proc. 3rd Cologne-Zermatt Symposium, ed. V. Ossenkopf, J. Stutzki, & Winnewisser, p 330
- Schilke, P., Benford, D.J., Hunter, T.R., Lis, D.C. Phillips, T. G. 2001, ApJS, 132, 281 (Sch01)
- Schloerb, F. P., Irvine, W. M., Friberg, P., Hjalmarson, A., Hoglund, B. 1983, ApJ, 264, 161
- Schulz, A., Güsten, R., Walmsley, C. M., Serabyn, E. 1991, A&A, 246, L55.
- Serabyn, E., Weisstein, E. W. 1995, ApJ, 451, 238
- Sutton, C., Blake, G., Masson, C., Phillips, T.G. 1985, ApJS, 58, 341 (Sut85)
- Sutton, E.C., Peng, R., Danchi, W.C., et al. 1995, ApJS, 97, 455 (Sut95)
- Turner, B.E. 1991, A&AS, 76, 617
- Turner, B.E. 1989, ApJS, 70, 539
- White, G. J., Griffin, M. J., Rainey, R. et al. 1986, A&A, 162, 253
- White G.J., Sandell G. 1995 A&A, 299, 179 (Whi95)
- White G.J., Padman, R. 1991 Nat, 354, 511
- Wilner, D.J., Wright, M.C.H., Plambeck, R.L. 1994, ApJ, 422, 642
- Wright, M. C. H., Plambeck, R. L., Vogel, S. N. et al. 1983, ApJ, 267, L41 (Wri83)
- Zeng, Q., Pei, C. C. 1995, Ap&SS. 229, 301 (Zen95)
- Ziurys, L. M., McGonagle, D. 1993, ApJS, 89, 155
- Xu, L.-H., & Lovas, F. J., 1997, J. Phys. Chem. Ref. Date, 26, 17.