THE X-RAY BINARY GRS 1741.9–2853 IN OUTBURST AND QUIESCENCE

M. P. MUNO, F. K. BAGANOFF, AND J. S. ARABADJIS

Center for Space Research, Massachusetts Institute of Technology, Cambridge, MA 02139;

muno@space.mit.edu Submitted to ApJ Letters

ABSTRACT

We report *Chandra* and *XMM-Newton* observations of the transient neutron star low-mass X-ray binary GRS 1741.9–2853. Chandra detected the source in outburst on 2000 October 26 at an X-ray luminosity of ~ 10^{36} erg s⁻¹ (2–8 keV; 8 kpc), and in quiescence on 2001 July 18 at ~ 10^{32} erg s^{-1} . The latter observation is the first detection of GRS 1741.9–2853 in quiescence. We obtain an accurate position for the source of 17h 45m 2.33s, $-28^{\circ}54'49.7''$ (J2000), with an uncertainty of 0.7''. GRS 1741.9–2853 was not detected significantly in three other *Chandra* observations, nor in three XMM-Newton observations, indicating that the luminosity of the source in quiescence varies by at least a factor of 5 between $(< 0.9 - 5.0) \times 10^{32}$ erg s⁻¹ (2–8 keV). A weak X-ray burst with a peak luminosity of 5×10^{36} erg s⁻¹ above the persistent level was observed with *Chandra* during the outburst on 2000 October 26. The energy of this burst, 10^{38} erg, is unexpectedly low, and may suggest that the accreted material is confined to the polar caps of the neutron star. A search of the literature reveals that GRS 1741.9-2853was observed in outburst with ASCA in Fall 1996 as well, when the BeppoSAX WFC detected the three previous X-ray bursts from this source. The lack of X-ray bursts from GRS 1741.9–2853 at other epochs suggests that it produces bursts only during transient outbursts when the accretion rate onto the surface of the neutron star is about 10^{-10} M_{\odot} y⁻¹. A similar situation may hold for other low-luminosity bursters recently identified from WFC data.

Subject headings: stars: individual: GRS 1741.9–2853— stars: neutron — X-rays: bursts

1. INTRODUCTION

In recent years, the *BeppoSAX* Wide-Field Camera (WFC) and the Rossi X-ray Timing Explorer All-Sky Monitor (RXTE ASM) have identified two classes of faint low-mass X-ray binary (LMXB) that may be closely related: low-luminosity transients and low-luminosity bursters. The low-luminosity transients consist of a rather inhomogeneous group of about 15 LMXBs with outbursts that last from a few days to several months, and get no brighter than a few times 10^{36} erg s⁻¹, which corresponds to an accretion rate of 10^{-10} M_{\odot} y⁻¹ (e.g., in 't Zand 2001). This distinguishes the low-luminosity transients from more easily detected transient LMXBs such as Aql X-1 that usually exhibit outbursts brighter than 10^{37} erg s⁻¹, and from which faint outbursts are less common (e.g., Simon 2002). The faintness of these outbursts has been attributed to average mass transfer rates of $\dot{M} \leq 10^{-11}$ M_{\odot} y⁻¹ (in 't Zand 2001; King 2000). The low-luminosity transients have attracted particular attention because they include the four known accreting millisecond X-ray pulsars (Wijnands & van der Klis 1998; Markwardt et al. 2002; Galloway et al. 2002; Markwardt et al. 2003). For this reason, it has been hypothesized that the low average accretion rates allow relatively strong (> 10^8 Gauss) magnetic fields to persist on the surfaces of the neutron stars among these LMXBs, whereas the surface field is buried in systems with higher accretion rates (Cumming, Zweibel, & Bildsten 2001).

The low-luminosity X-ray bursters are sources from which bright thermonuclear X-ray bursts (see Lewin, van Paradijs, & Taam 1993 for a review) have been observed with the BeppoSAX WFC, and yet there was no evidence

for X-ray emission from persistent accretion at the time of the burst (Cocchi et al. 2001; Cornelisse et al. 2002a,b). This sample of low-luminosity bursters may represent a large population of undiscovered neutron star X-ray binaries, depending upon how often these systems produce bursts. The intervals between X-ray bursts is a strong function of the accretion rate per unit area onto the neutron star, which determines how quickly a sufficient column of material is collected for helium burning to become unstable (e.g., Bildsten 2000). Since the nuclear energy in accreted material is at most 4% of the gravitational energy that it emitted during accretion (Lewin et al. 1993), sufficient nuclear energy to produce an easilydetectable 10^{39} erg burst is collected in 5 hours on the surface of a neutron star producing 10^{35} erg s⁻¹ of X-rays through accretion (corresponding to an accretion rate of 10^{-11} M_{\odot} y⁻¹), while such a burst could not occur for several years on a star emitting persistently at 10^{32} erg s^{-1} (10⁻¹⁴ M_☉ y⁻¹). Unfortunately, the *BeppoSAX* WFC could only place an upper limit of 10^{36} erg s⁻¹ on the luminosity of persistently faint bursters close to the Galactic center, so estimates of the frequency of bursts from the low-luminosity bursters are uncertain by several orders of magnitude. Therefore, the number of low-luminosity bursters can only be guessed within a factor of 1000, and it is still possible that many of them are also faint X-ray transients.

GRS 1741.9–2853 is both a faint transient system and a low-luminosity burster. It was discovered as a transient X-ray source located 10' from the Galactic Center with *Granat*, during two observations on 1990 March 24 and April 8 (Pavlinsky, Grebenev, & Sunyaev 1994). The flux during these observations remained approximately con-



FIG. 1.— Image in the 0.5-8.0 keV band of the ACIS-I field containing GRS 1741.9–2853 in outburst on 2000 October 26. GRS 1741.9–2853 lies in the upper right. The bright region at the center is the Sgr A complex.

stant at $2 \times 10^{-10} \text{ erg cm}^{-2} \text{ s}^{-1}$ (4–30 keV), but by the Fall of 1990 the source had decreased in flux by at least a factor of seven, making it undetectable with *Granat*. In the Fall of 1996, the *BeppoSAX* Wide-Field Camera (WFC) detected three thermonuclear X-ray bursts from the source, identifying the compact object in GRS 1741.9-2853 as a weakly-magnetized neutron star (Cocchi et al. 1999). The brightest of these bursts had a peak flux of 4×10^{-8} erg $cm^{-2} s^{-1}$, and exhibited a "radius expansion" episode in which the photosphere of the neutron star appeared to be driven from its surface by radiation pressure. Assuming that the peak intensity of this burst represents the Eddington limit for a neutron star with standard parameters $(\approx 2 \times 10^{38} \text{ erg s}^{-1}; \text{ see Lewin et al. 1993}), \text{ the system lies}$ near the Galactic Center, at a distance of approximately 8 kpc (since GRS 1741.9-2853 also is only 10' away from the Galactic center, we adopt this as its distance throughout the paper). At this distance, the luminosities of the transient outbursts were only $\approx 10^{36}$ erg s⁻¹. Repeated observations of the Galactic center by the BeppoSAX WFC each Spring and Fall from mid-1996 through 2001 did not detect any outbursts from GRS 1741.9-2853 above a sensitivity limit corresponding to 3×10^{36} erg s⁻¹ (2–28 keV) (Cornelisse et al. 2002a), and continuous monitoring with the RXTE All-Sky Monitor rules out outbursts brighter than $6 \times 10^{36} \text{ erg s}^{-1}$ (2–12 keV) lasting more than a month from 1996 until the present. However, the source was detected twice as part of the ASCA survey of the Galactic Plane, with luminosities between 10^{34} and 10^{36} erg s⁻¹ (1-10 keV) (see Table 1 and Sakano et al. 2002).

In this paper, we report several serendipitous *Chandra* and *XMM-Newton* observations of GRS 1741.9–2853. *Chandra* detected the source in outburst, which allowed

us to determine its location precisely (Muno et al. 2003). Using this location, we study the flux from this source in quiescence. We also report the detection of a thermonuclear X-ray burst while the source was in outburst.

2. OBSERVATIONS AND DATA ANALYSIS

$2.1. \ Chandra$

To date, the location of GRS 1741.9–2853 has been observed on five occasions with *Chandra*: three times as part of a program to monitor the super-massive black hole at the center of the Galaxy, Sgr A* (Baganoff et al. 2001, 2003; Muno et al. 2003), and twice as part of a shallow survey of the inner 300 pc of the Galaxy (Wang et al. 2002). The observations are listed in Table 2.

All five observations were taken using the Advanced CCD Imaging Spectrometer imaging array (ACIS-I). The ACIS-I is a set of four 1024-by-1024 pixel CCDs, covering a field of view of 17' by 17'. When placed on-axis at the focal plane of the grazing-incidence X-ray mirrors, the imaging resolution is determined by the pixel size of the CCDs, 0''.492. The CCDs measure the energy of incident photons with a resolution of 50-300 eV (depending on photon energy and distance from the read-out node) within a calibrated energy band of 0.5–8 keV.

We reduced the data starting with the level 1 event files provided by the *Chandra* X-ray Center (CXC). We first removed the pixel randomization applied by the default processing software, and corrected the pulse heights of each event for the position-dependent charge-transfer inefficiency caused by radiation damage early in the mission (Townsley et al. 2000). We excluded most events flagged as possible non-X-ray background, but left in possible cosmic ray after-glows because they are difficult to distinguish from the strong diffuse emission and numerous point sources near the Galactic center. We applied the standard ASCA grade filters to the events, as well as the good-time filters supplied by the CXC. We examined each observation for background flares, and removed 10 ks of strong flaring from ObsID 0242 when the count rate rose to 3 standard deviations above the mean. Finally, we corrected the pointing aspect for each observation as specified in the *Chandra* data caveats page.¹

To search for point sources in each observation, we employed the CIAO tool wavdetect (Freeman et al. 2002) in the manner described in Muno et al. (2003). We set the detection threshold to 10^{-7} , which corresponds to the probability that a background fluctuation in any given pixel will be identified as a source. The resulting source list for the Sgr A^{*} monitoring campaign, which covers ObsIDs 0242, 1561a, and 1561b, is presented in Muno et al. (2003). The source lists for ObsIDs 2284 and 2287 will be reported elsewhere. These lists were used to identify candidate counterparts to GRS 1741.9-2853 and to exclude point sources from the estimate of the background flux. We verified that the astrometry for each observation was accurate to within $0^{\prime\prime}_{...5}$ by cross-correlating the positions of the half-dozen sources that were both detected in the soft X-ray band (0.5–1.5 keV) and present in the USNO-B catalog.²

GRS 1741.9-2853 was evident by eye as a bright transient in the image from ObsID 1561a (2000 October 26; Figure 1). Since the source is located 10' from the aimpoint of the observation, its location is uncertain by about 5'' (see also Muno et al. 2003). Its position is within the 1' error circles of previous localizations of GRS 1741.9-2853 with GRANAT and ASCA. The transient source exhibited a thermonuclear burst during this Chandra observation (see below), so we can confidently associate it with GRS 1741.9–2853. We searched the source lists from the other observations, and identified a faint point source within our initial localization of GRS 1741.9-2853 at 6' from the aim point of ObsID 2287. Its position was 17h 45m 2.33s, -28°54′49.7″ (J2000), with an uncertainty of 0.7'' (compare Brandt et al. 2001). No point source was identified by wavdetect within 5" of GRS 1741.9-2853 in the other three observations.

established Having \mathbf{a} precise position for GRS 1741.9-2853, we estimated its flux in each of the five observations using the acis_extract routine from the Tools for X-ray Analysis (Broos et al. 2002). We extracted event lists for the source from a polygonal region chosen to match the contour of 90% encircled energy from the point-spread function (PSF) at an energy of 4.5 keV. Background event lists were extracted from circular regions centered on GRS 1741.9-2853, excluding from the event lists counts in circles circumscribing the 95% contours of the PSF around the point sources detected in each field. The number of total, background, and net counts from each observation are listed in Table 3. We computed



FIG. 2.— ACIS-I spectrum of GRS 1741.9–2853 while in outburst on 2000 October 26. The *top panel* displays the number of counts received as a function of energy, which depends both on the incident spectrum and the detector effective area. The spectrum is modeled with an absorbed power law continuum (solid line). The *bottom panel* displays the residuals after applying the model, normalized to the 1σ uncertainty on the counts in each spectral bin.

90% uncertainties and upper limits on the net number of counts using the expected Poisson distribution of the total number of counts given the background derived from the photometry (Kraft, Burrows, & Nousek 1991).

We extracted a spectrum for ObsID 1561a (2000 October 26), which contained 170,214 counts from 35,705 s of live time in an extraction region of 1563 pixels. We used response matrices appropriate for data with the chargetransfer inefficiency corrected, and an effective area function (ARF) corrected to account for the fraction of the PSF enclosed by the extraction region and for the hydrocarbon build-up on the detectors.³ The burst contributed only 250 counts to the spectrum, so we did not remove it from the persistent spectrum. Although a count rate this high (5 count s^{-1}) would generally cause serious pile-up if the source were observed on-axis with Chandra, the 10'offset angle spread the counts over a large enough area on the ACIS-I CCDs that pile-up is not an issue. The mean count rate per pixel is only 0.003 count s⁻¹ pixel⁻¹, which produces a pile-up fraction of 0.3%. The brightest pixel only contained 803 counts, for a rate of 0.02 count s^{-1} pixel⁻¹ and a pile-up fraction of 2%.

We fit the spectrum between 2–7 keV, because outside this energy range there were systematic residuals. These residuals may have been caused by the high count rate incident on a relatively small area of the CCD, which may have changed the efficiency of charge-transfer as the events were read out. We modeled the spectrum with a power-law continuum absorbed and scattered by interstellar gas and dust.⁴ We linked the optical depth of dust to the absorption using the relation $\tau = 0.485 \cdot N_{\rm H}/(10^{22} {\rm cm}^{-2})$ (Baganoff et al. 2001), and fixed the halo size to 100 times that of the PSF. The fit had a $\chi^2/\nu = 403/339$, and is displayed in Figure 2. The best-fit photon index was $\Gamma = 1.88 \pm 0.04$ and the column density $N_{\rm H} = (9.7 \pm 0.2) \times 10^{22}$

 $^{^1}$ http://cxc.harvard.edu/cal/ASPECT/fix_offset/fix_offset.cgi

² D. Monet et al.; see http://www.nofs.navy.mil/projects/pmm/catalogs.html

³ http://cxc.harvard.edu/cal/Acis/Cal_prods/qeDeg/

 $^{^4}$ We modified the dust model from the default version in XSPEC so that it did not assume that the dust was optically thin.

20

FIG. 3.— Light-curve of GRS 1741.9–2853 in outburst in the 0.5-8.0 keV energy band. An X-ray burst lasting about 100 s is observed 25,700 s into the observation. A typical error bar is displayed in the lower left.

cm⁻². We extrapolated the fit to compute an "observed" X-ray flux of $F_{\rm X} = 2 \times 10^{-10}$ erg cm⁻² s⁻¹between 2–8 keV, where we chose the energy range to be consistent with the catalog in Muno et al. (2003).⁵ The deabsorbed X-ray flux between 2–8 keV was 4×10^{-10} erg cm⁻² s⁻¹, which corresponds to a luminosity of 3×10^{36} erg s⁻¹ at 8 kpc.

For the remaining four observations, we converted the count rates and upper limits into fluxes using PIMMS, and list the results in Table 3. The source was detected significantly $(> 7\sigma)$ in ObsID 2287 on 2001 July 18, and marginally in ObsID 1561b on 2001 July 14 (2σ) and in ObsID 2284 on 2001 July 18 (4 σ). If we assume that the spectrum is the same as in outburst, the observed flux was $(1-5) \times 10^{-14}$ erg cm⁻² s⁻¹ (2-8 keV). Assuming a 0.3 keV blackbody typical for a quiescent neutron star LMXB (Brown, Bildsten, & Rutledge 1998) results in fluxes a factor of 3 lower. Including the uncertainty on the spectrum in quiescence, the deabsorbed luminosities range between $(1-10) \times 10^{32}$ erg s⁻¹ (2–8 keV, 8 kpc), which is typical for a LMXB in quiescence. There is evidence for variability in the quiescent flux, as the upper limit on 1999 September 21 is lower than the detection on 2001 July 18 at 17:38 with a significance greater than $3-\sigma$.

We extracted a light curve from the observation in which the transient was bright (ObsID 1561a on 2000 October 26), which we display in Figure 3. The bin size is 23 s, so that there are over 100 counts per bin. An X-ray burst lasting about 100 s is observed 25,500 s into the observation (Figure 4a). It decays exponentially on a time scale of 54 ± 6 s. We examined the burst in two energy bands, 0.5–4.5 keV and 4.5–8.0 keV, which provide approximately equal counts during the persistent periods before and after the burst. The burst is harder in the peak than in the tail (Figure 4b), and the profile decays faster at high energies $(\tau = 43 \pm 5 \text{ s})$ than at low energies $(\tau = 77 \pm 18 \text{ s})$. Thus, the burst appears to cool as it decays, which suggests that the burst is thermonuclear in origin, and not a sudden burst of accretion (see Lewin et al. 1993). However, the burst is weak, reaching a peak flux only 3.5 times that of

FIG. 4.— Light curve (top panel) and hardness ratio (4.5–8.0 keV:0.5–4.5 keV; bottom panel) from 600 s during which the 100 s burst was observed. The spectrum of the burst is harder in the peak than in the tail, which is consistent with the cooling that is expected in the tail of a thermonuclear X-ray burst. A typical uncertainty from the light curve is displayed in the upper left of the top panel.

the persistent emission. For a 2 keV blackbody (e.g., Kuulkers et al. 2002, see also below), the peak rate of 18 count s^{-1} is equivalent to 6×10^{-10} erg cm⁻² s⁻¹, or 5×10^{36} erg s⁻¹ at 8kpc (2–8 keV). The fluence of the burst is approximately 400 counts. This translates to 1×10^{-8} erg cm⁻² (2–8 keV, absorbed) assuming a 2 keV blackbody spectrum and a 1×10^{23} cm⁻² absorption column, for an energy output of 10^{38} erg (8 kpc).

We extracted a spectrum of the first 80 s of the burst in order to determine the average temperature and solid angle of the emitting area. We used the persistent emission from the entire observation prior to 20 s before the burst as background (e.g., Kuulkers et al. 2002), and modeled the burst as a blackbody absorbed by a 10×10^{22} $\rm cm^{-2}$ column of gas and dust using XSPEC. The spectrum contained only about 250 counts, so we grouped the spectrum so that each bin contained ≥ 20 counts. We can only place 90% confidence limits on the apparent temperature of 2-8 keV, and on the apparent radius of 0.3-1.5 km (at 8 kpc). The fit was acceptable, with $\chi^2 = 6$ for 12 degrees of freedom. The solid angles derived from spectra of X-ray bursts are usually within a factor of two of the expected radius of a neutron star (about 10 km; e.g., Kaptein et al. 2000; Cocchi et al. 2001; Cornelisse et al. 2002a; Kuulkers et al. 2002). The small solid angle from which the burst emission originates appears to be the main factor in its relative weakness compared to previous bursts from this source (compare Cocchi et al. 1999).

In order to search for variability in the light curve aside from the burst, we removed 140 s of data starting 20 s prior to the burst, filled the gap with the mean count rate,





 $^{^{5}}$ The lower bound of 2 keV was chosen because Galactic absorption limits the flux observed below this value. The upper bound of 8 keV is the limit above which the ACIS effective area is small, and its spectral response is not well calibrated.

and computed a Fourier Transform of the resulting light curve. We find broad-band low-frequency variability with a total rms power of 0.7% rms above the expected Poisson noise between 5.5×10^{-5} and 0.13 Hz. We find no evidence for periodic signals over this frequency range with a significance greater than $2-\sigma$, and place a 90% upper limit on any periodic signal in this range of 1.5%, after incorporating the expected distribution of noise power (Vaughan et al. 1994).

2.2. XMM-Newton

XMM-Newton observed GRS 1741.9–2853 several times as part of a survey of the Galactic Plane and observations of the bursting pulsar GRO J1744–28. We searched for GRS 1741.9–2853 in the three observations taken with the European Photon Imaging Camera (EPIC) that were in the public archive as of April 2003 (Table 4). Data from the two metal-oxide semiconductor (MOS) CCDs were available from all three observations, while data from the pn CCD camera were only available on 2001 September 04. The cameras record X-rays within a calibrated energy band of 0.2–12 keV with an energy resolution of $E/\Delta E \approx 50$ at 6.5 keV. The medium filters were used for all observations.

We reduced the data starting with the standard pipeline event lists for each EPIC camera using SAS version 5.4.1. We filtered the data to remove event grades higher than 12, events flagged as bad by the standard processing, and events below 0.3 keV (as these are likely to be background events). We then examined the light curve from each observation in order to check for soft proton flares. The entire exposure on 2001 March 31 was contaminated with such flares, limiting the usefulness of the data. Flares were also observed on 2001 September 04, so we removed all intervals for which the count rate was 3 standard deviations above the mean. We removed 6000 s of flaring from the pn data in this manner (resulting in the exposure listed in Table 4), and 3500 s from the MOS data.

The field of view of the EPIC cameras are 30' in radius, so GRS 1741.9-2853 was near the edge of the field in all three observations. GRS 1741.9–2853 was not detected by eve, and no source at its location was listed in the source catalogs that are produced in the standard pipeline processing. We were unable to use the technique applied to the *Chandra* observations to derive an upper limit on the flux from GRS 1741.9–2853 during the XMM-Newton observations for two reasons. First, although the full-width of the half-maximum of the XMM-Newton PSF is only 6''on-axis, off-axis it is significantly degraded. At an offset of 10', 90% of the photons at 5 keV are enclosed by the PSF within radii of 70'' for the pn CCDs and 85'' for the MOS CCDs. Second, the diffuse X-ray emission detected with Chandra from the "darkest" regions of the inner 10' of the Galaxy varies by 10% around a mean value of 2×10^{-13} erg cm⁻² s⁻¹arcmin⁻² (Muno et al., in preparation). We therefore used the fluxes from the faintest sources detected as part of the pipeline processing between 8' and 12' from the aim-point of each observation as estimates of the upper limit on the count rate from GRS 1741.9-2853. These count rates were 0.009 count s^{-1} (MOS, 2–12 keV) for 2001 March 31, 0.003 count s^{-1} (MOS, 2–12 keV) for 2001 April 01, and 0.002 count s^{-1}



FIG. 5.— The luminosity history of GRS 1741.9–2853 assembled from *GRANAT* (squares; 3–30 keV), *ASCA* (diamonds; 2–10 keV), *XMM-Newton* (circles; 2–12 keV), and *Chandra* (triangles; 2–8 keV) observations. The luminosity varies over four orders of magnitude on time scales of months, as is commonly observed from LMXB transients. The variability at low luminosities (< 10^{33} erg s⁻¹) is significant (see Table 3 for uncertainties), and has been observed from other sources on only a few occasions.

(pn, 2–12 keV) for 2001 September 04. We converted these to energy fluxes using PIMMS by assuming a $\Gamma = 1.9$ power law and $N_{\rm H} = 10 \times 10^{22}$ cm⁻², and listed them in Table 1. The resulting limits are similar to both the *Chandra* detections in quiescence and the variations in the background flux on arcminute scales.

3. DISCUSSION

We plot the long-term history of the persistent luminosity from GRS 1741.9-2853 in Figure 5 (deabsorbed, and assuming a distance of 8 kpc). It is clear that this source varies in luminosity by at least four orders of magnitude on time scales of months. GRS 1741.9-2853 was fainter than 10^{33} erg s⁻¹ in Fall 1999, and Fall and Spring 2001. The source has been observed to reach luminosities above 10^{36} erg s⁻¹ three times in the last 13 years: in Spring 1990, Fall 1996, and Fall 2001. This suggests that the bright states are transient outbursts driven by instabilities in the accretion disk (e.g., Lasota 2001). It is unlikely that there were any outbursts brighter than 6×10^{36} erg s⁻¹ lasting more than a month between 1996 and the present, because they would have been detectable with the BeppoSAX WFC and the RXTE ASM. Thus, GRS 1741.9–2853 appears to be a genuine low-luminosity transient. Under the disk instability model, the faintness of the outbursts can be explained if the mass transfer rate from the companion is low, either because it is due to a weak wind, or because the companion is a very low-mass star that fills its Roche-lobe (King 2000).

GRS 1741.9–2853 was also detected several times at luminosities below 10^{35} erg s⁻¹ (Figure 5). It was observed with *ASCA* at a luminosity of 10^{34} erg s⁻¹ Fall 1994. Several other sources have been observed with varying flux at this level, including the neutron star LMXBs SAX J1747.0–2853 (Wijnands et al. 2002), SAX J1808.4–3568 (Wijnands et al. 2001), 1RXS J1718.2–402934 (Kaptein et al. 2000), and SAX J1828.5–1037 (Cornelisse et al. 2002a). Such lowluminosity activity is not addressed by the simplest disk instability models, which generally produce $10^{37} - 10^{38}$ erg s^{-1} outbursts because a large fraction of the accretion disk is expected to be disrupted (e.g., Dubus, Hameury, & Lasota 2001). Furthermore, Chandra observations demonstrate that GRS 1741.9-2853 varies in X-ray intensity when its luminosity is only about 10^{32} erg s⁻¹(Table 3 and Figure 5). The fact that GRS 1741.9–2853 is significantly brighter after the Fall 2000 outburst could be attributed to the flux from the hot $(kT \sim 0.3 \text{ keV})$ surface of the neutron star, which may have been heated during the outburst (Brown et al. 1998). However, the high absorption column toward the source should make most of this thermal emission unobservable (see Figure 2). On the other hand, short time-scale variations have been observed in the flux from the LMXBs Cen X-4 (Campana et al. 1997; Rutledge et al. 2001) and Aql X-1 (Rutledge et al. 2002) that are inconsistent with a cooling neutron star. This emission has been explained as residual accretion either onto the neutron star surface or onto the magnetospheric boundary (Campana et al. 1998). The emission from GRS 1741.9-2853with luminosities between 10^{32} and 10^{34} erg s⁻¹ probably also represents continued accretion. These low luminosities are an important, and relatively unexplored, regime of accretion that challenge current disk instability models for LMXBs.

X-ray bursts were observed from GRS 1741.9-2853 with the *BeppoSAX* WFC during the bright state in Fall 1996 (Cocchi et al. 1999; Sakano et al. 2002), and with Chandra during the outburst in Fall 2000 (this work). The WFC observed the Galactic Center for approximately 500 ks each spring and fall from mid-1996 through the end of 2001, and detected no further bursts. This suggests that the bursts only occur frequently when the persistent X-ray emission reaches $\sim 10^{36}~{\rm erg~s^{-1}}$, so that accretion provides $10^{-10}~M_{\odot}~y^{-1}$ of fuel to the surface of the neutron star. It is likely that some other low-luminosity bursters are similar, and that sensitive observations at or just before the times that bursts were observed would have revealed persistent emission just below the sensitivity limit of the BeppoSAX WFC and RXTE ASM (~ 10^{36} erg s⁻¹; see also Cocchi et al. 2001). If this is the case, then the fraction of low-luminosity bursters that have been detected so far depends upon how often the transient outbursts occur, which is currently uncertain.

The weakness of the X-ray burst observed with Chandra is a bit puzzling. Its peak flux of 6×10^{-10} erg cm⁻² s⁻¹ (2– 8 keV) is a factor of 40 below those of Eddington-limited bursts observed previously from this source with the Bep-

poSAX WFC (10⁻⁸ erg cm⁻² s⁻¹). Faint bursts are not uncommon from LMXBs, but they are usually preceded by bright bursts tens of minutes before (e.g. Langmeier et al. 1987; Gottwald et al. 1986), and are thought to occur in fuel that was not burnt in the previous burst (Lewin et al. 1987; Fujimoto et al. 1987). However, no burst was observed from GRS 1741.9-2853 in the seven hours of Chandra observations prior to the weak burst, and the fluence of the burst (10^{38} erg) is an order of magnitude smaller than the amount of nuclear energy that could have been stored in accreted hydrogen during the *Chandra* observation prior to the burst $(0.04 \cdot 2.5 \times 10^4 \text{ s} \cdot 10^{36} \text{ erg s}^{-1} = 10^{39} \text{ erg}$; see Figure 3). On the other hand, this weak burst is reminiscent of a 10^{37} erg burst observed by Gotthelf & Kulkarni (1997) with ASCA from the globular cluster M28. The source that produced that burst had a persistent luminosity of at most 10^{33} erg s⁻¹. However, if material was accreted onto the entire surface of the neutron star at such a low accretion rate (< $10^{-13} M_{\odot} y^{-1}$), the recurrence time for bursts should be very long, and their fluence high. For instance, to collect the 10^8 g cm⁻² column necessary to trigger a burst over the entire star (an area of 10^{13} cm⁻²) would take several years, and the nuclear energy stored on the surface would be at least 10^{39} erg (e.g., Fujimoto, Hanawa, & Miyaji 1981). Therefore, Gotthelf & Kulkarni (1997) suggested that the accreted material in the M28 source is channeled onto a small area of the surface of the neutron star, such as its magnetic poles. This explanation may also apply for GRS 1741.9-2853, for which we infer that the burst emission originates from a solid angle that is only a few percent of the surface area of a 10 km neutron star. Moreover, the fact that four lowluminosity transients are accreting millisecond X-ray pulsars suggests that faint transients are more likely than the general LMXB population to have magnetic fields strong enough to channel the accretion flow onto the poles of the neutron star. Therefore, we suggest that future observations of GRS 1741.9-2853 in outburst with sensitive timing instruments could reveal millisecond pulsations.

We thank A. Levine, D. Galloway, R. Remillard, and C. Markwardt for useful discussions, and for providing information on the flux history and burst properties of GRS 1741.9–2853 and similar LMXBs. We thank the referee for the careful reading and useful comments. This work has been supported by NASA grants NAS 8-39073 and NAS 8-00128.

REFERENCES

- Baganoff, F. K., Bautz, M. W., Brandt, W. N., Chartas, G., Feigelson, E. F., Garmire, G. P., Maeda, Y., Morris, M., Ricker, G. R., Townsley, L. K., & Walter, F. 2001, Nature, 413, 45
 Baganoff, F. K. et al. 2003, ApJ, 591, 891
 Belloni, T., Colombo, A. P., Homan, J., Campana, S., & van der Klis, M. 2002, A&A, 390, 199

- Khs, M. 2002, A&A, 390, 199
 Bildsten, L. 2000, in Cosmic Explosions: Tenth Astrophysics Conference, eds. S. S. Holt and W. W. Zhang, AIP Conference Proceedings, Vol. 522, College Park, MD, 11-13 Oct 1999. American Institute of Physics, 2000, pg. 359, astro-ph/0001135
 Brandt, W. N. et al. 2001, AJ, 122, 2810
 Broos, P., Townsley, L., Getman, K. & Bauer, F. 2002, "ACIS Extract, An ACIS Point Source Extraction Package", Pennsylvania State University
- Pennsylvania State University
- Brown, E. F., Bildsten, L., & Rutledge, R. E. 1998, ApJ, 504, L95

- Campana, S., Mereghetti, S., Stella, L., & Colpi, M. 1997, A&A, 324, 94î
- 941
 Campana, S., Colpi, M., Mereghetti, S., Stella, L., & Tavani, M. 1998, Astronomy and Astrophysics Review, 8, 279
 Cocchi, M., Bazzano, A., N, L., Ubertini, P., Heise, J., Muller, J. M., & in' t Zand J. M M. 1999, A&A, 346, L45
- Cocchi, M., Bazzano, A., Natalucci, L., Ubertini, P., Heise, J., Kuulkers, E., Cornelisse, R., & in 't Zand, J. J. M. 2001, A&A, 378, L37
- Cornelisse, R., Verbunt, F., in 't Zand, J. J. M., Kuulkers, E., Heise, Cornelisse, R., Vorbin, Y., M. Cocchi, M., Natalucci, L., Bazzano, A., & Ubertini, P. 2002a, A&A, 392, 885
 Cornelisse, R., Verbunt, F., in 't Zand, J. J. M., Kuulkers, E., & Heise, J. 2002b, A&A, 392, 931
 Cumming, A., Zweibel, E., & Bildsten, L. 2001, ApJ, 557, 958

- Dubus, G., Hameury, J.-M., Lasota, J.-P. 2001, A&A, 373, 251
- Freeman, P. E., Kashyap, V., Rosner, R., & Lamb, D. Q. 2002, ApJS, 138, 185
- Fujimoto, M. Y., Hanawa, T., & Miyaji, S. 1981, ApJ, 246, 267
 Fujimoto, M. Y., Sztajno, M., Lewin, W. H. G., & Van Paradijs, J.

- rujimoto, M. Y., Sztajno, M., Lewin, W. H. G., & Van Paradijs, J. 1987, ApJ, 319, 902
 Galloway, D. K., Chakrabarty, D., Morgan, E. H., & Remillard, R. A. 2002, ApJ, 576, L137
 Gotthelf, E. V. & Kulkarni, S. R. 1997, ApJ, 490, L161
 Gottwald, M., Haberl, F., Parmar, A. N., & White, N. E. 1986, ApJ, 308, 213
- in 't Zard, J. J. M. 2001, in Exploring the Gamma-ray Universe, eds. A Gimenez, V. Regelo, & C. Winkler, ESA publication division,
- A Gimenez, V. Regelo, & C. Winkler, ESA publication division, p. 463. astro-ph/0104299 Kaptein, R. G., in 't Zand, J. J. M., Kuulkers, E., Verbunt, F., Heise, J., & Cornelisse, R. 2000, A&A, 358, L71 King, A. R. 2000, MNRAS, 315, L33 Kraft, R. P., Burrows, D. N., & Nousek, J. A. 1991, ApJ, 374, 344 Kuulkers, E., Homan, J., van der Klis, M., Lewin, W. H. G., & Méndez, M. 2002, A&A, 382, 947 Langmeier, A., Sztaino, M., Hasinger, G., Trümper, J., & Gottwald.

- Langmeier, A., Sztajno, M., Hasinger, G., Trümper, J., & Gottwald, M. 1987, ApJ, 323, 288 Lasota, J.-P. 2001, New Astronomy Reviews, 45, 449 Lewin, W. H. G., Pennix, W. Van Paradijs, J., Damen, E., Sztajno,
- M., Trümper, J., & Van der Klis, M. 1987, ApJ, 319, 893

- Lewin, W. H. G., Van Paradijs, J., & Taam, R. E. 1993, Space Science

- Lewin, W. H. G., Van Paradıjs, J., & Taam, R. E. 1993, Space Science Review, 62, 223
 Markwardt, C. B., Smith, E., & Swank, J. H. 2003, IAU Circ., 8080
 Markwardt, C. B., Swank, J. H., Strohmayer, T. E., in 't Zand, J. J. M., & Marshall, F. E. 2002, ApJ, 575, L21
 Muno, M. P., Baganoff, F. K., Bautz, M. W., Ricker, G. R., Morris, M., Garmire, G. P., Feigelson, E. D., Brandt, W. N., Townsely, L. K., & Broos, P. S. 2003a, ApJ, 589, 225
 Pavlinsky, M. N., Grebenev, S. A., & Sunyaev, R. A. 1994, ApJ, 425, 110
- 110
- Rutledge, R. E., Bildsten, L., Brown, E. F., Pavlov, G. G., & Zavlin, V. E. 2001, ApJ, 551, 921 Rutledge, R. E., Bildsten, L., Brown, E. F., Pavlov, G. G., & Zavlin,
- V. E. 2002, ApJ, 577, 346 Simon, V. 2002, A&A, 381, 151 Sakano, M., Koyama, K., Murakami, H., Maeda, Y., & Yamaushi, S.
- 2002, ApJS, 139, 19
- Townsley, L. K., Broos, P. S., Garmire, G. P., & Nousek, J. A. 2000, ApJ, 534, L139

- Vaughan, B. A. et al. 1994, ApJ, 435, 362
 Wang, Q. D., Gotthelf, E. V., & Lang, C. C. 2002, Nature, 415, 148
 Wijnands, R. & van der Klis, M. 1998, Nature, 394, 344
 Wijnands, R., Méndez, M., Markwardt, C., van der Klis, M., Chakrabarty, D., & Morgan, E. 2001, ApJ, 560, 896
 Wijnanda, P. Miller, J. M. & Wang, O. D. 2002, ApJ, 570, 422
- Wijnands, R., Miller, J. M., & Wang, Q. D. 2002, ApJ, 579, 422

Date	Mission	$F_{\rm X}{}^{\rm a}$	References
1990 Mar 24	Granat	2000	(1)
1990 Apr 08	Granat	2000	(1)
1993 Sep 30	ASCA	$<\!\!12$	(2)
1993 Oct 01	ASCA	$<\!\!40$	(2)
1993 Oct 03	ASCA	$<\!20$	(2)
1993 Oct 04	ASCA	$<\!19$	(2)
$1994 { m Sep } 15$	ASCA	18	(2)
1996 Sep 11	ASCA	1000	(2)
1997 Mar 16	ASCA	$<\!\!8$	(2)
1999 Sep 21	Chandra	< 0.09	(3)
2000 Oct 26	Chandra	2000	(3)
2001 Mar 31	XMM	$<\!\!2$	(3)
2001 Apr 01	XMM	< 0.8	(3)
2001 Jul 14	Chandra	0.1	(3)
2001 Jul 18	Chandra	0.3	(3)
2001 Jul 18	Chandra	0.5	(3)
$2001~{\rm Sep}~04$	XMM	< 0.2	(3)

TABLE 1 Flux History of GRS 1741.9–2853

^aX-ray flux in units of 10^{-13} erg cm⁻² s⁻¹. The energy ranges are: 3–30 keV for *Granat*, 2–10 keV for *ASCA*, 2–8 keV for *Chandra*, and 2–12 keV for *XMM*.

Note. — The *BeppoSAX* WFC observed the Galactic center every Spring and Fall from mid-1996 through the end of 2001. The WFC detected three thermonuclear bursts during Fall 1996, but never detected persistent emission from accretion with $F_{\rm X} > 2 \times 10^{-10}$ erg cm⁻¹ s⁻¹ (Cocchi et al. 1999).

References. -(1) Pavlinsky et al. (1994); (2) Sakano et al. (2002); (3) This work.

	Aim Point					
Start Time (UT)	ObsID	$\begin{array}{c} \text{Exposure} \\ \text{(s)} \end{array}$	RA (degrees	DEC J2000)	$\begin{array}{c} \text{Roll} \\ \text{(degrees)} \end{array}$	$\begin{array}{c} \text{Offset}^{\mathrm{a}} \\ (\operatorname{arcmin}) \end{array}$
1999 Sep 21 02:43:00 2000 Oct 26 18:15:11 2001 Jul 14 01:51:10 2001 Jul 18 14:25:48	0242 1561a ^b 1561b 2284	40,872 35,705 13,504 10,625	$\begin{array}{c} 266.41382\\ 266.41344\\ 266.41344\\ 266.40415 \end{array}$	-29.0130 -29.0128 -29.0128 -28.9409	$268 \\ 265 \\ 265 \\ 284$	$10 \\ 10 \\ 10 \\ 8$
2001 Jul 18 17:37:38	2287	10,625	266.21412	-28.8391	284	$\overline{5}$

TABLE 2 Chandra Observations of GRS 1741.9–2853

^aThe offset listed is from the nominal position of GRS 1741.9–2853 from Pavlinsky et al. (1994).

^bThis observation was taken during two epochs because the ACIS-I camera was powered off at the beginning of the first epoch. It is listed in the archive under a single ObsID, 1561.

Counts						
Date	Tot	Bkgd	Net	Flux		
1999 Sep 21 02:43	30	27.6	< 11.9	< 0.09		
2000 Oct 26 18:15	170214	87	$170127 {\pm} 412$	2000 ± 120		
2001 Jul 14 01:51	14	8.6	$5.4^{+6.6}_{-5.0}$	$0.11^{+0.15}_{-0.10}$		
2001 Jul 18 14:26	11	2.0	$9.0^{+6.4}_{-4.6}$	$0.23_{-0.13}^{+0.19}$		
2001 Jul 18 17:38	18	0.8	$17.2_{-6.1}^{+7.9}$	$0.46_{-0.18}^{+0.23}$		

 $\label{eq:Table 3} TABLE \ 3 \\ Chandra \ Photometry \ of \ GRS \ 1741.9-2853 \ (2-8 \ {\rm keV})$

Note. — Confidence limits are 90%, based on the observed number of counts (see text). Fluxes are absorbed in units of 10^{-13} erg cm⁻² s⁻¹ (2–8 keV).

TABLE 4XMM-Newton Observations of GRS 1741.9-2853

Start Time (UT)	Sequence	Aim PointExposureRADECRollOff(s)(degrees J2000)(degrees)(arcs)				
2001 Mar 31 11:24:26 2001 Apr 01 13:02:43 2001 Sep 04 01:19:34	$\begin{array}{c} 0112971601\\ 0112971901\\ 0112972101 \end{array}$	$3830 \\ 9191 \\ 19,040$	$\begin{array}{c} 266.40146\\ 266.12667\\ 266.43463\end{array}$	-28.9820 -28.7313 -29.0313	$95 \\ 95 \\ 271$	10 10 10

Note. — The offset listed is from the nominal position of GRS 1741.9–2853 from Pavlinsky et al. (1994). A fourth observation, 0112972001, is listed in the archive, but contains no usable data.