

The Properties and the Evolution of the Highly Ionized Gas in MR 2251–178

Shai Kaspi, Hagai Netzer, Doron Chelouche

*School of Physics and Astronomy, Raymond and Beverly Sackler
Faculty of Exact Sciences, Tel-Aviv University, Tel-Aviv 69978, Israel*

Ian M. George, T. J. Turner

*Joint Center for Astrophysics, Physics Department, University of
Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD
21250, USA, and Laboratory for High Energy Astrophysics, NASA/
Goddard Space Flight Center, Code 662, Greenbelt, MD 20771, USA.*

Kirpal Nandra

*Astrophysics Group, Imperial College London, Blackett Laboratory,
Prince Consort Road, London SW7 2AZ, UK.*

Abstract. We present the first *XMM-Newton* observations of the radio-quiet quasar MR 2251–178. We model the X-ray spectrum with two power laws, one at high energies with a slope of $\Gamma = 1.6$ and the other to model the soft excess with a slope of $\Gamma = 2.9$, both absorbed by at least two warm absorbers (WAs). The high-resolution grating spectrum shows emission lines from N VI, O VII, O VIII, Ne IX, and Ne X, as well as absorption lines from the low ionization ions O III, O IV, and O V. A study of the spectral variations in MR 2251–178 over a period of 8.5 years yields that all X-ray observations can be fitted with the above model. Luminosity variations over timescales of years seem to correlate with the soft excess variations but not with the WA properties variations. The overall picture is that of a stratified WA that enters and disappears from the line-of-sight on timescales of several months. We also present the first *FUSE* spectrum of MR 2251–178. The general characteristics of the UV and X-ray absorbers seem to be consistent.

1. Introduction

MR 2251–178 ($z = 0.06398 \pm 0.00006$, $V \approx 14$) is the first quasar detected by X-ray observations (using *Ariel V* and *SAS-3* in 1977) and also the first quasar where a warm absorber (WA) was suggested to explain the X-ray spectrum (Halpern 1984). It was observed by practically all X-ray missions since *Einstein*. Previous X-ray data of MR 2251–178 were modeled using a power-law with photon index $\Gamma \approx 1.6$ –1.7 and a WA with column density in the range $10^{21.3-22.2} \text{ cm}^{-2}$ (e.g., Mineo & Stewart 1993; Orr et al. 2001; Morales & Fabian 2002). The X-ray flux of the source is variable on timescales of ~ 10




Figure 1. (a) Ratio of the 2002 *XMM-Newton* EPIC-pn data to a power law model (including Galactic absorption of $10^{20.45} \text{ cm}^{-2}$) fitted to the 3–11 keV band. Excess absorptions is evident near the O VII and the O VIII absorption edges and at energies below 0.5 keV. (b) Ratio of the 2000 *XMM-Newton* EPIC-pn data to the scaled 2002 EPIC-pn model. The additional soft excess in the 2000 spectrum is evident.

days. The observed 2–10 keV flux of MR 2251–178 covers the range of $(1.7\text{--}5.1) \times 10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ which translates to a 2–10 keV luminosity of $(1.7\text{--}5.1) \times 10^{44} \text{ erg cm}^{-2} \text{ s}^{-1}$.

UV spectra of MR 2251–178 were obtained by the *Hubble Space Telescope* at three epochs and show clear Ly α and C IV absorption. Ganguly et al. (2001) found the C IV doublet absorption to vary with time, suggesting an intrinsic origin for this absorption.

In this contribution we present new *XMM-Newton* and *FUSE* observations of MR 2251–178. We also carry out an in-depth analysis of the 10 available *ASCA* observations and the two *BeppoSAX* observations.

2. The X-ray Spectra and Past variations

MR 2251–178 was observed with *XMM-Newton* in 2000 and in 2002. The flux in the 2002 observation was about 2.5 times lower than the 2000 observation. The EPIC-pn data of the 2002 observation clearly show a power law with a photon index of $\Gamma \approx 1.6$ at energies above 3 keV. Extrapolating this power law to lower energies reveals the presence of a WA around 0.8 keV, an additional absorber below 0.5 keV, and some excess emission around 0.5 keV (Figure 1a). Our best fitted model for these data yields a WA with a column density, N_{H} , of $10^{21.51 \pm 0.03} \text{ cm}^{-2}$, ionization parameter, U_{OX} , of $10^{-1.78 \pm 0.05}$ and a line of sight covering factor of 0.8. Assuming gas with the same properties produces the emission, we find a global covering factor of 0.3. For the less ionized absorber we find that it can be fitted by a neutral absorber (in addition to the galactic one) with $N_{\text{H}} \approx 10^{20.3} \text{ cm}^{-2}$. This absorber can also be modeled as a combination of low-ionization absorber with $\log(U_{\text{OX}}) \approx -4$ and $N_{\text{H}} \approx 10^{20.3} \text{ cm}^{-2}$ and a neutral absorber of $N_{\text{H}} \approx 10^{20.06} \text{ cm}^{-2}$. Both cases give equally good fits. Fitting the above model to the 2000 observation clearly reveals the presence of a soft excess (Figure 1b) which can be fitted with an additional power law with a photon index of $\Gamma \approx 2.9$ at energies less than ~ 1 keV.




Figure 2. Combined RGS1 and RGS2 spectrum of MR 2251–178 binned to ~ 0.04 Å. The spectrum has been corrected for Galactic absorption and for the redshift of the source. The strongest emission lines are due to the O VII triplet and O VIII Ly α . Other suggested absorption and emission lines are marked. Gaps in the spectrum due to chip gaps are marked as ‘art’. The two absorbers model described in the text is shown (gray line).

The high-resolution grating spectrum of the 2002 observation is shown in Figure 2. The spectrum shows emission lines from N VI, O VII, O VIII, Ne IX, and Ne X, as well as absorption lines from the low ionization ions O III, O IV, and O V. The data were fitted with a detailed photoionization model which includes the components described above and is shown in Figure 2.

We used the model with the two power laws and two absorbers described above to fit historical data of MR 2251–178 obtained over 8.5 years. The WA properties during the 6 weeks of *ASCA* observations in 1993 were consistent with those of an absorber with a column density of $10^{21.5}$ cm $^{-2}$. These data are consistent with the scenario in which the decrease in flux caused a corresponding decrease in the ionization parameter. However, the MR 2251–178 observations are not frequent and detailed enough to infer on the location of the gas.

On timescales of years, the WA properties change in time but are not correlated with luminosity variations. Our only successful model requires that the absorbing material is changing in time. For example, the *ASCA* 1996 observations clearly indicate a larger column density ($> 10^{21.8}$ cm $^{-2}$ vs. $10^{21.5}$ cm $^{-2}$) and a smaller ionization parameter ($\log U_{\text{OX}} \sim -2.3$ vs. ~ -2.0) of the WA compared to 1993. We suggest that a physical motion of gas into and out of our line-of-sight can cause these changes in the absorber properties. In the 2002 *XMM-Newton* observation, the flux is smaller by a factor of 2 compared with 1993 *ASCA* observations, yet the column density and the ionization parameter are similar. Comparing the two *XMM-Newton* observations yields a similar conclusion. This might mean that the absorbing material has changed between




Figure 3. Selected bands from the *FUSE* spectrum of MR 2251–178. Identified intrinsic emission lines are marked at their theoretical position with vertical lines and identified airglow and Galactic lines are marked below the spectrum with \oplus .

the epochs since the SED is very similar but the luminosity decrease between the epochs was not accompanied by a corresponding decrease in ionization parameter. An alternative explanation can be that the absorbing material is far from the central source and does not respond to the luminosity variations.

3. The High Resolution *FUSE* spectrum

The *FUSE* spectrum (Figure 3) shows broad emission lines of O VI and C III which also show significant blueshifted absorption. We also detect blueshifted absorption from at least 10 lines of the H I Lyman series. We identify at least 4 absorption systems, one at -580 km s^{-1} and at least 3 others which are blended together and form a wide trough covering the velocity range of 0 to -500 km s^{-1} . The trough profiles are consistent with the absorption in the high resolution X-ray spectrum (though the later does not have sufficient resolution) and suggest that the UV and X-ray absorptions may arise from the same region in the AGN.

Acknowledgments. We acknowledge a financial support by the Israel Science Foundation grant no. 545/00. S. K. also acknowledge financial support by Colton Foundation.

References

- Ganguly, R., Charlton, J. C., & Eracleous, M. 2001, *ApJ*, 556, L7
Halpern, J. P. 1984, *ApJ*, 281, 90
Mineo, T. & Stewart, G. C. 1993, *MNRAS*, 262, 817
Morales, R. & Fabian, A. C. 2002, *MNRAS*, 329, 209
Orr, A. et al. 2001, *A&A*, 376, 413