

Variable stars in the Magellanic Clouds: II. The data and infrared properties

Yoshifusa Ita^{1*}, Toshihiko Tanabé¹, Noriyuki Matsunaga¹, Yasushi Nakajima², Chie Nagashima², Takahiro Nagayama³, Daisuke Kato³, Mikio Kurita³, Tetsuya Nagata³, Shuji Sato³, Motohide Tamura², Hidehiko Nakaya⁴ and Yoshikazu Nakada^{1,5}

¹*Institute of Astronomy, School of Science, The University of Tokyo, Mitaka, Tokyo 181-0015, Japan*

²*National Astronomical Observatory of Japan, Mitaka, Tokyo 181-8588, Japan*

³*Department of Astrophysics, Nagoya University, Chikusa-ku, Nagoya 464-8602, Japan*

⁴*Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'ohoku Place, Hilo, HI 96720, U.S.A.*

⁵*Kiso Observatory, School of Science, The University of Tokyo, Mitake, Kiso, Nagano 397-0101, Japan*

Received – / Accepted –

ABSTRACT

The data of 8,852 and 2,927 variable stars detected by OGLE survey in the Large and Small Magellanic Clouds are presented. They are cross-identified with the SIRIUS *JHK* survey data, and their infrared properties are discussed. Variable red giants are well separated on the period- $J - K$ plane, suggesting that it could be a good tool to distinguish their pulsation mode and type.

Key words: galaxies: Magellanic Clouds – stellar content – stars: AGB and post-AGB – variables – infrared: stars – surveys

1 INTRODUCTION

Because of their proximity and comparatively well-known distances, the Large and Small Magellanic Clouds (LMC and SMC, respectively) are the ideal places to study stellar evolution. To search for the gravitational lensing events, several projects are now underway monitoring the sky toward the Magellanic Clouds and also the Galactic bulge, where there are a large number of background stars that are potential targets for microlensing. As the natural by-products of these surveys, many variable stars were newly found. The Optical Gravitational Lensing Experiment (OGLE) project is one of such surveys and its I band time-series data in the Magellanic Clouds (Žebruň et al. 2001) obtained during four years of OGLE-II run (Udalski, Kubiak & Szymanski 1997; January 1997 - November 2000) is now available over the Internet (OGLE homepage; <http://sirius.astrouw.edu.pl/~ogle/>).

To detect dust-enshrouded variables that have been possibly escaped from the ongoing/previous optical projects, we started the *JHK* monitoring survey toward the Large and Small Magellanic Clouds using the InfraRed Survey Facility (IRSF) at the South African Astronomical Observatory Sutherland station. The IRSF consists of a dedicated 1.4m telescope to which is attached a near infrared (NIR) camera SIRIUS. Details of the instrument are found in Nagashima et al. (1999) and Nagayama et al. (2002). This campaign was initiated in December 2000, and so far, many dust-

enshrouded variables were found and their light variation data has been steadily accumulating (see Ita et al. 2002 for the first results).

In the previous paper (Ita et al. 2003, hereafter referred to as Paper I), we cross-identified the OGLE data and the single-epoch SIRIUS *JHK* data from the monitoring survey, and studied the pulsation properties and metallicity effects on period- K magnitude relations by comparing the variable stars in the LMC and SMC. In this paper, we present the data of the variable stars in the Magellanic Clouds and discuss their infrared properties.

2 DATA

Details of the cross-identifications and period findings are described in Paper I. We did not analyze multi-periodic stars (e.g., Bedding et al. 1998) and/or variable stars that show too complex light curves to find a predominant period. We made two catalogs presenting data of variable stars in the LMC and SMC. They contain following information: OGLE name, pulsation period in days, statistical parameter θ calculated by Phase Dispersion Minimization technique (Stellingwerf 1978), pulsation amplitude ($\Delta I = I_{\max} - I_{\min}$), intensity mean I magnitude $\langle I \rangle$, *JHK* magnitudes and positional difference r between the SIRIUS coordinate and the OGLE coordinate in arcsec. Table 1 is a sample that shows the first five records of the LMC catalog. The full version of the data, including SMC catalog are available in the on-line version of this paper. In cases the *JHK* measurements are saturated or below the de-

* E-mail: yita@ioa.s.u-tokyo.ac.jp

Table 1. The first 5 records in the LMC catalog. This is a sample of the full version, which is available in the electronic version of Monthly Notices.

Name	Period [days]	θ	ΔI	$\langle I \rangle$	J	H	K	r
					[mag]			[arcsec]
OGLE050437.94-692217.2	0.460	0.178	1.025	19.250	99.999	16.056	16.030	2.260
OGLE050438.43-691527.2	49.892	0.837	0.089	13.951	13.559	10.786	11.337	0.306
OGLE050440.26-693039.7	42.783	0.866	0.160	14.189	12.786	11.880	11.599	0.466
OGLE050440.29-692708.3	39.200	0.661	0.096	14.734	13.423	12.617	12.333	0.465
OGLE050440.76-692416.0	96.365	0.358	0.317	14.060	12.294	11.491	11.063	0.574

Table 2. Colour mapping table

Label	Colour	Population
A ⁻	Cyan	RGB variables & metal poor and old AGB variables
A ⁺	Yellow	less regularly pulsating AGB variables
B ⁻	Orange	RGB variables & metal poor and old AGB variables
B ⁺	Green	less regularly pulsating AGB variables
C'	Blue	Mira variables pulsating in the 1st overtone mode
C	Red	Mira variables pulsating in the fundamental mode
D	Steel-blue	Some obscured variables & Unknown variables
F	Magenta	Cepheid variables pulsating in the fundamental mode
G	Purple	Cepheid variables pulsating in the 1st overtone mode

tection limit, the corresponding columns contain the value 99.999. The JHK magnitudes (note that SIRIUS measures K_s) in the catalog are referred to the Las Campanas Observatory (LCO) system ones based on observations of a few dozen stars from Persson et al. (1998) and are not corrected for the interstellar absorption. The θ is a measure of the regularity of the light variation, being near 0 for the regular variation and near 1 for the irregular variation.


3 DISCUSSION

In the following discussions, we use the JHK magnitudes that are corrected for the interstellar absorption based on the relations in Koornneef (1982), assuming $R = 3.2$. We adopted $(A_J, A_H, A_K) = (0.129, 0.084, 0.040)$ and $(0.082, 0.053, 0.025)$, corresponding to the total mean reddening of $E_{B-V} = 0.137$ and 0.087 as derived by Udalski et al. (1999) for the OGLE's observing fields in the LMC and SMC, respectively. The reddening of the Magellanic Clouds are still somewhat controversial (e.g., Westerlund 1997), and it is certain that this type of statistical average approach can not account for individual corrections. However, the adopted reddening should not affect any of the conclusions of this paper. The coefficients we calculated in equations 1 to 4 depend on reddening, but the effects should be small.

3.1 Classification of the variable stars on the period- K magnitude plane

In Paper I, we classified variable stars into nine (LMC) and eight (SMC) prominent groups based on their locations on the period- K magnitude plane. Refer to Paper I for the details of the grouping. For the easier reading and completeness of this paper, we show the period- K magnitude diagram with classification boxes in figure 1, and also for conciseness, we tabulated the colour mapping chart in table 2. Stars outside the classification boxes are represented by black tiny dots. Throughout this paper, we use the same labels and colours for each group. In the LMC, there is a clump of variables around $\log P \lesssim -0.1$ and $K \lesssim 15.0$. This is a group of RR Lyrae variables, which we do not analyze in this paper.

Roughly speaking, five types of variable stars constitute the

**Figure 1.** Variable stars in the LMC (upper panel) and SMC (lower panel) are classified into several groups according to their location on the period- K magnitude plane.

majority of the catalog: Cepheids, Miras, Semi-regulars, Irregular variables and Eclipsing binaries. However, telling Semi-regulars from Mira variables is rather difficult. Even recently, several authors (e.g., Alard et al. 2001; Cioni et al. 2001) use a simple criterion which is based on a certain limiting pulsation amplitude to separate Semi-regulars and Miras. However, this simple criterion could be rather artificial. In this paper, we do not separate Semi-regulars and Miras explicitly. Instead, we distinguish “regularly pulsating variables, ($\theta \leq 0.55$)” from “less regularly pulsating variables, ($\theta > 0.55$)” relying on the parameter “ θ ”.

3.2 Colour-magnitude and colour-colour diagram

The NIR data from the SIRIUS survey enabled us to study the infrared properties of the OGLE variables. Figure 2 and 3 show the NIR colour-magnitude and colour-colour diagram of the OGLE variables. In the inset of the figure 2, we show the colour-magnitude diagram of the SIRIUS data (including non-variables), closing-up around the tip of the first giant branch (TRGB). By comparing the insets and the main diagrams, it is clear that the stars of group A⁻ and B⁻ accumulate at the exact location of the TRGB.




Figure 2. NIR colour-magnitude diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel). The inset shows the plot of the all SIRIUS data (including non-variables) around the K -band luminosity of the TRGB.




Figure 3. NIR colour-colour diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel).




Figure 4. Period-colour diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel). The dashed and thick solid lines indicate the least-square fits of linear relation to each distinct sequence. The thin solid lines labeled by I to III in the upper panel are the period- $J - K$ colour relations for Mira variables obtained by Feast et al. (1989) and Whitelock et al. (2000) (see text). The insets show the closeup of less regularly pulsating variables.

Most of the stars from group *D* are found at the place a bit brighter than the TRGB. On the other hand, some of the *D* variables are found in very red ($J - H > 1.8$, $H - K > 1.5$ and $J - K > 3.2$) and relatively bright area ($11.0 \lesssim K \lesssim 12.5$), where usually the carbon-rich stars and dusty AGB stars dominate (Nikolaev & Weinberg 2000). A possible explanation for these red stars in group *D* is that, these stars could in fact be very dusty AGB stars that are subjected to severe circumstellar extinctions, even in the K band. Whitelock et al. (2003) studied obscured variables in the LMC and showed that the K magnitudes of obscured stars cover a wide range at a given pulsation period. For instance, their data indicates that the difference can be more than two magnitudes in the case of the carbon-rich stars. It is interesting to note that their red stars (those with long periods and K magnitudes fainter than ~ 11 mag) clearly locate the brighter part of sequence *D* on the period- K magnitude plane.

3.3 Period-colour diagram

Figure 4 shows the $J - K$ colours of OGLE variables in the Magellanic clouds as a function of their periods. The insets are the close-ups for the sake of clarity. A general trend that the colours get redder with increasing period can be seen.

The dashed lines are the least-square fit of a linear relation to the Cepheid variables (*F* and *G* stars), which yields the period- $J - K$ colour relations of

$$J - K[\text{mag}] = 0.147(\pm 0.010) \log P[\text{days}] + 0.248(\pm 0.006) \quad (1)$$

for the LMC ($\sigma = 0.062$) and

$$J - K[\text{mag}] = 0.167(\pm 0.009) \log P[\text{days}] + 0.272(\pm 0.004) \quad (2)$$

for the SMC ($\sigma = 0.074$), respectively. The quoted errors are the 1σ errors of each coefficient. Although the fundamental (group *F*) and first overtone (group *G*) Cepheids are clearly separated in the period-*K* magnitude plane, we don't see any significant difference between them in this diagram. This means, if we compare the fundamental and first overtone Cepheids with the same pulsation periods, the stellar radii of first overtone Cepheids should be larger than those of fundamental Cepheids because their stellar temperatures are nearly the same but the first overtone Cepheids are brighter than the fundamental ones. Theories predict exactly the same conclusion (e.g., Bono et al. 2001), and if the stellar radius of the Cepheid variables could be measured accurately, it will help to identify the pulsation modes.

The less regularly pulsating red giants (group $A^\pm$ and $B^\pm$) also follow a tight sequence on the period-*J* - *K* colour plane. The thick solid lines in the figure indicate the least-square fit of a linear relation to the stars from the four groups, whose *J* - *K* colours and periods are within the ranges, $0.7 < J - K < 1.3$ and $1.2 < \log P < 2.1$. Their period-*J* - *K* colour relations are also calculated and they are,

$$J - K[\text{mag}] = 0.231(\pm 0.004) \log P[\text{days}] + 0.671(\pm 0.007). \quad (3)$$

for the LMC ($\sigma = 0.058$) and

$$J - K[\text{mag}] = 0.272(\pm 0.011) \log P[\text{days}] + 0.560(\pm 0.019). \quad (4)$$

for the SMC ($\sigma = 0.068$), respectively.

It is obvious that most of the *C* and *C'* stars in the LMC, and also some B^+ stars and most of *C* stars in the SMC do not follow the extension of these relations. In Paper I, we concluded that stars on sequences *C* and *C'* are Mira variables pulsating in the fundamental and first overtone mode respectively, putting an end to the longstanding problem over the pulsation mode of Mira variables. Figure 4 clearly shows that *C* and *C'* stars behave very similarly on the period-*J* - *K* colour plane. This corroborates the idea that the *C* and *C'* stars are the same type of stars (i.e., Miras), but only different in the pulsation modes. Also, it is likely that the "some B^+ stars" in the SMC are the counterparts of the *C'* stars in the LMC.

In earlier work, Feast et al. (1989), Glass et al. (1995) and Whitelock, Marang & Feast (2000) obtained period-*J* - *K* colour relations of Mira variables in the LMC, Sgr I Baade window of the Galactic Bulge and solar neighborhood, respectively. These relations are all based on the SAAO system. The atmospheres of Mira variables are rather complex, making their quantitative comparisons difficult between different filter systems. Thus here we make only qualitative discussions. Just to get a rough idea, we indicated in the upper panel of figure 4 the period-*J* - *K* colour relations obtained by Feast et al. (1989) for carbon- (labeled I) and oxygen-rich (labeled II) Miras and by Whitelock et al. (2000) (labeled III) after referring to the LCO system by assuming $(J - K)_{\text{LCO}} = 0.928(J - K)_{\text{SAAO}} - 0.004$ (Carpenter 2001). Of course we can not tell exactly which of our stars are carbon- or oxygen-rich, but it is statistically fair to say that stars with *J* - *K* colours redder than 1.4 are primarily carbon-rich stars (Nikolaev & Weinberg 2000). In this point of view, we confirmed the result of Feast et al. (1989) that oxygen- and carbon-rich Miras follow different period-*J* - *K* colour relations.

Now that we see the *C* and *C'* stars separate on the period-




Figure 5. Period- θ diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel).

J - *K* colour plane, it will be interesting to search for the Galactic counterpart of the first overtone Miras (*C'* stars) by using this tool. Then we can redetermine their distances using their exclusive period-*K* magnitude relation, which we obtained in Paper I. It will help to understand the differences between the fundamental and first overtone Miras through more detailed studies that are technically difficult in the Magellanic Clouds (i.e., spectroscopic work and mid-infrared observations etc.). Because not only *C* and *C'* stars but also the other variables separate on the period-*J* - *K* colour plane, the same argument is true for them, such as *D* stars.

However, before applying this tool to different environments, we should know whether it depends on chemical abundance. Wray, Eyer & Paczynski (2003) showed that small amplitude red giant variables in the Galactic Bar can be separated into two groups, and they follow different period-colour relations. Also, they suggest that their *A* and *B* stars correspond to type A^- and B^- variables, respectively of Paper I. The insets of the figure 4 clearly confirm their result, showing that A^- and B^- are well separated on the period-*J* - *K* color plane. This infers that the period-*J* - *K* relations we derived in equations 3 and 4 are merely the general trends of the variable red giants, and in fact there are exclusive period-colour relations for each population. Rejkuba et al. (2003) studied infrared properties of long period variables in Cen A. Although they did not mention it, it is likely that there are two groups in their period-colour diagrams (see their figure 10). We suggest that their short-period ($\log P \sim 2.45$) and long-period ($\log P \sim 2.65$) groups could correspond to type *C'* and *C* stars in the LMC, respectively. These facts strengthen the idea that, period-colour diagram is a good and universal tool, at least to tell variability type.

Lebzelter et al. (2002) studied the AGAPEROS/DENIS variables in the LMC and showed that so-called Semi-regular variables




Figure 6. Period-amplitude(ΔI) diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel).

follow a period-colour relation but the “regular” variables do not (see their figure 12). To see the regularity of the light variation quantitatively, we show the relationship between the statistic parameter θ and period in figure 5. The figure clearly confirmed the Lebzelter’s suggestion and shows that stars that follow period- $J-K$ relation ($A^\pm$ and $B^\pm$ stars with $\log P \lesssim 2.1$) are less regularly pulsating, and those that don’t (B^+ stars with $\log P \gtrsim 2.1$, C' and C stars) are regularly pulsating.

3.4 Period-amplitude (ΔI) and colour-amplitude diagram

Figure 6 is a plot to show the relationship between the pulsation period and amplitude (ΔI). The pulsation amplitude is determined from the OGLE’s I band data, given as $\Delta I = I_{\max} - I_{\min}$. Note that all of the original OGLE light curves were carefully eye-inspected, and we calculated the ΔI after eliminating the obvious spurious data points.

The figure shows that the fundamental Cepheids show larger amplitude variations than the first overtone Cepheids. Also, the upper envelope of Cepheid amplitude goes down with increasing period. This trend contrasts markedly with that of red giants, because their pulsation amplitudes generally go up with increasing period.

Variables stars with K magnitudes below the TRGB (A^- and B^- stars) pulsate with very small amplitude. With a few exceptions, most of the stars from D group also pulsate with relatively small amplitude. The large amplitude pulsation ($\Delta I \gtrsim 0.9$) seems to occur only among the long-period ($\log P \gtrsim 2.3$) variables. If one had to separate Semi-regular from Mira variables based on pulsation amplitudes, the threshold would be about $\Delta I \approx 0.9$, because one can see a discernible gap around there. However, we again emphasize that this type of criteria could be artificial.




Figure 7. Colour-amplitude(ΔI) diagram of OGLE variables in the LMC (upper panel) and the SMC (lower panel).

Figure 7 shows the relationship between the $J-K$ colour and pulsation amplitude (ΔI). The variable red giants show the clear trend that the amplitude get larger with increasing colour. According to Nikolaev & Weinberg (2000), obscured carbon-rich AGB stars have the $J-K$ colours redder than about 2.0 mag. Almost all of such dusty carbon-rich stars in the Magellanic Clouds have large pulsation amplitudes. Some of the relatively blue stars ($0.8 \lesssim J-K \lesssim 1.2$) on the sequence C also show large amplitude pulsation. Judging from their $J-K$ colour and being on the sequence C , they are likely to be oxygen-rich Mira variables. It is remarkable that the carbon-rich Miras tend to have greater I -band amplitudes at redder $J-K$ colour, but the amplitudes of oxygen-rich Miras are almost independent of the colour. This might reflect the difference in the molecular opacities at I -band between the two groups.

3.5 Similarities and differences between the LMC and SMC samples

In a large sense, variable stars in the Magellanic Clouds behave similarly in the figures we discussed so far. However, on closer view, there are some differences between variable stars in the two galaxies. The first thing to note is that we can see a deficiency of short period (hence faint) Cepheids in the LMC. This is not the selection effect. The comparison studies of Cepheids in various environments (e.g., Antonello, Fugazza & Mantegazza 2002) can shed light on this matter, but this issue is beyond the scope of the present paper. Next thing to note is that, especially in the period-colour diagram, we see smaller number of oxygen-rich Mira variables in the SMC (counterpart of LMC Miras with $J-K$ colours bluer than 1.4) compared to the LMC. This is probably because the difference in

average chemical abundance and/or in age between the two galaxies.

4 SUMMARY

We presented the data of 8,852 and 2,927 variable stars in the Large and Small Magellanic Clouds. Based on these data, we discussed the infrared properties of the variable stars. We showed that period- $J - K$ colour diagram is a good tool to tell the pulsation modes of the Mira variables. A follow-up campaign searching for the Galactic counterpart of the first overtone mode Miras would tell us the difference between them and fundamental mode Miras.

ACKNOWLEDGMENTS

We are grateful to people involved in OGLE project for making their data so easy to access and use. We would like to thank Dr. Michael Feast and Dr. Ian Glass for the valuable and helpful comments on the manuscript. This research is supported in part by the Grant-in-Aid for Scientific Research (C) No. 12640234 and Grant-in-Aids for Scientific on Priority Area (A) No. 12021202 and 13011202 from the Ministry of Education, Science, Sports and Culture of Japan. The IRSF/SIRIUS project was initiated and supported by Nagoya University, National Astronomical Observatory of Japan and University of Tokyo in collaboration with South African Astronomical Observatory under a financial support of Grant-in-Aid for Scientific Research on Priority Area (A) No. 10147207 of the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

REFERENCES

- Alard C. et al., 2001, *ApJ*, 552, 289
 Antonello E., Fugazza D., Mantegazza L., 2002, *A&A*, 388, 477
 Bedding T. R., Zijlstra Albert A., Jones A., Foster G., 1998, *MNRAS*, 301, 1073
 Bono G., Gieren W. P., Marconi M., Fouqué, 2001, *ApJ*, 552, L141
 Carpenter J. M., 2001, *AJ*, 121, 2851
 Cioni M.-R. L., Marquette J.-B., Loup C., Azzopardi M., Habing H. J., Lasserre T., Lesquoy E., 2001, *A&A*, 377, 945
 Feast M. W., Glass I. S., Whitelock P. A., Catchpole R. M., 1989, *MNRAS*, 241, 375
 Glass I. S., Whitelock P. A., Catchpole R. M., Feast M. W., 1995, *MNRAS*, 273, 383
 Ita Y. et al., 2002 *MNRAS*, 337, L31
 Ita Y. et al., 2003 *MNRAS*, in press. [astro-ph/0310083] (Paper I)
 Koornneef J., 1982, *A&A*, 107, 247
 Lebzelter T., Schultheis M., Melchior A. L., 2002, *A&A*, 393, 573
 Nagashima C. et al., 1999, in Nakamoto T., ed., *Star formation 1999*, Nobeyama Radio Observatory, p. 397
 Nagayama T. et al., 2002, *Proc. SPIE*, 4841, 459
 Nikolaev S., Weinberg M. D., 2000, *ApJ*, 542, 804
 Persson S. E., Murphy D. C., Krzeminski W., Roth M., Rieke M. J., 1998, *AJ*, 116, 2475
 Rejkuba M., Minniti D., Silva D. R., Bedding T. R., 2003, *A&A*, in press. [astro-ph/0309358]
 Stellingwerf B. F., 1978, *ApJ*, 224, 953
 Udalski A., Kubiak M., Szymański M., 1997, *AcA*, 47, 319
 Udalski A., Szymański M., Kubiak M., Pietrzyński G., Soszyński I., Woźniak P., Żebruń K., 1999, *AcA*, 49, 201
 Westerlund B. E., 1997, *The Magellanic Clouds*. Cambridge Univ. Press, Cambridge
 Whitelock P. A., Marang F., Feast M. W., 2000, *MNRAS*, 319, 728
 Whitelock P. A., Feast M. W., van Loon J. Th., Zijlstra A. A., 2003, *MNRAS*, 342, 86
 Wray J. J., Eyer L., Paczynski B., 2003, *MNRAS*, submitted. [astro-ph/0310578]
 Żebruń K., et al., 2001, *AcA*, 51, 317