THE DEEP2 GALAXY REDSHIFT SURVEY: EVOLUTION OF CLOSE GALAXY PAIRS AND MAJOR-MERGER RATES UP TO z $\sim 1.2^{-1}$

LIHWAI LIN ^{2,3}, DAVID C. KOO ³, CHRISTOPHER N. A. WILLMER ^{3,4}, DAVID R. PATTON ⁵, CHRISTOPHER J. CONSELICE ⁶, RENBIN YAN ⁷, ALISON L. COIL ⁷, MICHAEL C. COOPER ⁷, MARC DAVIS ^{7,8}, S. M. FABER ³, BRIAN F. GERKE ⁸, PURAGRA GUHATHAKURTA ³, AND JEFFREY A. NEWMAN ⁹

ABSTRACT

We derive the close, kinematic pair fraction and merger rate up to redshift $z \sim 1.2$ from the initial data of the DEEP2 Redshift Survey. Assuming a mild luminosity evolution, the number of companions per luminous galaxy is found to evolve as $(1+z)^m$, with $m=0.51\pm0.28$; assuming no evolution, $m=1.60\pm0.29$. Our results imply that only 9% of present-day L^* galaxies have undergone major mergers since $z \sim 1.2$ and that the average major merger rate is about 4×10^{-4}

¹ Some of the data presented herein were obtained at the W. M. Keck Observatory, which is operated as a scientific partnership among the California Institute of Technology, the University of California, and the National Aeronautics and Space Administration. The Observatory was made possible by the generous financial support of the W. M. Keck Foundation.

²Department of Physics, National Taiwan University, No. 1, Sec. 4, Roosevelt Road, Taipei 106, Taiwan; d90222005@ntu.edu.tw

³University of California Observatories/Lick Observatory, Department of Astronomy and Astrophysics, University of California at Santa Cruz, 1156 High Street, Santa Cruz, CA 95064.

⁴On leave from Observatório Nacional, Brazil.

⁵Department of Physics, Trent University, 1600 West Bank Drive, Peterborough, ON K9J 7B8, Canada.

⁶Department of Astronomy, California Institute of Technology, Pasadena, CA 91125.

⁷Department of Astronomy, University of California at Berkeley, 601 Campbell Hall, Berkeley, CA 94720-3411.

⁸Department of Physics, University of California at Berkeley, 366 LeConte Hall, Berkeley, CA 94720-7300.

⁹Hubble Fellow; Institute for Nuclear and Particle Astrophysics, Lawrence Berkeley National Laboratory, Berkeley, CA 94720.

 $h^3~{\rm Mpc^{-3}~Gyr^{-1}}$ for $z\sim 0.5-1.2$. Most previous studies have yielded higher values.

Subject headings: galaxies:interactions - galaxies:evolution - large-scale structure of Universe

1. INTRODUCTION

Galaxy interactions and mergers are an integral part of our current paradigm of the hierarchical formation and evolution of galaxies. Such processes are expected to affect the morphologies, gas distributions, and stellar populations of galaxies (e.g., Mihos & Henquist 1994, 1996; Dubinski, Mihos, & Henquist 1996). Although mergers are rare today (Patton et al. 2000, hereafter P2000), cold dark matter N-body simulations show merger rates of halos increasing with redshift as $(1+z)^m$, with $2.5 \le m \le 3.5$ (Governato et al. 1999; Gottlöber, Klypin, & Kravtsov 2001).

Observations, however, have yielded results with $0 \le m \le 4$ (Zepf & Koo 1989; Burkey, Keel, & Windhorst 1994; Carlberg et al. 1994; Woods, Fahlman, & Richer 1995; Yee & Ellingson 1995; Patton et al. 1997; Neuschaefer et al. 1997; Le Fèvre et al. 2000; Carlberg et al. 2000; Patton et al. 2002; Bundy et al. 2004). The diverse results are likely due to different pair criteria, observational techniques, selection effects, and cosmic variance (Abraham 1999; P2000). To identify close pairs, the most secure method is via spectroscopic redshifts for both galaxies to find kinematic pairs (Patton et al. 2002, hereafter P2002). This Letter adopts the approach of P2002 and uses the early data of the DEEP2 (Deep Extragalactic Evolutionary Probe 2) Redshift Survey (Davis et al. 2003; S. M. Faber et al. 2004, in preparation) to derive the pair fraction and merger rates up to z = 1.2. In §2, we describe the sample and selection functions. In §3, we use both the projected separation and the relative velocity to select close galaxy pairs and to determine their evolution. Major merger rates out to $z \sim 1.2$ are computed in §4. The results are discussed in §5. Throughout this Letter, we adopt a cosmology of $H_0 = 70$ km s⁻¹ Mpc⁻¹, $\Omega_m = 0.3$, and $\Omega_{\Lambda} = 0.7$.

2. DATA AND SELECTION FUNCTIONS

The DEEP2 Redshift Survey (DEEP2 for short) will measure redshifts for $\sim 50,000$ galaxies at $z \sim 1$ (Davis et al. 2003) using DEIMOS (DEep Imaging Multi-Object Spectrograph; Faber et al. 2003) on the 10 m Keck II telescope. The survey convers four fields, with Field 1 (Extended Groth Strip) being a strip of $0.25 \times 2 \text{ deg}^2$ and Fields 2, 3, and 4

each being $0.5 \times 2 \text{ deg}^2$. The photometry is based on BRI images taken with the $12k \times 8k$ camera on the Canada-France-Hawaii Telescope (see Coil et al. 2004 for details). Galaxies are selected for spectroscopy using a limit of $R_{AB} = 24.1$ mag. Except in Field 1, a two-color cut is also applied to exclude galaxies with redshifts z < 0.75. A 1200 line mm⁻¹ grating (R ~ 5000) is used with a spectral range of 6400 - 9000 Å, where the [O II] $\lambda 3727$ doublet would be visible at $z \sim 0.7 - 1.4$. The data are reduced using an IDL pipeline developed at UC-Berkeley (J. A. Newman et al. 2004, in preparation). The K-correction is derived from spectra of local galaxies (C. N. A. Willmer et al. 2004, in preparation). The data used here are from Fields 1 and 4, cover $\sim 0.4 \text{ deg}^2$, and contain ~ 5000 galaxies.

To measure the spectroscopic selection functions (see L. Lin et al. 2004, in preparation, for details), we compared the sample with successful redshifts to galaxies in the full photometric catalog that satisfy the limiting magnitude and two-color cuts. Following analogous approaches by Yee, Ellingson, & Carlberg (1996) and P2002, we calculate the spectroscopic weight w of each galaxy as 1/S, where S is the spectroscopic selection function derived from the R magnitude, B-R and R-I colors, R—band surface brightness μ_R , and local galaxy density of the galaxy itself. To correct for bias due to slit collisions (Davis, Gerke, & Newman 2004), we also compute the angular weight, $w(\theta)$, as a function of angular separation θ (P2002). For $\theta \leq 20$ ", we find $0.95 \leq w(\theta) \leq 1.5$, a result confirmed with the DEEP2 mock catalog of Yan, White, & Coil (2004).

3. PAIR STATISTICS

A major limitation on the direct measurement of merger fractions for galaxies is the difficulty in identifying on-going mergers, especially for distant galaxies. One alternative is to count only the pairs with projected separations Δr and relative line-of-sight heliocentric velocities Δv less than r_{max} and v_{max} , respectively. A large fraction of pairs with physical separations less than 20 h^{-1} kpc and velocity difference less than 500 km s⁻¹ appear to have disturbed morphologies or signs of interactions, and these galaxies are expected to merge within 0.5 Gyr (P2000). In our work, close pairs are defined such that their projected separations satisfy 10 h^{-1} kpc $\leq \Delta r \leq r_{max}$ and their rest-frame relative velocities Δv are less than 500 km s⁻¹. We adopt an inner cutoff with a projected distance 10 h^{-1} so as to avoid the ambiguity between very close pairs and single galaxies with multiple star-forming knots. We choose values of $r_{max} = 30$, 50, and 100 h^{-1} kpc, where 30 h^{-1} kpc is most likely to include genuine merger pairs, while the two larger separations provide larger samples and thus better statistics. To ensure the selection of the same types of galaxies at different redshifts in the presence of luminosity evolution, P2002 adopted a specific range

in evolution-corrected absolute magnitude M_B^e , defined as $M_B + Qz$, where the evolution is parameterized as M(z) = M(z=0) - Qz. Following P2002, we adopt Q=1 as a primary model, and we also study the effect of different models on the pair statistics. We restrict the analysis to galaxies with luminosities $-21 \le M_B^e \le -19$ for z=0.45-1.2. Since there is only a 2 mag range in our sample, and assuming a constant M/L ratio, most observed pairs are thus major mergers, i.e., mass ratios between 1: 1 and 6: 1. Using data from the DEEP2 Fields 1 and 4 helps us reduce the effects of cosmic variance. Following P2000 and P2002, we compute the average number of companions per galaxy

$$N_c = \frac{\sum_{i=1}^{N_{tot}} \sum_j w_j w(\theta)_{ij}}{N_{tot}},\tag{1}$$

where N_{tot} is the total number of galaxies within the chosen magnitude range, w_j is the spectroscopic weight for the jth companion belonging to the ith galaxy, and $w(\theta)_{ij}$ is the angular weight for each pair as described in Section 2. Details of the weighting scheme can be found in Section 5 of P2002 and references therein. This quantity N_c is similar to the pair fraction when there are few triplets or higher order N-tuples in the sample, which is the case here. In this work, N_c will sometimes simply be referred to as the pair fraction.

Figure 1 shows N_c versus redshift z for a sample with $v_{max} = 500 \text{ km s}^{-1}$ and $10 h^{-1} \text{ kpc}$ $\leq \Delta r \leq r_{max}$ with $r_{max} = 30$, 50, and $100 h^{-1}$ kpc, respectively. In the case of using $r_{max} = 50 h^{-1}$ kpc, we find 79 paired galaxies out of 2547 galaxies. The derived N_c is $\sim 8\%$ at $z \sim 0.6$ and increases to 10% at $z \sim 1.1$. Figure 1 also shows results from the SSRS2 (Southern Sky Redshift Survey; P2000) at $z \sim 0.015$ and CNOC2 (Canadian Network for Observational Cosmology; P2002) at $z \sim 0.3$, after corrections that adopt the same cosmology and the same luminosity range.

The DEEP2 sample is R-band-selected and hence, for redshifts greater than 0.8, is biased toward galaxies that are bright in the rest-frame UV, i.e., against faint red galaxies, especially at redshifts z > 1.0. To avoid this bias, we divide our sample into blue and red galaxies using the color bimodality feature and then repeat the calculation of N_c using $r_{max} = 50 \ h^{-1}$ kpc for blue galaxies at z = 0.3 (CNOC2), 0.6, and 0.85 bins (DEEP2). When using only blue galaxies, the values of the pair fraction N_c^{blue} decrease at z = 0.3 and 0.85 but increase at z = 0.6 compared to the results using the original sample. The changes in the measured value of N_c could be up to a factor of 2, depending on the field and redshift. To mitigate the underestimation of the pair fraction in the redshift bin 1.0 < z < 1.2, we calculate a corrected N_c at z = 1.1 as

$$N_c^{cor} = \max[N_c, 2N_c^{blue}], \tag{2}$$

shown as open triangles in Figure 1. Parameterizing the evolution of N_c as $Nc(0)(1+z)^m$, we find that the best fit of $(N_c(0), m)$ is $(0.029 \pm 0.005, 1.08 \pm 0.40)$ with $\chi^2 = 0.27$ for

 $r_{max}=30\ h^{-1}\ \mathrm{kpc},\ (0.068\pm0.008,0.51\pm0.28)$ with $\chi^2=0.89$ for $r_{max}=50\ h^{-1}$ kpc, and $(0.177\pm0.014,0.47\pm0.18)$ with $\chi^2=1.2$ for $r_{max}=100\ h^{-1}$ kpc. For the highest redshift bin, N_c^{cor} instead of N_c is used for the above fitting procedure. The best fits are shown as long-dashed curves in Figure 1. For three different choices of r_{max} , we find only a small increase of N_c from redshifts z=0 to z=1.2. In figure 1, we also list the upper limits of errors from the cosmic variance for each redshift bin calculated by assuming a spherical geometry of the survey volume. Incorporating the cosmic variance into the fitting procedure slightly raises m to 0.62 ± 0.49 for $r_{max}=50\ h^{-1}$ kpc.

The adopted choice of Q=1 in the luminosity evolution model sets the luminosity range of the sample at each redshift and thus affects our results. To assess this effect, we repeat the pair analysis for other choices of Q. The luminosity ranges are chosen such that they are locked to $-22 \le M_B \le -20$ at z=1. Using $r_{max}=50$ h^{-1} kpc and an upper redshift limit of z=1, we find that m varies from 1.60 ± 0.29 for Q=0, 0.86 ± 0.29 for Q=0.5, 0.41 ± 0.30 for Q=1, to -0.24 ± 0.35 for Q=2. In figure 2, we plot N_c as a function of absolute B-band magnitude (M_B) for three redshift samples. A clear dependence of N_c on M_B is evident. The trend that m decreases with Q can be understood as the result of including fainter galaxies and thus having larger N_c at lower redshifts when adopting higher Q values. Nevertheless, even with the lowest choice of Q=0 to maximize m, models with m>3.5 are ruled out at a 3 σ level of confidence for all three choices of separation limits.

4. MAJOR MERGER RATES

The comoving merger rate, usually defined as the number of mergers per unit time per comoving volume, can be estimated as

$$N_{mg} = 0.5n(z)N_c(z)C_{mg}T_{mg}^{-1}, (3)$$

where T_{mg} is the time-scale for physically associated pairs to merge, C_{mg} denotes the fraction of galaxies in close pairs that will merge in T_{mg} , and n(z) is the comoving number density of galaxies. The factor 0.5 is to convert the number of galaxies into the number of merger events. The merger rate calculated above, however, is not suitable for comparison to the merger rate derived from morphological approaches. The reason is that while we have restricted the luminosity range of companions to compute N_c , the morphological approach does not. To correct for this restriction, equation (3) can be modified:

$$N_{mg} = (0.5 + G)n(z)N_c(z)C_{mg}T_{mg}^{-1},$$
(4)

where the added parameter G accounts for the excess number of companions failing to fall into our sample. Assuming that the maximum mass ratio needed to yield significant

morphology distortions is 4: 1, we calculate the value of G as 1.24 for $-21 \le M_B^e \le -19$ using the local luminosity function (Driver & De Propris 2003). T_{mg} depends on the relative mass ratio, dynamical orbit, and detailed structure of the two merging galaxies, and thus it varies from case to case. Here we adopt $T_{mg} = 0.5$ Gyr, a value suggested by N-body simulations and simplified models (Mihos 1995; P2000). The value C_{mg} is approximately 0.5 for close pairs with 5 h^{-1} kpc $\le \Delta r \le 20$ h^{-1} kpc and $v_{max} = 500$ km s⁻¹; this estimate is based on morphological studies of local close pairs (P2000). Nevertheless, both T_{mg} and C_{mg} remain uncertain.

The derived merger rates are displayed in Figure 3. Here we have applied the pair fraction evolution of m=0.51 derived for the $r_{max}=50$ h^{-1} kpc case to the $z\sim 0$ (SSRS2) N_C result for pairs with 5 h^{-1} kpc $\leq \Delta r \leq 20$ h^{-1} kpc. The parameter n(z) is calculated as the sum of the number of galaxies in the adopted magnitude range, each weighted by its spectroscopic weight and divided by the comoving volume occupied by the included sources. The errors shown for DEEP2 measurements represent 40% variations that are typical for close pair counts in our sample, and they do not include the uncertainties of T_{mg} and C_{mg} . Also shown are the results from Conselice et al. (2003), who relied on morphologically identified mergers for $M_B < -19$ using Hubble Space Telescope (HST) data in the Hubble Deep Field-North (filled triangles). Clearly, the comoving merger rate in our data changes little with redshift. From $z\sim 0.5$ to $z\sim 1.2$, the average is $N_{mg}\sim 4\times 10^{-4}~h^3$ Mpc⁻³ Gyr⁻¹ for $-21\leq M_B^e\leq -19$. This value is about 1 order of magnitude lower than the average merger rate for galaxies with $M_B\leq -19$ derived by Conselice et al. (2003).

Finally, we calculate the merger remnant fraction, f_{rem} , defined as the fraction of present-day galaxies that have undergone major mergers (P2000). Adopting the merger fraction C_{mg} to be 0.5 and T_{mg} to be 0.5 Gyr, we estimate, using equation (32) in P2000, that about 9% of present L^* luminous galaxies have undergone major mergers since $z \sim 1.2$.

5. DISCUSSION

Studies up to $z \sim 1$ using pair counts have found a wide range in the evolution of merger rates. For example, based on roughly 300 Canada-France Redshift Survey galaxies measured with HST, Le Fèvre et al. (2000) concluded that the pair fraction evolves with $m = 2.7 \pm 0.6$ while the fraction of merger candidates evolves with $m = 3.4 \pm 0.6$. In contrast, Carlberg et al. (2000) measured the mean fractional pair luminosity from z = 0.2 to 1 and found no evolution with $m \sim 0 \pm 1.4$, consistent with our result. Since the pair fraction depends on adopted luminosity limits, differences among studies may depend on the choice of luminosity evolution models. Le Fèvre et al. (2000), e.g., applied no corrections while Carlberg et al.

(2000) adopted the Q=1 model, as we do here. The key advantages of our pair sample over previous surveys include having measurements at z > 1, a larger sample, and more restrictive pair criteria than those adopted by Le Fèvre et al. (2000) and Carlberg et al. (2000).

The discrepancy in the derived merger rates at $z \sim 1$ between our estimates and those from Conselice et al. (2003) is more difficult to reconcile. Besides the choice of luminosity ranges and the uncertainties in C_{mq} and T_{mq} , other factors may explain the discrepancy. First, cosmic variance can always come into play, since most morphological studies have been forced to use the few small fields covered deeply by HST imaging, whereas pair-count surveys cover much larger areas. Second, the two approaches may sample interacting and merging galaxies at different stages, with morphological approaches identifying very advanced mergers and merger remnants, while the pairs detect some of the same systems before distortions are discernible. Matching the derived merger rates from these two approaches requires knowledge of the precise timescales of close pairs and of the duration time for the appearance of distorted morphologies. However, neither is well understood yet. Third, while our close pairs identify only major mergers, morphological criteria may be sensitive to minor mergers as well. Finally, both the pair-count and morphology methods are subject to systematic uncertainties that require detail modeling (Bell 2004). Given the current discrepancy in results, it behooves us to study the connection between kinematic pairs and morphologically disturbed galaxies at various redshifts, both via observational approaches (P2000; J. Lotz et al. 2004, in preparation) and through numerical simulations (T. J. Cox et al. 2004, in preparation).

The merger rates from pair counts are also an excellent test for those estimated from N-body simulations, since the former reveal the behavior of luminous baryons while the latter reflect the behavior of dark halos. Gottlöber et al. (2001) defined merger rates in N-body simulations as equivalent to the major merger fraction per gigayear by tracing the formation and evolutionary history of each halo. These fractions will differ from the pair fractions roughly by a constant, and therefore their redshift evolution is amenable to direct comparisons. They found merger rates evolving as $(1+z)^3$ up to $z\sim 2$. This theoretical value of m=3 is significantly higher than the $m\sim0.5$ we find from observed pair counts. We should, however, be wary about this comparison for the following reasons. First, definitions of merger rates in observations and simulations may not be consistent with each other, since the halos and visible galaxies span different size scales and since N-body simulations also suffer limitations in resolution. Second, the merger rates/fractions of halos are likely to be a function of halo mass, which is suggested by our finding that the galaxy pair fraction is a function of luminosity. Pair count works using K-band-selected samples (e.g., Bundy et al. 2004) can also provide another avenue for testing the mass dependence of merger rates since the K-band luminosity is more representative of any underlying stellar mass and suffers less from evolution corrections. Although preliminary, mock catalog simulations by E. Van Kampen et al. (2004, in preparation) predict flat slopes for the pair-count fractions (see Fig. 3 of Bell 2004), just as seen in our observations. We expect a dramatic improvement in our understanding of merger histories via pair counts after more realistic comparisons to simulations are possible and especially after a 10-fold increase in sample size when DEEP2 is complete in 2005.

We thank the referee for helpful comments, J. Primack, J. Lotz, and B. Allgood for useful discussions, and the Keck staff for dedicated assistance. L. L. acknowledges support from Taiwan via the COSPA project and NSC grant NSC92-2112-M-002-021, D. R. P. from NSERC of Canada, and the DEEP2 team from NSF grants AST00-71048, AST00-71198, and KDI-9872979. We close with thanks to the Hawaiian people for allowing us to use their sacred mountain.

REFERENCES

Abraham, R. G. 1999, IAU Symp. 186, Galaxy Interactions at Low and High Redshifts (astro-ph/9802033)

Bundy, K., Fukugita, M., Ellis, R. S., Kodama, T., & Conselice, C. J. 2004, ApJ, 601, L123

Bell, E. F. 2004, preprint (astro-ph/0408023)

Burkey, J. M., Keel, W. C., & Windhorst, R. A. 1994, ApJ, 429, L13

Carlberg, R. G., Pritchet, C. J., & Infante, L. 1994, ApJ, 435, 540

Carlberg, R. G., et al. 2000, ApJ, 532, L1

Coil, A. L., et al. 2004, ApJ, submitted (astro-ph/0403423)

Conselice, C. J., Bershady, M. A., Dickinson, M., & Papovich, C. 2003, AJ, 126, 1183

Davis, M., et al. 2003, SPIE, 4834, 161 (astro-ph/0209419)

Davis, M., Gerke, B., & Newman, J. A. 2004, preprint(astro-ph/0408344)

Driver, S., & DePropris, R. 2003, Ap&SS, 285, 175

Dubinski, J., Mihos, J. C., & Hernquist, L. 1996, ApJ, 462, 576

Faber, S. M., et al. 2003, SPIE, 4841, 1657

Gottlöber, S., Klypin, A., & Kravtsov, A. V. 2001, ApJ, 546, 223

Governato, F., Gardner, J. P., Stadel, J., Quinn, T., & Lake, G. 1999, AJ, 117, 1651

Le Fèvre, O., et al. 2000, MNRAS, 311, 565

Mihos, J. C., & Hernquist, L. 1994, ApJ, 425, L13

Mihos, J. C. 1995, ApJ, 438, L75

Mihos, J. C., & Hernquist, L. 1996, ApJ, 464, 641

Neuschaefer, L. W., Im, M., Ratnatunga, K. U., Griffiths., R. E., & Casertano, S. 1997, ApJ, 480, 59

Patton, D. R., Pritchet., C. J., Yee, H. K. C., Ellingson, E., & Carlberg, R. G. 1997, ApJ, 475, 29

Patton, D. R., Carlberg, R. G., Marzke, R. O., Pritchet., C. J., da Costa, L. N., & Pellegrini,
 P. S. 2000, ApJ, 536, 153

Patton, D. R., et al. 2002, ApJ, 565, 208

Woods, D., Fahlman, G. G., & Richer, H. B. 1995, ApJ, 454, 32

Yan, R., White, M., Coil, A. L. 2004, ApJ, 607, 739

Yee, H. K. C., & Ellingson, E. 1995, ApJ, 445, 37

Yee, H. K. C., Ellingson, E., & Carlberg, R. G. 1996, ApJS, 102, 269

Zepf, S. E., & Koo, D. C. 1989, ApJ, 337, 34

This preprint was prepared with the AAS LATEX macros v5.2.




Fig. 1.— Pair fraction, N_c , as a function of redshift z. The filled squares represent measurements from DEEP2 Field 1, and the open squares represent those from Field 4. The crosses mark the results from SSRS2 and CNOC2. The open triangles show the corrected pair fraction N_c^{cor} for the highest redshift bin. The long-dashed curves are best fits by the form $(1+z)^m$, using DEEP2, SSRS2, and CNOC2 data (see text); the dot-dashed curves represent evolution with m=2.5 for reference. From the top to the bottom, $r_{max}=100$ h^{-1} kpc, $r_{max}=50$ h^{-1} kpc, and $r_{max}=30$ h^{-1} kpc. Note that the bottom panel has a different vertical scale. The error bars shown in the plot and used for fitting are calculated by bootstrapping. The numbers appearing in the bottom panel indicate the upper limits of errors (in units of percentage) from the cosmic variance for each redshift bin. Those in the parentheses are for open symbols.




Fig. 2.— Pair fraction, N_c , as a function of absolute B-band magnitude (M_B) using $r_{max} = 50$ h^{-1} kpc for different redshift samples as indicated. For each data point, N_c is calculated using galaxies with absolute B-band magnitude between $M_B - 1$ and $M_B + 1$.




Fig. 3.— Comoving volume merger rate, N_{mg} , as a function of redshift z. Estimates from the DEEP2 fields using equation (4) are marked with symbols as indicated; the data from Conselice et al. (2003) are represented by the filled triangles.