

Resolving the Steep Spectrum Emission in the Central Radio Source in ZwCl 0735.7+7421

A. S. Cohen¹, T. E. Clarke^{1,2}, L. Feretti³ and N. E. Kassim¹

ABSTRACT

It has long been known that an extremely steep spectrum radio source lies at the center of ZwCl 0735.7+7421, a galaxy cluster with high X-ray luminosity and a cooling core. In this Letter, we present VLA observations of this radio source at both 1425 and 325 MHz. With a resolution below $21''$ (75 kpc) for both 1425 and 325 MHz, we show the morphology of the central source, and find that it is most likely a large (400 kpc) radio galaxy rather than diffuse cluster emission. We estimate a steep spectral index of $\alpha_{325}^{1400} = -1.54$ for the core (although it may be contaminated by lobe emission), while the outer lobes are extremely steep objects, both with $\alpha_{325}^{1400} < -3.1$. We also find evidence for restarted core activity in the form of a set of inner lobes oriented at a somewhat different angle from the outer lobes. A spectral analysis extending the frequency range down to 74 MHz appears to show a turnover at very low frequencies. Comparison of the minimum energy radio pressures with the average thermal pressure surrounding the radio cavities from McNamara et al. (2005) shows that the radio lobes appear to be roughly in pressure balance with the thermal gas.

Subject headings: galaxy clusters: individual(ZwCl 0735.7+7421)

1. Introduction

The hot X-ray emitting gas in clusters of galaxies cools through thermal bremsstrahlung emission for which the cooling time is $t_{cool} \propto T^{0.5} n_e^{-1}$, where T is the gas temperature and n_e is the thermal electron density. In the absence of a major cluster merger, the intracluster medium (ICM) relaxes toward hydrostatic equilibrium. In the classical cooling flow picture (Fabian 1994), the relaxed cluster system develops an inflow of gas toward the cluster center in order to maintain pressure support against the cluster atmosphere. Early estimates of the mass inflow rate from *Einstein* and *ROSAT* suggested cooling rates of 10 to 100's $M_\odot \text{ yr}^{-1}$ (Fabian 1994). Recent X-ray spectroscopic observations from *XMM-Newton*, however, show a greatly reduced amount

¹Naval Research Laboratory, Code 7213, Washington, DC 20375; aaron.cohen@nrl.navy.mil

²Department of Astronomy, University of Virginia, P. O. Box 3818, Charlottesville, VA 22903; tclarke@virginia.edu

³Istituto di Radioastronomia CNR, Via P. Gobetti, 101, I-40129 Bologna, Italy

of gas at intermediate X-ray temperatures (a few 10^6K), and generally find that the gas is cooling to only one-half to one-third of the ambient cluster temperature (Peterson et al. 2003). To avoid the expected rapid cooling to very low temperature, there must be some form of energy input into the ICM to balance the cooling. Currently one of the best candidates for supplying the energy is cluster-center radio sources. Birzan et al. (2004) show that in at least some clusters, the energy input from the central radio source is sufficient to balance cooling, at least for short time periods.

We present an analysis of the radio properties of the exceptionally steep-spectrum radio source (4C+74.13) in the core of the galaxy cluster ZwCl 0735.7+7421. This cluster is a high X-ray luminosity ($L_x = 3.7 \times 10^{44} \text{ ergs s}^{-1}$ in the 0.2–2.5 keV band; Donahue & Stocke (1995), adjusted for Wilkinson Microwave Anisotropy Probe [WMAP] parameters) system at the intermediate redshift of $z = 0.216$. It was identified as a “cooling flow” candidate by Donahue, Stocke, & Gioia (1992) that was later confirmed by Donahue & Stocke (1995). This cluster has been long known to contain an extremely steep spectrum radio source at the center (Slingo 1974). A 408 MHz image (Slingo 1974) hinted at extended structure but did not have the resolution to characterize this, while a snapshot image in the Very Large Array (VLA) A-configuration at 1.4 GHz showed only a very small ($\sim 5''$) faint source (Donahue & Stocke 1995).

In order to investigate the nature of the radio source in this unusual system, we have conducted VLA observations at both 1425 and 325 MHz. Throughout this Letter we use WMAP cosmological parameters (Spergel et al. 2003) that give a comoving radial distance of 870 Mpc, and an angular scale of $3.47 \text{ kpc arcsec}^{-1}$.

2. Observations

We observed 4C+74.13 at 325 MHz with the VLA in B-configuration on 2003 December 4, and at 1425 MHz with the VLA C-configuration on 2004 April 15. For both frequencies, we used the multichannel continuum mode to reduce the bandwidth smearing of bright sources throughout the field of view. Also, both frequencies used 3C48 as a flux and bandpass calibrator. Phase calibration was done using B0735+743 at 1425 MHz and B0739+703 at 325 MHz. The entire primary beam area was imaged in order to subtract confusing sources and the uv -data were weighted with even balance between uniform and natural weighting (robust = 0). Table 1 summarizes the parameters of each data set.

The final image at 1425 MHz (Figure 1) has an rms noise level of $48 \mu\text{Jy/beam}$ and a synthesized beam size of $19.06'' \times 12.39''$ with $\text{PA} = -56.1^\circ$. The final image at 325 MHz (Figure 2) has an noise level of $550 \mu\text{Jy/beam}$ and a synthesized beam size of $20.99'' \times 15.37''$ with $\text{PA} = 13.07^\circ$.

3. Results

3.1. The 1425 MHz Morphology

Our 1425 MHz image reveals a morphology consisting of three major peaks. The brightest peak is near the center and co-incident with the cD galaxy position listed by Bauer et al. (2000), and thus we take this to be the core of the radio galaxy. The other two peaks are consistent with radio lobes and are located roughly $58''$ (200 kpc) to the north and south of the core. The lobes are not co-linear with the core, possibly as a result of interactions with the ICM. We determined the positions of these three peaks with single-Gaussian fits. These three positions are listed in Table 2 and are labeled in Figures 1 - 4.

At the core, there is a significant extension of emission to the south west, which we also label in Figures 1 - 4 as the southwest extension. This structure may be the result of the radio jet interacting with the thermal ICM, or it may represent an inner radio lobe from renewed activity in the core. Unfortunately, the current radio data do not have sufficient resolution to separate these possibilities. In Figure 3 we show the 1425 MHz image obtained after subtracting a point source located at the core from the $u - v$ data. This image reveals excess emission on both sides of the core, suggestive of a set of inner radio lobes from renewed activity. Although higher resolution data are required to confirm the possibility of a recent outburst, the data suggest that either the jet axes have been deflected from their original directions into the outer lobes, or the renewed activity has begun at a different position angle from the original outburst that created the outer radio lobes. We note that another cooling core cluster, A2199, contains the famous renewed radio source 3C 338, which also has a steep spectrum. (Giovannini et al. 1998; Owen & Eilek 1998).

3.2. The 325 MHz Morphology

The 325 MHz image (Figure 2) shows a similar overall extent to the 1425 MHz image but significantly different morphology. The core is significantly fainter relative to the rest of the radio galaxy, indicating that the outer lobes are significantly steeper than the core. Also, the outer lobes are significantly more extended (particularly to the eastern edge) than at 1425 MHz. The eastern extension of the lobes at low frequencies may suggest relative motions between 4C +74.13 and the ICM.

3.3. Spectral Index Measurements

We measure a total flux density for each frequency of $S_{325} = 820.0 \pm 41.1$ mJy and $S_{1425} = 21.94 \pm 0.54$ mJy. These measurements are similar to the Westerbork Northern Sky Survey (Rengelink et al. 1997) and NRAO VLA Sky Survey (Condon et al. 1998) values of $S_{325} = 827.0$ mJy and $S_{1425} = 22.8$ mJy respectively. The corresponding spectral index ($S_\nu \propto \nu^\alpha$) of $\alpha_{325}^{1425} = -2.45 \pm 0.04$

confirms the extremely steep spectrum of this object. In comparison, previous observations of this source reported a spectral index of $\alpha_{408}^{1407} = -1.7$ (Slingo 1974). A typical extragalactic radio source has a spectral index of -0.7 (Kellermann 1964) and a sample of 97 cluster radio sources has shown a typical spectral index of $\alpha_{325}^{1400} \sim -1$ with the steepest at $\alpha_{325}^{1400} = -2.08$ (Edge et al. 2003).

We also measure the flux density at 74 MHz from an image (Figure 4) taken from the VLA Low-Frequency Sky Survey (Cohen et al. 2005) to be 10.4 ± 0.6 Jy. This implies a spectral turnover toward low frequency with $\alpha_{74}^{325} = -1.71 \pm 0.05$. In Figure 5 we plot these flux measurements along with those from the 4C (178 MHz; Gower, Scott, & Wills 1967), 6C (151 MHz; Hales, Mayer, Warner, & Baldwin 1991) and 8C (38 MHz; Rees 1990) surveys. While the 4C and 6C measurements seem consistent with the overall trend of a spectral turnover, the 8C measurement at 38 MHz is surprisingly high, indicating that the spectrum turns up again ($\alpha_{38}^{74} = -2.2 \pm 0.1$). An examination of all images shows the contamination of the 8C measurement to be unlikely, and so the low frequency spectral steepening appears real, although it needs to be confirmed by more sensitive data.

Using the 4C flux measurement, we estimate the monochromatic radio luminosity at 178 MHz to be $P_{178} = 5.2 \times 10^{26} \text{ W Hz}^{-1} \text{ sr}^{-1}$. This is roughly 4 times larger than the monochromatic power of Hydra A (Taylor et al. 1990), and is typical of FR-II radio galaxies (Fanaroff & Riley 1974).

We have also measured the spectral indices of individual components of the radio galaxy. First, we measured positions for these components by fitting a Gaussian to the region immediately surrounding the peak of each component. For the core and the north and south outer lobes, we used the 1425 MHz image to measure the positions. But, as the southwest extension is not a peak in the 1425 MHz image, we fitted the peak in the 325 MHz image to determine its position. We then convolved both images to a $21.0''$ circular beam and measured the surface brightness at the location of each component. The results are summarized in Table 2. The outer lobes have extremely steep spectral indices of $\alpha_{325}^{1425} = -3.17 \pm 0.05$ and -3.13 ± 0.05 for the north and south outer lobes respectively. As discussed in Section 3.2, the outer lobes appear to show an extension toward the east at 325 MHz that suggests spectral steepening in this region. We have estimated the spectral indices on the eastern and western sides of each outer lobe and find a spectral gradient of 0.20 ± 0.08 for the north lobe and 0.15 ± 0.08 for the south lobe. This emission may represent a region of older radio plasma that is displaced to east by relative motions between the radio source and the thermal ICM. The southwest extension has a somewhat flatter spectral index of -2.32 ± 0.04 , which is expected for a presumably younger particle population. The spectral index of the core is much flatter at $\alpha_{325}^{1425} = -1.54 \pm 0.04$. As the flux density measurement for the core at 325 MHz is probably significantly contaminated by the surrounding lobe emission its spectral index is likely to be even flatter than this.

3.4. Radio/X-Ray Comparison

Recent *Chandra* images of ZwCl 0735.7+7421 by McNamara et al. (2005) reveal large (~ 200 kpc diameter) cavities in the thermal gas in this system. They show that the 1.4 GHz radio emission of 4C +74.13 appears to fill the X-ray cavities. An X-ray temperature map shows that the gas in the region of the cavities is significantly hotter than the surrounding cluster, indicating that the radio outburst is driving shocks into the ICM (McNamara et al. 2005). They estimate that a total energy of $\sim 6 \times 10^{61}$ ergs was required to drive the shocks.

We have used our new radio observations to estimate the minimum-energy synchrotron pressure in the outer lobes of 4C +74.13 for comparison with the thermal pressure. We assume that the emission from the radio lobes comes from a uniform prolate cylinder with a filling factor of unity, and that there is equal energy in relativistic ions and electrons. Using a model with the magnetic field perpendicular to the line of sight, lower and upper cutoff frequencies of 10 MHz and 100 GHz, and spectral indices given in Table 2, we estimate minimum energy pressures of 3.9×10^{-11} and 2.4×10^{-11} dyn cm $^{-2}$ for the north and south outer lobes, respectively. Based on the *Chandra* data, McNamara et al. (2005) estimate an average pressure surrounding the lobes to be $P \simeq 6 \times 10^{-11}$ dyne cm $^{-2}$, roughly 1.5 – 2.5 times higher than the minimum-energy pressures of the lobes. This system appears to be much closer to pressure balance than most other bubble systems that are generally an order of magnitude different (Blanton 2003).

Using the spectral index of the full source ($\alpha_{325}^{1425} = -2.45$), we have calculated the total radio luminosity of 4C +74.13 over the frequency range of 10 MHz - 10 GHz to be $L_{radio} = 38.2 \times 10^{42}$ ergs s $^{-1}$. Based on the spherical hydrodynamic shock model, McNamara et al. (2005) estimate a required average radio jet power of $P_s \simeq 1.7 \times 10^{46}$ ergs s $^{-1}$ for the radio outburst that created the cavities. This suggests that the jet power is a few hundred times the observed radio luminosity, similar to what is found by Birzan et al. (2004) for radio filled cavities. McNamara et al. (2005) conclude that the radio outburst released sufficient energy to balance the central cooling flow for several gigayears. In comparison, Birzan et al. (2004) found that roughly one-half of the 16 systems they studied had sufficient mechanical luminosities to balance cooling in the cores. Although the Birzan et al. (2004) sample specifically targetted clusters displaying X-ray surface brightness depressions, and thus the results are not generally representative of cluster systems, we notice that the objects balancing the central cooling show a large spread in radio luminosity and in spectral index.

4. Conclusions

From these 1425 and 325 MHz data sets, we find that the central radio source in the galaxy cluster ZwCl 0735.7 + 7421 is an extended (400 kpc) radio galaxy with a clearly defined core and outer lobes. The overall spectrum is very steep between these two frequencies ($\alpha_{325}^{1425} = -2.45 \pm 0.04$) but turns over somewhat at 74 MHz with $\alpha_{74}^{325} = -1.71 \pm 0.05$, with an apparent steepening to lower frequencies. The spectral index of the core ($\alpha_{325}^{1425} = -1.54 \pm 0.04$) is very steep although

much flatter than that of the outer lobes, both with $\alpha_{325}^{1425} < -3.1$. We find evidence of “inner” lobes, which may be the result of a recent radio outburst. The spectral index at the location of the southwest inner lobe (the southwest extension of the core) is -2.32 ± 0.04 , which is steeper than the core but flatter than the outer lobes.

Minimum energy estimates of the pressures of the radio lobes are within a factor of 2.5 of the average thermal pressure in the surrounding regions as determined by McNamara et al. (2005). Using estimates of the total radio luminosity, we find that the jet power must be a few hundred times the observed radio luminosity.

Basic research in radio astronomy at the US Naval Research Laboratory is supported by the Office of Naval Research. T. E. C. was supported in part by *Chandra* awards GO2-3160X, GO3-4155X, GO3-4160X, GO4-5133X, GO4-5149X, and GO4-5150X issued by the Chandra X-ray Observatory, which is operated by the Smithsonian Astrophysical Observatory for and on behalf of NASA under contract NAS8-39073.

REFERENCES

- Bauer, F. E., Condon, J. J., Thuan, T. X., & Broderick, J. J. 2000, *ApJS*, 129, 547
- Birzan, L., Rafferty, D. A., McNamara, B. R., Wise, M. W., & Nulsen, P. E. J. 2004, *ApJ*, 607, 800
- Blanton, E. L. 2003, in *The Riddle of Cooling Flows in Galaxies and Clusters of Galaxies*, Eds. T. H. Reiprich, J. C. Kempner, & N. Soker, astro-ph/0402342
- Cohen, A. S., Lane, W. M., Lazio, T. J. W., Kassim, N. E., Cotton, W. D., Perley, R. A., Condon, J. J., & Erickson, W. C. 2005, American Astronomical Society Meeting #205, paper #91.05
- Condon, J. J., Cotton, W. D., Greisen, E. W., Yin, Q. F., Perley, R. A., Taylor, G. B., & Broderick, J. J. 1998, *AJ*, 115, 1693
- Donahue, M., Stocke, J. T., & Gioia, I. M. 1992, *ApJ*, 385, 49
- Donahue, M. & Stocke, J. T. 1995, *ApJ*, 449, 554
- Edge, A. C., Ebeling, H., Bremer, M., Röttgering, H., van Haarlem, M. P., Rengelink, R., & Courtney, N. J. D. 2003, *MNRAS*, 339, 913
- Fabian, A. C. 1994, *ARA&A*, 32, 277
- Fanaroff, B. L. & Riley, J. M. 1974, *MNRAS*, 167, 31P
- Gower, J. F. R., Scott, P. F., & Wills, D. 1967, *MmRAS*, 71, 49
- Hales, S. E. G., Mayer, C. J., Warner, P. J., & Baldwin, J. E. 1991, *MNRAS*, 251, 46

- Giovannini, G., Cotton, W. D., Feretti, L., Lara, L., & Venturi, T. 1998, ApJ, 493, 632
- Kellermann, K. I. 1964, ApJ, 140, 969
- McNamara, B. R., Nulsen, P. E. J., Wise, M. W., Rafferty, D. A., Carilli, C., Sarazin, C. L., & Blanton, E. L. 2005, Nature in press (astro-ph/0411553)
- Owen, F. N. & Eilek, J. A. 1998, ApJ, 493, 73
- Peterson, J. R., Kahn, S. M., Paerels, F. B. S., Kaastra, J. S., Tamura, T., Bleeker, J. A. M., Ferrigno, C., & Jernigan, J. G. 2003, ApJ, 590, 207
- Rees, N. 1990, MNRAS, 244, 233
- Rengelink, R. B., Tang, Y., de Bruyn, A. G., Miley, G. K., Bremer, M. N., Roettgering, H. J. A., & Bremer, M. A. R. 1997, A&AS, 124, 259
- Slingo, A. 1974, MNRAS, 168, 307
- Spergel, D. N., et al. 2003, ApJS, 148, 175
- Taylor, G. B., Perley, R. A., Inoue, M., Kato, T., Tabara, H., & Aizu, K. 1990, ApJ, 360, 41

This preprint was prepared with the AAS L^AT_EX macros v5.2.

Table 1. Summary of Radio Observations

Observation:	90cm	20cm
Configuration	VLA B	VLA C
Central Frequency	321.56 (IF 1)	1385.1 (IF 1)
(MHz)	328.50 (IF 2)	1464.9 (IF 2)
Bandwidth (MHz)	6.25	12.5
Time on Source (min.)	215	205
Resolution (")	15.4 × 21.0	12.4 × 19.1
RMS noise (μJy/b)	550	48




Fig. 1.— Central radio source (4C+74.13) in the core of ZwCl 0735.7+7421 at 1425 MHz. The contours begin at $120 \mu\text{Jy}/\text{beam}$ and increase in factors of $\sqrt{2}$.




Fig. 2.— Radio contours of the 325 MHz observations of 4C+74.13. The contours begin at 1.5 mJy/beam and increase in factors of $\sqrt{2}$. This image is shown at the same scale as that in Figure 1.




Fig. 3.— Radio contours of the central region of ZwCl 0735.7+7421 at 1425 MHz with the core subtracted. The contours begin at 120 $\mu\text{Jy}/\text{beam}$ and increase in factors of $\sqrt{2}$. This image is shown at the same scale as that in Figure 1.




Fig. 4.— ZwCl 0735.7+7421 at 74 MHz. The contours begin at 150 mJy/beam and increase in factors of $\sqrt{2}$.




Fig. 5.— Total flux density of 4C+74.13 as a function of frequency.

Table 2. Flux Densities and Spectral Index Measurements

Component	α (J2000) (h, m, s)	δ (J2000) ($^{\circ}$, $'$, $''$)	S_{1425} (mJy)	S_{325} (mJy)	α_{325}^{1425}
Core	07 41 44.54	+74 14 39.6	9.79 ± 0.20	94.9 ± 4.8	-1.54 ± 0.04
S Inner Lobe	07 41 40.88	+74 14 25.1	2.86 ± 0.075^a	88.4 ± 4.5^a	-2.32 ± 0.04
N Outer Lobe	07 41 45.63	+74 15 36.9	0.952 ± 0.054^a	102.8 ± 5.2^a	-3.17 ± 0.05
S Outer Lobe	07 41 39.02	+74 13 41.0	0.978 ± 0.054^a	99.4 ± 5.0^a	-3.13 ± 0.05
Full Source	-	-	21.94 ± 0.54	820.0 ± 41.1	-2.45 ± 0.04

^aThese values represent surface brightness in mJy/beam, with circular beam FWHM = $21''$.