Star Clusters with Primordial Binaries: II. Dynamical Evolution of Models in a Tidal Field

Michele Trenti^{1,2*}, Douglas C. Heggie^{3*} and Piet Hut^{4*}
¹Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

- ² Yukawa Institute for Theoretical Physics, Kyoto University, 606-8502 Kyoto, Japan
- ³ School of Mathematics, University of Edinburgh, King's Buildings, Edinburgh EH9 3JZ, Scotland, U.K.
- ⁴Institute for Advanced Study, Princeton, NJ 08540, USA

Accepted; Received; in original form

ABSTRACT

We extend our analysis of the dynamical evolution of simple star cluster models, in order to provide comparison standards that will aid in interpreting the results of more complex realistic simulations. We augment our previous primordial-binary simulations by introducing a tidal field, and starting with King models of different central concentrations. We present the results of N-body calculations of the evolution of equal-mass models, starting with primordial binary fractions of 0 - 100 %, and N values from 512 to 16384. We also attempt to extrapolate some of our results to the larger number of particles that are necessary to model globular clusters.

We characterize the steady-state 'deuterium main sequence' phase in which primordial binaries are depleted in the core in the process of 'gravitationally burning'. In this phase we find that the ratio of the core to half-mass radius, r_c/r_h , is similar to that measured for isolated systems (Heggie et al. 2005). In addition to the generation of energy due to hardening and depletion of the primordial binary population, the overall evolution of the star clusters is driven by a competing process: the tidal dissolution of the system. If the primordial binary fraction is greater than 5% and the total number of particles $N \ge 8192$, we find that primordial binaries are not fully depleted before tidal dissolution, in systems initially described by a King model with a self-consistent tidal field.

We compare our findings, obtained by means of direct N-body simulations but scaled, where possible, to larger N, with similar studies carried out by means of Monte Carlo methods (Fregeau et al. 2003, 2005). We find significant qualitative and quantitative differences with the results in the earlier paper. Some of these differences are explicable by the different treatment of the tidal field in the two studies. Others, however, confirm the conclusion of Fregeau et al (2005) that the efficiency of binary burning in the earlier Monte Carlo runs was too high. There remain unexplained differences, however. In particular, the binary population appears to be depleted too quickly, even in the most recent Monte Carlo results.

Key words: globular clusters – methods: N-body simulations – stellar dynamics.

1 INTRODUCTION

Several observational surveys of globular clusters have highlighted the presence of binaries, which are found to constitute up to 50% of the mass of the core (e.g., see Rubenstein & Bailyn 1997; Albrow et al. 2001; Bellazzini et al. 2002; Pulone et al. 2003). Such a high number of objects is extremely unlikely to have been

ate a significant binary population due to tidal capture and three body encounters, the most probable outcome from these processes is a merger (Chernoff & Huang 1996; Portegies Zwart & McMillan 2002), which could eventually lead to the formation of a central intermediate mass black hole (Portegies Zwart et al. 2004). Therefore these binaries are regarded as being primordial. The primordial binary fraction appears to be a key parameter for determining the dynamical evolution of a star cluster, as binaries are an efficient heating source which can halt core collapse. In addi-

formed dynamically, as, at the density required to cre-

^{*} E-mail addresses: trenti@stsci.edu (MT); piet@ias.edu (PH); d.c.heggie@ed.ac.uk (DCH)

2

tion, interesting astronomical objects can be produced due to three- and four-body encounters, such as blue stragglers, X-ray binaries and binaries containing pulsars.

The evolution of star clusters with primordial binaries still presents open issues, mainly due to the intrinsic complexity of numerical simulations of a system where the local dynamical timescale may be many orders of magnitude smaller than the global relaxation timescale. (Hard binaries have an orbital period of a few hours, while the half mass relaxation time t_{rh} may be of order a few billion years.) Numerical simulations have thus been performed either using approximate algorithms such as Fokker Planck or Monte Carlo methods (Gao et al. 1991; Giersz & Spurzem 2000; Fregeau et al. 2003), which often have to rely on a knowledge of the relevant interaction cross sections; or using direct Nbody codes but employing only a modest number of particles of order $N \approx 10^3$ (McMillan et al. 1990; McMillan & Hut 1994; Heggie & Aarseth 1992). Both approaches could potentially lead to misleading results. On the one hand Monte Carlo methods rely on input physics that may not accurately reflect the realistic interactions in a densely populated and fluctuating core (e.g. the velocity distribution may be assumed to be isotropic), while interaction cross sections in the case of unequal masses are not well known. On the other hand, extrapolation from direct simulations may prove to be non trivial, as it is not clear how to scale the results obtained with the number of particles.

We have thus recently started (Heggie et al. 2005, hereafter Paper I) a program to survey the basic properties of the evolution of an idealised star cluster with a population of primordial binaries, with the aim of placing a stepping stone between simplified models, such as those of Fokker-Planck type, and realistic complex numerical simulations, such as those recently performed by Portegies Zwart & McMillan (2004). In the first paper of our series we have focused on the evolution of isolated, equal-mass models, with a primordial binary fraction in the range 0-100 % and with a number of particles N in the range from 256 to 16384. We have compared our results not only with those of the study carried out by Gao et al. (1991) (who used a Fokker Planck code), and also to some extent with Giersz & Spurzem (2000) and with Fregeau et al. (2003), but also with a theoretical model which predicts the evolution of the core radius (Vesperini & Chernoff 1994). We have shown that significant differences arise between direct N-body simulations and the Fokker Planck calculations of Gao et al. (1991). In particular we found that, starting from the same initial conditions, the number of binaries in the core after the core collapse is significantly lower in our simulations than the number reported by Gao et al. (1991). This could have important consequences if one attempted to infer the original primordial binary fraction by observations of the current number of binaries in the cores of globular clusters. Another interesting finding is that the ratio of core radius r_c to half mass radius r_h in the post collapse phase seems not to follow the theoretical expectation from the work of Vesperini & Chernoff (1994), as, by varying the number of particles N, we observed a steeper decrease of r_c/r_h than that expected ($\propto 1/\log(0.4N)$).

In this paper we extend our simulations to include the effect of the tidal field of the galaxy. We consider the evolution of King models with different values of N, the con-

centration index W_0 and the primordial binary ratio f. We compare our direct simulations to the Monte Carlo simulations performed by Fregeau et al (2003, 2005), hereafter Fregeau et al. They studied the evolution of isolated and tidally truncated star clusters with a population of primordial binaries in the range 2-20 % and a realistic particle number $(N = 3 \ 10^5)$. Their two-dimensional Monte Carlo Method is expected to offer significant advantages over the one-dimensional code used by Gao et al. (1991) as the influence of anisotropy in the velocities can be taken into account, and it is unnecessary to assume that the distributions of energy and position (for binaries) are independent. The work of Fregeau et al. led to some important conclusions. For tidally limited clusters they noted for the first time the possibility of an initial expansion of the core radius of the cluster in the presence of a significant population of primordial binaries, when starting from models with a high central concentration (i.e. King profiles with $W_0 \gtrsim 7$). Fregeau et al. also showed that, in general, primordial binaries delay the deep core collapse phase that is observed in clusters with only single stars, so that the system can be tidally dissolved before collapsing. The comparison of our results with their simulations is mainly focused on fundamental quantities like the disruption rate of the binaries, the core radius and the dissolution time-scale. It must be noted, however, that Fregeau et al improved their code between the 2003 and 2005 papers by the inclusion of direct numerical integration of encounters. They note (Fregeau et al. 2005) that this alters some of their earlier results, as we mention at appropriate points in the present paper. In particular Fregeau et al. (2005) note that binary burning was too efficient in their earlier models.

The paper is organized as follows: in the next Section we present the parameters of our simulation dataset and the numerical method used. In Sec. 3 we give a physical picture of the evolution of a star cluster with primordial binaries and tidal field that will guide our interpretation of the simulations. In Sec. 4 and 5 we present our results for runs with a tidal field and a tidal cutoff, respectively, including comparison with the results of Fregeau et al. The last section of the paper provides a summary and discussions.

2 SIMULATIONS: SETUP AND ANALYSIS

2.1 Specification of the models

The models considered in this paper are tidally limited, with stars of equal mass m. The initial distribution is a King model with scaled central potential $W_0=3,7,11$. Our standard models have a primordial binary ratio of 10-20 %, although we have also performed some runs with fewer or no primordial binaries (0,2,5%) as well as with higher ratios, 50 % and 100 %. As in Paper I we define the primordial binary fraction as:

$$f = n_b/(n_s + n_b) \tag{1}$$

with n_s and n_b being the number of singles and binaries respectively. This implies that the fraction of the total mass in binary stars, in the case of equal masses, is larger in the following way:

$$f_m = 2n_b/(n_s + 2n_b)$$

For example, for a run with 10% primordial binaries, f = 10% whereas $f_m = 2/11 \sim 18\%$.

All our results are presented using standard units (Heggie & Mathieu 1986) in which

$$G = M = -4E_{\mathrm{T}} = 1$$

where G is the gravitational constant, M is the total mass, and $E_{\rm T}$ is the total energy of the system of bound objects. In other words, $E_{\rm T}$ does not include the internal binding energy of the binaries, only the kinetic energy of their centre-of-mass motion and the potential energy contribution where each binary is considered to be a point mass. We denote the corresponding unit of time by

$$t_d = GM^{5/2}/(-4E_T)^{3/2} (2)$$

in general. For the relaxation time, we use the following expression

$$t_{rh} = \frac{0.138Nr_h^{3/2}}{\ln(0.11N)},\tag{3}$$

where r_h is the half-mass radius, and N denotes the number of original objects, i.e. $N=n_s+n_b$. When we discuss a run with N=4096 and 50% primordial binaries, we are dealing with 6144 stars. We have considered runs with N in the range 512-16384.

Our runs follow the evolution of the system until the number of particles drops below 10. However, to avoid possible biases introduced by long-living small-N configurations, we define the dissolution time t_{dis} , as the moment when 98 % of the initial mass of the system is no longer bound.

Following Fregeau et al., our initial distribution of the binaries' binding energies is flat in log scale in the range from ϵ_{min} to $133\epsilon_{min}$, with $\epsilon_{min} = m\sigma_c(0)^2$. Here $\sigma_c(0)$ is the initial central velocity dispersion and this choice, if applied to an isolated Plummer model, corresponds approximately to the standard range adopted in Paper I, i.e. to $\epsilon_{min} \approx 5.1~kT$ and $133\epsilon_{min} \approx 680~kT$, where the mean kinetic energy per particle (over the entire system, with binaries replaced by their centres of mass) is 3kT/2.

To follow the evolution of the system we have used Aarseth's NBODY6 (Aarseth 2003), which has been slightly modified to provide additional runtime diagnostics on the spatial distributions of single and binary stars. A King model with $W_0 = 7$ required almost one month of CPU time on a Pentium4 3Ghz PC before tidal dissolution for N = 16384 and f = 10%; the use of GRAPE hardware would not help significantly, as the computational bottleneck is in the treatment of the local dynamics of binaries (however see Makino & Hut 1990, for a discussion on the scaling of the computational time with the number of particles in the presence of primordial binaries).

The galactic tidal field is treated in our standard sample of simulations as that due to a point mass, and the tidal force acting on each particle is computed using a linear approximation of the field. The tidal radius r_t is defined as (Spitzer 1987):

$$r_t^3 = \frac{M}{3M_{Gal}}R_{Gal}^3, \qquad (4)$$

where M_{Gal} is the galaxy mass, and R_{Gal} is the distance of the centre of the star cluster from the centre of the galaxy.

We assume R_{Gal} is constant, i.e. the cluster describes a circular orbit. Thus the equations of motion of a star in the cluster are

$$\begin{vmatrix}
\ddot{x} + 2\omega\dot{y} + (\kappa^2 - 4\omega^2)x & = -\phi_x \\
\ddot{y} - 2\omega\dot{x} & = -\phi_y \\
\ddot{z} & + \nu^2 z & = -\phi_z,
\end{vmatrix}$$
(5)

where the axes are orientated conventionally with respect to the direction to the centre of the galaxy and the galactic motion of the star cluster, ω is its angular velocity, κ is its epicyclic frequency, ν is its vertical frequency, and the right sides are the components of the gravitational acceleration due to the other stars. The value of $r_t(0)$ is set at the beginning of the simulation to match the cut-off given by the adopted initial King model. In our standard runs particles are removed from the system when they reach a distance from the density centre greater than twice the instantaneous tidal radius r_t , and the value of r_t is updated during the simulation according to the decrease of M, the total mass of the system.

Note that this method of treating the influence of the galactic field is different from the one adopted by Fregeau et al., as they imposed a radial "cutoff", i.e. they did not consider the tidal force acting on individual particles, and simply removed particles whose distance exceeded r_t (see the description of their code in Joshi et al. 2000). At variance with our tidal field treatment, the simulations with tidal cutoff do not include the Coriolis and centrifugal contributions to the particles acceleration. Therefore the two approaches have important physical differences. The effects of this different treatment are discussed in Sec. 5, where we describe a subset of simulations ($W_0 = 7$ and $W_0 = 3$, f = 10%, with N from 512 to 16384) using the tidal radial cutoff treatment adopted by Fregeau et al.

Our results, unless otherwise noted, are presented by applying a triangular smoothing filter with width 2.5 $t_{rh}(0)$ (see Fig. 3 and Sec. 3.2 in Paper I for further details).

A summary of the runs is presented in Table 3.

2.2 Core radius: definition

In Paper I we adopted the following definition for the core radius:

$$\tilde{r}_c = \sqrt{\frac{\sum_{i=1,\tilde{N}} r_i^2 \rho_i^2}{\sum_{i=1,\tilde{N}} \rho_i^2}},$$
(6)

where the sum is made over the particles within the half mass radius and the density ρ_i around each particle is computed from the distance to the fifth nearest neighbour (Casertano & Hut 1985). This definition presents some systematic differences (see Fig. 1) from the one adopted by Fregeau et al.:

$$r_c = \sqrt{\frac{3\sigma_c^2}{4\pi G\rho_0}},\tag{7}$$

where σ_c^2 is the central velocity dispersion (mass weighted as in Vesperini & Chernoff 1994) and ρ_0 the central density (Spitzer 1987). For a proper comparison between our set of simulations and the runs discussed by Fregeau et al. it is important to refer to the same quantities, and in this paper we adopt as definition for the core radius r_c the choice taken




Figure 1. Core radius measured using the method in Paper I (Eq. 6) and in Fregeau et al. (Eq. 7) for runs with N=16384 and f=10%. Note that the density-averaged definition of the core radius (Paper I) underestimates the core radius at the beginning of the simulation compared with the theoretical value for the corresponding King model (i.e. 0.67 and 0.21 for $W_0=3$ and 7, respectively). In this paper we adopt the definition used by Fregeau et al. for the core radius.

by Fregeau et al., i.e. Eq. (7). We estimated the velocity dispersion σ_c^2 using all the particles within the radius defined by Eq. (6), while ρ_0 has been obtained from the Lagrangian radius containing 1 % of the instantaneous mass of the system. For the purpose of comparison with Paper I, we also report in Table 3 the value of the alternative definition for the core radius, \tilde{r}_c , at the end of the core collapse.

2.3 Time-scales

In the presence of primordial binaries and a tidal field, several dynamical processes operate, each with its own time scale. Here we summarize them:

- Dynamical time $(t_d, \text{cf. eq.}(2))$: within a small numerical factor, this is the typical timescale for a particle to cross the system. It also represents the timescale on which approximate dynamical equilibrium is established in the case of initial conditions starting out of equilibrium. (Note however that all our simulations start from approximate dynamical equilibrium.)
- Half-mass relaxation time $(t_{rh}, \text{ cf. eq.}(3))$: this is the timescale for energy diffusion in phase space, i.e. for establishing thermal equilibrium. It is of order $N/\log(0.11N)$ t_d .
- Total-mass relaxation time (t_{rt}) : this is the timescale for relaxation of the cluster as a whole and is defined using the tidal radius as characteristic length scale for estimating the relaxation time. t_{rt} is proportional to t_{rh} (defined above): $t_{rt}/t_{rh} \propto (r_t/r_h)^{3/2}$.
- Time for initial core contraction (t_{cc}) : this is the timescale on which the core of the cluster contracts (or ex-

pands) to reach a quasi-steady configuration in which the energy released in the core fuels the expansion of the half mass radius. The typical value of t_{cc} is of the order of $10t_{rh}(0)$.

- Tidal dissolution time (t_{dis}) : Time needed to lose 98% of the initial mass of the system due to escape across the tidal boundary. Depending on the tidal field strength, the initial concentration of the system and the initial number of particles, this timescale ranges from a few relaxation times to a few tens of relaxation times.
- Tidal half-mass dissolution time (t_{Half}) : Similar to t_{dis} , but defined as the time needed for 50% of the mass of the system to escape.
- Binary burning time (t_{bb}) Time needed to deplete 80% of the initial number of binaries. Around this time gravothermal oscillations set in, provided that N is large enough. For initial binary fractions above 10%, t_{bb} is of the order of several tens to hundreds of initial relaxation times.

3 THE PHYSICAL PICTURE: PRIMORDIAL BINARIES AND THE TIDAL FIELD

The evolution of an isolated star cluster made of equal mass stars is driven initially by the gravothermal instability. Two-body encounters drive a heat flow from the central region of the star cluster, which behaves like a self gravitating system with negative specific heat, to the halo. This triggers a thermal "collapse" on the timescale of the heat flow process.

The "collapse" phase lasts until an efficient form of energy generation in the centre can stop the process by providing an energy production rate equal to the energy loss rate by two body relaxation from the region of the core. In the case of systems consisting of single stars only, the collapse is stopped only after one or more binaries are formed due to three body encounters in the dense central region. The collapse typically takes a time t_{cc} of order 15 $t_{rh}(0)$, where $t_{rh}(0)$ is the initial half-mass relaxation time. At this point, i.e. at the end of core collapse (sometimes called "core bounce"), the central density can exceed 10⁴ times the average density inside the half-mass radius r_h (for N=16384, or far more for larger N). After core bounce, the generation of energy in the core is accompanied by a steady expansion of the half mass radius. If N is large enough, however, binary activity in the core may cause a temporary temperature inversion which drives the gravothermal instability in reverse. This suppresses binary activity, and further collapses may recur, in a succession of "gravothermal oscillations" (Sugimoto & Bettwieser 1983).

This picture is strongly modified when a population of primordial binaries is present, as the generation of energy due to existing binaries is more efficient than that due to dynamically formed pairs (see Sec. 2 of Paper I). For this reason, if the primordial binary fraction is above a few percent, the initial contraction of the core is more gentle, and it is halted at a lower core density (and a larger core radius). The timescale for initial core contraction (t_{cc}) remains of the same order of magnitude as in the case where no binaries are initially present (see Fig. 17 in Paper I). From then until the time when the population of binaries becomes heavily depleted, the evolution of the system proceeds with an almost constant ratio of core radius to half mass radius. This phase lasts for a time proportional to the initial binary ratio,

and thereafter a phase of gravothermal oscillations sets in, much as for the case where there are no primordial binaries (see Figs. 5-6 in Paper I). For an isolated system, provided that N is large enough (i.e. $N \gtrsim 10^4$), the onset of gravothermal oscillations occurs when the mass of binaries has fallen to roughly 10–15% of its initial value (Fregeau et al. 2003, Fig.4; Paper I, Figs. 5,7). These are empirical results, however, and may change for primordial binary fractions smaller than the lowest used in these simulations, i.e. 2%.

The presence of a tidal field introduces a new timescale for the evolution of the system, i.e. t_{dis} . In the presence of a tidal field, the star cluster steadily loses mass at a rate that is almost independent of the properties of the central parts of the system, where binaries accumulate (because of mass segregation) and act as an energy source for the system. In principle the tidal dissolution can be so fast $(t_{dis} < t_{bb})$ that the system does not have enough time to deplete its reservoir of primordial binaries before dissolving. On the other hand, if the initial primordial binary fraction is low or if the tidal field is weak, then the system can burn almost all of the binaries before being dissolved, and it then undergoes a phase of gravothermal oscillations, as already described for an isolated system, provided that N is large enough.

In an isolated system with an initial binary ratio of 10%, approximately $100 \, t_{rh}(0)$ is the typical timescale t_{bb} required for disrupting 80% of the binary population considered in our runs (e.g., see Fig. 7 in Paper I). Note, however, that the range of binding energy in these runs is $5-700 \, kT$, where 3/2NkT is the total kinetic energy of the cluster (when the binaries are replaced by their barycentres). The time to disrupt most of the binaries is certainly dependent on the initial distribution.

The time to disrupt most of the binaries can also be estimated with the following argument. A primordial binary in the range of $5-700\ kT$ gives off, on average, around $100\ kT$ before leaving the system (or being dissolved; see e.g., Hut et al. 1992). Thus if we start with a 10% primordial binary population each single star receives around $10\ kT$. The energy loss at the half mass radius due to the temperature gradient is of the order of $0.1\ kT$ per star per relaxation time, so we have to wait an order of $100\ relaxation$ times before depleting the binaries. The same argument suggests that the time taken to burn most of the binaries is proportional to f.

4 TIDAL FIELD RUNS: RESULTS AND COMPARISONS

In this Section we discuss the properties of our simulations, our main purpose being to set out the essential empirical facts about the evolution of systems with primordial binaries in a galactic tidal field. We shall also compare our results with those obtained by Fregeau et al., though they treated the tidal field as a cutoff. To elucidate this comparison, we were motivated to carry out further N-body runs with the same treatment (i.e. a cutoff). We defer presentation of those calculations to the next section, however; in the present section we consider only runs that include our most realistic treatment of the tidal force (eq.(5)).

4.1 Total mass and dissolution timescale

Our results are summarized by the values of t_{dis} in Tab. 3 and Tab.2, and details for a subset of the runs are presented in the top panel in Figs. 2-5; these are intended also to facilitate comparison with corresponding plots in Fregeau et al., as specified in the captions. The evolution of the total mass in our different runs depends on three initial parameters: (i) the initial fraction of primordial binaries, f; (ii) the scaled central potential of the initial King model, W_0 ; and (iii) the initial number of objects, N. We consider the dependence on each in turn.

4.1.1 Dependence on f

The initial fraction of primordial binaries has only a small influence on the rate of mass loss of the system and on t_{dis} . The dependence is, however, systematic, as can be seen from Fig. 6. (Though this figure shows runs with $W_0 = 7$, the trend shown is representative of all concentration indexes that we have studied.) The simulation with the longest dissolution time is the one starting with 100% of primordial binaries. This may not be surprising, as this run has the highest total number of stars and so, after disruption of a given proportion of binaries, it has the longest relaxation time. On the other hand it is closely followed by the simulations with low binary content (2 % and 5 %) and then by the run with single stars only, which is dissolved in 90% of the time needed in the 100 % case. This may be due to the fact that simulations with $f\lesssim 5\%$ deplete their primordial binaries before dissolution $(t_{bb} < t_{dis})$; this leads to a deep core collapse, which creates a strongly bound core which is rather resilient to dissolution. Curiously, simulations with 20% of primordial binaries are subject to the fastest tidal dissolution, and this could be due to effects related to mass segregation, which is absent if f = 0 or 1, and which tends to expel single stars.

4.1.2 Dependence on W_0

Consider, as an example, runs with N = 4096 and f = 10%. A low concentration King model with $W_0 = 3$ is tidally dissolved in approximately 13 $t_{rh}(0)$ (Table 3), while the system can survive for approximately 50 $t_{rh}(0)$ for $W_0 = 7$; but $W_0 = 11$ is intermediate between the two $(t_{dis} \approx 25 \ t_{rh}(0))$. This non-monotonic dependence on concentration index W_0 is a general feature, independent of the number of particles. It may be understood in terms of the non-monotonic dependence of the tidal radius on W_0 , at fixed total mass and total energy (Tab.1; see also Fig. 8.3 in Heggie & Hut 2003). (Another way of looking at this is that, in the units we adopt, the strength of the tidal field varies non-monotonically with W_0 .) Though the difference in r_t between the cases $W_0 = 7$ and $W_0 = 11$ may not seem great, two further factors may be relevant: (i) r_h also increases slightly between $W_0 = 7$ and $W_0 = 11$ (Table 1), which would increase the vulnerability of the more concentrated system, even if the tidal field were the same; (ii) high-concentration models begin with a short period of core expansion (Fig.5), and the associated burst of energy generation enhances tidal overflow.

If the tidal dissolution time is expressed in units of the total-mass relaxation time (t_{rt}) , then the differences be-

Table 1. Initial tidal and half-mass radius $r_t(0)$, $r_h(0)$ (in natural units) as functions of the scaled central potential W_0

W_0	$r_t(0)$	$r_h(0)$		
3	3.15	0.84		
7	7.00	0.81		
11	6.62	1.01		

tween the different King models are greatly reduced. By accounting for the factor $(r_t/r_h)^{3/2}$, so that the time is in units of t_{rt} , the difference in tidal dissolution time between the various King models considered here is within 20% for N = 4096 and within 10% for N = 16384.

4.1.3 Dependence on N

In our set of simulations we observe a marked dependence of the dissolution time-scale t_{dis} on the number of particles used (see Tab. 3 and Fig. 7). This is not surprising, as a similar effect is well known in simulations with the tidal force modeled as in our eq. (5), though using single particles only: the time for the loss of half of the mass of a cluster is approximately proportional to $t_{rh}(0)^{3/4}$, i.e.

$$t_{Half} \propto \left(\frac{\ln\left(0.11N\right)}{N}\right)^{1/4} t_{rh}(0),\tag{8}$$

(Baumgardt 2001). Besides empirical support from N-body simulations, this formula can be understood on the basis of the combined effect of the tidal field and of two body relaxation (Baumgardt 2001). We find (Fig.7) that it also provides a reasonable fit to the dissolution time t_{dis} of models with primordial binaries (for the parameters specified in the caption of the figure), though the observed dependence on N is slightly stronger. Therefore we can take advantage of eq.(8) for the approximate extrapolation of results on the dissolution time t_{dis} to larger numbers of particles N. Note, however, that eq.(8) is derived in Baumgardt (2001) using simulations with $N \leq 16384$; it is expected to be valid up to $N \lesssim 10^6$ at most, as for larger N the equation eventually leads to $t_{Half} < t_{rh}(0)$, a clearly unphysical result.

Assuming, then, that this scaling applies to the dissolution time t_{dis} , we can compare our results with those in Fregeau et al. If we extrapolate to $N=3\ 10^5$, we find that t_{dis} decreases by a factor ≈ 2.6 from the results obtained for the simulations with N=4096 and by a factor ≈ 1.9 from the runs with N=16384. This would mean that a simulation starting from $W_0 = 7$ with 10 % binaries and $N = 3 \times 10^5$ would take $\approx 18 \ t_{rh}(0)$ to dissolve. For comparison Fregeau et al. measure $t_{dis} \gtrsim 38t_{rh}(0)^1$ for these initial conditions. This and other comparisons are given in Table 2 (second and fourth columns). An understanding of such differences requires consideration of the treatment of the tide, and so we defer further discussion to Sec.5.




Figure 2. Time dependence of the total mass and of the mass in binaries (upper panel) and of the half mass and core radius (bottom panel). The dotted line is the half mass radius for singles while the dashed line in the lower panel is the half mass radius for binaries. The lowest curve is the core radius of the system (in units of the initial half mass radius). The simulation has been performed with 16384 particles and 10% of binaries starting from $W_0 = 3$. It is the equivalent of Fig 13 in Fregeau et al. (2003)




Figure 3. Like Fig. 2 for a simulation starting from $W_0 = 7$ with 16384 particles and 10% of binaries. It is the equivalent of Fig 10 in Fregeau et al. (2003)

Note that our definition of $t_{rh}(0)$ differs by a factor $\ln(0.11N)/\ln(0.4N)$ from that used by Fregeau et al.. This has been taken into account in this paper to ensure a proper comparison.




Figure 4. Like Fig. 2 for a simulation starting from $W_0 = 7$ with 16384 particles and 20% of binaries. It is the equivalent of Fig 11 in Fregeau et al. (2003) or Fig.3 in Fregeau et al. (2005). The results from a new, unpublished run by Fregeau (Fregeau 2006, private communication) are overplotted in red.




Figure 5. Like Fig. 2 for a simulation starting from $W_0 = 11$ with 16384 particles and 10% of binaries. It is the equivalent of Fig 12 in Fregeau et al. (2003)

4.2 Core and half mass radius

In the presence of a tidal field, the system evolves toward conditions in the centre very similar to those that we observed for isolated systems. The core radius evolves, usually in a few t_{rh} , to reach a ratio r_c/r_h that is close to that




Figure 6. Evolution of the total mass for a King models with $W_0 = 7$ and different ratios of primordial binaries. The simulations have been performed with 4096 particles.

Table 2. Comparative results on the dissolution timescale $t_{dis/t_{rh}(0)}$ for simulations with f=10~%

W_0	THH field	THH cutoff	Fregeau et al. cutoff
3	5	15	13
7	18	20 - 40	$\gtrsim 38$
11	11	N/A	27

Notes: The values for t_{dis} for our runs (THH) have been extrapolated to $N=3\cdot 10^5$ from N=16384 (see discussion in the text). In the second column we give t_{dis} for our standard tidal field runs, extrapolated by using eq. (8). In the third column we report our results from runs with a tidal cut-off, with constant extrapolation for $W_0=3$ and a range of values for $W_0=7$ (see Sec.5). The fourth column contains the results from Fregeau et al.

attained at the end of core collapse during the evolution of isolated Plummer models (see Sec. 4.1 of Paper I) with a similar fraction of primordial binaries. As is depicted in Fig. 8, the evolution of the core to half mass radius, after a first adjustment phase, is largely independent of the initial conditions considered. Thereafter this ratio is almost constant, as long as there are binaries to burn or until tidal dissolution has reduced the total mass below 10% of the initial value.

We discuss the value of the core radius in further detail below, but summarise here the behaviour of the half mass radius. This depends on the strength of the tidal field, which depends on W_0 (Sec.4.1). For $W_0 = 7$, r_h remains almost constant, up to the final stages of the life of the system, when it eventually falls to zero. For stronger tidal fields (models with $W_0 = 3$ and 11) the half mass radius decreases steadily. Our interpretation is that, for $W_0 = 7$, the tendency toward expansion (due to the energy generated in the core)




Figure 7. Tidal dissolution t_{dis} time for models with $W_0 = 7$, f = 10% and a galactic tidal field, for different values of N, compared to the result by Fregeau et al. (square point; obtained using a tidal cut-off). The points associated with runs using a tidal field are compared with the theoretical scaling (red solid line) given by Baumgardt (2001) for tidal field runs.

is balanced by the mass loss due to the tidal field, while the tide dominates in the other two cases.

Now we return to the evolution of the core radius, and consider first the initial phase of contraction or expansion. The evolution in the first few relaxation times depends on how the initial value of r_c compares with the quasiequilibrium value attained in the intermediate steady binary burning phase. Thus runs starting initially with a relatively big core, such as $W_0 = 3$ $(r_c(0) \approx 0.6r_h(0), \text{ Fig. 2})$ have a rather deep collapse; after about $8t_{rh}(0)$ in our simulation with N = 16384 and f = 10% we measure $r_c/r_h(0) \simeq 0.05$. For $W_0 = 7$, f = 10% and N = 16384 (see Fig. 3) the duration of the initial contraction is shorter, and thereafter the core radius is somewhat larger. On the other hand runs that start with a small, concentrated core $(W_0 = 11)$, have a first rapid expansion of the core radius (see also Fig. 5) up to $r_c/r_h(0) \approx 0.09$ for our N = 16384 run with f = 10%. This can be understood in terms of the energy balance of the core: if the density is too high, the binary heating exceeds the cooling by heat transport to the halo, so that the core has to expand to lower density to reduce the rate of energy generation. Incidentally, though we have focused on numerical values for runs with f = 10%, we find that the effect of primordial binaries saturates by around f = 10% (see Fig. 9), much as we have noted in the evolution of isolated clusters in Paper I (see also Vesperini & Chernoff 1994).

Now we turn to the value of the core radius during the phase of steady binary burning, and, in particular, its dependence on the number of particles. Our results are summarised in Fig. 10. At variance with Paper I, here the Vesperini & Chernoff (1994) model provides an excellent fit to the dependence on N of the ratio of the core to half mass




Figure 8. Evolution of the ratio of the half mass to core radius (upper panel) and of the concentration parameter $c = \log r_t/r_c$ for different King models with f = 10%, 20%. The solid line refers to simulations starting from $W_0 = 7$, the dotted line to $W_0 = 11$ and the dashed line to $W_0 = 3$; the number of particles used is 16384. The evolution quickly erases differences due to initial conditions, and during binary burning c and r_h/r_c evolve very similarly. This figure can be directly compared with Fig. 16 in Fregeau et al. Note that we kept the same range in the y-axis to highlight the differences between our runs and theirs.

radius, provided that in the Coulomb logarithm the value 0.11 is used for the coefficient of N. Thus the formula used for the fit is (see Vesperini & Chernoff 1994 and Paper I):

$$\frac{r_c}{r_h} = \frac{\alpha}{\log_{10}(0.11N)} \frac{\phi_b(1 - \phi_b)\mu_{bs} + \phi_b^2\mu_{bb}}{(1 + \phi_b)^4},\tag{9}$$

where the various quantities are defined as in Paper I. Thus the quantity α is a parameter depending on γ and v_c/v_h , where γ is the ratio of the expansion timescale r_h/\dot{r}_h to t_{rh} , and v_c and v_h are the velocity dispersion in the core and at the half-mass radius, respectively; we have assumed typical values for these parameters: $\gamma = 10.5$ and $v_c/v_h = \sqrt{2}$ (Heggie & Aarseth 1992; Goodman & Hut 1989). The quantity ϕ_b is the binary fraction in the core defined as

$$\phi_b = \frac{n_b}{n_s + n_b}.$$

Finally, μ_{bs} and μ_{bb} are coefficients for the efficiency of binary-single and binary-binary burning and depend on the distribution of binding energy of the binaries. They have been computed as in Paper I assuming a flat distribution in the logarithm of the binding energy from 12 kT to 300 kT. α , μ_{bs} and μ_{bb} do not significantly depend on the number of particles N (see Sec. 5.1.2 in Paper I).

With regard to the core radius there are many quantitative and even some qualitative differences between our results and the earlier results of Fregeau et al. (2003). But they point out (Fregeau et al. 2005) that the core radius is one parameter which has turned out to behave rather differently following the code improvements summarised in their




Figure 9. Dependence of the ratio r_c/r_h on time (in units of $t_{rh}(0)$) for simulations starting from a King model with $W_0 = 7$ and different ratios of primordial binaries. The simulations have been performed with 4096 particles.




Figure 10. Dependence of the ratio r_c/r_h , averaged for $10\ t_{rh}(0)$ after core collapse, on the number of particles. Simulations start from a King model with $W_0=7$ and f=10% using a tidal field (crosses) or isolated runs (squares). Points associated with N from 512 to 2048 have been obtained by averaging over a sample of 16 simulations for each N. A comparison with the Vesperini & Chernoff (1994) model is drawn: the agreement (using log(0.11N) instead of log(0.4N)) is excellent for the tidal field runs. The adopted values for the parameters used are given in the text.




Figure 11. Like Fig. 2 for a simulation starting from $W_0 = 7$ with 4096 particles and 5% of binaries.

later paper. Indeed the one case for which a direct comparison with their most recent results is possible is shown in Fig.4 ($W_0 = 7, f = 20\%$). Both models (our N-body model and their Monte Carlo model) show an initial contraction followed by a phase of steady binary burning. The comparison with the new results by Fregeau et al. (2005) is quantitatively satisfactory, except that, in the phase of steady binary burning, our core radius is about twice as large as theirs. However, it is known that the core radius in this phase is N-dependent (see above, and Paper I), and the sense of the difference is consistent with this.

As already mentioned in Sec.3, the *late* evolution of the core radius depends on whether the binary population can be heavily depleted by the time of dissolution, t_{dis} . The evolution of the binary population is discussed in the following section, and so we defer further discussion until then.

4.3 Evolution of the primordial binary population

The destruction of the primordial binary population is depicted in the top panel of Figs. 2-5. The number of binaries decreases due to both destruction of the softer pairs in the centre of the system and the ejection of stars across the tidal boundary. The changing relative contribution of these two processes can be inferred from Fig. 12, upper panel. At first the ratio of binaries to singles decreases, as at the beginning of the simulation relatively many soft binaries (i.e. those with $E_b \leq 15kT$) are present. However at later times the relative number of binaries to singles starts to rise, as most of the surviving binaries are hard to disrupt in three- or four-body encounters. In addition tidal ejection is more probable for singles, as binaries have sunk toward the core of the cluster by mass segregation (see bottom panel in Figs. 2-5 and Fig. 13).

There are some quantitative differences between our results and those of Fregeau et al., even after the recent im-

provements to their technique. For example they point out (Fregeau et al. 2005) that, for a model with $W_0 = 7$ and f = 20%, the binary fraction shrinks to about 10% before recovering to about its initial value shortly before t_{dis} . This differs considerably from our result in Fig.12 (upper panel). Another example is illustrated in the upper panel of Fig.4, where it is clear that the disruption of binaries in the most recent available run by Fregeau et al. appears to be rather faster than in our run starting from similar initial conditions.

Now we consider models in which the population of binaries can be strongly depleted before dissolution of the cluster, i.e. $t_{bb} < t_{dis}$. We have already remarked (Sec.4.1) that t_{dis} increases as N decreases, and is longest for $W_0 = 7$ (out of the three values we adopted). It has also been argued (Sec.3) that the time taken to burn most binaries is roughly proportional to f. Based on these considerations, it may be expected that the only runs that may burn most of their binaries within the lifetime of the system are those in which f or N is sufficiently small, and that intermediate values of W_0 are most favourable. Indeed, we found that the only runs which did so were those with $W_0 = 7$ and $N \lesssim 4096$ and $f \lesssim 5\%$. (Note, however, that $W_0 = 7$ is the only value of W_0 for which we carried out runs with N < 4096, see Table 2.) At the point where the binaries become heavily depleted, there is a deep core contraction, similar to what happens in runs with single stars only; however, here N is too low for gravothermal oscillations to occur. In Fig 11 we depict the evolution of a $W_0 = 7$ model starting from f = 5% and N = 4096: by about 30 $t_{rh}(0)$ the fraction of binaries has fallen below 20 % of its initial value and the core radius contracts sharply.

Next we comment in particular on the evolution of the binary population in the core. At the end of our simulations starting from f = 10 - 20% we find that the fraction of binaries in the core has increased by at least a factor 4. In general the density of binaries in the core is in broad agreement with the numbers observed in Paper I and does not depend significantly on the details of the initial conditions if f is in this range (see bottom panel of Fig. 12, where runs starting from $W_0 = 7$ and $W_0 = 11$ show a rather similar behaviour). However there are differences in detail in the time evolution. In an isolated cluster the proportion of binaries in the core, after the sharp initial increase caused by mass segregation, declines slowly; in these tidally limited models, on the other hand, there is a slow increase. The $W_0 = 3$ King model is an extreme example, as there is an enhanced rate of ejection of singles relative to the rate of disruption and ejection of binaries, because of the relatively small tidal radius; in this situation the binary fraction in the core increases rapidly throughout the evolution.

The evolution of the internal binding energy (see Fig. 14) is similar to that observed for isolated systems, with a preferential decrease in the number of less bound binaries. Furthermore, as expected, there is a significant correlation of the internal binding energy with the radius (see Fig. 15). After approximately 15 $t_{rh}(0)$ the survival probability in the core for binaries with binding energy below $\approx 20~kT$ is low, as they have either been destroyed or hardened. We thus see that softer binaries are mainly present around the half mass radius and in the halo, while the core is dominated by hard binaries (with occasionally some short-lived low energy binaries that have been formed dynamically). At later




Figure 12. Dependence of the binary fraction on time (in units of $t_{rh}(0)$) for simulations starting from King models with $W_0 = 3$ (dotted lines), $W_0 = 7$ (solid lines), and $W_0 = 11$ (dashed lines) for f = 10%, 20%. The upper panel refers to the total number of binaries, the lower to the binaries in the core. The simulations have been performed with 16384 particles.

stages of the evolution (see bottom right panel of Fig. 15), the softer binaries have almost completely disappeared.

This picture is in qualitative agreement with the results already known in the literature (see e.g., McMillan & Hut 1994; Giersz & Spurzem 2000 and Fregeau et al.). As our resolution is limited by the modest number of binaries that are present in our runs, it is hard to attempt any quantitative comparison.

5 TIDAL CUTOFF RUNS: RESULTS AND COMPARISON

It was pointed out in Sec.4.1.3 that our result on the lifetime differed significantly (at least in the one case considered) from that of Fregeau et al. This conclusion, however, required a knowledge of the N-dependence of the lifetime, and it is known that this in turn depends on the treatment of the tide, which differs between the work of Fregeau et al. and the N-body simulations which we have presented so far. For the case of a tidal cutoff, as adopted by Fregeau et al., it was found by Baumgardt (2001) that the half-mass dissolution time-scale t_{Half} (expressed in relaxation times) does not depend on the number of particles of the system (as we assumed for the purpose of our comparison) if $N \gtrsim 4096$. Note, however, that Baumgardt (2001) used a fixed tidal radius and, as we have mentioned, measured the timescale of mass-loss by the half-mass time t_{Half} , whereas our discussion has focused on the time for loss of 98% of the mass, t_{dis} .

Because of these complications, for a better comparison with Fregeau et al.we have run a second series of simulations,




Figure 13. Evolution of the distribution of binary radius in units of the instantaneous core radius. The ordinate is the number of binaries per bin, which are of length 0.5 in \log_e . Curves have unit normalization, and are shown for $t/t_{rh}(0) = 0, 8, 16, 24$. The dashed line is the position of the half mass radius. The simulation has been performed with 16384 particles and 20% of binaries starting from a King model with $W_0 = 7$.




Figure 14. Evolution of the distribution of binary binding energy (given in units of kT). The ordinate is the number of binaries per bin, which are of unit length in \log_2 . Curves have unit normalization, and are shown for $t/t_{rh}(0)=0,8,16,24$. The simulation has been performed with 16384 particles and 20% of binaries starting from a King model with $W_0=7$.




Figure 15. Evolution of the distribution of binaries in radius and binding energy. The radius is given in units of the instantaneous core radius; the energy in units of kT. The panels refer to $t/t_{rh}(0) = 0, 8, 16, 24$. The dashed line is the position of the half mass radius. The simulation has been performed with 16384 particles and 20% of binaries starting from a King model with $W_0 = 7$. It can be compared to Fig. 15 in Fregeau et al. (2003)

in which the tidal field we have used hitherto has been replaced by a tidal cutoff. We have ensured that the cutoff radius decreases in the correct manner (eq.(4)) as the mass inside this radius decreases, exactly as adopted by Fregeau et al. The initial parameters of these models are $W_0 = 7$ and $W_0 = 3$, f = 10%, with N in the range 512 to 16384.

For the series starting from $W_0=3$ with f=10 % (i.e. with a strong tidal field, Sec.4.1.2), we do not observe a significant N-dependence of the half-mass dissolution time t_{Half} in units of the relaxation time, except for a modest ($\approx 10\%$) variation from N=4096 to N=8192. This is consistent with the experiments (without primordial binaries) by Baumgardt (2001), despite his use of a constant tidal field. Moreover, it is also consistent with unpublished results by one of us using the initial conditions of the collaborative experiment summarised in Heggie (2003), except for replacement of the tidal field by a tidal cutoff; in this case the cutoff was evolved in the correct manner as mass was lost. For the $W_0=3$ runs with tidal cut-off the agreement with the dissolution time t_{dis} of Fregeau et al. (2003) is good: we extrapolate $t_{dis}\approx 15~t_{rh}(0)$ and they have $t_{dis}=13~t_{rh}(0)$.

By contrast, for the series with $W_0 = 7$ and f = 10 % and a cutoff, we observe an N-dependence similar to that found when a tidal field is used, as in our standard runs (see Fig. 16). On the other hand it is not known why the N-dependence should differ for the cases $W_0 = 3$ and 7 when a cutoff is used. One possibility is that it is a "small-N" effect which is more important in a system with a small core; if so, the N-dependence of the dissolution time t_{dis} will eventually flatten out for larger N. The presence of primordial binaries does not seem to be the issue; we have run a few simulations

with a tidal cut-off, starting from a $W_0=7$ model and no binaries, and find that the half-mass dissolution time t_{Half} is almost equal to that measured when binaries are present. Tanikawa & Fukushige (2005) have also observed differences (in the scaling of the half-mass time t_{Half} with N) between models with $W_0=3$ and 7, though their models adopted a tidal field, and the potential was softened. They tentatively attribute the difference to the time taken for a particle to escape.

Whatever the explanation, this (unexpected) N dependence of t_{dis} in our tidal cutoff runs with $W_0 = 7$ makes the extrapolation of our results obtained with the tidal cut-off prescription highly uncertain. While the dissolution time of our model with $W_0 = 7$, f = 10% and N = 16384 in a tidal cutoff is $t_{dis} \simeq 41.2t_{rh}(0)$ (Table 3), the inferred dissolution time for a $W_0 = 7$ f = 10% model with $N = 3 \cdot 10^5$ may be in the range of $20-40t_{rh}(0)$ (Table 2, col.3), depending on whether t_{dis} is proportional to $t_{rh}(0)$ or scales in accordance with eq.(8). If the actual value is close to the upper limit, then the tidal dissolution time t_{dis} in our $W_0 = 7$ cutoff models would compare marginally well with Fregeau et al., as they find $t_{dis} \gtrsim 38$. Furthermore, recent improvements to their code appear to have reduced the lifetime slightly for a model with $W_0 = 7, f = 20\%$, and a similar reduction for f = 10% would improve the agreement between our N-body and their Monte Carlo results.

We close this section with a brief comparison between other properties of our runs with a tidal field and those with a tidal cutoff. Because of the limited scope of our runs with a tidal cutoff, we are concerned less here with general trends; but these results will be useful for comparison with simplified treatments in which it is necessary to use a cutoff, such as Fokker-Planck models; they also illustrate the kinds of results which may be sensitive to the manner in which the tidal boundary conditions are treated.

For our run with $W_0 = 3$, f = 10%, N = 16384 and a cutoff (Fig. 17), the initial contraction of the core radius is slower than that observed in the run with a tidal field (see Fig. 2): after $8t_{rh}(0)$, r_c is approximately twice as large in the cutoff run. However just before tidal dissolution the values of the core radius in the two runs are quite similar. Actually, if the time is normalized by the dissolution time t_{dis} , the evolution of r_c looks very similar, except at the very last moment, where r_c decreases for the run with a cutoff.

By contrast, the run with $W_0=7$, $f=10\,\%$, N=16384 and a cutoff (Fig. 18) is much more similar to the corresponding run with a tidal field (Fig. 3). The dissolution time t_{dis} scales for the two tidal treatments differ by only $\approx 25\%$, and the core radius evolves in a similar way. However the half mass radius for the run with a cutoff initially increases in much the same way as in the run reported by Fregeau et al., while in our run with a tidal field it remains almost constant.

6 DISCUSSION AND CONCLUSIONS

In this paper we have continued our investigation of the dynamical evolution of stellar systems with primordial binaries. In Paper I we considered simple isolated models with equal mass stars and a primordial binary population in the range 0-100 %. Then the main parameters that




Figure 16. Tidal dissolution time t_{dis} for a $W_0 = 7 f = 10\%$ model with a tidal cutoff, with different values of N, compared to the result by Fregeau et al. (square point; obtained using a tidal cut-off). The lower series (3-sided stars) gives the half mass dissolution time t_{Half} , compared to some runs starting from $W_0 = 7$ and no binaries (triangles). The scaling with N of both t_{dis} and t_{Half} appears similar.




Figure 17. Like Fig. 2, but for a tidal cutoff simulation starting from $W_0 = 3$ with 16384 particles and 10% of binaries. It is the equivalent of Fig 13 in Fregeau et al. (2003)




Figure 18. Like Fig. 3 for a tidal cutoff simulation starting from $W_0 = 7$ with 16384 particles and 10% of binaries. It is the equivalent of Fig 10 in Fregeau et al. (2003)

control the evolution of the system are the primordial binary fraction f and the number of bodies N. However, for $f\gtrsim 10\%$ the efficiency of energy generation due to three-and four-body encounters saturates, so that little difference was observed in the size of the core radius for runs starting with the same number of particles and $f\gtrsim 10\%$. This behaviour had already been predicted by the theoretical model of Vesperini & Chernoff (1994), and our runs confirmed this expectation. Interestingly, however, the scaling of the core radius with the number of particles was not quite the same as that predicted by Vesperini & Chernoff (1994).

Here we have added an important new ingredient to our simulations: a tidal field. The presence of a tidal field introduces a new time-scale in the simulation, the time needed for the dissolution of the system, which is usually shorter than the time needed to deplete the primordial binary population for simulations starting from $f\gtrsim 10\%$ and $N\gtrsim 4096$. We have considered runs starting from King models with initial central potential $W_0=3,7,11$ (and thus a different intensity of the tidal field, in the units adopted) with 0 to 100 % primordial binaries, and N from 512 to 16384.

By comparison with the results of Paper I, our Fig. 8 clearly confirms that the evolution of the core to half mass radius proceeds much as for isolated models, after an initial transient; the models starting with $W_0 = 7$ and 11 evolve very similarly, for initial binary fractions of f = 10%, 20%. The King model with $W_0 = 3$, however, nicely illustrates the effects of a strong tidal field: the evolution of r_c/r_h tries to go toward the steady value reached by the other runs with higher W_0 , but the tidal field manages to dissolve the system just when this common r_c/r_h value is about to be attained.

One objective of this paper was also to compare the results obtained by means of our direct N-body simulations with the outcome of simulations able to employ realistic number of particles by using approximate methods,

like Monte-Carlo and/or Fokker Planck approaches. Just as in Paper I we compared our work with the milestone study of Gao et al. (1991), here we took as comparison the recent work carried out by Fregeau et al. There are considerable differences with the results reported in Fregeau et al. (2003). but they have since reported that several results change significantly following an improved treatment of binary interactions (Fregeau et al. 2005). As a result of these improvements they have discovered that the emission of energy in encounters with binaries was too high in the earlier models. Therefore we have focused our comparison on one case which they illustrated in some detail in their later paper. While it is promising to report that most differences have been cleared up by the improvements, there are exceptions. One is that the binaries still appear to be depleted too rapidly, even in the most recent runs (Fig.12; cf. Fregeau et al. (2005), Fig.3). There remain uncertainties also about the lifetime, but this depends on perplexing problems in the scaling with N: we find evidence that the tidal cutoff scaling derived by Baumgardt (2001) for single stars with a fixed cutoff radius does not seem to be applicable to the self-consistent prescription of a cutoff radius prescription used by us (for some simulations) and by Fregeau et al. Therefore the best approach to understanding the remaining differences between our runs and those of Fregeau et al. would be to directly compare a set of Monte Carlo simulations starting from the same initial conditions as we have adopted.

We finally note that, in the presence of a tidal field, the prediction of the Vesperini & Chernoff (1994) model, on the size of the core as a function of the number of particles used in the simulation, is in detailed quantitative agreement with the values measured in our set of simulations with $W_0 = 7$. This fact comes as a surprise, because, as we discussed in Paper I, the N dependence given by Vesperini & Chernoff (1994) was not verified for isolated clusters: however, the model does not make any assumption about the presence or absence of a tidal field. The presence of a tidal field seems thus to introduce a difference in the scaling with N of the ratio r_c/r_h in the steady burning phase, which it would be interesting to understand better.

ACKNOWLEDGEMENTS

We are indebted to Sverre Aarseth for providing his code NBODY6. We thank John Fregeau, Fred Rasio and Atakan Gürkan for interesting discussions and for critical comments on a draft of the manuscript; John Fregeau also kindly supplied us with some of his unpublished data. We are very grateful to the referee for his help in making us see places where improvements were needed, and in particular for his observation that the half-mass time scales well with the total-mass relaxation time. Finally we thank Andreas Ernst for useful suggestions. MT thanks Prof. Mineshige for his kind hospitality at the Yukawa Institute at Kyoto University, through the Grant-in-Aid 14079205 of the Ministry of Education, Culture, Sports, Science and Technology, Japan. P.H. thanks Prof. Ninomiya for his kind hospitality at the Yukawa Institute at Kyoto University, through the Grantsin-Aid for Scientific Research on Priority Areas, number 763, "Dynamics of Strings and Fields", from the Ministry of Education, Culture, Sports, Science and Technology, Japan.

REFERENCES

- Aarseth S., 2003, Gravitational N-body Simulations. Cambridge University Press
- Albrow M. D., Gilliland R. L., Brown T. M., Edmonds P. D., Guhathakurta P., Sarajedini A., 2001, Ap.J., 559, 1060
- Baumgardt H., 2001, MNRAS, 325, 1323
- Bellazzini M., Fusi Pecci F., Messineo M., Monaco L., Rood R. T., 2002, A.J., 123, 1509
- Casertano S., Hut P., 1985, Ap.J., 298, 80
- Chernoff D. F., Huang X., 1996, in IAU Symp. 174: Dynamical Evolution of Star Clusters: Confrontation of Theory and Observations, ed. P. Hut & J. Makino (Dordrecht: Kluwer), 263
- Fregeau J. M., Gürkan M. A., Joshi K. J., Rasio F. A., 2003, Ap.J., 593, 772
- Fregeau J. M., Gürkan M. A., Rasio F. A., 2005, to appear in Few-Body Problem: Theory and Computer Simulations, ed C. Flynn, Annales Universitatis Turkuensis; also astro-ph/0512032
- Giersz M., Spurzem R. 2000, MNRAS, 317, 581
- Gao B., Goodman J., Cohn H., Murphy B., 1991, Ap.J., 370, 567
- Goodman J., Hut P., 1989, Nature, 319, 40
- Goodman J., Hut P., 1993, Ap.J., 403, 271
- Heggie D.C., 1975, MNRAS, 173, 729
- Heggie D. C., 2003, IAUS 208, 103
- Heggie D. C., Aarseth S. J., 1992, MNRAS, 257, 513
- Heggie D.C., Hut P., 2003, The Gravitational Million-Body Problem, Cambridge, Cambridge University Press
- Heggie D. C., Mathieu R. D., 1986, in LNP 267: The Use of Supercomputers in Stellar Dynamics (Springer: Berlin), p.233
- Heggie D.C., Trenti M., Hut P., 2006, MNRAS, 368, 677
- Hut P., McMillan S., Goodman J., Mateo M., Phinney E. S., Pryor C., Richer H. B., Verbunt F., Weinberg M., 1992, PASP, 104, 981
- Joshi K. J., Rasio F. A., Portegies Zwart S., 2000, Ap.J., 540, 969
- Makino J., Hut P., 1990, Ap.J., 365, 208
- McMillan S., Hut P., Makino J., 1990, Ap.J., 362, 522
- McMillan S., Hut P., 1994, Ap.J., 427, 793
- Portegies Zwart S.F., McMillan S. L. W., 2002, Ap.J., 428, 724
- Portegies Zwart S. F., McMillan S. L. W., 2004, ASP Conference Series, 322, 459
- Portegies Zwart S.F., Baumgardt, H., Hut, P., Makino, J., McMillan, S. L. W. 2004, Nature, 428, 724
- Pulone L., De Marchi G., Covino S., Paresce F., 2003, A&A, 399, 121
- Rubenstein E. P., Bailyn C. D., 1997, Ap.J., 474, 701
- Spitzer L., Jr, 1987, Dynamical evolution of globular clusters, (Princeton University Press: Princeton)
- Sugimoto D., Bettwieser E., 1983, MNRAS, 204, 19P
- Tanikawa A., Fukushige T., 2005, PASJ, 57, 155
- Vesperini E., Chernoff D. F., 1994, Ap.J., 431, 231
- This paper has been typeset from a TEX/ LATEX file prepared by the author.

Table 3. N-body models with Fregeau et al.-like initial conditions. In this table we report summary information for all runs with $N \geqslant 4096$. For runs with smaller N we report averaged values over 16 simulations for each entry line. The entries at the bottom of the table refer to simulations with tidal cut-off as in Fregeau et al. Column entries are: the number of bodies N, the initial value of W_0 , the primordial binary ratio f, time of core collapse t_{cc} , time of dissolution t_{dis} (computed at the point where only 2% of the initial mass of the system is left), core radius at core collapse r_c , and core to half mass radius after core collapse $r_c/r_h(t_{cc})$ (averaged for 5 $t_{rh}(0)$ after core collapse), fraction of binaries to singles in the core after core collapse $(N_b/N_s)_c(t_{cc})$ (same average as for core to half mass radius), and relative mass of the system at core collapse $M(t_{cc})/M(0)$.

N	W_0	f	t_{cc}	$t_{dis}(98\%)$	$\tilde{r}_c(t_{cc})$	$r_c(t_{cc})$	$r_c/r_h(t_{cc})$	$(N_b/N_s)_c(t_{cc})$	$M(t_{cc})/M(0)$
512	7.0	10	1.6	81.2	0.147	0.161	0.179	0.35	0.99 (averaged)
1024	7.0	10	4.0	66.8	0.129	0.147	0.157	0.44	0.97 (averaged)
2048	7.0	10	5.6	59.7	0.108	0.128	0.143	0.47	0.92 (averaged)
4096	3.0	0	11.9	17.3	0.019	0.023	0.081	0.04	0.26
4096	3.0	0	11.7	16.8	0.018	0.022	0.080	0.03	0.27
4096	3.0	10	11.1	13.8	0.071	0.075	0.210	2.84	0.15
4096	3.0	10	10.5	13.0	0.073	0.080	0.175	2.53	0.17
4096	3.0	20	9.5	13.4	0.085	0.097	0.230	4.48	0.23
4096	3.0	20	10.0	13.0	0.066	0.074	0.235	6.39	0.17
4096	3.0	50	9.8	13.9	0.075	0.090	0.190	8.84	0.24
4096	3.0	100	14.1	16.5	0.062	0.072	0.274	34.05	0.09
4096	7.0	0	11.5	50.3	0.015	0.022	0.030	0.02	0.79
4096	7.0	0	10.8	47.7	0.014	0.021	0.031	0.02	0.77
4096	7.0	2	20.8	51.8	0.035	0.045	0.041	0.03	0.52
4096	7.0	2	17.4	48.4	0.036	0.046	0.053	0.06	0.61
4096	7.0	5	4.1	52.2	0.091	0.110	0.129	0.24	0.95
4096	7.0	5	14.0	53.4	0.063	0.078	0.089	0.23	0.69
4096	7.0	10	3.3	48.0	0.100	0.121	0.132	0.61	0.95
4096	7.0	10	7.6	45.8	0.096	0.117	0.126	0.57	0.84
4096	7.0	20	2.0	41.4	0.132	0.158	0.132	1.24	0.98
4096	7.0	20	6.6	40.2	0.097	0.118	0.130	1.44	0.83
4096	7.0	50	7.5	44.0	0.113	0.141	0.163	4.24	0.84
4096	7.0	100	3.4	55.1	0.147	0.189	0.203	11.08	0.93
4096	11.0	0	0.0	28.7	0.019	0.028	0.034	0.02	1.00
4096	11.0	0	0.0	32.2	0.021	0.030	0.031	0.02	1.00
4096	11.0	10	0.0	25.1	0.082	0.103	0.121	0.54	1.00
4096	11.0	10	0.0	30.9	0.079	0.097	0.122	0.56	1.00
4096	11.0	20	0.0	23.8	0.074	0.094	0.128	1.08	1.00
4096	11.0	20	0.0	26.7	0.082	0.102	0.137	1.03	1.00
4096	11.0	50	0.0	28.4	0.080	0.101	0.158	2.86	1.00
4096	11.0	100	0.0	32.4	0.082	0.107	0.179	11.93	1.00
8192	7.0	10	7.1	38.7	0.074	0.092	0.108	0.77	0.82
16384	3.0	10	9.4	9.6	0.034	0.038	0.080	6.05	0.02
16384	3.0	20	8.5	9.2	0.036	0.043	0.241	6.75	0.05
16386	7.0	10	7.2	32.4	0.069	0.086	0.097	1.16	0.76
16384	7.0	20	6.2	28.5	0.076	0.096	0.095	2.53	0.81
16384	11.0	10	0.0	20.1	0.056	0.074	0.102	0.78	1.00
16384	11.0	20	0.0	19.3	0.069	0.090	0.105	1.77	1.00
512	7.0	10	1.8	83.9	0.137	0.153	0.196	0.35	0.99 (averaged + tidal cut-off)
1024	7.0	10	3.5	75.4	0.124	0.142	0.172	0.39	0.96 (averaged + tidal cut-off)
2048	7.0	10	4.7	67.2	0.111	0.132	0.146	0.53	0.94 (averaged + tidal cut-off)
4096	7.0	10	3.5	56.3	0.095	0.118	0.129	0.64	0.94 (tidal cut-off)
8192	7.0	10	10.9	50.4	0.077	0.096	0.108	0.74	0.78 (tidal cut-off)
16384	7.0	10	7.6	41.2	0.065	0.084	0.088	0.88	0.83 (tidal cut-off)
4096	3.0	10	12.9	17.0	0.061	0.068	0.275	2.30	0.22 (tidal cut-off)
8192	3.0	10	13.5	14.8	0.047	0.048	0.248	5.53	0.09 (tidal cut-off)
16384	3.0	10	14.2	15.3	0.041	0.046	0.272	8.31	0.07 (tidal cut-off)
512	7.0	10	1.6	N/A	0.141	0.157	0.193	0.32	1.00 (averaged, isolated)
1024	7.0	10	3.2	N/A	0.131	0.152	0.155	0.40	0.99 (averaged, isolated)
2048	7.0	10	4.9	N/A	0.112	0.134	0.128	0.50	0.99 (averaged, isolated)
4096	7.0	10	5.6	N/A	0.104	0.128	0.107	0.59	0.98 (averaged [5 sim.], isolated)
8192	7.0	10	9.7	N/A	0.082	0.104	0.094	0.64	0.97 (isolated)