Properties of ⁸Be and ¹²C deduced from the folding-potential model

P. Mohr, H. Abele, V. Kölle, G. Staudt Physikalisches Institut, Univ. Tübingen, D-72076 Tübingen, Germany

H. Oberhummer and H. Krauss

Institut für Kernphysik, Technische Universität Wien, A-1040 Vienna, Austria

Abstract: The α - α differential cross sections are analyzed in the optical model using a double-folded potential. With the knowledge of this potential bound and resonance-state properties of α -cluster states in 8 Be and 12 C as well as astrophysical S-factors of 4 He(α , γ) 8 Be and 8 Be(α , γ) 12 C are calculated. Γ_{γ} -widths and B(E2)-values are deduced.

PACS: 25.55.Ci, 25.70.Ef, 24.10.Ht

In order to determine reaction rates for the triple–alpha process [1], we have calculated in the potential model the $\alpha+\alpha$ differential cross section as well as the bound and resonance–state properties of α –cluster states in ⁸Be and ¹²C. The potentials are calculated using the folding procedure [2], [3].

In this approach the nuclear densities are derived from nuclear charge distributions [4] and folded with an energy and density dependent NN-interaction $v_{\rm eff}$ [2]. By means of a normalization factor λ the depth of the potential can be adjusted to elastic scattering data and to bound and resonant state energies of nuclear cluster states. In this work the folding potentials were determined using the computer code DFOLD [5]. The imaginary part of the optical potential can be neglected at the low energies considered.

Using the folding procedure for the $\alpha+\alpha$ scattering potential we fitted the experimental differential cross section for all projectile energies below the break-up threshold ($E_{\rm lab}\approx 7\text{--}35\,{\rm MeV}$) with a renormalization factor $\lambda=1.656$. An excellent agreement between the calculated and experimental cross section has been found [6]. With the same potential we reproduce the resonance energies and widths of the ⁸Be ground–state: $E_{\rm R}^{\rm th}=92.1\,{\rm keV}$ and of the 2^+- and 4^+ –state in ⁸Be (see Table 1). Especially for the ground state the calculated values agree excellently with the experimental data. This result confirms that this nucleus can be well described in a two-cluster approach [7]. This marked α –cluster structure is also reflected by the values of the α –particle spectroscopic–amplitudes for these ⁸Be–states [8]: $S(g.s.)=1.48, S(2^+)=1.46, \text{ and } S(4^+)=1.39.$

The s-wave function for the relative motion of the two α -clusters in the ⁸Be ground state can be folded with the nucleon densities of both α -particles to calculate the ⁸Be density [6]. In our calculation we used an upper cut-off radius of 20 fm. Averaging over 4π we obtain the nucleon density distribution. The result is shown in Fig. 1. With the knowledge of the ⁸Be density distribution we are able

Table 1. Relevant data of resonant states in ⁸Be

J^{Π}	$E^{\rm calc} \ [{ m MeV}]$	E^{\exp} [MeV] [9]	$\Gamma_{\alpha}^{\mathrm{calc}} \ [\mathrm{eV}]$	$\Gamma_{\alpha}^{\text{exp}}[\text{eV}] [9]$
0 ⁺	0.0921	0.09212(5)	$6.86 \\ 1.3 \cdot 10^{6} \\ \stackrel{>}{\scriptstyle \sim} 3 \cdot 10^{6}$	6.8 ± 1.7
2 ⁺	2.78	3.13(3)		$1.5(2) \cdot 10^{6}$
4 ⁺	11.02	11.5(3)		$\approx 3.5 \cdot 10^{6}$

Table 2. Relevant data of bound and resonant states in ¹²C

J^{Π}	E [MeV]	λ	$\Gamma_{\alpha}^{\mathrm{calc}} \ [\mathrm{eV}]$	$\Gamma_{\alpha}^{\text{exp}}[\text{eV}] [10]$
0_{1}^{+}	0.0	1.144	_	_
$0_{1}^{+} \ 2_{1}^{+} \ 0_{2}^{+} \ 2_{2}^{+}$	4.437	1.010		
0_{2}^{+}	7.654	1.286	7.5	8.3 ± 1.0
2_{2}^{+}	10.3	1.152	$\approx 2.0 \cdot 10^6$	$(3.0\pm0.7)\cdot10^6$

to calculate the optical potential for the α^{-8} Be system using the folding procedure again. Since no experimental phase shifts for α^{-8} Be are available, we have to fit the normalization factors λ of the folding potentials to the bound and resonance–state energies of 12 C.

The ground state (0_1^+) , the first (2_1^+) and second (0_2^+) excited, and the (2_2^+) state in are those states which are of interest in the triple–alpha process (see Table 2). With respect to a $\alpha \otimes^8$ Be clustering in 12 C, the 0_1^+ and 2_1^+ bound states belong to the ground state band with a harmonic oscillator quantum number Q=4, whereas for the 0_2^+ – and 2_2^+ –state we assume that they can be interpreted as members of a band with Q=6. We assume the broad 10.3 MeV level in 12 C to be the 2_2^+ state. The adopted spin assignment of this level is 0^+ [10]. However, an assignment of 2^+ for this state is strongly suggested by experimental results [11] as well as by microscopic calculations [12].

Table 3. Γ_{γ} -widths and B(E2)-values

	$E_{\rm f} \ [{ m MeV}]$	J_f^π	E_i [MeV]	$J_{ m i}^\pi$	$\Gamma_{\gamma} \ [{ m meV}]$	B(E2) [W.u.]
⁸ Be ¹² C	0.0 4.437 7.654	0^{+} 2_{1}^{+} 0_{2}^{+}	3.04 7.654 10.3	2^{+} 0_{2}^{+} 2_{2}^{+}	8.4 4.1 17	75 8.8 100

Using the double–folded α^{-8} Be potential we have fitted the energies of these 12 C states. These states have quite a different $\alpha \otimes ^8$ Be(g.s.) cluster structure as can be seen from the values of the spectroscopic factors: S(g.s.)=0.557 [8], S(2₁⁺)=0.154 [13], and S(12 C, 0₂⁺)=1.8 [14]. Therefore, the α^{-8} Be folding potentials have to be renormalized separately to the ground and excited states in 12 C. The normalization factors λ for the potential obtained in this way are listed in Table 2. Using the optical model we also calculated the alpha widths Γ_{α} of the resonant 0₂⁺ and 2₂⁺ states. The good agreement between the experimental and the calculated α -width for the 0₂⁺-state confirms the assumption that this state has a marked $\alpha \otimes ^8$ Be(g.s.)-structure.

Furthermore, we calculated the cross sections and astrophysical S-factors of the reactions ${}^4\text{He}(\alpha,\gamma_0){}^8\text{Be}$ and ${}^8\text{Be}(\alpha,\gamma_1){}^{12}\text{C}$ in the direct capture model [3], [15]. These cross sections are essentially determined by the overlap of the scattering wave functions in the entrance channel, the bound state wave functions in the exit channel, and the electric quadrupole operator. The wave functions were calculated using the α - α and α - ^8Be potential, respectively. In order to compute the reaction ${}^4\text{He}(\alpha,\gamma_0){}^8\text{Be}$ the resonance energy of the ${}^8\text{Be}$ ground state was lowered by about 100 keV. This means that this state was assumed to be weakly bound. For the description of the bound states the α -particle spectroscopic factors given above are used. In Fig. 2 the cross section of the resonant E2-capture into the ${}^8\text{Be}$ ground state is shown. Excellent agreement is found with the result of a calculation in a bremsstrahlung model [16]. Approximating the $\sigma(E)$ cross sections near the resonance energy by a Breit-Wigner parametrization, Γ_{γ} -widths and B(E2)-values can be deduced. These quantities are given in Table 3.

Acknowledgments: We want to thank the DFG-project Sta290/2, and the Austrian Science Foundatio (FWF) (project P8806-PHY).

References

- [1] Oberhummer, H., Krauss, H., Grün, K., Rauscher, T., Abele, H., Mohr, P., Staudt, G.: this volume
- [2] Kobos, A. M., Brown, B. A., Lindsay, R., Satchler, G. R.: Nucl. Phys. A425, 205 (1984)
- [3] Oberhummer, H., Staudt G.: in Nuclei in the Cosmos. Ed. H. Oberhummer,p. 29. Berlin: Springer-Verlag 1991
- [4] De Vries, H., Jager, C. W., de Vries, C.: At. Data and Nucl. Data Tables 36, 495 (1987)
- [5] Abele H.: computer code DFOLD, Univ. Tübingen, unpublished

- [6] Abele H.: Ph.D. thesis, Univ. Tübingen 1992
- [7] Wildermuth, K., Tang, Y. C.: A Unified Theory of the Nucleus. Braunschweig: Vieweg & Son 1977
- [8] Kurath, D.: Phys. Rev. C7, 1390 (1973)
- [9] Ajzenberg-Selove, F.: Nucl. Phys. **A490**, 1 (1988)
- [10] Ajzenberg-Selove, F.: Nucl. Phys. **A506**, 1 (1990)
- [11] Jacquot, C., Satamoto, Y., Jung, M., Girardio, L.: Nucl. Phys. **201**, 247 (1973)
- [12] Fujiwara, Y. et al.: Prog. Theor. Phys. Suppl. **68**, 29 (1980)
- [13] Kwaśniewicz, E., Jarczyk, L.: J. Phys. **G12**, 697 (1986)
- [14] Fukushima, Y., Kamimura, N.: J. Phys. Soc. Japan 44, 225 (1978)
- [15] Mohr, P., Abele, R. Zwiebel, G., Staudt, G., Krauss, H., Oberhummer, H. Denker A., Hammer J. W., Wolf, G.: Phys. Rev. C, in press
- [16] Langanke, K.-H., Rolfs, C.: Phys. Rev. C33, 790 (1986)

Captions to figures

- Fig. 1: Nucleon density distribution of 8 Be calculated in the potential model Fig. 2: Capture cross section of 4 He $(\alpha,\gamma_0)^8$ Be near the 2^+ resonance