

DISTANCE MEASUREMENT OF GALAXIES TO REDSHIFT OF ~ 0.1 USING
THE CO-LINE TULLY-FISHER RELATION

Yoshiaki SOFUE^{1,2}, Franz SCHÖNIGER¹, Mareki HONMA¹, Yoshinori TUTUI¹,
Takashi ICHIKAWA², Ken-ichi WAKAMATSU³,
Ilya KAZES⁴, and John DICKEY⁵

E-mail: sofue@mtk.ioa.s.u-tokyo.ac.jp

(1) Institute of Astronomy, University of Tokyo, Mitaka, Tokyo 181, Japan

(2) Kiso Observatory, IoA, University of Tokyo, Kiso-gun, Nagano 397-01, Japan

(3) Dept. Physics, Gifu University, Gifu, 501-11, Japan

(4) Observatoire de Paris, F-92190 Meudon, France

(5) Dept. Astronomy, Univ. Minnesota, 116 Church St. SE,

Minneapolis, MN 55455, USA

(to appear in PASJ, 1996)

Abstract:

We report on the first results of a long-term project to derive distances of galaxies at cosmological distances by applying the CO-line width-luminosity relation. We have obtained deep CO-line observations of galaxies at redshifts up to $cz \sim 29,000 \text{ km s}^{-1}$ using the Nobeyama 45-m mm-wave Telescope, and some supplementary data were obtained by using the IRAM 30-m telescope. We have detected the CO line emission for several galaxies, and used their CO line widths to estimate the absolute luminosities using the line-width-luminosity relation. In order to obtain photometric data and inclination correction, we also performed optical imaging observations of the CO-detected galaxies using the CFHT 3.6-m telescope at high resolution. The radio and optical data have been combined to derive the distance moduli and distances of the galaxies, and Hubble ratios were estimated for these galaxies. We propose that the CO line width-luminosity relation can be a powerful method to derive distances of galaxies to redshift of $z \sim 0.1$ and to derive the Hubble ratio in a significant volume of the universe.

Key words: Cosmology – Galaxies: general – Distance scale – Radio lines: CO molecular

1. Introduction

The HI-line width - luminosity relation (Tully-Fisher relation) is one of the most powerful tools to measure distances to galaxies (Tully & Fisher 1977; Aaronson et al. 1986; Pierce & Tully 1988; Kraan-Korteweg et al. 1988; Fouqué et al. 1990; Fukugita et al. 1991). However, distances to galaxies so far reached by HI observations are limited to around 100 Mpc, or $cz \sim 10,000$ to $15,000 \text{ km s}^{-1}$, even with the use of the world's-largest telescopes (Schöniger and Sofue 1993). We have no routine method to determine distances to galaxies beyond this distance, at which beam sizes of a few arc-minutes at $\lambda 21 \text{ cm}$ become too large to resolve individual galaxies in a cluster. Interferometers like the VLA are not very useful for this purpose because of the limited number of spectral channels (velocity resolution). Furthermore, the red-shifted HI frequency results in increases in beam size as well as increased interference, which also makes observations of distant cluster galaxies difficult. Moreover, HI line profiles are easily disturbed by interactions among galaxies, which is inevitable in the central region of a cluster, causing uncertainty in the interpretation of HI line profiles for the Tully-Fisher relation. On the other hand, molecular gas is tightly confined to the luminous stellar disk, so that it is less affected by tidal interactions and by ram-pressure distortion due to the intra-cluster gas. The molecular gas is distributed to a radius of several to ten kpc, so that the integrated line profiles manifest the maximum velocity part of the rotation curve (Sofue 1992).

Molecular-line observations at millimeter wavelengths, particularly in the CO-line emission at 115 GHz, can be achieved with much sharper beams. Therefore, we will be able to resolve individual member galaxies

in a cluster more easily, which makes it possible to avoid contamination by other member galaxies in a beam. Moreover, the larger the redshift of an object, the lower the CO frequency, which results in a decrease in the system noise temperature due to the atmospheric O₂ emission at 118 GHz: the more distant is a galaxy, the lower is the noise temperature.

Dickey and Kazes (1992) have addressed the use of CO line widths instead of and/or supplemental to HI observations, and proposed the CO-line Tully-Fisher relation as an alternative to the HI Tully-Fisher relation. The CO-line width - luminosity relation has been established for the local distance calibrators and tens of nearby galaxies (Sofue 1992; Schöniger and Sofue 1993, 1996). In these works, we have shown that CO-line measurements can be used as an alternative to HI by deriving a good linear correlation between CO and HI linewidths for the galaxies in the sample. The only disadvantage of the use of CO line is the sensitivity. Actually, we need a few mK rms noise for line-width measurements at a velocity resolution of 10 km s⁻¹ for normal galaxies beyond $cz \sim 10,000$ km s⁻¹, which requires long integration times. Such observations are possible only by a long-term project with the use of the world's-largest mm-wave telescopes.

On the basis of these studies, we have conducted a long-term project to observe ¹²CO($J = 1 - 0$) line profiles for distant cluster galaxies at redshift to $z \sim 0.1$ using the Nobeyama 45-m telescope. The targets are so chosen that they include galaxies in clusters of galaxies at $cz \sim 10,000$ to 30,000 km s⁻¹. Before proceeding to measuring CO in normal galaxies at these distances, we chose far-infrared (FIR) luminous galaxies, so that the CO detection is easier at this first trial. We selected the targets using the NED (Nasa Extragalactic Database) searching facility. In this paper, we present results for the most FIR luminous galaxies with 60 μ m and 100 μ m fluxes greater than 0.2 and 0.5 Jy, respectively, as well as with redshifts within 10,000 and 30,000 km s⁻¹. This sample selection may include some danger in applying the Tully-Fisher relation because of possible peculiarities in optical absolute magnitudes for the FIR luminous galaxies: We may need to evaluate the accuracy of the relation for such FIR luminous galaxies, which remains problem for the future. The result presented here should be taken as a demonstration of the possible usage of the CO Tully-Fisher relation for measuring distances of galaxies of these large distances. However, we mention that such a FIR luminous starburst galaxy NGC 253 is included in the small number of standard calibrators of the Tully-Fisher relation. It is also known that IRAS galaxies obey a standard Tully-Fisher relation (van Driel et al 1995).

We obtained additional CO-line spectra for some galaxies using the IRAM 30-m mm-wave telescope. We also obtained high-quality optical imaging of detected galaxies in CO in order to measure the inclination and magnitude using the Canada-France-Hawaii 3.6-m telescope. In this paper, we report a first-step result of the measurement of CO line profiles in order to demonstrate that the line profiles can be obtained in routine

work within a reasonable integration time, and that distances can be obtained by applying the Tully-Fisher relation. We also report on optical imaging observations of several CO-detected galaxies, and try to obtain a possible value of the Hubble ratios.

2. CO-Line Observations

2.1. Observations

We have selected spiral galaxies which satisfy the following criteria using NASA Extragalactic Database (NED); 1) galaxies with red-shift between $10,000 \text{ km s}^{-1}$ and $30,000 \text{ km s}^{-1}$, 2) with position error less than $10''$, 3) with relatively strong far infrared emission at $60 \mu\text{m}$ and $100 \mu\text{m}$. The first criterion was set to select distant galaxies for which the HI Tully-Fisher relation cannot reach. The highest redshift galaxy was IRAS 08344+5105 which had $cz = 29029 \text{ km s}^{-1}$. The second criterion was set so that the position error is small enough compared to the beam size of CO-line observation. Accurate positions of the sources were measured on the Palomar Sky Survey Prints using the position measuring facilities at the Kiso Observatory, as well as using the STScI Digitized Sky Survey images. The third criterion was to select galaxies which are expected to be bright in the CO-line emission. One might suspect that galaxies with bright infrared emission could deviate from the Tully-Fisher relation for normal galaxies. However, van Driel et al.(1995) studied so-called IRAS Minisurvey galaxies and showed that there is no significant difference between Tully-Fisher relation for IRAS selected galaxies and that for normal galaxies. Among galaxies which satisfy these criteria, interacting galaxies and Seyfert-type galaxies are excluded. Galaxies with larger error of recession velocity than 100 km s^{-1} are also excluded, though galaxies for which the error of recession velocity is unknown are included. There are some galaxies for which types are not known through NED. For such galaxies, we have confirmed that they are disk galaxies and do not show interaction on the Palomar Sky Survey Prints.

Observations of the $^{12}\text{CO}(J = 1 - 0)$ line of distant cluster galaxies were made on 1994 January 14 to 23, 1994 December 9 to 12, and 1995 January 6 to 10, March 13 to 17, 1995 December 17, 18, 21, 22, and 1996 February 18 to 22, using the 45-m telescope of the Nobeyama Radio Observatory. The antenna had a HPBW of $15''$ at the CO line frequency, and the aperture and main-beam efficiencies were $\eta_a = 0.35$ and $\eta_{mb} = 0.50$, respectively. We used two SIS (superconductor-insulator-superconductor) receivers with orthogonal polarization which were combined with 2048-channel acousto-optical spectrometers. The total channel number corresponds to a frequency width of 250 MHz, and, therefore, to a velocity coverage in the rest frame at the galaxy of $c \ 250\text{MHz}/\nu_{\text{obs}} = c(1+z)(250 \text{ MHz}/115.2712 \text{ GHz})=650(1+z) \text{ km s}^{-1}$. The center frequency was so tuned that the center channel 1024 corresponds to $115.2712(1+z) \text{ GHz}$ for each galaxy.

After combining every 32 channels in order to increase the signal-to-noise ratio, we obtained spectra with a velocity resolution of 10.2 km s^{-1} . The system noise temperature (SSB) was 300 to 400 K at the observing frequencies. The calibration of the line intensity was made using an absorbing chopper in front of the receiver, yielding an antenna temperature (T_A^*), corrected for both the atmospheric and antenna ohmic losses. We used an on-off switching mode, and the on-source total integration time was 2 to 6 hours for each galaxy. After careful flagging and subtraction of linear baselines, the rms noise of the resultant spectra at velocity resolution of 10 km s^{-1} was typically 2 mK in T_A^* . The pointing of the antenna was tested by observing nearby SiO maser sources at 43 GHz every 1 to 1.5 hours, and was typically within $\pm 3''$ during good weather conditions which were attained for about one fourth of the allocated observing time.

2.2. CO-line Profiles and Results

Among the observed galaxies, sufficient quality data have been obtained for fifteen galaxies including eight galaxies with CO-line detection. Good CO line profiles were obtained for several galaxies with a sufficient signal-to-noise ratio for determining the velocity width. CO-detected galaxies are NGC 6007, IZw 23, IC2846, CGCG 1448.9+1654, CGCG 1113.7+2936, CPG 60451, CGCG 1417.2+4759, and IRAS 17527+6422. We obtained possible (marginal) detection for two galaxies, IRAS 14210+4829 and IRAS 23420+2227. These galaxies are listed in Table 1, and the spectra are shown in Fig. 1. The intensity scale used in this paper is the antenna temperature (T_A^*). The abscissa in Fig. 1 is relative recession velocity V_0 with respect to the optical redshift (cz) given in table 1. We have also performed CO line observations using the IRAM 30-m Telescope, and obtained detection for CGCG 1417.2+4759. The observation with the IRAM 30-m telescope was made on July 9, 1994, equipped with an SIS receiver of system temperature of 270 K and combined with a 256-channel filter-bank spectrometer of a velocity coverage 1400 km s^{-1} . The CO profile from the IRAM 30-m telescope is also shown in Fig. 1.

The profiles for the detected galaxies show a nearly double-peaked emission, which is characteristic of a rotating disk. From these data we have measured the line width, peak antenna temperature, and line intensity for individual galaxies. The results are listed in Table 2. The farthest galaxies for which the CO-line has been detected in the present program are IRAS 23420+2227 and IRAS 17527+6422 at $cz = 26022$ and 26151 km s^{-1} , respectively.

–Fig. 1; Tables 1, 2–

Our program included farther galaxies with redshift to $cz = 29000 \text{ km s}^{-1}$, and good data were obtained for several galaxies after integration of a few hours. Some examples are shown in Fig. 2, which are for IRAS 16305+4823 with $cz = 26327 \text{ km s}^{-1}$, showing a slight enhancement at relative velocity of $V_0 \sim -180$ to 0

km s⁻¹. IRAS 07243+1215 with $cz = 28204$ km s⁻¹, showing marginal detection at $V_0 \sim +70$ to $+200$ km s⁻¹, while not conclusive; IRAS 08344+5105 with $cz = 29029$ km s⁻¹, the farthest galaxy in our program showing a slight sign of enhancement at $V_0 \sim -100$ to $+180$ km s⁻¹.

–Fig. 2–

3. Optical Observations

3.1. Observations

In order to obtain photometry, morphology and inclination of the CO detected galaxies, we have performed high-quality optical imaging observations of three galaxies, CGCG 1417.2+4759, CGCG 1448.9+1654, and CPG 60451. The galaxies were so chosen that they are among galaxies showing the best quality CO line profiles observed during the 1994/1995 missions, and are within a right ascension range observable during the allocated time of optical observation. The observations were made using the Canada-France-Hawaii 3.6-m Telescope (CFHT) on June 30 - July 1, 1995. Additional images have been taken with the 105-cm Schmidt telescope of the Kiso Observatory in January 1994. Using these data, we have determined the apparent magnitudes and inclinations of the galaxies. We summarize the optical data in Table 3.

— Table 3 —

Imaging observations were made in Johnson V and Kron-Cousins R_C and I_C bands at the Canada-France-Hawaii 3.6-m Telescope with the SIS (Subarcsecond-Imaging-Spectrograph). The Loral3 CCD had an effective imaging area 2048×2048 pixels with a pixel size of $15\mu\text{m} \times 15\mu\text{m}$. The camera covered a sky area of $3' \times 3'$ with a resolution of $0.086''$ per pixel. The exposure times in the V , R_C , and I_C bands were 900, 900, and 300 sec, respectively. The observations were performed under photometric sky conditions on 30 June, 1995. Standard stars from Landolt (1992) were also observed for flux calibration. The FWHM seeing size was $0.6 - 0.8''$, which is small enough to determine the morphology of the observed galaxies. The dome screen was exposed to obtain flat frames.

Standard data processing (bias subtraction and flat fielding) was performed with the IRAF software package. For further image processing and surface photometry (e.g., sky subtraction and flux calibration) we used the SPIRAL package developed at Kiso Observatory and installed into the IRAF system. The large field of view gives a sufficiently large sky area around the galaxies to allow an accurate sky-subtraction. The flux calibration using the standard star frames resulted in a photometric accuracy of about $\pm 3\%$. The transformation to Johnson and Kron-Cousins' standard photometric systems and the atmospheric extinction corrections were determined by analyzing the standard star frames.

3.2 Surface Photometry

In Fig. 3 we present R_C band images of three galaxies. The contours are drawn at an interval of 1 mag./square arcsec, starting from 26.5 mag./square arcsec. The maps were fitted with ellipses to obtain the isophotal magnitudes and the axial diameters. The second column of Table 3 gives morphological types of the galaxies. We stress that the galaxies show quite normal morphology as spirals, and are isolated galaxies. The isophotal magnitudes of the galaxies above 25.5, 26.0, 24.5 mag arcsec⁻² levels in the V , R_C , and I_C bands are presented, respectively. The major A and minor B axial diameters at 25.5 mag arcsec⁻² in the V band are given in the last two columns. The typical error in the total magnitude estimate was about 0.03 magnitude.

– Fig. 3 –

We measured the intensity of the galaxies in elliptic annuli of one pixel width centered on the galactic nucleus, after subtracting the sky brightness and stellar images, thus obtaining a growth curve. Since the integrated magnitude in the tail of each growth curve approximately follows an exponential law, asymptotically reaching a constant value, we define here the total magnitudes as the extrapolation of this curve to infinity.

To evaluate the Tully-Fisher relation we need the velocity width to be corrected for the inclination i and for redshift effect to an edge-on value at the rest frame. We obtain an intrinsic velocity width in the rest frame referred to the galaxy corrected for the inclination by

$$W_i = W/\sin i, \quad (1)$$

where W is the rest-frame velocity width given by

$$W = W_{\text{rest}} = c\Delta\nu_{\text{obs}}/\nu_{\text{obs}} = c(1+z)\Delta\nu_{\text{obs}}/(115.2712 \text{ GHz}), \quad (2)$$

where $\Delta\nu_{\text{obs}}$ is the observed line width in frequency. We obtain the inclination i of a galaxy from the conventional formula given by Hubble (1926) for oblate spheroid,

$$\cos^2 i = (q^2 - q_0^2)/(1 - q_0^2) \quad (3)$$

with $q = b/a$ and $q_0 = c/a$, where a , b , and c are the lengths of three axes of the spheroid. Here, we adopt $q_0 = 0.20$ for the present analysis. The measured result is given in Table 4. The typical error in the estimate of inclination was $\pm 3^\circ$.

The Galactic absorption A_B is taken from Burstein & Heiles (1984). The internal absorption correction A_i are computed by the methods given in Pierce & Tully (1992). For the K-correction, we take the correction

of Fukugita, et al (1995). The corrected total magnitude is written as

$$m_{\text{T}}^{b,i} = m_{\text{T}} - A_B - A_i - K. \quad (4)$$

Obtained results of the optical measurements are summarized in Table 4. Typical errors in the magnitude estimate was about ± 0.1 magnitude except for the systematic errors which might be present in the applied corrections.

– Table 4 –

4. Discussion

4.1. Distances and Hubble Ratios

On the basis of our previous study that the CO-line width of a galaxy is equivalent to the HI-line width within the error of the Tully-Fisher relation for local calibrators, we try to derive the distances to the three galaxies for which the optical photometry and imaging have been obtained with the CFHT. We adopt the zero point of the TF relation given by Pierce & Tully (1992) for R and I and by Shimasaku & Okamura (1992) for V band.

$$M_V^{b,i} = -20.43 - 6.21(\log W_i - 2.5) \quad (5)$$

$$M_R^{b,i} = -20.40 - 8.23(\log W_i - 2.5) \quad (6)$$

$$M_I^{b,i} = -20.94 - 8.72(\log W_i - 2.5) \quad (7)$$

We adopt the same scheme for their extinction and inclination corrections. However, the distance moduli of the common local calibrators exhibit large discrepancies. The differences are -1.0 to $+0.1$ mag and 0.38 mag on average. (Shimasaku & Okamura’s moduli are larger). Four calibrators (M31, M33, NGC300, and NGC2403) have distances determined well by Cepheid variables. The Cepheid distances are in favor of Shimasaku & Okamura’s values. Therefore, we adopt the Shimasaku & Okamura’s calibration and add 19% to the distance moduli obtained by the Pierce & Tully calibration in R and I_C bands.

Using the Tully-Fisher relation for the CO-line widths measured in section 2 we derive distance moduli, distances and Hubble ratios V_C/r for the three galaxies. Here the recession velocity V_C is the value with respect to the rest frame referred to the cosmic background radiation (Smoot et al. 1991). Distance modulus, distance, and Hubble ratio obtained in individual optical bands (V , R_C and I_C) are shown in Table 5 for each object. Error in the absolute magnitude ΔM arises from the error in CO line-width measurement ΔW as well as from that due to the error in inclination ΔW_i , and is given by

$$\Delta M \simeq K \sqrt{(\Delta W/W)^2 + (\Delta W_i/W_i)^2}, \quad (8)$$

where K is the coefficient of the Tully-Fisher relation in equations (5) to (7), and

$$\Delta W_i = W \frac{\cot i}{\sin i} \Delta i = W_i \cot i \Delta i. \quad (9)$$

Here, Δi is the error in inclination measurement, which was typically $\pm 3^\circ$. The distance error is then given by

$$\frac{\Delta D}{D} \simeq \frac{\Delta(m - M)}{5 \log e} = \frac{1}{5 \log e} \sqrt{\Delta m^2 + \Delta M^2}. \quad (10)$$

Here Δm is the error in the photometric measurement of apparent magnitude. In the present case the error in the apparent magnitude (~ 0.1 mag.) is much smaller than that of the estimate of absolute magnitude arising from the line-width measurement and inclination correction. Hence, we have

$$\frac{\Delta D}{D} \simeq \frac{\Delta M}{5 \log e} \simeq \frac{K}{2.17} \Delta W_i / W_i \simeq \frac{K}{2.17} \sqrt{(\Delta W / W)^2 + (\cot i \Delta i)^2} \quad (11)$$

The measured values in the V , R_C , and I_C band are shown in Tables 4 and 5. The errors shown here do not include systematic error in the adopted photometric corrections or the internal scatter within the Tully-Fisher relation itself. As to the CO line width of CGCG 1417.2+4759 we adopted the CO data from IRAM 30-m telescope. The Hubble ratio obtained for CGCG 1417.2+4759 from the NRO 45-m telescope data is as large as $110 \text{ km s}^{-1}/\text{Mpc}$, while the data from IRAM 30-m gives a value around 80, and we here adopt the value from IRAM data. The larger value of Hubble ratio from the NRO data appears to be due to an underestimate of the velocity width because of the smaller velocity coverage in the spectrometer at NRO than the IRAM 30-m telescope. The obtained Hubble ratios in table 5 are scattered from ~ 50 to $\sim 80 \text{ km s}^{-1}/\text{Mpc}$, while the values are within a reasonable range of current determinations.

– Table 5 –

4.2. Remarks

We have underway a long-term program using the Nobeyama 45-m mm-wave telescope, and partly using the IRAM 30-m telescope, to study the CO-line width-luminosity (CO Tully-Fisher) relation. In this program, we have shown that the present method can be a useful technique to derive distances of galaxies at redshifts greater than $cz \sim 10,000 \text{ km s}^{-1}$, at which the HI-line Tully-Fisher relation becomes more difficult to be applied. Although the number of CO detection is not sufficient for conclusive determination of the Hubble ratio, we have obtained reasonable values for a few galaxies. We indeed tried to observe more distant galaxies, whose redshifts were as large as $z \sim 0.1$, in January and February of 1996. We obtained marginal detection for a few galaxies as listed in Table 1 and 2, and their CO line profiles are shown in Fig. 1 and 2. In order to derive the Hubble ratios of these galaxies, however, high-resolution optical imaging as shown in Fig. 3 is required, which will be performed in the near future.

Our present sample selection may include some danger in applying the Tully-Fisher relation because of possible peculiarities of FIR luminous galaxies, although the optical imaging (Fig. 3) has shown that the three galaxies used here are of normal morphology. However, we mention that FIR luminous galaxies from the IRAS minisurvey follow the Tully-Fisher relation (van Driel et al 1995). We also note that such a FIR luminous starburst galaxy as NGC 253 is included in the small number of standard calibrators of the Tully-Fisher relation, indicating that the FIR luminosity does not necessarily disturb this relation.

Acknowledgment: The observations were performed under a long-term project at the Nobeyama Radio Observatory. The authors would like to express their sincere thanks to Prof. N. Ukita and N. Nakai of the NRO for their kind and efficient help during the observations as well as for their invaluable suggestions.

References

- Aaronson M., Bothun G., Mould J., Shommer R. A., Cornell, M. E., 1986, ApJ 302, 536
- Aaronson, M., Mould, J., Huchra, J. 1980, ApJ, 237, 655
- Burstein, D., Heiles, C. 1984, ApJS, 54, 33
- Dickey J., Kazes I., 1992, ApJ 393, 530
- Fukugita M., Okamura S., Tarusawa K., et al., 1991, ApJ 376, 8
- Fukugita, H., Shimasaku, K., Ichikawa, T. 1995, PASP, 107, 945.
- Fouqué P., Bottinelli L., Gouguenheim L., 1990, ApJ 349, 1
- Hubble, E. P. 1926, ApJ, 64, 321
- Kraan-Korteweg R. C., Cameron L. M., Tammann G. A., 1988, ApJ 331, 610
- Landolt, A. U. 1992, AJ, 104, 340
- Pierce M. J., Tully R. B., 1988, ApJ 330, 579
- Pierce, M. J., Tully, R. B. 1992, ApJ, 387, 47
- Schöniger, F., Sofue, Y. 1993, AA, 283, 21.
- Schöniger, F., Sofue, Y. 1996, AA, in press.
- Shimasaku, K., Okamura, S. 1992, ApJ., 398, 441
- Smoot, G. F., et al. 1991, ApJ,371, L1
- Sofue Y., 1992, PASJ 44, L231
- Tully B., Fisher J. R., 1977, AA 64, 661
- van Driel, W., van den Broek, A, C., Baan, W. 1995 ApJ 444, 80.

Table 1. Galaxies detected in the CO line emission[†].

Name	RA(1950) (h m s)	Dec(1950) (° ' ") (posi. err. ")	cz (km s ⁻¹) (err. km s ⁻¹)	$S_{60\mu\text{m}}$ (Jy)	$S_{100\mu\text{m}}$ (Jy)	Type	detection
(1993/94 program)							
CGCG 1113.7+2936	11 13 46.9	+29 35 59(4)	13880(33)	0.63	2.03	?	D
CGCG 1448.9+1654	14 48 54.4	+16 54 02(4)	13700(51)	0.14	1.13	Sp	D
CGCG 1417.2+4759	14 17 14.8	+47 59 00(2)	21465	0.62	1.54	SBb	D
(1994/95 program)							
NGC 6007	15 51 01.6	+12 06 27(10)	10547(6)	0.69	2.03	SBbc	D
CPG 60451	17 30 00.6	+20 09 48.6(5)	14989(?)	0.48	1.36	Scd	D
IC 2846	11 25 24.804	+11 26 00.5(1)	12294(33)	4.21	6.72	Sp	D
(1995/96 program)							
IRAS 17527+6422	17 51 44.99	+64 22 14.1(3)	26151(24)	2.22	3.25	?	D
IZw 23	09 56 01.0	+52 29 48.0(1)	12224(57)	0.62	1.74	Sp	D
(Marginal detection)							
IRAS 14210+4829	14 21 06.2	+48 29 59.0(1)	22690(?)	0.38	0.88	Sp	?
IRAS 23420+2227	23 42 00.60	+22 27 49.8(3)	26022(?)	1.41	2.07	?	?

[†] Sources of the positions and redshifts is the NED (NASA Extragalactic Database). These velocities were used for the center frequencies of the CO-line detectors.

Table 2. CO-line data for detected galaxies.

Name	cz (km s ⁻¹)	cz_{CO} (km s ⁻¹)	$\Delta\nu_{\text{obs}}$ (GHz)	W † (km s ⁻¹)	T_{A}^* (mK)	$I_{\text{CO,rest}}$ (K km s ⁻¹)	Type
NGC 6007	10547	10542±5	0.129	347±10	23±3	5.1±0.2	SBbc
IZw 23	12224	12364±5	0.0391	106±10	37±5	2.4±0.2	Sp
IC 2846	12294	12302±10	0.0898	243±15	25±4	5.5±0.2	Sp
CGCG 1448.9+1654	13700	12693±5	0.0989	269±10	20±3	3.0±0.1	Sp
CGCG 1113.7+2936	13880	13910±5	0.117	318±5	16±2	2.5±0.1	SBb
CPG 60451	14989	14984±5	0.168	460±15	25±3	6.3±0.2	Scd
CGCG 1417.2+4759	21465	21527±10	0.111	308±15	12±3	1.6±0.2	SBb
" (IRAM 30m)			—	380±15	16±3	2.5±0.3	—
IRAS 17527+6422	26151	26165±20	0.145	410±20	15±5	4.1±0.4	?

† $W = W_{\text{rest}} = c\Delta\nu_{\text{obs}}/\nu_{\text{obs;center}} = c(1+z)\Delta\nu_{\text{obs}}/(115.2712 \text{ GHz})$.

Table 3: Optical Imaging of three galaxies.[†]

Galaxy	Type	$V_{25.5}$ (mag)	$R_{26.0}$ (mag)	$I_{24.5}$ (mag)	$A_{V_{25.5}}$ ($''$)	$B_{V_{25.5}}$ ($''$)
CGCG 1417.2+4759	SBcd	15.34	14.82	14.39	32.2	23.1
CGCG 1448.9+1654	Sab	15.00	14.46	13.98	33.5	29.6
CPG 60451	Sc	15.97	15.31	14.72	35.2	13.7

[†] The errors in the magnitudes are typically 0.03 mag.

Table 4. Data for the CO-Line TF Relation for measured galaxies.[†]

Galaxy	V_T mag	R_T mag	I_T mag	V_C km s ⁻¹	i deg	W_i km s ⁻¹
CGCG 1417.2+4759	15.31	14.80	14.33	21502	45.3	433±27
" " using (IRAM 30-m)						534±33
CGCG 1448.9+1654	14.96	14.43	13.93	13483	28.7	560±54
CPG 60451	15.92	15.26	14.66	14840	70.1	489±18

[†] The errors in the magnitudes are typically 0.1 mag, and inclination errors are ±3 degrees.

Table 5. Distance modulus, Distance, and Hubble ratio for the detected galaxies. †

Galaxy	$m - M$	D	$H = \langle V_C/r \rangle$
	mag	Mpc	km s ⁻¹ Mpc ⁻¹
CGCG 1417.2+4759 (IRAM 30-m Telescope)			
(V-band)	37.15±0.40	270±50	80±15
(R _C -band)	37.07±0.53	260±62	83±20
(I _C -band)	37.25±0.57	282±74	76±20
(Avrg.)	37.16±0.51	271±66	80±19
CGCG 1448.0+1654			
(V-band)	36.93±0.64	243±71	55±16
(R _C -band)	36.87±0.84	237±92	57±16
(I _C -band)	37.03±0.89	255±103	53±21
(Avrg.)	36.95±0.79	245±92	55±21
CPG 60451			
(V-band)	37.53±0.23	320±34	46±5
(R _C -band)	37.22±0.31	278±39	53±7
(I _C -band)	37.25±0.33	282±43	53±8
(Avrg.)	37.33±0.34	293±66	51±11

† The errors do not include systematic errors in photometric corrections and internal scatter within the Tully-Fisher relation itself.

Figure Captions

Fig. 1: CO line profiles for detected galaxies and marginal detection taken with the Nobeyama 45-m Telescope. The CO line profile of galaxy CGCG 1417.2+4759 obtained with the IRAM 30-m telescope is also presented. All data are presented in V_0 vs T_A^* plane, where V_0 is radial velocity referred to the galaxy's optical cz .

Fig. 2: The same as Fig. 1, but non-detection or marginal for the farthest galaxies up to $cz = 29000$ km s^{-1} in the present program obtained under good conditions. Some show a sign of detection, while not conclusive as yet.

Fig. 3: Optical images taken with the CFHT 3.6-m telescope in R_C band of galaxies CGCG 1417.2+4759, CGCG 1448.9+1654, and CPG 60451 in contour map and gray-scale representations. Contours are drawn at an interval of 1 mag./square arcsec, starting at 26.5 mag./square arcsec.




Fig. 1a




Fig. 1b




Fig. 1c




Fig. 1d




Fig. 1e




Fig. 1f




Fig. 1g




Fig. 1h




Fig. 2a




Fig. 2b




Fig. 2c




Fig. 1i




Fig. 1j






Fig. 1k


PLot file version 4 created 18-JUN-1996 18:53:31
CGCG 144 CG1448.CONV.1




Peak flux = 6.1529E+01
Levs = 5.0000E-02 * (1.000, 2.000, 4.000,
10.00, 20.00, 40.00, 60.00, 100.0, 200.0)




PLot file version 2 created 18-JUN-1996 18:49:13
CPG 6045 CPG.CONV.1



Peak flux = 1.8015E+01
Levs = 5.0000E-02 * (1.000, 2.000, 4.000,
10.00, 20.00, 40.00, 60.00, 100.0, 200.0)



PLot file version 4 created 18-JUN-1996 18:52:13
cgcg 141 CG1417.CONV.2



Peak flux = 2.2494E+01
Levs = 5.0000E-02 * (1.000, 2.000, 4.000,
10.00, 20.00, 40.00, 60.00, 100.0, 200.0)

