

# Ultraviolet Signatures of Tidal Interaction in the Giant Spiral Galaxy, M101

William H. Waller<sup>1</sup>, Ralph C. Bohlin<sup>2</sup>, Robert H. Cornett<sup>1</sup>, Michael N. Fanelli<sup>1</sup>, Wendy L. Freedman<sup>3</sup>, Jesse K. Hill<sup>1</sup>, Barry F. Madore<sup>4</sup>, Susan G. Neff<sup>5</sup>, Joel D. Offenberg<sup>1</sup>, Robert W. O’Connell<sup>6</sup>, Morton S. Roberts<sup>7</sup>, Andrew M. Smith<sup>5</sup>, and Theodore P. Stecher<sup>8</sup>

## ABSTRACT

We present new evidence for tidal interactions having occurred in the disk of M101 in the last  $10^8 - 10^9$  years. Recent imaging of the far-ultraviolet emission from M101 by the Shuttle-borne Ultraviolet Imaging Telescope (UIT) reveals with unprecedented clarity a disk-wide pattern of multiple linear arm segments (“crooked arms”). The deep FUV image also shows a faint outer spiral arm with a (“curly tail”) feature that appears to loop around the supergiant HII region NGC 5471 — linking this outlying starburst with the rest of the galaxy. These FUV-bright features most likely trace hot O & B-type stars along with scattered light from associated nebular dust. Counterparts of the outermost “crooked arms” are evident in maps at visible wavelengths and in the 21-cm line of HI. The inner-disk FUV arms are most closely associated with H $\alpha$  knots and the outer (downstream) sides of CO arms. Comparisons of the “crooked arm” and “curly tail” morphologies with dynamical simulations yield the greatest similitude, when the non-axisymmetric forcing comes from a combination of *external interactions* with one or more companion galaxies and *internal perturbations* from massive objects orbiting within the disk. We speculate that NGC 5471 represents one of these “massive disturbers” within the disk, whose formation followed from a tidal interaction between M101 and a smaller galaxy.

*Subject headings:* galaxies: evolution; — galaxies: interactions — galaxies: individual (M101) — galaxies: interactions — galaxies: spiral — ultraviolet: galaxies

---

<sup>1</sup>Hughes STX Corporation, NASA Goddard Space Flight Center, Laboratory for Astronomy and Solar Physics, Code 681, Greenbelt, MD 20771

<sup>2</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

<sup>3</sup>Carnegie Observatories, 813 Santa Barbara St., Pasadena, CA 91101-1292

<sup>4</sup>Infrared Processing and Analysis Center, Caltech, M/S 100-22, 770 So. Wilson Ave., Pasadena, CA 91125

<sup>5</sup>NASA Goddard Space Flight Center, Laboratory for Astronomy and Solar Physics, Code 681, Greenbelt, MD 20771

<sup>6</sup>University of Virginia, Department of Astronomy, P. O. Box 3818, Charlottesville, VA 22903

<sup>7</sup>National Radio Astronomy Observatory, 520 Edgemont Rd., Charlottesville, VA 22903-2475

<sup>8</sup>NASA Goddard Space Flight Center, Laboratory for Astronomy and Solar Physics, Code 680, Greenbelt, MD 20771

## 1. Introduction

Recent HST measurements of Cepheid variables in M101 (Kelson et al. 1996) have shown that this Sc(s)I galaxy is  $\sim 50\%$  more distant than was previously inferred from groundbased measurements of the galaxy's brightest supergiant stars (Humphreys & Aaronson 1987). At the new distance of  $7.4 \pm 0.60$  Mpc, M101 has a Holmberg radius of 30.1 kpc (Holmberg 1958), while its outermost giant HII regions lie more than 25 kpc from its nucleus. A galaxy of such enormous extent (70% larger than the giant ScI galaxy M100) presents an unusually large cross-section for interactions with its galaxian neighbors. The presence of the companion dwarf galaxies NGC 5477 and NGC 5474 with projected radii of 47 kpc and 99 kpc, respectively, lends further credence to the idea of tidal interactions having recently modified the disk of M101 (cf. van der Hulst & Sancisi 1988; Kenney et al. 1991; Combes 1991).

Evidence for interaction events during the last Gyr include (1) the lopsided distribution of HI gas (towards the northeast) with respect to the galaxy's dynamical nucleus (Bosma et al. 1981), (2) deviations from circular rotation in the outer galaxy (Bosma et al. 1981; Kamphuis 1993) (3) massive ( $10^7 - 10^8 M_{\odot}$ ) high-velocity clouds of HI gas to the east of the nucleus along with contiguous "holes" of comparable size (van der Hulst & Sancisi 1988), (4) a non-axisymmetric distribution of O/H abundances (Kennicutt & Liu 1996), and (5) outer spiral arms with nearly linear morphologies to the east and with bifurcating non-logarithmic curvatures to the west. The unusual arm morphology to the east has been described in a variety of terms, including "crooked arms" (Garcia-Burillo et al. 1991), "kinks" (Elmegreen et al. 1992), "spurs" (Byrd 1983; Byrd et al. 1984), "squared-off arm sections" and "intermediate-scale structure" (Roberts 1992; Roberts 1993).

Similar non-logarithmic spiral structure has been noted in the eastern-most arm of M51, whose obvious interaction with NGC 5195 has prompted the development of dynamical models, some of which successfully predict the generation of distorted spiral structure (Garcia-Burillo et al. 1991; Elmegreen et al. 1993; Byrd & Salo 1995). Other numerical simulations involving massive objects orbiting *within* the disks of spiral galaxies yield transient crooked arms, kinks, and spurs (Byrd et al. 1984; Roberts 1992).

In this paper, we present new evidence for dy-

namical interactions having occurred in the disk of M101 during the last  $10^8 - 10^9$  years. Recent imaging of the far-ultraviolet emission from M101 by the Shuttle-borne *Ultraviolet Imaging Telescope* reveals a remarkable pattern of multiple linear arm segments ("crooked arms") throughout the disk and a faint outer spiral arm with a ("curly tail") feature that appear to be of tidal origin. These FUV-prominent features most likely trace hot O & B-type stars along with scattered light from associated concentrations of dust. The morphological results presented herein build upon those obtained at lower resolution in a previous rocket flight, where the larger-scale UV emission is described (Stecher et al. 1982).

Section 2 briefly describes the *Astro-2/UIT* observations and reductions. The FUV-emitting features are presented in Section 3. The interpretation of these features as physically significant structures is evaluated in Section 4, and the dynamical implications are discussed in Section 5. Subsequent papers will present more detailed analyses of the FUV emission features in M101 (Waller et al. 1996) and of the dynamical processes which these features trace (Combes et al. 1996).

## 2. Observations and Reductions

The Ultraviolet Imaging Telescope imaged M101 in the far-ultraviolet ( $\lambda_0 = 1521 \text{ \AA}$ ,  $\Delta\lambda = 354 \text{ \AA}$ ) on 11, March 1995 as part of the 16-day Spacelab/*Astro-2* mission aboard the Space Shuttle Endeavour. The 1310-sec exposure was obtained with a dual-stage image intensifier with CSI photocathodes and was recorded on carbon-backed IIAO Kodak film. After processing of the film and scanning of the emulsion, the resulting digitized "density image" was fog-subtracted, flat-fielded, linearized to "exposure units," and calibrated to flux units using IUE observations of standard stars (cf. Stecher et al. 1992; Waller et al. 1995). Correction for image distortion produced by the magnetically focused image intensifiers was carried out according to the procedures described by Greason et al. (1995). The resulting corrections amounted to a few arcsec in the field center increasing to 10-20 arcsec near the edge of the 40 arcmin field of view. The astrometry of the distortion-corrected image is good to  $\sim 3$  arcsec, and the spatial resolution is of similar magnitude.

### 3. Morphology of FUV Emission

**Figure 1** (Plate xxxx) shows the distortion-corrected FUV image of M101. The emission morphology is dominated by spiral arms and star-forming knots. Most of the FUV-bright knots coincide with H $\alpha$ -bright giant HII regions (cf. Waller 1990; Scowen et al. 1992). However, the FUV knots are connected by brighter lanes of emission. Some of this FUV emission can be attributed to scattering of starlight by dust along the spiral arms (Stecher et al. 1982; Hill et al. 1995), although the balance of scattered vs. direct FUV emission remains uncertain (see next Section).

The structure of the FUV arms deviates markedly from that of logarithmic spirals. Instead, multiple “crooked arms” are evident throughout the disk of M101. The “crooked arm” morphology consists of linear arm segments that intersect one another — often at angles of  $\sim 120^\circ$ . These patterns are identified in **Figure 2a** (Plate xxx), where the FUV image of M101 has been spatially filtered to highlight the intermediate-scale structure.

The face-on locations, pitch angles, lengths, surface brightnesses, and contrasts of the 12 arm segments are listed in **Table 1**, where an inclination of  $18^\circ$  and major axis position angle of  $39^\circ$  have been assumed for the rectification (Bosma et al. 1981). A wide range of mean position angles and radii are represented in the list, further highlighting the ubiquitous nature of the linear arm segments. The large and varying pitch angles indicate arm morphologies that strongly differ from logarithmic spirals. The background-subtracted FUV surface brightnesses of the arm segments average to  $(5.6 \pm 3.2) \times 10^{-18} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2} \text{ \AA}^{-1}$ , which corresponds to  $2.9 \times 10^{37} \text{ erg s}^{-1} \text{ \AA}^{-1} \text{ kpc}^{-2}$  in the disk of the galaxy, or the equivalent of 300 Orion nebulae per square kpc (Bohlin et al. 1982). The contrast (ratio) of arm-interarm median intensities ranges from  $1.3 \pm 0.2$  for the innermost arm segment (#5) to  $\geq 5.0 \pm 0.2$  for the prominent arms in the east — where the uncertainties refer to the rms dispersion of contrasts derived from an arbitrary sequence of strip scans across the galaxy.

In addition to the crooked arms, a faint arm evident to the south is seen to extend all the way to the outermost giant HII region, NGC 5471, where it appears to loop around and ultimately connect with this kpc-size starbursting region (see **Figure 2b** [Plate xxx]). Previous UV imaging (Stecher et al. 1982; Blecha et al. 1990) detected the faint southern arm

(see features B&C in Stecher et al. 1982) but were of insufficient resolution ( $\sim 20''$ ) to delineate the “curly tail” structure which appears to connect with NGC 5471.

### 4. Physical Interpretation of FUV Features

Through FUV imaging of M101 at  $\sim 3$ -arcsec resolution, an anomalous pattern of non-logarithmic arms can now be traced with unprecedented clarity. The physical significance of this FUV morphology, however, is less obvious. We note that the physical interpretation of other spatial patterns in the sky has sometimes led to spurious conclusions. Well-known examples include the interpretation of dark patterns on Mars as evidence for life-sustaining canals (cf. Hoyt 1976) and the more recent controversy over stellar ring configurations seen on the Palomar Sky Survey images (cf. Vidal & Bern 1973). Because of the human eye-brain’s propensity to “connect the dots” and see patterns where none actually exist, the “crooked arm” and “curly tail” features noted in the previous section are subject to similar misinterpretation. Fortunately, these interpretive difficulties can be overcome by comparing the FUV features in M101 with stellar and interstellar tracers at other wavelengths.

#### 4.1. Arms and Knots at other Wavelengths

“Kinks” and “spurs” in the spiral arms of M101 have been previously noted from optical images (Bosma et al. 1981; Elmegreen et al. 1992). These descriptions invariably referred to the *outermost arms* to the east and northeast (arm segments #10,11,12 in **Figure 2a** and **Table 1**), where confusion with the underlying disk is minimal. At smaller galactocentric radii, emission from the underlying disk and absorption by lanes of dust complicate the observed morphology at blue–red wavelengths.

Comparison of the FUV image with the blue photograph of M101 printed in the Atlas of Galaxies (Sandage & Bedke 1988) shows that the FUV linear arm segments in the inner galaxy *do not* coincide with the spiral lanes of dust that are seen in silhouette against the brighter disk. Better correspondence is evident between the FUV-bright knots making up the arm segments and the young star clusters revealed in the blue print. Continuum-subtracted H $\alpha$  emission images show the star-forming knots in the inner galaxy with equivalent contrast to that observed in

the FUV image (Waller 1990; Scowen et al. 1992). However, the diffuse inter-knot emission along the arms is less evident at  $H\alpha$  than in the FUV, making the “crooked-arm” morphology less compelling. The connecting emission seen in the FUV but not at  $H\alpha$  is probably tracing *non-ionizing* populations of B-type stars along with associated reflection nebulosity.

The FUV-bright knots along the arms have similar sizes at  $H\alpha$  and in the FUV. By contrast, the sizes of the knots in the red-continuum are smaller by factors of  $\sim 2$  (Waller 1990). Since the red-continuum is probably tracing direct starlight from the underlying OB clusters, while the  $H\alpha$  emission arises from nebular ionized gas, the larger FUV and  $H\alpha$  sizes indicate that the FUV emission from the knots is at least partly nebular (e.g. reflection nebulosity associated with the HII regions). For HII regions and reflection nebulae in the Milky Way, scattering efficiencies by nebular dust are  $\sim 3$  times greater in the FUV than at visible wavelengths (Witt et al. 1992), thus explaining the larger FUV sizes. By comparing the peak and total emission from 3 bright knots in the R and FUV images (at matching resolution), we estimate that at least 30% of the FUV emission is scattered by nebular dust. UV images of the Orion nebula (Bohlin et al. 1982) and the giant HII regions in M81 (Hill et al. 1995) show evidence for similar amounts of scattering.

Comparison of the UIT’s FUV image with maps of 21cm HI emission at 45” resolution (Bosma et al. 1981), 80” resolution (van der Hulst & Sancisi 1988), and  $13'' \times 16''$  resolution (Kamphuis et al. 1991; Kamphuis 1993) shows that the outermost straight-arm segments to the east and northeast (#9,10,11,12) coincide with linear ridges of atomic gas. The inner-disk HI distribution is dominated by clumps and holes with little resemblance to the FUV arms.

Complementary information on the morphology of gas in the inner  $4 \times 4$  arcmin ( $8.6 \times 8.6$  kpc) is obtained from a recent mapping of 2.6mm CO emission at  $9'' \times 7''$  resolution (Kenney et al. 1996). **Figure 3** (Plate xxx) shows a contour map of the CO emission superposed on a greyscale representation of the FUV emission. Besides resolving the central gaseous bar into discrete features, the CO map reveals spiral structure beyond the bar, some of which coincides with prominent FUV emission. *These well-correlated FUV–CO emission features are present despite likely concentrations of associated dust* — further confirming the utility of FUV emission as a tracer of high-

mass star formation in disk galaxies of low inclination (Chen et al. 1992; Hill et al. 1995).

The most extensive FUV–CO correlations are evident in an arc of shared emission 1–1.5 arcmin to the W–SW of the nucleus, in linear arm segment #5 which adjoins the SW end of this arc, and in a large complex of knots 1-arcmin to the E–SE. A significant fraction of the correlated FUV emission is located on the outer (downstream) sides of the CO arms, thus suggesting *a spatio-temporal sequence of molecular-cloud aggregation, massive star formation, and cloud disruption* that can be modeled in terms of density-wave theory. Similar spatio-temporal relations between the CO and  $H\alpha$  arms are discussed by Kenney et al. (1996).

Along the W–SW arc (which shows the most coherent sequencing), the azimuthal displacements between the FUV and CO emission amount to  $0.35 \pm 0.1$  kpc — which when divided by the expected time lag between cluster formation and peak FUV brightness ( $\sim 3$  Myr [Hill et al. 1995]) — yields migration velocities through the spiral density wave of  $115 \pm 30$  km/s. Differencing this velocity with that of the orbiting gas at these radii (Bosma et al. 1981; Kenney et al. 1991) results in a pattern speed for the density-wave front of  $19 \pm 5$  km s $^{-1}$  kpc $^{-1}$  with a co-rotation radius of  $12 \pm 3$  kpc ( $5.5 \pm 1.5$  arcmin) — remarkably close to the co-rotation radius determined by Elmegreen et al. (1992) from their multi-mode analysis. The fact that coherent sequencing of the CO and FUV arms is not universal to the inner galaxy indicates that other dynamical processes are probably competing with the larger-scale density waves.

#### 4.2. Faint Southern Arm and NGC 5471

The faint FUV southern arm and the supergiant HII region (NGC 5471) at the arm’s NE tip are well correlated with other tracers of stellar and interstellar matter. The blue photograph of M101 in the Atlas of Galaxies (Sandage & Bedke 1988) shows the arm to contain many blue star clusters at the limit of detection. The supergiant HII region, NGC 5471, is beyond the field of the Atlas print, but appears in the deep blue Schmidt plate obtained by Sandage & Tammann (1974) along with diffuse emission to the NW and  $\sim 15$  discrete sources to the E and NE (2 of which appear to be background galaxies).

The compilation of  $H\alpha$  sources by Hodge et al. (1990) indicates  $\sim 25$  sources associated with the faint

southern arm. Near NGC 5471, their H $\alpha$  image reveals counterparts to the 4 FUV knots that delineate the tightly curled part of the “curly tail” structure (T1, T2, T3, & T4 in **Fig. 2a**). Beginning 1.8 arcmin to the east of NGC 5471 and looping northward and westward, the corresponding H $\alpha$  sources are catalogued as #1258, 1252, 1248, and 1231. The supergiant HII region, NGC 5471, is the most luminous HII region in M101. Spectroscopic studies indicate that the underlying stellar population is very young ( $\leq 3$  Myr) resulting in a high H $\alpha$  equivalent width, some evidence for an embedded supernova remnant, but little evidence for Wolf-Rayet or red supergiant stars (Skillman 1985; Rosa & Benvenuti 1994). That NGC 5471 is just now undergoing a massive starburst appears to be fortuitous.

In HI emission, the faint southern arm shows up as a significant distortion in the HI radial velocity field, deviating from circular rotation by 10–25 km/sec along the line of sight (Bosma et al. 1981). The  $13'' \times 16''$  HI mapping by Kamphuis et al. (1991) reveals an extended “hole” of low HI column density interior to the faint southern arm. The “hole” terminates near NGC 5471, where an expansive ( $\sim 1$  arcmin  $\approx 2$  kpc) region of HI emission is concentrated. In the following section, we contend that the velocity distortions and HI deficits associated with the faint southern arm are both caused by tidal interaction with one or more interloping galaxy companions.

## 5. Dynamical Implications

The anomalous morphological and kinematic features in the disk of M101 pose a serious challenge to dynamicists concerned with the evolution of giant spiral galaxies. Can a single model explain the “crooked arm” and “curly tail” morphologies seen in the FUV, along with the lopsided HI distribution, extended HI holes, and contiguous high-velocity HI features? Anticipating numerical simulations specific to the M101 system (Combes et al. 1996), we have examined existing simulations of M101 (Combes 1991) and of “generic” disk galaxies to see which ones best replicate the observed features in M101.

### 5.1. External Interactions

We find that tidal interactions with one or more companion galaxies are able to produce lopsided distributions in the gas and stars, large-scale spiral structure with some cusp-like corners in the gas, and high-

velocity features with contiguous “holes” in the gas (cf. Combes 1991; Garcia-Burillo et al. 1991; Howard et al. 1993; Elmegreen et al. 1993; Byrd & Salo 1995). Simulations with direct encounters relative to the direction of galactic rotation — along with large gas fractions and strong self-gravity within the disks — yield the clearest spiral structure (Howard et al. 1993). Characteristic interaction time scales are of order  $10^8$ – $10^9$  years. For M101, prime candidates for interaction within this timescale are the dwarf galaxies NGC 5477 and NGC 5474, whose masses and projected radii are  $0.5 \times 10^9 M_\odot$  (47 kpc) and  $19 \times 10^9 M_\odot$  (99 kpc), respectively (Allen et al. 1978; Combes 1991). Such interactions can induce the formation of massive condensations at the ends of the tidal tails (Elmegreen et al. 1993). According to this scenario, the kinematically distinct FUV arm to the south may represent a tidally induced tail, and the starburst NGC 5471 at the end of the faint arm may have resulted from a gravitational instability in the tidal tail.

Further insight to the relevant interaction timescales can be gleaned from the geometric and kinematic configuration of the tidally induced features. In a theoretical study of tidal perturbations, gravitational amplification, and galaxy spiral arms, Byrd (1995) derives the time since the perturber’s closest approach to be

$$t = R (\cot p) / v_{rot}$$

where  $t$  is the time since closest approach,  $p$  is the pitch angle, and  $v_{rot}$  is the rotation velocity (see also Binney & Tremaine 1987). Using this simple formulation, we obtain for the faint southern arm a mean interaction timescale of  $(4.4 \pm 3.3) \times 10^8$  yrs along its 19–28 kpc radius. If we adopt a  $27^\circ$  inclination (Kamphuis 1993) instead of the assumed  $18^\circ$  (Bosma 1981), the timescale would increase to  $(6.5 \pm 4.8) \times 10^8$  yrs. Numerical simulations of the interaction between NGC 5474 and M101 yield very similar timescales (Combes et al. 1996).

### 5.2. Internal Perturbations

Simulations involving *external interactions* with interloping galaxies do not usually generate “crooked arms” with *linear* arm segments (cf. Howard et al 1993). Multi-mode spiral density waves seem to do a better job of creating breaks and kinks in the spiral structure, but the arm segments tend to be curved

(Elmegreen et al. 1992). Such superpositions of two and three-arm density-wave modes are more successful at explaining the broken but symmetric spiral-arm structure evident in the “grand design” spiral NGC 628 (Chen et al. 1992; Elmegreen et al. 1992).

We find that *internal perturbations* by “massive disturbers” orbiting *within* the disk seem to do the best job at replicating the crooked FUV arms observed in M101. Simulations explicitly including massive disturbers in the disk (Byrd 1983; Byrd, Smith, & Miller 1984), as well as simulations which implicitly include them through large gas/star mass ratios and strong spiral forcing (leading to massive agglomerations) (Roberts 1992; Roberts 1993) yield spiral arm patterns with linear arm segments and with opening angles of approximately 120°.

### 5.3. Summation

Based on the above comparisons, we speculate that M101 has been subject to tidal perturbations from one or more companion galaxies in the past  $10^8$ – $10^9$  years. The interaction(s) have reshaped the large-scale spiral structure in the disk of M101, including the generation of a tidal tail to the south and a massive condensation (NGC 5471) at the end of the tail to the east that is currently undergoing a major starburst. Lower levels of ongoing star formation are evident along the length of the tidal tail. Further perturbations by NGC 5471 and other “massive disturbers” orbiting in the disk of M101 have induced a disk-wide pattern of crooked arms with linear arm segments.

The prominence of the crooked-arm pattern at FUV wavelengths indicates that it traces the most recent epoch of star formation and hence is a transient phenomenon. Consideration of the shear flow in the disk of M101 as approximated by

$$\Delta v = \Delta R R d\Omega/dR \approx -v_{rot} \Delta R/R,$$

where  $d\Omega/dR$  is the differential rotation in units of  $\text{km s}^{-1} \text{kpc}^{-2}$  (cf. Waller 1990; Binney & Tremaine 1987), indicates that significant evolution of the crooked-arm pattern will occur in a time less than 20 Myr — thus explaining the exceptional clarity of the FUV arms compared to the more diffuse spiral structures evident at longer wavelengths.

The proposed tidal scenario for explaining the unusual morphological and kinematic features in M101 can be tested via numerical simulations of the M101

system of galaxies (cf. Combes et al. 1996) and by evaluating the properties of other giant spiral galaxies that display similar features (e.g. NGC 1232, NGC 2805, & NGC 4303).

We thank Jeff Kenney for access to his CO map in advance of publication. Gene Byrd, Françoise Combes, Kevin Olson, and Bill Roberts generously provided insights on the dynamics of interacting disk galaxies and assistance in diagnosing M101’s unusual properties. UIT research is funded through the Space-lab Office at NASA Headquarters under Project number 440-51. We are deeply grateful to the crew of STS-67 and to the many people who helped make the Spacelab/*Astro-2* mission such a great success.

### REFERENCES

- Allen, R. J., van der Hulst, J. M., Goss, W. M., & Huchtmeier, W. 1978, *A&A*, 64, 359
- Binney, J. & Tremaine, S. 1987, *Galactic Dynamics* (New Jersey: Princeton University Press), pp. 122, 378
- Blecha, A., Golay, M., Huguenin, D., Reichen, D., & Bersier, D. 1990, *A&A*, 233, L9
- Bohlin, R. C., Hill, J. K., Stecher, T. P., & Witt, A. N. 1982, *ApJ*, 255, 87
- Bosma, A., Goss, W. M., & Allen, R. J. 1981, *A&A*, 93, 106
- Byrd, G. G. 1983, *ApJ*, 264, 464
- Byrd, G. G., Smith, B. F., & Miller, R. H. 1984, *ApJ*, 286, 62
- Byrd, G. G. & Salo, H. 1995, in *Proceedings of the World of Galaxies II*, *Ap. Lett. & Comm.*, 31, 193
- Byrd, G. G. 1995, in *Waves in Astrophysics*, eds J. H. Hunter, Jr. & R. E. Wilson, *Annals of the New York Academy of Science*, 773, p. 302
- Chen, P. C., et al. 1992, *ApJ*, 395, L41
- Combes, F. 1991, *A&A*, 243, 109
- Combes, F., Olson, K., & Waller, W. H. 1996, in preparation
- Elmegreen, B. G., Elmegreen, D. M., & Montenegro, L., 1992, *ApJS*, 79, 37
- Elmegreen, B. G., Kaufman, M., & Thomasson, M. 1993, *ApJ*, 412, 90
- Garcia-Burillo, S., Combes, F., & Gerin, M. 1991, in *Dynamics of Galaxies and Their Molecular Cloud*

TABLE 1  
LINEAR ARM SEGMENTS

No.	$\langle PA \rangle$ (deg)	$\langle R \rangle$ (kpc)	$\langle Pitch \rangle$ (deg)	$\Delta Pitch$ (deg)	$Length$ (kpc)	$\langle I(FUV) \rangle$	$Contrast$
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
1	312	22.3	2	-22	8.6	323	4.7
2	331	21.0	24	-18	6.9	342	5.4
3	353	7.8	7	-80	10.2	594	3.1
4	195	6.5	30	-51	6.6	534	1.7
5	168	7.5	16	106	12.5	365	1.3
6	156	9.3	21	27	4.6	943	3.5
7	50	5.8	22	-43	4.7	498	1.9
8	42	8.8	27	-47	8.9	918	2.7
9	31	15.6	35	-42	13.6	100	2.6
10	120	11.0	10	45	8.5	1030	5.0
11	40	21.8	23	-57	22.6	228	5.2
12	81	14.6	34	34	10.4	965	6.7

NOTE.—All quantities are measured in the plane of M101's disk, assuming an inclination of  $i = 18^\circ$  and major axis position angle of  $PA = 39^\circ$  (Bosma et al. 1981). (1) — Linear arm segment number (see **Figure 2a**). (2) — Mean position angle (in degrees) of arm segment, where the major axis has  $PA = 39^\circ$ , as observed on the sky. (3) — Mean galactocentric radius (in kpc) of the arm segment, where the distance to M101 is assumed to be 7.4 Mpc. For reference, the Holmberg radius is 30.1 kpc, and the radius of NGC 5471 is 25.1 kpc. (4) — Mean pitch angle (in degrees) of arm segment with respect to a circle corresponding to the mean radius. (5) — Range of pitch angles (in degrees) along arm segment as measured from west to east. (6) — Length of arm segment (in kpc). (7) — Average FUV intensity (surface brightness) in units of  $10^{-20} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ arcsec}^{-2} \text{ \AA}^{-1}$ , based on strip scans of  $45''$  (1.6 kpc) width on and off each arm. (8) — Arm–interarm contrast from the ratio of median intensities on and off each arm segment.

- Distributions, IAU Symposium #146, eds. F. Combes & F. Casoli (Dordrecht: Kluwer Academic Publishers), p. 351
- Greason, M. R., Offenberg, J. D., Cornett, R. H., Hill, R. S., & Stecher, T. P. 1994, *PASP*, 106, 1151
- Hill, J. K. et al. 1995, *ApJ*, 438, 181
- Hodge, P. W. Gurwell, M., & Goldader, J. D. 1990, *ApJS*, 73, 661
- Holmberg, E. 1958, *Medd. Lunds. Obs. Ser. II*, 136
- Howard, S., Keel, W. C., Byrd, G., & Burkey, J. 1993, *ApJ*, 417, 502
- Hoyt, W. G. 1976, *Lowell and Mars* (Tucson: University of Arizona Press)
- Humphreys, R. M. & Aaronson, M. 1987, *ApJ*, 318, L69
- Kamphuis, J. 1993, Ph.D. Dissertation, Kapteyn Astronomical Institute
- Kamphuis, J., Sancisi, R., & van der Hulst, T. 1991, *A&A*, 244, L29
- Kelson, D. D. et al. 1996, *ApJ*, 463, 26
- Kenney, J., Scoville, N., & Wilson, C. 1991, *ApJ*, 366, 432
- Kenney, J., et al. 1996, *ApJ*, submitted
- Kennicutt, R. C. & Garnett, D. R. 1996, *ApJ*, 456, 504
- Roberts, W. W. 1992, in *Astrophysical Disks*, eds. S. F. Dermott, J. H. Hunter, & R. E. Wilson (New York: New York Academy of Sciences), p. 93
- Roberts, W. W. 1993, *PASP*, 105, 670
- Rosa, M. R. & Benvenuti, P. 1994, *Å*, 291, 1
- Sandage, A. & Tammann, G. A. 1974, *ApJ*, 194, 223
- Sandage, A. & Bedke, J. 1988, *Atlas of Galaxies*, NASA SP-496, (Washington, DC: U.S. Govt. Printing Office), panel 12
- Scowen, P. A., Dufour, R. J., & Hester, J. J. 1992, *AJ*, 104, 92
- Skillman, E. D. 1985, *ApJ*, 290, 449
- Stecher, T. P., Bohlin, R. C., Hill, J. A., & Jura, M. A. 1982, *ApJ*, 255, L99
- Stecher, T. P. et al. 1992, *ApJ*, 395, L1
- van der Hulst, T. & Sancisi, R. 1988, *AJ*, 95, 1354
- Vidal, N. V. & Bern, K. Å 1973, *A&A*, 29, 277
- Waller, W. H. 1990, Ph.D. Dissertation, University of Massachusetts
- Waller, W. H., et al. 1995, *AJ*, 110, 1255
- Waller, W. H. et al. 1996, in preparation
- Witt, A. N. et al. 1992, *ApJ*, 395, L5

Fig. 1.— (Plate xxx) Far-ultraviolet ( $\lambda_0 1521, \Delta\lambda 354$ ) image of M101 as obtained by the Ultraviolet Imaging Telescope during the Spacelab/*Astro-2* mission of March 1995. The image has been corrected for residual field distortion. North is up and East is to the left. The astrometry is based on comparison of point sources with corresponding sources in the Guide Star Catalog. After rotation and centering of the target, the unmasked field of view has a diameter of 32.3 arcmin (69.5 kpc).

Fig. 2.— (Plate xxx) **(a)** — Far-ultraviolet image of M101 after spatial filtering to highlight the intermediate-scale structure. FUV arm segments described in the text are delineated and labeled (see **Table 1**). **(b.)** High-contrast presentation of the FUV emission in M101. The image has been Gaussian smoothed to a resolution of  $10''$  (FWHM). This was done to highlight the faint southern arm (“Arm”) and its looping connection with NGC 5471 to the east. The FUV knots (“T1, T2, T3, T4”) that help delineate the “curly tail” feature have  $H\alpha$  counterparts as described in the text.

Fig. 3.— (Plate xxx) Superposition of contoured CO emission (Kenney et al. 1996) on a grey-scale image of the FUV emission in the central  $4' \times 4'$  ( $8.6 \times 8.6$  kpc) of M101.

---

This 2-column preprint was prepared with the AAS L<sup>A</sup>T<sub>E</sub>X macros v4.0.

This figure "m101fig1.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9612165v1>

This figure "m101fig2a.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9612165v1>

This figure "m101fig2b.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9612165v1>

This figure "m101fig3.gif" is available in "gif" format from:

<http://arxiv.org/ps/astro-ph/9612165v1>