

Pulsar Braking Indices Revisited

Simon Johnston¹ & David Galloway^{2,1}

¹*Research Centre for Theoretical Astrophysics, University of Sydney, NSW 2006, Australia*

²*School of Mathematics and Statistics, University of Sydney, NSW 2006, Australia*

29 August 2021

ABSTRACT

Using the standard equation for the slowdown of a neutron star, we derive a formula for the braking index via integration rather than the conventional differentiation. The new formula negates the need to measure the second time derivative of the rotation frequency, $\ddot{\nu}$. We show that the method gives similar braking indices for PSR B1509–58 and the Crab pulsar to those already in the literature. We point out that our method is useful for obtaining the braking indices of moderate aged pulsars without the need for long, phase-connected timing solutions. We applied the method to 20 pulsars and discuss the implications of the results. We find that virtually all the derived braking indices are dominated by the effects of (unseen) glitches, the recovery from which corrupts the value of $\dot{\nu}$. However, any real, large, positive braking index has implications for magnetic field decay and offers support to recent models of pulsar evolution.

Key words:

pulsars: timing noise – pulsars: glitches

1 INTRODUCTION

Neutron stars are powered by rotational kinetic energy and lose energy by accelerating particle winds and by emitting electromagnetic radiation at their rotation frequency, ν . The rotation frequency thus decreases with time and this slowdown is usually described by the relation

$$\dot{\nu} = -K\nu^n \quad (1)$$

Here, K is a positive constant which depends on the moment of inertia and the magnetic dipole moment of the neutron star and n is the braking index. Conventionally, the braking index is derived by *differentiation* of equation 1, yielding

$$n = \frac{\nu \ddot{\nu}}{\dot{\nu}^2} \quad (2)$$

In a highly simplified model in which the spin-down torque arises from dipole radiation at the rotation frequency, one expects $n = 3$.

Only 4 pulsars have had their braking indices measured and all have $n < 3$. The Crab pulsar has a value 2.509 ± 0.001 (Lyne, Pritchard & Smith 1988; Lyne, Pritchard & Smith 1993), PSR 1509–58 has a braking index 2.837 ± 0.001 (Kaspi et al. 1994), PSR 0540–69 has $n = 2.04 \pm 0.02$ (Manchester & Peterson 1989; Nagase et al. 1990; Gouiffes, Finley & Ögelman 1992) and, finally, the Vela pulsar has the low value 1.4 ± 0.2 (Lyne et al. 1996). Melatos (1997) has shown that a modification of the simple model, which involves treating the neutron star and the inner magnetosphere as one entity,

allows him to derive values of braking index very close to those observed (except for the Vela pulsar).

Braking indices are very difficult to measure in all the other pulsars. For a typical ‘old’ pulsar with $\nu = 1$ Hz, $\dot{\nu} = 10^{-15}$ Hz/s, the expected $\ddot{\nu}$ from equation 2 is only $\sim 10^{-30}$ Hz/s², much too small to measure even over hundreds of years of timing - the second derivative only contributes one extra phase rotation every 600 yr! There are a number of pulsars with ages ~ 20 kyr for which one might expect to be able to measure n . However, two different effects tend to dominate the value of $\ddot{\nu}$ over that expected from spin-down alone. First, these young pulsars glitch often (Shearer & Lyne 1996). These glitches lead to discontinuities in both ν and $\dot{\nu}$ making it very difficult to phase connect (i.e. count the exact number of rotations of the pulsar) over the glitch. Furthermore, the recovery from a glitch can last many hundreds of days and the measurement of $\dot{\nu}$ reflects the recovery rather than the intrinsic spin-down. Finally, young pulsars have large random variations in arrival times known as ‘timing noise’. Cordes & Helfand (1980) recognised that the timing noise dominates over the intrinsic $\ddot{\nu}$ by a factor ~ 100 in these pulsars and that many early published values of braking indices, based on $\ddot{\nu}$, were spurious.

2 BRAKING INDEX BY INTEGRATION

Instead of differentiating equation 1, we integrate from a time t to $t + T$ to obtain

$$\left[\frac{\nu^{1-n}}{1-n} \right]_{\nu_1}^{\nu_2} = -KT = \frac{\dot{\nu}_1}{\nu_1^n} T \quad (3)$$

from which follows

$$\frac{\nu_2^{1-n} - \nu_1^{1-n}}{1-n} = \frac{\dot{\nu}_1 T}{\nu_1^n} \quad (4)$$

Hence

$$1-n = \frac{1}{\dot{\nu}_1 T} \left[\nu_2 \left(\frac{\nu_1}{\nu_2} \right)^n - \nu_1 \right] \quad (5)$$

But, $(\nu_1/\nu_2)^n = \dot{\nu}_1 / \dot{\nu}_2$ and so

$$n = 1 + \frac{\nu_1 \dot{\nu}_2 - \nu_2 \dot{\nu}_1}{\dot{\nu}_1 \dot{\nu}_2 T} \quad (6)$$

This allows the braking index to be computed without the need to measure $\ddot{\nu}$. The advantage of this method is that, in principle, ν and $\dot{\nu}$ can be measured over a short interval of time and then re-measured 20 yr later without the need for a phase connected solution over the whole 20 yr time span.

2.1 Error analysis

We can rewrite equation 6 as

$$n = 1 + \frac{1}{T} \left(\frac{\nu_1}{\dot{\nu}_1} - \frac{\nu_2}{\dot{\nu}_2} \right) \quad (7)$$

Thus, for $n \sim 3$, the value of the expression inside the brackets must be $\sim 2T$. Typically, the fractional error in $\dot{\nu}$ is much larger than that in ν . Let $\dot{e}_1$ and $\dot{e}_2$ be the error in $\dot{\nu}_1$ and $\dot{\nu}_2$. The absolute error, E , on the braking index is thus given by

$$E \simeq \frac{\nu}{T \dot{\nu}^2} \sqrt{\dot{e}_1^2 + \dot{e}_2^2} \quad (8)$$

Taking a young pulsar such as the Crab as an example, the parameters for which are $\nu \sim 30$ Hz, $\dot{\nu} \sim -3.8 \times 10^{-10}$ Hz/s and $\dot{e} \sim 10^{-15}$ Hz/s over one month (Lyne, Pritchard & Smith 1988). If measurements are made ~ 1 yr apart then the error in the braking index is ~ 0.01 .

For older pulsars, those near 100 kyr, one can obtain a reasonably low error on the braking index, provided the time span is large enough. For example, PSR B0540+23 was discovered more than 25 yr ago. Its parameters are $\nu \sim 4$ Hz, $\dot{\nu} \sim -2.5 \times 10^{-13}$ Hz/s and $\dot{e} \sim 10^{-18}$ Hz/s (Arzoumanian et al. 1994). Equation 8 yields an estimated error in the braking index of ~ 0.1 , accurate enough to distinguish between various slow-down models.

3 DISCUSSION

3.1 Pulsars with known braking index

For the Crab pulsar, Lyne et al. (1988) have tabulated ν , $\dot{\nu}$ and their associated errors at monthly intervals over a period of five years. In order to determine the braking index, they first compute $\ddot{\nu}$ by performing a straight line fit to the $\dot{\nu}$ versus time graph. They derive a braking index of 2.509 ± 0.001 based on the value of $\ddot{\nu}$ over this interval. We can use their tabulated values and derive a braking index from equation 6 above, without the need to compute $\ddot{\nu}$ explicitly.




Figure 1. Braking index for the Crab pulsar as a function of time over the period 1982-1987. The braking index is computed using equation 6 above and parameters for ν and $\dot{\nu}$ from Lyne et al. (1988). 2278 data points are shown.

We can do this for every pair of values in the table (i.e. 2278 pairs in total). The results are displayed in Fig. 1.

It can clearly be seen that for the shortest intervals, both measurement error and short-term timing noise dominate the value of the braking index. For time intervals greater than ~ 500 days, the braking index is stable near 2.5. This is consistent with the ~ 600 days periodicity seen in the timing noise by Lyne et al. (1988). The weighted average of all the measurements is 2.502 (however, the braking indices are not truly independent); the braking index measurement from the largest time interval is 2.516 ± 0.003 . Both these measurements are consistent with the Lyne et al. (1988) result. It is not possible to compute a braking index for the Crab over a longer period of time. Lyne et al. (1993) have shown that the glitches permanently alter the value of $\dot{\nu}$; however the inter-glitch value of the braking index remains roughly constant.

Kaspi et al. (1994) published a braking index of 2.837 ± 0.001 for PSR B1509-58 based on a phase-connected solution across 11 yr of timing data. This pulsar has not been observed to glitch in that interval, and the timing noise is surprisingly low for such a young pulsar. We applied our method by finding a local fit to ν and $\dot{\nu}$ over a period of 3 months in 1993 and 1997. Over this interval of 1700 days we find $n = 2.80 \pm 0.03$ consistent with the Kaspi et al. (1994) result but without the need for a phase connected solution.

3.2 Young, Vela-like pulsars

There are 21 pulsars with $|\dot{\nu}| > 10^{-12}$ Hz/s. Of these, 4 have their braking indices measured (see above), and 11 of the remaining 17 have had one or more glitches and are expected to glitch every ~ 10 yr or so (see e.g. Shemar & Lyne 1996). For the Vela pulsar, for example, the intrinsic $\ddot{\nu}$ is completely dominated by the recovery of $\dot{\nu}$ following the regular glitches. Lyne et al. (1996) solved this problem by taking $\dot{\nu}$ at a fixed interval after each glitch and deriving $\ddot{\nu}$ and hence the braking index from the changing $\dot{\nu}$. Presumably, if such a method is valid, one could apply equation 6

over the largest time interval available (25 yr) to obtain an error on the braking index of ~ 0.1 .

The six pulsars which are not known to have glitched are PSRs B1951+32, B1853+01, B1930+22, B1643-43, B0906-49 and J0631+1036. Taking PSR B1951+32 as an example, Foster et al. (1994) have shown that this pulsar suffers from excessive timing noise. By fitting for higher orders in ν they obtain a value of $\ddot{\nu}$ of 2.7×10^{-22} Hz/s². This is about a factor ~ 50 higher than that expected from the simple spin-down, making it virtually impossible to compute a braking index. PSR B0906-49 was discovered 12 yr ago, and has not glitched since discovery. Using recent timing data, the value of $\ddot{\nu}$ from timing noise is $\sim 1.8 \times 10^{-23}$ Hz/s², more than 30 times the expected spin-down value. Thus, timing noise is dominating the value of $\ddot{\nu}$ in the literature (D’Amico et al. 1988) and a braking index cannot be computed. In particular the formal fitting errors given by TEMPO (the least-squares pulsar timing package) are seriously underestimating the systematic effects of the timing noise.

For these young pulsars, the new method suffers from similar problems to the standard calculation of the braking index. In Vela-like pulsars glitches dominate the overall spin-down behaviour, masking the true spin-down $\ddot{\nu}$. In slightly older pulsars, glitches are less frequent; however, the true $\ddot{\nu}$ is hidden in large timing noise residuals.

3.3 Pulsars with moderate $\ddot{\nu}$

As described above, it should be possible in principle to obtain an error on the braking index as low as ~ 0.2 for pulsars of age $\sim 10^5$ yr discovered more than 20 yr ago (especially those with low timing noise). We select pulsars in the following way. We construct an ‘expected’ value of $\ddot{\nu}$ for every pulsar, assuming a braking index of 3 and select all those with $\ddot{\nu} > 10^{-27}$ Hz/s². We reject all those pulsars which are known to have glitched in the past and the 21 pulsars described in 3.2 above. We then searched the literature for at least two timing solutions for the remaining pulsars and with a $\ddot{\nu}$ of sufficient accuracy to allow the error on the computed braking index to be less than 20. We discovered, however, that in some publications, the epoch of ν and $\dot{\nu}$ corresponded not to the middle of the data span, but rather to the start. If there is a significant value of $\ddot{\nu}$ in the data span which is unaccounted for (either related to timing noise or to the intrinsic spin-down), this leads to an error in the quoted $\dot{\nu}$ of $\sim \frac{1}{2} \ddot{\nu} T$ which, for large T , can be significantly larger than the formal fitting error. This led to the rejection of a further five pulsars.

Table 1 gives the braking index for 20 pulsars which survive the above selection criteria. Column 1 of the table gives the pulsar name. Columns 2 and 3 give the pulsar’s rotation frequency and its first derivative, and column 4 the time interval between the epochs of measurement. Column 5 gives the braking index and associated error according to equations 6 and 8. The final column gives the references for the timing solutions. Note that for PSR B0656+14 the value of $\dot{\nu}$ given in Ashworth & Lyne (1981) is clearly in error. Hence, the time interval between timing solutions is rather shorter than it might have been.

The values of the braking indices listed in the table raise the question as to how realistic the quoted errors are. Typically (but not always) in timing solutions the errors are

twice the formal standard error given by the TEMPO least-squares timing package. However, the errors are computed assuming that the each TOA has uncorrelated residuals (i.e. white noise), which is clearly not the case in pulsars with significant (red) timing noise. Thus the error on $\dot{\nu}$ is likely to be larger than that given from the fit alone. Also, as described above, not fixing the epoch in the middle of the data span leads to errors in the value of $\dot{\nu}$.

	PSR B	ν (Hz)	$\dot{\nu} \times 10^{-15}$ (Hz/s)	Interval (days)	n	Refs
1	0114+58	9.86	-562.3	2271.1	-9.6 $\pm$ 1.5	1,2
2	0136+57	3.67	-144.2	4492.0	-81 $\pm$ 4.7	5,2
3	0154+61	0.43	-34.2	4336.5	28 $\pm$ 14	5,2
4	0540+23	4.06	-255.0	5543.5	11.1 $\pm$ 8.6	8,2
				5990.5	11.81 $\pm$ 0.12	3,2
5	0611+22	2.99	-530.8	5541.5	20.1 $\pm$ 1.1	3,2
6	0656+14	2.60	-371.5	2163.3	14.7 $\pm$ 1.4	1,2
7	0740-28	6.00	-604.9	4245.2	17.7 $\pm$ 1.4	9,7
				5827.2	25.6 $\pm$ 0.8	9,2
8	0919+06	2.32	-74.0	4521.7	28.9 $\pm$ 4.1	5,2
9	1221-63	4.62	-105.7	6661.3	18.7 $\pm$ 12.3	6,10
10	1356-60	7.84	-389.9	7050.3	6.3 $\pm$ 6.8	6,10
11	1719-37	4.23	-194.6	4824.0	-183 $\pm$ 10	6,2
12	1742-30	2.72	-79.0	1581.0	-132 $\pm$ 5	7,2
13	1829-08	1.54	-151.4	1541.0	2.5 $\pm$ 0.9	4,2
14	1907+10	3.52	-32.8	5842.5	24 $\pm$ 17	3,2
15	1914+09	3.70	-34.5	5556.6	-15 $\pm$ 16	3,2
16	1915+13	5.14	-190.0	6080.5	36.08 $\pm$ 0.48	3,2
17	2000+32	1.44	-216.7	1381.0	-226 $\pm$ 4.5	4,2
18	2002+31	0.47	-16.7	6076.5	23.3 $\pm$ 1.0	3,2
19	2148+52	3.01	-91.2	2307.2	49.6 $\pm$ 3.5	1,2
20	2334+61	2.02	-776.2	2347.1	8.60 $\pm$ 0.13	1,2

[Table 1 references: 1-Dewey et al. (1988), 2-Arzoumanian et al. (1994), 3-Gullahorn & Rankin (1978), 4-Clifton et al. (1992), 5-Backus, Taylor & Damashek (1982), 6-Newton, Manchester & Cooke (1981), 7-Siegmán, Manchester & Durdin (1993), 8-Helfand et al. (1980), 9-Manchester et al. (1983), 10-Parkes timing programme.]

Figure 2 displays the braking index for these 20 pulsars with error bars a factor of 5 larger than in the table. We believe these error bars are conservative and more accurately reflect the contributions from the underlying timing noise to the intrinsic value of $\dot{\nu}$. From the figure, 14 of the 20 pulsars have significant values of braking index. The eight pulsars with the smallest error bars all have positive braking indices (PSRs B0540+23, B0611+22, B0656+14, B0740-28, B1915+13, B2002+31, B2148+52 and B2334+61). The four pulsars with large negative braking indices are PSRs B0136+57, B1719-37, B1742-30, B2000+32 and B2255+58.

3.4 Implications

The presence of glitches in pulsars can lead to spurious values of the braking index and we surmise that (unseen) glitches are the main cause for $n \neq 3$. If we assume that glitches cause $\Delta \dot{\nu} / \dot{\nu}$ to be positive in all cases and that the value of $\dot{\nu}$ recovers to nearly its original value, then large




Figure 2. Braking index for 20 pulsars from Table 1. The error bars are a factor of 5 larger than in the table.

negative braking indices can be caused by glitches between timing solutions and positive braking indices arise when the glitch occurs before the epoch of the first timing solution. Alpar & Baykal (1994), in a statistical survey of the frequency of pulsar glitches, showed that the number of glitches, g , in a sample of pulsars is given by

$$g = \left(\frac{\delta\nu}{\nu} \right)^{-1} \sum T \frac{\dot{\nu}}{\nu} \quad (9)$$

where T is the time interval over which the pulsar is monitored and $\delta\nu/\nu$ is 1.74×10^{-4} (Alpar & Baykal 1994).

Using the values of ν , $\dot{\nu}$ and T from the table, we derive $g = 2.8$, i.e. one expects there to have been glitches in ~ 3 pulsars in between their timing solutions. We have four pulsars with large negative braking indices which is consistent with this estimate. The values of $\Delta \dot{\nu} / \dot{\nu}$ for these four pulsars are 1.3, 3.6, 0.5, 4 and 0.6×10^{-3} respectively, values typical of glitches (Shemar & Lyne 1996). It is thus possible that these pulsars have all glitched between the epochs of the observations. However, none of the 4 appear to have suffered any change in ν , i.e. any step change at the time of the glitch would have been recovered in the intervening time period. Alternative explanations are hard to find - for PSR B2000+32, for example, one of the two timing solutions would have to be in error by more than 100σ to obtain a ‘real’ braking index of 3.

For the positive values of braking index, it is harder to compute the number of glitches expected *before* the timing solutions. Both the recovery time and the form of the recovery (linear or exponential) for the change in $\dot{\nu}$ are unclear. Shemar & Lyne (1996) estimate that the recovery time is greater than 3 yr for the exponential case or tens of years if the recovery is linear. Indeed, in some pulsars, a permanent alteration in the value of $\dot{\nu}$ is observed. If we let $T = 25$ yr, then we would expect 6.4 glitches prior to the first timing solution for these 20 pulsars. We have 8 positive braking indices, broadly in line with this estimate (and we also note that the 2 pulsars with the highest glitch probability are PSRs B2334+61 and B0611+22, both of which have positive braking indices).

A number of other factors may contribute to $n \neq 3$. It is possible that the different methods of obtaining the

pulsar position could affect the results (however, for PSR B0540+23, for example, the positions are identical within the errors). Another possibility is that the different fitting software, clock corrections, ephemeris changes etc can affect the result but it is difficult to believe this could have an effect at the 500σ level. These possibilities could be overcome by recovering the original TOAs from the 1970s and reducing them with the same software as the 1990s data.

The glitch interpretation put forward above, is statistical in nature. It may be that in one or more cases the braking index is indeed significantly larger than 3. What are the implications of this? Blandford & Romani (1988) showed that if the value of K in equation 1 is time variable then one can write

$$\frac{\dot{K}}{K} \frac{\nu}{\dot{\nu}} = n_{\text{obs}} - n. \quad (10)$$

$K / \dot{K}$ then gives the timescale for variation, if we assume that $n = 3$. For the cases in which $n_{\text{obs}} > 3$, $\dot{K} < 0$, and the timescale is surprisingly constrained in the range ~ 14 to ~ 56 kyr. $\dot{K} < 0$ can arise from magnetic field decay, alignment of the spin and magnetic axis, or an increase in the moment of inertia. Evidence for and against magnetic field decay and axis alignment have raged in the literature for 25 yr without any clear consensus emerging. Recently, Chen, Ruderman & Zhu (1998) have developed a model in which the magnetic field increases by a factor of ~ 10 over the first 10 kyr of a pulsar’s life (which may explain the *low* braking index of the Vela pulsar) and then decreases thereafter. In their model, they expect a braking index of ≥ 5 for middle aged pulsars before it settles in older pulsars back to 3. If any of the above braking indices are thus real, this implies a magnetic field decaying by a factor of ~ 100 in ~ 200 kyr, lending support to the Chen et al. (1998) model.

In any case, one is forced to conclude that the initial starting premise that the spin-down of a pulsar can be described as a simple power law in ν is highly questionable. In virtually all young pulsars, and in most older ones, complex glitch behaviour appears to be dominating the spin-down behaviour. This glitch behaviour, coupled with changes in the magnetic field strength over time, indicates that the entire concept of a ‘braking index’ must be treated with some caution.

Verification of the glitch hypothesis or the presence of a true braking index could be obtained by deriving additional timing solutions for these pulsars now. This would give a 7 yr span back to the Arzoumanian et al. (1994) data and, in some cases, more than 20 yr back to earlier data. One might then expect to observe the exponential recovery of $\dot{\nu}$ in the case of a glitch.

4 CONCLUSIONS

We have derived a new method for computing pulsar braking indices based on integration of the standard slow-down equation. We show that the method replicates the known values for both the Crab pulsar and PSR B1509–58. The new method conveys no advantage over the previous method for pulsars with regular glitches and/or large timing noise. The advantage of equation 6 is that for pulsars with ages $\sim 10^5$ yr where both the timing noise and the glitch frequency are

relatively low, it is not necessary to obtain a phase connected solution across 25 yr or more in order to obtain a braking index. Rather, local fits over 1-2 yr at large intervals may be sufficient to yield the braking index to an error of ~ 0.1 .

We computed braking indices for 20 pulsars based on timing solutions available in the literature. Four pulsars have large negative braking indices which we believe have been caused by glitches occurring in between the epochs of the timing solutions. A further eight pulsars have moderate positive braking indices which appear to be robust to the effects of timing noise but are possibly due to glitches which occurred before the start of the timing observations. If any of the large, positive braking indices are real, it may provide evidence for magnetic field decay in moderate aged pulsars. However, the entire concept of a smooth pulsar spin-down and of a constant braking index must be treated with some caution.

ACKNOWLEDGMENTS

We thank the ATNF pulsar group for providing TOAs for PSRs B1221–63, B1356–60 and B1509–58. We also thank L. Ball, R. Manchester and M. Wardle for comments on the manuscript. We acknowledge use of the TEMPO software package for pulsar timing.

REFERENCES

- Alpar M. A., Baykal A., 1994, MNRAS, 269, 849
 Arzoumanian Z., Nice D. J., Taylor J. H., Thorsett S. E., 1994, ApJ, 422, 671
 Ashworth M., Lyne A. G., 1981, MNRAS, 195, 517
 Backus P. R., Taylor J. H., Damashek M., 1982, ApJ, 255, L63
 Blandford R. D., Romani R. W., 1988, MNRAS, 234, 57P
 Chen K., Ruderman M., Zhu T., 1998, ApJ, 493, 397
 Clifton T. R., Lyne A. G., Jones A. W., McKenna J., Ashworth M., 1992, MNRAS, 254, 177
 Cordes J. M., Helfand D. J., 1980, ApJ, 239, 640
 D’Amico N., Manchester R. N., Durdin J. M., Stokes G. H., Stinebring D. R., Taylor J. H., Brissenden R. J. V., 1988, MNRAS, 234, 437
 Dewey R. J., Taylor J. H., Maguire C. M., Stokes G. H., 1988, ApJ, 332, 762
 Foster R. S., Lyne A. G., Shemar S. L., Backer D. C., 1994, AJ, 108, 175
 Gouffes C., Finley J. P., Ögelman H., 1992, ApJ, 394, 581
 Gullahorn G. E., Rankin J. M., 1978, AJ, 83, 1219
 Helfand D. J., Taylor J. H., Backus P. R., Cordes J. M., 1980, ApJ, 237, 206
 Kaspi V. M., Manchester R. N., Siegman B., Johnston S., Lyne A. G., 1994, ApJ, 422, L83
 Lyne A. G., Pritchard R. S., Smith F. G., 1988, MNRAS, 233, 667
 Lyne A. G., Pritchard R. S., Smith F. G., 1993, MNRAS, 265, 1003
 Lyne A. G., Pritchard R. S., Smith F. G., Camilo F., 1996, Nat, 381, 497
 Manchester R. N., Peterson B. A., 1989, ApJ, 342, L23
 Manchester R. N., Newton L. M., Hamilton P. A., Goss W. M., 1983, MNRAS, 202, 269
 Melatos A., 1997, MNRAS, 288, 1049
 Nagase F., Deeter J., Lewis W., Dotani T., Makino F., Mitsuda K., 1990, ApJ, 351, L13

- Newton L. M., Manchester R. N., Cooke D. J., 1981, MNRAS, 194, 841
 Shemar S. L., Lyne A. G., 1996, MNRAS, 282, 677
 Siegman B. C., Manchester R. N., Durdin J. M., 1993, MNRAS, 262, 449