Deep10and18 µmImagingoftheHR4796ACircumstellarDisk: TransientDustParticles&TentativeEvidenceforaBrigh tnessAsymmetry¹

C.M.Telesco²,R.S.Fisher²,R.K.Piña²,R.F.Knacke³, S.F.Dermott²,M.C.Wyatt²,K.Grogan²,E.K.Holmes²,A.M.Ghez⁴,L.Prato⁴, L.W.Hartmann⁵,andR.Jayawardhana⁵

Received:_____;Accepted:_____

¹BasedonobservationsattheW.M.KeckObservatory,w hichisoperatedasa scientificpartnershipbetweenCaltech,theUniversityo fCalifornia,andNASA, andmadepossiblebythegenerousfinancialsupportoftheW.M .Keck Foundation.

²DepartmentofAstronomy,UniversityofFlorida,Gainesvil le,FL32611; telesco,fisher,rpina,dermott,grogan,hol mes,wyatt@astro.ufl.edu

³SchoolofScience,PennsylvaniaStateUniversityatErie ,StationRoad,Erie,PA 16563;rfk2@psu.edu

⁴DepartmentofAstronomy,UniversityofCalifornia,Los Angeles,CA90095; ghez@athena.astro.ucla.edu,lprato@minerva.astro .ucla.edu

⁵Harvard-SmithsonianCenterforAstrophysics,60GardenStr eet,Cambridge,MA 02138;hartmann@fuor.harvard.edu,rayjay@shakti.harvard.edu

ABSTRACT

Wepresentnew10.8and18.2 µmimagesofHR4796A,ayoungA0Vstarthatwas recentlydiscoveredtohaveaspectacular, nearlyedgeon, circumstellardisk prominentat ~20 µm(Jayawardhanaetal.1998;Koerneretal.1998).Thesene wimages, obtained withOSCIRatKeckII.showthatthedisk'ssizeat 10 umiscomparabletoitssizeat 18 μm. Therefore, the 18 μm-emitting dust may also emitsome, or all, of the 10 μm radiation.Usingthesemulti-wavelengthimages,wedeter minea"characteristic"diameter of 2-3 µmforthemid-infrared -emittingdustparticlesiftheyaresphericalandcompose d ofastronomicalsilicates.Particlesthissmallare expectedtobeblownoutofthesystem retheseparticlesareunlikelyto byradiationpressure in a few hundred years, and there fo bepromordial.Rather,asinferredinacompanionpaper(W yattetal.2000),theyare probablyproductsofcollisionsthatdominateboththecrea tionanddestructionofdustin theHR4796Adisk.Dynamicalmodelingofthedisk(Wyattet al.2000)indicatesthat thedisksurfacedensityisrelativelysharplypeakednear 70AU, which agrees with the meanannularradiusdeducedbySchneideretal.(1999)fromthei rNICMOSimages. Interiorto70AU, the model density drops steeply by af actoroftwobetween70and60 AU, falling to zeroby 45 AU, which corresponds to the edge of thepreviously discovered central hole; in the context of the dynami calmodels,this"soft"edgeforthe centralholeresultsfromthefactthatthedustparti cleorbitsarenon-circular.Theoptical depthofmid-infrared -emittingdustintheholeis -3% of the optical depth in the disk, and the hole is therefore relatively very empty. We pr esentevidence(-1.8σ significance) forabrightnessasymmetrythatmayresultfromthe presenceoftheholeandthe gravitationalperturbationofthediskparticleorbitsbyt helow-massstellarcompanionor aplanet. This "pericenterglow," which must still be confirmed, results from avery small (afewAU)shiftofthedisk'scenterofsymmetryrel ativetothecentralstarHR4796A; onesideoftheinnerboundaryoftheannulusisshiftedto wardsHR4796A,thereby becomingwarmerandmoreinfrared-emitting. The possible det ectionofpericenterglow implies that the detection of even complex dynamical effectsofplanetsondisksiswithin reach.

1. INTRODUCTION

Planetsformindustycircumstellardisks, and so thee xplorationofcircumstellar diskscontributesfundamentallytoabroaderunderstandi ngoftheoriginandevolutionof ourownandotherplanetarysystems.Vega-typedisks,o ftencalleddebrisdisks, are those thatsurroundmainsequencestars(seethereviewbyBackm an&Paresce1993).Unlike thediskssurroundingpre-main-sequencestars, the dust masse sofdebrisdisksare ,whichisroughlytheminimummassrequiredfor generallymuchsmallerthan0.01M theformationofaplanetarysystemlikeourown(see ,e.g.,Beckwith&Sargent1993; Strometal. 1993). The lower mass of debrisdisks confor mswiththeexpectationthat disksdispersewithtimeasthematerialaccreteson tothestars,formsplanets,and/oris blownoutofthesystems(e.g., Strometal. 1993). Thee lucidationoftheseevolutionary processesisakeychallengeofobservationalastronomy.

High-angular-resolution, multi-wavelengthimaging of cir cumstellardiskscan provideuniqueinsightintotheseprocesses, since a disk's de tailedstructureshouldbear tracesofitshistoryandofanydevelopingormatureplane ts.Untilrecently,thedisk surroundingtheA5star BPictoris(Smith&Terrile1984;Telescoetal.1988;Lagage & Pantin1994;Kalas&Jewitt1995)wastheonlydebrisdisktha twasspatiallywellresolvedatopticalorinfraredwavelengths.However,ma nynewresultshaveemerged overthepastfewyearsincluding:possibleopticalimaging ofadiskaroundthebinary starBD+31°643(Kalas&Jewitt1997);submillimeterimagingofimpressive structurein severaldebrisdisks, including the class archetype Vega(Hollandetal.1998;Greaveset al. 1998); and ~20 µmimaging of the nearly edge-ondisk around the AOV starHR 4796A(Jayawardhanaetal.1998[J98];Koerneretal.1998[K98]), resultsthatwere recentlycomplementedwithNICMOSimagesrevealingwhat appearstobeasharply definedring(Schneideretal. 1999).

Herewepresentnew10.8and18.2 umimages, obtained at the Keck II telescope, of theHR4796Adebrisdisk.Jura(1991)firstdrewattentiontoH R4796, pointing out that itsinfraredspectralenergydistributionresemblestha tof **BPicbutwithtwicetheIR** excess(L_{IR}/L*).TheIRASdatashowednodiscernableemissionfromdust hotterthan ~110K, implying that there is little dust closer to the starthan40AU(Juraetal.1993). Theexistenceofoneormoreplanetswassuggestedasone ofthepossiblecausesofthis centralholeinthedustdistribution(Juraetal.1995;Jura etal. 1998). J98 and K98 determinedthatthedisk'sdiameteris -3", which corresponds to 200 A Uat the star's distanceof67pc, and they confirmed directly the existence ofthecentralhole.An especiallysignificantaspectofthediscoveryisthat ,basedonisochronefittingandthe nionHR4796B,located7.7" stronglithiumabsorptionline, the presumably coeval compa awayatPA=226 °(Juraetal.1993),isanM2-M3pre-mainsequencestarwith anage ± 3) $\times 10^{6}$ yr(Staufferetal. 1995; Juraetal. 1998). tightlyconstrainedtobe(8 Observationalandtheoreticalstudiesofoursolarsystem, includingchronologicaldata $\sim 10^7$ yrisareasonable estimate of the timescale fort formeteorites, suggest that he durationofthesolarnebulaandthegrowthoftheplanets tonearlytheirfinalmasses(see reviewsbyLissauer1993andPodosek&Cassen1994).Thus,theco nstraintsontheage ofHR4796Aplaceitwithinacriticalstageofplanetary diskevolution.

2

Ournewimagesclarifythestructureofthecentralcl earingzone, orhole, which may beindirectevidenceforaplanetorbitingHR4796A(see,e.g. ,Roquesetal.1994).They mayalsorevealanasymmetryinthedisk,possiblyconf irmedbytheNICMOSimages (Schneideretal. 1999), that could reflect the influence of thestellarcompanionora hypotheticalplanet.Finally,thesemulti-wavelengthdat aprovideareasonablebasisfor estimatingacharacteristicsizeforthemid-infrared -e mittingdustparticles, an estimate ution.Ourresultsanddiscussion withimplicationsforourunderstandingofthediskevol complementthoseinacompanionpaperbyWyattetal.(2000, hereafterW2000).

2.MID-INFRAREDIMAGINGWITHOSCIRATKECKII

WeimagedHR4796AintheN(10.8 μ m)andIHW18(18.2 um)bandsattheKeckII Telescopein1998May7-12withOSCIR, the University of Flo ridamid-infrared imager/spectrometer.OSCIRcontainsa128 ×128-pixel,Si:AsBlockedImpurityBand (BIB)arraydevelopedbyBoeing.Theplatescalewas0.062ar csec/pixelwiththef/40 choppingsecondaryattheKeckIIvisitorbent-cassegrai nport, which gave an array field of view of 7.9" \times 7.9". We used a guidest arand the off-axis autoguider, which ensured excellenttrackingandimagestability, and the standardc hop/nodtechnique,withachop frequencyof4Hzandathrowof8"indeclination.Thes tars α Sco, γ Aql,and δ Vir wereusedforfluxcalibration. These eingwasgenerally verygoodforourobservations. Typicalvaluesofthefullwidthathalf-maximuminten sity(FWHM)oftheobserved pointspreadfunctions(PSFs)were0.32"at10.8 µmand0.45"at18.2 µm,comparedto the diffraction-limited values (λ /D) of 0.22" and 0.37", respectively. Quadratic µm"seeing" of -0.2"-0.3" (which may include other subtractionoftheseimplies10-20 contributions, e.g., from autoguider jitter). The diffracti onlimitsat10and18 um correspondtofourandsixpixels, respectively, and thus OSCIRspatiallysamplesthe diffraction-limitedPSFmorefinelythanconventional Nyquistsampling.Weencountered lightcirrusthroughoutmuchofourKeckrun.Bycomparingo bservationswithinand amongthenights, webelievethat we have been ablegen erallytoremovethedatafor which the photometry is most uncertain. However, becaus eofthecirrus,some unquantifiableuncertaintyremainsinthephotometry.Key properties of the observations, including flux densities and noise estimates, are listed i nTable1.Theimagesare presentedascontourplotsinFigures1aand1b.Figure1cshow stheunsmoothed18.2 µmimage, which indicates the quality of the original data.

3.DISKMORPHOLOGYINTHETHERMALINFRARED

 $\label{eq:holest} The discovery images clearly showed the disk around HR4796A to be extended at 18 \ \mu m, but both teams only marginally resolved it at 10 \ \mu m (J98;K98). Figure 1 \ demonstrates that the emission ridge corresponding to the edge-ondisk is well resolved at both 10 and 18 \ \mu m in the senew images. For comparison, the PSFs determined from nearby stars are presented in Figure 2 on both logarithmica and linear brightness scales. At 10 \ \mu m, the disk is evident as very low-level wings protruding from or the strong stellar points ource (which, for convenient display, is there for the strong stellar or the s$

1a). Theridge is much more prominent at 18 μ m (plotted linearly in Figure 1b). The line connecting the two outer peaks is oriented along PA=26 ° ±2°, and the maximum extent of the lowest 18 μ m contour displayed in Figure 1 is about 3.1", or 208 AU. The maximum extent at 10 μ m is about 2.8", or 85 -90% that of the 18 μ m emission, and thus the 10 μ m and 18 μ m sources are roughly equal insize along the ridge. We also resolved the 18 μ m disk along the minor axis, as indicated by the scansth rough the lobes and a comparison starshown in Figure 3.

Threepeaksofcomparablebrightnessareembeddedinthe18 umridge.Eachofthe twoouter18 µmpeaks, which were fer to as the lobes, is separated fr omthecentralpeak by0.8"(54AU). The brightest region within the central 18 umcontourisindicatedin Figure1bbyafilledcircle.Thatpointisslightlyoff setfromthecentroidofthehighest centralcontour; it lies on the line connecting the tw olobesandhalfwaybetweenthem. ByimagingsequentiallyatthetwowavelengthswiththeK eckIIvisitor-portautoguider edeterminedthatthecentral18 inoperation, which facilitated accurate registration, w μm peakcoincides with the 10 µmpeaktowithin0.3". Asdescribedbelow, mostof the 10 µmemissionisdirectstarlight, and therefore the co incidencebetweenthecentral10 and18 µmpeaksimpliesthatthecentral18 umpeakalsocoincides with the star, which wethereforeassumeiscoincidentwiththefilledci rcleinFigure1b.Inthe10 umcontour plotinFigure1aweindicate(crosses)thelocationo fthepeaksofthe18 umlobes.The 18 µmlobesclearlycoincidewiththe10 umwings, and therefore much, possibly most, ofthe10 µm-emittingdustmustbemixedwith, or identical to, the 18 um-emittingdust. The18 µmbrightnessatthelobesiscomparabletothatatth estar, whereas their 10 μm brightnessisonlv4 -5% that at the star. The disk morphology that we see at18 µmis consistentwiththatseenbyK98intheirKeckimages.K 98didnotresolvethediskat 10 µm.

Themid-infraredradiationfromHR4796Aoriginatesbothin thestellarphotosphere and indust that is heated by the starlight. The stella rphotosphericfluxdensitiesat10and 18 µmareabout80% and5%, respectively, of the total observ edfluxdensities.Although the18 µmstellarfluxisasmallfractionofthetotal, it contributesalargefractionofthe emissionofthecentralpeakitself.Todeterminetheco ntributionofthestellar photospheretotheimage, we subtracted the stellar PSFf romtheimageofHR4796A afterregisteringthepeaksandnormalizingthecomparison star'sfluxtothatinferredfor HR4796A(Table1).ThePSFstarwaslocatednearHR4796Aand observedjustbefore and just after it. Due to the hexagonal shape of the Keckpri maryandsecondarymirrors, thePSFishexagonallyshaped, so that, for the most ac curatepoint-sourcesubtractionor deconvolution, one cannot assume circular symmetry oft hePSF.Duringthe18 μm observationsofHR4796A,thePSFrotatedabout23 °relativetotheimagerfieldofview (arotationinherentinthealt-azmount).Fromtheob servedPSFsweconstructedatimeaveraged PSF appropriate to the rotational range of the PSFduringourobservationsof HR4796A.

Theslightlysmoothed(threepixels)originalandstar showninFigures4aand4b,respectively.Thestarwasas brightestpart(filledcircle,Fig.1b)ofthecentral ma fluxremovesthecentralpeak,leavingaholeatthatp

tar -subtracted18 μmimagesare as sumedtobecoincidentwiththe maximum.Subtractionofthestellar osition.Becauseoftheapparent

absenceinHR4796ofsignificantdusthotterthanabout110K, Juraetal.(1995) concludedthatthediskhasaroughly40AU-radiuscentralho leinit.Thisconclusion wassubsequentlyconfirmedbythefirstdirectimagesof -20 µm(J98;K98) thediskat and, more recently, by NICMOS images at 1-2 µm(Schneideretal.1999;seebelow). Weseetheholedirectlyinour18 umimage(Figures1b&4a)asadepressionbetween thecentralpeakandeachofthetwolobes; it is obvi ousintheimagewiththestarlight subtracted(Figure4b).Sometenuousemissionmayalsobe presentintheinnerregion, i.e., the region within -0.5" of the star. The appearance of this residual emissio ndepends onhowwecharacterizeandpositionthesubtractedstel larPSF.IntheAppendixwe addressbrieflysomeaspectsofhowthecomputedresiduali seffectedbyslighterrorsin thepositionandwidthofthePSFusedtosubtractHR4796A's starlight.

Wealsosubtractedthestellarphotospheric10 µmfluxfromthe10 µmimage.The centralholeisevidentinthatsubtractedimage,butth starthatthesubtractionprocedureisveryuncertain. confidenceintheaccuracyofthisresultinsofaras partoftheemittingregion.However,those10 µmresultsareusefulinthecontextof estimatingthecharacteristicsizeofthemid-infrared -e mittingparticles(§6).

Thereissomeevidencethatthelobesarenotofequal maximumbrightness:theNE lobeis 5.9 $\pm 3.2\%$ brighter than the SW lobe, where this asymmetry c orrespondstothe differenceinthelobepeakbrightnessesdividedbytheir mean.Thisvalueforthe asymmetryincludesa0.8% correction for an asymmetry int hePSF.Theformal statistical significance of the brightness asymmetry o fthelobesistherefore 1.8 σ . The brightnessasymmetryitselfwasdeterminedbyfittingapo lvnomialtoa 10-pixel $\times 10$ -pixel (0.6" $\times 0.6$ ") region approximately centered on each peak. The uncertaintyassociated with this asymmetry was establish edbyconvolvingthebest-fit model(§4)withagaussianapproximationtothePSF,applyingga ussiannoiselikethat observed across the array, and fitting the polynomiala smentionedabove.Afterrepeating thisprocedure 50,000 times, we computed therms deviation, which constitutedthe formaluncertainty. This asymmetry may also be apparent inthe10 umimage (particularlyinthescanshowninFigure8b,discussedbe low)andintheNICMOSimage (Schneideretal. 1999). The asymmetry is discussed in a muc hbroadercontextinthe companionpaperbyW2000.Webrieflycommentonitbelow.

4.MODELINGTHEBRIGHTNESSDISTRIBUTION

J98andK98modeledtheir ~20 µmimagesofHR4796Atoinferstructuralproperties ofthedisk.Bothmodelsassumedopticallythin,passived isksinwhichthedustparticles ngthas λ^{-1} .J98foundthatan haveaninfraredemissionefficiencyvaryingwithwavele herange40 -80AU edge-ondustydiskwithaninnerholeradiussomewhereint providesanadequatefittothespectralenergydistribution andtheimage.Thehigher instheirthin-diskmodel, resolutionKeckimageobtainedbyK98moretightlyconstra fromwhichtheyinferaninnerholeradiusof55 0 ±15AUandadiskinclinationof18 $(+9^{\circ},-6^{\circ})$ from edge on.

Inthecompanionpaper, W2000presentresultsofmodelingthe18 umbrightness distributionofHR4796A.Sinceitisrelevanttooures timateofthegrainsizeandtothe wesummarizethemodel comparisonofourresultstotheHSTNICMOSobservations umstellardiskandastellaror here.W2000considerthedynamicalinteractionofacirc planetarycompanion, astudy initially motivated by the de siretounderstandhowsuch interactionsgenerateasymmetriesandotherstructures incircumstellardisks.A distribution of orbiting dust particles is gravitationally perturbedbyacompanion, and W2000examinetheresultantorbitalanddensitydistributions fordustgrainsofagiven sizeandcomposition.W2000ignoretheeffectsofradiati onpressure, assuming, ineffect, $\beta \approx 0, \text{orbitHR4796A}; \beta \text{ is the ratio of the}$ thatlarge, dynamically stabled ust grains, with aparticle(called δ , the overpressure radiationpressureforcetothegravitationalforceon ratio,byBackman&Paresce1993).However,atleastin thecontextofMietheory, we knowthattheselargeparticlesarenottheonesthat weobserveinthemid-infrared.As discussedin§6,themid-infrared -emittingparticles, if Miespheresofastronomical silicates(Draine&Lee1984), are ~2-3 µmindiameter, with the corresponding value **β** ≈ 2. Radiation pressure will drive particles with such a high valueof Boutofthesystemon hyperbolicorbitsinafewhundredyears.Basedonthisne gligiblelifetimeandon ratesinthedisk,W2000(§6) estimatesoftheparticlenumberdensityandcollision conclude that these grains must be collision fragments ofmuchlargerparticles.The spatialdistributionoflargerparticlesintheHR4796Adi skisunknown,so,asan alternative, W2000 assume that larger particles are distr ibutedessentiallylikethatofthe smaller, 2-3 µm-size, Miegrainsemitting the observed mid-infrared rad iation.By accountingforthetemperaturedependenceofthedustasaf unctionofdistancefromthe star, they determine the spatial distribution of them i d-infrared-emittinggrainsfromour 18 µmimage.

Themodeldiskisnotflat, butflared, with a distribut ionoforbitalproperinclinations andpropereccentricitiesidenticaltothoseofthemai n-beltasteroidsinourownsolar system. The distribution of semi-major axes for the relationn(a) $\propto \alpha^{\gamma}$, wheren(a) is the number of orbits per unit semi-major particleorbitsthatarecircular, which is approximately surfacenumberdensityasafunctionofdistancerfromt tor γ^{-1} . The largest value, a max, used for the semi-major axis of the dust or bits is 130 AU, althoughtheresultsofthemodelingarenotsensitiveto fluxarisesneartheouteredgeofthedisk.Keymodelpar theobservationsare:(1)theexponent γ ;(2)theminimum value, a majoraxis, which controls the diameter of the centra thedensitydistribution;and(4)theinclinationofthed sight.Inaddition,observedasymmetriesinthebrightne adjustingthecommonforcedeccentricityandthedirec diskparticleorbits.

Figure4cshowsthemodelbrightnessdistributionat18 star-subtractedimageinFigure4b.Theresidual(observed Figure4d.Themodelfitsthelobeswell,withnearlyal within 0.4" of the star. We are not certain of the nat

particlesisassumedtofollowthe axis.For trueforthemodelorbits,the hestaristhereforeproportional thisvalue, sincerelatively little ametersthatarevariedtomatch min, of the dust semilclearing;(3)thenormalizationof isksymmetryplanetothelineof ssdistributionaremodeledby tionoftheforcedpericenterofthe

umthatisthebestfittothe minusmodel)isshownin loftheresidualfluxlocated ureoftheresidualflux.Someofit

mayrepresentrealfluxfromHR4796A'sinnerdisk,butsom eofitmayresultfrom imperfectsubtractionofthestellarphotosphericflux stemmingfromvariationsinthe seeingnotmanifestedinthePSF-starimages(whichwere ofshorterduration).We brieflyaddressthisissueintheAppendix.

Themodeldiskinclinationis13 ° ±1 ° fromedge-on. Aflatdiskwouldbemodeledas moreface-onthanthis, since the disk flaring that we assumeinthemodelaccountsfor theminoraxis(Figure3).Forthe someoftheresolveddiskextensionthatweseealong distribution of semi-majoraxes, $\gamma = -2 \pm 1$. The resultant model distribution of surface numberdensity(i.e., integrated normal to the disk) of 2. 5 µmgrainsisshowninFigure5 intheformofaradialcutthroughtheazimuthallyaver ageddisk(solidcurve).The surfacedensityisrelativelysharplypeakednear70AU. Exteriortothisposition, the surfacedensitydecreaseswithdistanceapproximatelyas r⁻³.Interiortothisposition,the and60AU, falling to zeroby surfacedensitydropssteeplybyafactoroftwobetween70 $min=62 \pm 2AU$, but because the dust particle orbits are 45AU.Themodelfitimpliesa assumedtobenon-circular,theholeisnotsharplyde finedinthemodels; i.e., the orbital eccentricityresultsinasmallrangeinaparticle' sradiallocation, which softens the hole'sedge.Flat-diskmodelsbasedonthedisk-discoveryi magesindicatecomparable valuesfortheholeradius (J98;K98), but the dynamical models considered here provide an apriori basisforunderstandingwhyaholeshouldhaveanindisti nct,orsoft,edge.

Howemptyistheholeinthedisk?Wecanestimatet heopticaldepthintheholeby comparing the brightness of the residual emission there (assumed to represent a nupper limittoemissionfromdust)totheobservedbrightness ofthelobes.Considerfirstthe opticaldepthinthedisklobes.TheSWlobeis54AUin projectionfromthestar. The modeltemperaturethereis113K(comparedtotheblackbody temperatureof81K)and, ignoringprojectioneffects, the corresponding edge-onop ticaldepthat18 µmis -0.002. Theresidualemission(observedminusmodel;Fig.4d)at20A U(0.3")fromthestaris typically0.2thatofthelobes.Themodeltemperatureat 20AUis163K, and thus the -5×10^{-5} , which is about 3% that of the lobes. 18 µmedge-onopticaldepththereis theedge-onopticaldepthinthe Assuming that the ratio of the face-on optical depth to holeiscomparabletothatratioatthelobes, thent heaverageopticaldepthof18 umemainpartofthediskat60emittingdustintheholeappearstobeabout3%thatinth 100AUfromthestar.Ifanyoftheresidualemissionres ultsfrompoorPSFsubtraction and is not intrinsic to the source, the optical depth of thedustintheholecouldbeeven lessthanthis. The central cleared region appears to b erelativelyveryempty.

Therehasbeenspeculationaboutthecausesofholesi nthecentersofcircumstellar disks(Lagage&Pantin1994;Juraetal.1998;Strom,Edwards,& Skrutskie1993: Roquesetal. 1994; J98; K98), and the disk in HR4796 Aislikely tobecomeastrong focusforthesediscussions(e.g., Juraetal. 1995; W2000). We onlywishtoemphasize herethatthepossiblediskbrightnessasymmetryisan interestingandperhapsunexpected manifestationofthecentralhole. The brightness as ymmetryinthemodelresultsfromthe gravitationalperturbationofthediskparticleorbitsbyt helow-masscompanionHR tric(W2000).Tofirst 4796Bevenifthecompanion'sorbitisonlymoderatelyeccen order, the effect of the perturbation is to shift the c enterofsymmetryofthecirculardisk fromHR4796Atowardsthecompanion'sapocenter, where the companionspendsmost

ofitstime; i.e., the long-termaverage of the pertur bationonthediskparticlesisdirected towardsthecompanion'sapocenter.Thesideofthedisk,a ndthereforetheholeedge,that isclosesttothecompanion'spericenterisshiftedclose rtotheprimarystarandis thereforewarmer("pericenterglow"). Thepericentergl owisadirectresultoftherebeing aholeinthedisk.Ifthismodelisvalidandthesepertur bationsdoarisefromHR4796B, thenHR4796B'spericentermustbelocatedNEofHR4796A, and HR4796Bis currentlylocatednearitsapocenterSWofHR4796A.Thes caleoftheshiftofthedisk's centerofsymmetryisindicatedbycomparingthedashedan ddottedcurvesinFigure5, whichshowsthedensitydistributionthroughthedisktowa rdsandawayfromthestellar which, whiles mall, leads to companion'spericenter.ThediskisshiftedonlyafewAU, anarguablymeasurablebrightnessasymmetry. Theperturber responsiblefortheshiftof thediskneednotbeexteriortothedisk.Astarorplanet withintheholecouldcausethe sameeffect(W2000);however,basedonnear-infraredspeckl eobservations(Juraetal. 1995), the mass of any stellar companion located between 11a nd120AUfrom HR4796Amustbelessthan0.13M .Thefactthatwemaybedetectingabrightness asymmetryresultingfromafew-AUshiftinthediskcent roidimpliesthatthedeted roidimplies that the detection of significantdynamicaleffectsoncircumstellardisks bystellarorplanetarycompanionsis nowwithinreach.TheHST/NICMOSimagespresentedbySchn eideretal.(1999)may confirmthispericenterglow($\S5$).

5.COMPARISONOFMID-INFRAREDANDHST/NICMOSIMAGES

Schneideretal.(1999)haveimagedtheHR4796Adiskat1.1and1. 6 µmwith NICMOS. They used a 0.6"-diameter coronographic spot, but, duetoavarietyof instrumentaleffects, the dataina ~1.3"-diameterregionareunusable. Theirimages show clearlywhatappearstobearelativelynarrowringincl inedtothelineofsight.Assuming thatthering is intrinsically circular, they determine amajor-axispositionangleof26.8 ° ± 0.6° and an inclination to the line of sight of 73.1 $^{\circ} \pm 1.2^{\circ}$, in a greement with our estimates(W2000;§§3&4)andK98's.Themeanradiusofthe ringis70AU, and, after quadratic subtraction of the PSF, the full width of the annulusisestimatedtobe17AU (Schneideretal. 1999). Therelatively sharp maximum in th edensitynear70AU determinedfromourmodelingoftheOSCIRobservationsagree spreciselywiththevalue derivedbySchneideretal.(1999)forthemeanradiusofthe ringbasedontheNICMOS observations. We now consider a more detailed comparison

InFigure6weshowour10and18 µmcontoursoverlaidonthe1.1 **µmNICMOS** image.Keepinginmindthedifferencesintheresolutions oftheimages(FWHM:0.12"at 1.1 μm;0.32"at10.8 μm;0.45"at18.2 μm), we see that the overall coincidence of the mid-infraredandnear-infraredemissionisexcellent. Howe ver,the18 µmlobesare locatednoticeablyclosertothestarthanarethebr ight1.1 µmpeaks.Thepeaksofthe 18 µmlobesareseparatedby1.73", comparedto2.10" for the peak-i ntensitypositionsat 1.1 µm(Schneideretal.1999).Toassessthenatureofthisoff set.weconsidereda perfectlythinring(annulus)ofdustwithuniformnumberdensi tyandthetemperature distributionderived in the W2000 models. We assumed a mean annularradiusof70AU and an annular width of 17 AU and computed the 18.2 *µmbrightnessateachpointinthe*

ring, which we convolved with the observed 18.2 µm PSF. In Figure 7 we compare the brightness distributional ong the majoraxis expected f or this convolved model to the observed brightness distribution.

Thiscomparisonimplies that ~60% of the 0.37" difference in the separation of the peaksat1.1and18.2 umresultsfromthelowerresolutionofthe18 µmimage.Figure7 indicatesthattheuniform-annulusmodelresultsinan18 umdistributionwithlobes hanactuallyobserved.The displacedtolargerradiiandappearingsomewhatnarrowert discrepancyinslopeisprimarilyontheinsidesofthe lobes.Theobservedouterslopeof theNElobeiswellmatchedbythemodel,whilethemat choftheSWlobeisnoticeably NICMOS-basedmodeland poorer.Wedonotthinkthatthesedifferencesbetweenthe the18 µmobservationsresultfrominaccuratesubtractionofthe stellarPSFfromthe 18 µmimage.Figure7givessomeindicationofthesmalldepe ndenceoftheshape inferredfortheinsideofthedisklobesontheassumed shapeofthePSF;weshowthe fluxdistributionsresultingfromuseofthenominal,o bservedPSF(solidcurve)andthe samePSFwithanadditional0.35" of seeing (dashedline; se eAppendix).Rather,the differenceintheobservationsandthebrightnessexpect edfromthemodelmayresult ilvbecauseofthelower from the annulus having a somewhat soft inneredge. Primar resolutionofthe18 µmobservations, amore gradual density fall of fon the ins ideofthe annuluswillbothshifttheobservedpeaksinwardandlower theinnerslopeofthelobes. Theexcellentagreementinthelocationofthepeakdensi tyinthedisk,asdetermined from models of the OSCIR images, and the average radiusoftheannulusdetermined frommodelsoftheHST/NICMOSimages, suggests that it i sthisinnerdensityrolloff,or asimilarrefinementsuchasexcessemissionintheh ole, that accounts for the relative locationsoftheobservedemissionpeaksinthetwodat asets.Aslowerdensityfalloffon the outside of the annulus would elevate the 18 µmwings, although we see this only on theSWsideofthedisk.AsshownbyW2000anddiscussedin theprevioussection, one mayaccountforasoftdensityfalloffontheinsideo ftheannulusbythekindsof dynamicalpropertiesanticipated(andassumedintheW2000mo dels)fortheorbiting dustparticles.

Figure6showsthatat1.1 μ mtheNEsideoftheannulusisnoticeablybrighterthan theSWside.Thisasymmetryisnotevidentinthe1.6 µmNICMOSimage.This asymmetrymaysupportour, albeit marginal, detection oft hepericenterglowat18 μm. However, one isseeing emission from dust particles at 18 µm, butscattering from dust µm.Wespeculatethatthemassshiftthat (and probably a different population) at 1-2 mayresultintheenhancedlobebrightnessat18 µmcouldalsoenhancethescatteringof lightthere.Detailedmodelsthatincorporaterealisti cscatteringfunctionsmustbecarried outtoexplorethisinterestingpossibility.

6. THEMID-INFRARED -EMITTINGDUST

6.1TheCharacteristicParticleSize

Circumstellardustspansarangeofcompositionsandsizes. However,foragiven composition,onecanderivefromobservationsattwowav elengthsauniqueparticlesize

thatisarguablycharacteristicofthedustresponsiblef orthatemission. The basis for this estimateisthattheemissionefficiency, and theref orethetemperature, of a dust particle dependsonitssize.Wemadeaninitialestimateofthe sizeofdustparticlesinHR4796A umfluxesinthelobes(aftercorrectingforthe bytakingtheratioofthe10and18 differentPSFsatthetwodifferentwavelengths)andc omparingitwiththatexpected vedifferentemissionefficiencies from Miesilicates pheres of different sizes, which ha and consequently different temperatures. The formal value ofthisinitiallyestimated diameterwas2.5 µm.(Miesilicategrainsofthisdiameterhaveinfrared emission efficiencies varying approximately as $\lambda^{-1.9}$.) This initial estimate was then used to do the modelingdescribedin§4, which was repeated using 2 and 3 um-diameterparticles.In Figure8aweshowtheresultantscanalongthemajoraxi softhe18 µmimagefrom which the stellar photospheric emission (using the nominal PSF;seeAppendix)hasbeen subtracted.Themodelprovidesagoodfittothe18 µmlobestructureforarangeof assumeddustsizes. However, Figure 8bshowshow the 10 umbrightnessconstrainsthe diameterto2.5 µm, consistent withour original estimate. We concludet hat, under the assumptionthattheparticlesweseeinthelobesare astronomicalsilicateMiespheres,2-3 µm-diametergrainsprovideareasonablefittothelobem id-infraredcolors.Ofcourse, basedonstudiesofinterplanetarydustparticlesinours olarsystem(e.g.,Gustafson 1994), it is evident that the dust particles in HR4796A's disk areunlikelytobesimple spheres, the implications of which we are in the process ofconsidering.

6.2ParticleEvolution

Wecanconsidertheeffectsofthestellarradiation on2.5 µm-sizegrainsby β , the ratio of the radiation pressure force to the estimatingthevalueoftheparameter gravitational force on a particle. Assummarized in Art ymowicz(1988) and elsewhere, ρ is its density, $\langle Q_m \rangle$ is the radiation $\beta \propto \langle Q_{pr} \rangle (L_*/M_*)/R\rho$, where Ristheparticle radius, pressureefficiencyaveragedacrossthestellarspectral energydistribution, and L */M*is theratioofthestellarluminosityandmass.W2000have computedthevalueof βfor2.5 µm-diametersilicatespheresnearHR4796A;theyassumedL *=21L °, M *=2.5M °, and $\rho = 2.5$ gcm ⁻³, and they compute $\langle Q_{pr} \rangle$ using a stellar spectral energy distribution for theA0VstarVega(M.Cohen, private communication).T heyderive the value $\beta = 1.7$ for the grainst hat characterize the mid-infrared -emittingparticlesinHR4796A'sdisk.If onetakesintoaccountthedifferencesinspectraltype and assumed grain density, this valueagreesreasonablywellwiththeonethatcanbede rivedfromtheresultsin Artymowicz(1988, seehisFigure1), who computed βforthestar βPicandforavariety ofgrainmaterialsincludingtheastronomicalsilicates ofDraine&Lee(1984).Dust grainsthatarecreatedincollisions, and for which β is greater than -0.5, have hyperbolic orbits;theyareblownbeyond ~100AUandoutofthesystemonessentiallyfree-fall time 2-3 µmgrainsweseeorbitingHR scales, on the order of a few hundred years. Thus, the 4796Aarenotprimordial; one expects very few of the min thisdisk(seealsoJuraetal. 1995). The detailed analysis by W2000 (see also Artymowicz 1997) i mpliesthatthe ⁴yracrossthedisk.ThemidcollisionallifetimeofthedustinHR4796Aisroughly10

infrared -emittingparticlesareprobablyrelativelyrecentfragm entsofcollisionsthat dominateboththecreationanddestructionofdustthere.

7. SUMMARYANDCONCLUSIONS

1.Ournew10and18 μmimagesofthecircumstellardiskaroundtheA0Vega-typ estar HR4796Aimagesshowextended,wellresolvedemissionat10 μminadditiontothe previouslydiscovered18 μmemission(J98;K98).Theresolutionofbothimagesis close totheKeckdiffractionlimit.Thesizeofthe10.8 μm-emittingregioniscomparableto thatat18 μm,implyingthatmostofthe10 μm-emittingparticlesaremixedwith,or identicalto,thoseemittingat18 μm.

2. Modelingofthediskbrightnessdistribution(W2000) impli esthatapronounced maximuminthesurfacedensityoccursatapproximately70AUf romthestar.Thisvalue equalsthemeanannularradiusdeducedbySchneideretal.(1999) fromtheir HST/NICMOSimages.Thebest-fitmodelsurfacedensitye xteriorto70AUfallssteeply withdistanceasr ⁻³.Interiorto70AU,thedensitydropssteeplybyafactor oftwo between70and60AU,fallingtozeroby45AU,whichcorresponds totheedgeofthe hedynamicalmodels,this"soft" previously discovered central hole; in the context of t edge for the central hole results from the fact thatthedustparticleorbitsarenon-circular.

3.Basedontheimages,theopticaldepthinthecentra lclearingzone,orhole,is -3%, or less,thatinthemainpartofthedisk.Thus,thehole isindeedrelativelyempty.

4.Wepresenttentativeevidence(1.8 umlobesareofunequalbrightness. σ)thatthe 18 µmbrightnessdistributionbasedondynamical Detailedmodelingofthe18 considerationsandpresentedinacompanionpaper(W2000)indica testhatsucha brightnessasymmetrycouldresultfromgravitationalper turbationofthediskparticle orbitsbythelow-massstellarcompanionoraplanet.Th eperturbationshiftsthedisk centerofsymmetryawayfromthecompanion'spericent er:tofirstorder.thisshift correspondstoaspatialshiftofthecenterofthecirc ularlysymmetricdisk.Thisshift bringstheinsideedgeoftheannulusclosertoHR4796A, which makesthatedgewarmer andmoreinfrared-emitting("pericenterglow")thantheot hersideoftheannulus. This measurablebrightnessasymmetrycorrespondstoaspatials hiftofthediskofonlyafew AU, and so the detection of even fairly complex dynamical effectsofplanetsondisks appearstobewithinreach.Therecent1.1 µmNICMOSimage(Schneideretal.1999) mayshowevidenceinsupportofthepericenterglow.

5.Wecompareour10and18 µmimageswithrecentNICMOSimagesofHR4796Aat 1.1and1.6 µmthatshowwhatappearstobearelativelynarrowri ng,orannulus,inclined tothelineofsightandwithameanradiusof70AUanda nannularwidthof ~17AU. Theoverallcoincidenceofthenear-infraredandmid-infr aredstructureisexcellent. However,evenafteraccountingfordifferencesinangu larresolution,wefindthatthe18 µmdistributionexpectedfromtheNICMOS-basedmodeldif ferssomewhatfromthe observeddistribution. These differences sugges that the annulus has a "soft" inneredge, which may correspond to the density fall offex pected from the considerations of particle dynamics.

6.Usingthe10and18 µmimages, we estimate that the mid-infrared-emitting dust particlesinthemainpartofthediskhaveacharacte risticdiameterof -2-3 µm. This areMiespherescomposedof estimateisbasedontheassumptionthattheparticles astronomical silicates. The corresponding value of the parameter β , the ratio of the radiation-pressure forcetogravitational force on the particle, is 1.7. Particles with $\beta > 0.5$ are expected to be quickly blow nout of the system. We co ncludethatthemidinfrared -emittingparticlesintheHR4796Adiskarenotprimordia l, butrather they are relativelyrecentandtransientcollisionfragments.

ACKNOWLEDGEMENTS

WewishtoacknowledgeKevinHannaforhisdevelopmentofthesuperbOSCIRelectronics;KevinHanna,JeffJulian,andChrisSingerfo
for,andcarryingout,theKeckobservingrun;MartinCohen
Vegaspectrum;andtheKeckObservatorypersonnelfortheirtheirassistanceinpreparing
forprovidinguswiththe
routstandingsupportofVegaspectrum;andtheKeckObservatorypersonnelfortheiroutstandingsupportof
Keckwassupportedinpartby
PennsylvaniaStateUniversity

APPENDIX

Wepointedoutin§3thatsomeoftheinnertenuousemiss ion(within0.5" of the star) apparentinthestar-subtracted18 umimageofHR4796A(Figure4b)andasresidual emissioninFigure4dmaybeanartifactresultingfromt heuseofanunsuitablePSF.This mismatchcouldoccuriftherearelow-frequencyseeingco mponentssampledduringthe ~1 hofchoppedintegrationtimeonHR4796A(Table1)thatare notsampledduringthe muchshorter(~1 m)integrationtimeonthenearbystarusedtodeterm inethePSF. (Giventhelimitedamountofavailableobservingtime, it isunrealistictoobservethePSF starforthesameamountoftimethatweobservethe programobjects.)Herewebriefly examinethisissue.

Figure9apresentsthestar-subtracted18 µmimagepresentedpreviouslyin§3 (Figure4b). That image resulted from the subtraction ofHR4796A'sdirectstarlight (photosphericemission)usingtheobservedstellarPSF(r eferredtoasthenominalPSF) andassumingthatthestellarpeakiscoincidentwiththe locationofmaximum18 um brightness(filledcircle,Figure1).Figure9bshowsthe resultofinsteadsubtractinga somewhatbroaderPSFobtainedbyconvolvingthenominalPSF withagaussianprofile withaFWHMof0.35". This might represent some additiona lseeingthatcouldhave affectedtheimageofHR4796AbutnottheobservedPSFst arimage.InFigures9cand 9d, were peat this exercise but with the star assumed to b eatthecentroidofthehighest centralcontourindicatedinFigure1b, i.e., at0.1"WN WofthefilledcircleinFigure1b. Theintentofthisexerciseistoillustratethesen sitivitvoftheresidualinnerstructureto theassumedstellarlocation.

Weseethateventhismodestamountofextra" seeing" re sultsinastar-subtracted imageofHR4796Athatisnoticeablylessstructuredatth ediskcenter.Thiscomparison isalsoshowninthescansinFigure7.Likewise,slig hterrorsinpositioningthestarresult insomedifferencesintheinferredinner-diskstructure. However, although the emission structuredifferssomewhat, the integrated residual fluxi svirtuallyunchanged, since the stellarfluxisthesameinbothcases. Aswehaven oted.thelightcirrusevident throughoutmuchofourKeckrunmayhaveresultedinpoorphoto metricconditions, but wedonotthinkthisisthesourceoftheresiduals:thepho tometryisgenerallyconsistent towithin ~20%, and even an error this large in the PSF flux nor malizationwouldonly marginallyeffecttheresidual.

We conclude that the detailed structure of the inner disk inferred from the star-subtracted image or the resultant (e.g., Figure 4d), must be viewed with caution. However, th more robust and may be real. If so, it may correspond t within the central clearing. (within 0.5" of the star), as image of the residual emission eintegrated residual flux is olow-level dust emission from

13

REFERENCES

Artymowicz, P.1988, ApJ, 335 , L79				
Artymowicz, P.1997, AREarthPlanet. Sci, 25, 175				
Backman, D., Gillett, F., & Witteborn, F. 1992, ApJ, 385, 680				
Backman, D., & Paresce, F. 1993, in Protostars and Planet	sIII,eds.E.Levy&J.Lunine			
(Tucson:U.ArizonaPress),1253				
Beckwith, S.V.W., & Sargent, A.I. 1993, in Protostarsa	ndPlanetsIII,eds.E.Levy&J.			
Lunine(Tucson:U.ArizonaPress),521				
Boss,A.,&Yorke,H.1996,ApJ,469,36				
Draine,B.T.,&Lee,H.M.1984,ApJ,285,89				
Gillett,F.1986,inLightonDarkMatter,ed.F.Israel (De	ordrecht:Reidel),61			
Greaves, J.S., et al. 1998, ApJ, 506, L133				
Grün, E., etal. 1985, Icarus, 62, 244				
Gustafson, B. Å.S. 1994, AREPS, 22, 553				
Holland, W.S., et al. 1998, Nature, 392, 788				
Jayawardhana, R., Fisher, S., Hartmann, L., Telesco, C., F	Piña,R.,&Fazio,G.1998,ApJ,			
503,L79				
Jura,M.1991,ApJ,383,L79				
Jura, M., Ghez, A.M., White, R.J., McCarthy, D.W., S	mith,R.C.,&Martin,P.G.			
1995,ApJ,445,451				
Jura, M., Malkan, M., White, R., Telesco, C., Piña, R., &	Fisher,R.1998,ApJ, 505,8	97		
Jura,M.,Zuckerman,B.,Becklin,E.E.,&Smith,R.C.	1993,ApJ,418,L37			
Kalas,P.,&Jewitt,D.1995,AJ,110,794				
Kalas,P.,&Jewitt,D.1997,Nature,386,52				
Koerner, D.W., Ressler, M.E., Werner, M.W., &Ba ck	man,D.E.1998,ApJ,503,L83			
Kurucz,R.L.1979,ApJS,40,1				
Lagage,P.,&Pantin,E.1994,Nature,369,628				
Larwood,J.D.,Nelson,R.P.,Papaloizou,J.C.B.,& Te	erquem,C.1996,MNRAS,282	,		
Lissauer.J.J.1993.ARAA.31.129				
Papaloizou, J.C.B., & Terguem, C. 1995, MNRAS, 274, 987				
Podosek, F.A., & Cassen, P. 1994, Meteoritics, 29,6				
Roques, F., Scholl, H., Sicardy, B., & Smith, B.A. 1994,	Icarus, 108, 37			
Schneideretal. 1999, ApJL, 513, L127				
Smith, B.A., & Terrile, R.J. 1984, Science 226, 1421				
Stauffer, J.R., Hartmann, L.W., & Navascues, D.B. 1995	,ApJ, 454,910			
Strom, S.E., Edwards, S., & Skrutskie, M.F. 1993, in Proto	starsandPlanetsIII,eds.E.			
Levy&J.Lunine(Tucson:U.ArizonaPress),837				
Strom,K.M.,Strom,S.E.,Edwards,S.,Cabrit,S., &Sk	rutskie, M.F. 1989. AJ, 97, 1	1451		
Telesco, C.M., Becklin, E.E., Wolstencroft, R.D., & De	cher,R.1988,Nature335,51			
Wyatt,M.C.,Dermott,S.F.,Telesco,C.M.,Fishe r,R.S	.,Grogan,K.,Holmes,E.,&			
Piña,R.K.2000,ApJ,inpress(W2000)	-			

FIGURECAPTIONS

- Figure 1. Images of HR4796A made with OSCIR. Northisup, and east i stotheleft. Theimagesfromwhichthetwocontourplotswereconst ructedweresmoothed witha3-pixel(FWHM).or0.19arcsec.gaussian.Forboth contourplots,the lowestcontourlevelisaboutfourtimesthesmoothedn oise. (a) 10.8 µmimage withcontours(mJyperarcsec ²)spacedlogarithmicallyat6.4,9.9,15,23,36, 55,85,132,203,and312.Thefilledcircleisthelocationof thepeak ². The crosses indicate the brightness, with the value 480 mJy perarcsec positionsofthe18.2 µmlobes; (b) 18.2 µmimagewithcontours(mJyper arcsec²)spacedlinearlyat58,93,128,163,198,234,269,304,339,and374. Thefilledcircleindicatesthepresumedpositionofthe star, where the ²;(c) Unsmoothedversion, with linear brightnessis362mJyperarcsec brightnesslevels, of the 18.2 µmimage.
- Figure 2. Plotsofthepointspreadfunctions (PSFs) at 10.8 and 18.2 μ mbased on observations of comparison stars. Northis up, and east is to the left. All plots are normalized to a maximum value of 100 units located at the filled circle. (a) Linear 10.8 μ mlevels: 1, 12, 23, 34, 45, 56, 67, 78, and 89; (b) Logarithmic 10.8 μ mlevels: 1.3, 1.9, 2.8, 4.2, 6.2, 9.2, 13.7, 20.2, 29.9, 44.3, an d65.5; (c) Linear 18.2 μ mlevels: 8.0, 18.2, 28.4, 38.6, 48.8, 59.0, 69.2, 79.5, and 89.7; (d) Logarithmic 18.2 μ mlevels: 8.0, 10.6, 13.9, 18.4, 24.3, 32.1, 42.3, 55.9, and 73.7.
- **Figure3.** Comparisonofminor-axisscansthroughthe18.2 star,whichshowsthatthelobesareresolvedalongth Thenorthwestsideofthescanistotheright(posit iveoffsets).
- Figure4. Comparisonofthe18.2 μmimagesofHR4796Awiththemodelbrightness distribution.Eastisupandnorthistotheleft.Overl aidcontourlevels(mJyper arcsec²)arelinearlyspaced. (a) Observedimagesmoothedwith3-pixel gaussian,withcontourlevelssameasinFig.1b; (b) Observedimageafter subtractionofstellarphotosphericemission.Contourl evels:58,93,128,163, 198,234,269,304,339,and374; (c)Modelbrightnessdistribution; (d) Residualemission,correspondingtothedifferencebetw eenthestar-subtracted andmodelbrightnessdistributions.Contourlevels:20,40, and60.
- **Figure5.** Plotsofthesurfacenumberdensityofthe2.5 µmdustgrainsinthedisk derivedfrommodelingthe18 µmbrightnessdistribution.Thesolidcurveis theazimuthalaverageofthesurfacenumberdensity,the dottedlineandthe dashedlineindicatethedensitythroughthediskinthedire ctionofandaway from,respectively,thestellarcompanion'spericenter. Therelative

displacementofthedottedanddashedlinesindicatesthe scaleofthe displacementofthediskcenterfromthestar.

- **Figure6.** Overlayof (a)10.8and (b)18.2 μmcontoursonthe1.1 μmHST/NICMOS imageofHR4796AdiskfromSchneideretal.(1999).Theangula rresolutions (FWHM)oftheimagesare:0.12"at1.1 μm;0.32"at10.8 μm;0.45"at 18.2 μm.Inbothimages,eastisapproximatelydown,andnorth istotheleft (seeSchneideretal.1999).Inthisorientation,thecompa nionstarHR4796Bis locatedtotheupperright.
- Figure7. Comparisonofmajor-axiscutthroughthelobesofour18 µmstar-subtracted image(solidcurve)tothat(dottedcurve)expectedfromt hesimplestdisk modelinferredbySchneideretal.(1999)fromtheirNICMOS image.Theleft sideofthescanisnortheastofthestar. The model diskisathinannuluswitha andthe meanradiusof70AU,awidthof17AU,auniformdensity, temperaturefalloffwithradiusdeterminedbyW2000.TheSchnei deretal. modelhasbeenconvolvedwiththeobserved18 umPSF.Thedashedcurve showstheresultofsubtractingfromtheobservations astellarPSFthathasan additional0.35" seeing, as discussed in the Appendix.
- **Figure8.** Comparison (a) the star-subtracted 18.2 µm and (b) 10.8 µm scans, indicated assolid lines, along the disk major axis of the HR4796Adisk.The leftsideofeachscanisnortheastofthestar.In the18 umscan.thefeatures within ~0.3" of the stellar position are probably artifacts res ultingfrom imperfectPSFsubtraction.Onlythelobesofthe10 umscanareshown(see text).The18 µmbrightnessdistributionhasbeenmodeled,asdescrib edinthe text, using spherical Miesilicate particles that are 2,2.5,and3 µmindiameter. The corresponding scanof the 10.8 *µmfluxdistributionexpectedfromadisk* composed of each of these particles indicates that the 10.8 µmbrightnessinthe lobesisbestfitby2.5 um-diameterparticles. Thus, in the context of Mie theory, the grains are constrained to be a few microns indiameter.
- µmdiskemissionresulting from the subtraction from t Figure9. Comparisonofthe18 he observedimage(Figure1b)ofamodestrangeofstellarpoi ntspreadfunctions normalizedtotheexpectedfluxofthestarHR4796A.Nort hisup, and east is totheleft.AsdiscussedintheAppendix,theintento fthisexerciseisto illustratetheeffectontheinferredinnerdiskstructure oftheuseofan inaccurate, orslightlymis-positioned, PSF. (a)NominalobservedPSF (FWHM=0.45") subtracted at brightest spot(filled circle inFigure1b); (b) Seeing-convolvedPSF(seeingFWHM=0.35")subtracted atbrig htestspot; (c)NominalobservedPSFsubtractedatcentroidofhighest centralcontour; (d) Seeing-convolvedPSFsubtractedatcentroidofhighestcen tralcontour.

Table1:OSCIRObservationsofHR4796A		
Ν	IHW18	
10.8	18.2	
5.3	1.7	
2784	3480	
0.188 ±0.047	0.905 ±0.130	
0.014	0.15	
0.150	0.048	
	817	
	26	
142		
	DbservationsofH N 10.8 5.3 2784 0.188 ±0.047 0.014 0.150 142	

^a Integrationtimescorrespondtobothhalvesofthecho ppercycle.Theyarethesumof skypositionwereprojectedonto thetimesforwhichthesourceitselfandthereference thedetector.

^b Theuncertainties for the 10.8 and 18.2 μ mfluxdensities, which are based on the reproducibilityofstandard-starfluxes, are principally due tothepresenceoflightcirrus clouds. This problem was more severe for the 10.8µmthanforthe18.2 µmimage.

с The10.8 µmphotosphericfluxdensitywasderivedbyJuraetal.(1998) by extrapolatingtheK-band(2.2 µm)fluxdensityto10.8 µmusingtheKurucz(1979)

This figure "figure1.gif" is available in "gif" format from:

This figure "figure2.gif" is available in "gif" format from:

This figure "figure3.gif" is available in "gif" format from:

This figure "figure4.gif" is available in "gif" format from:

This figure "figure5.gif" is available in "gif" format from:

This figure "figure6.gif" is available in "gif" format from:

This figure "figure7.gif" is available in "gif" format from:

This figure "figure8.gif" is available in "gif" format from:

This figure "figure9.gif" is available in "gif" format from: