

# Aubry-Mather Theory and Idempotent Eigenfunctions of Bellman Operator\*

A.N. Sobolevskii

Moscow State University  
e-mail: ansobol@idempan.phys.msu.su

## Abstract

We establish the connection between the Aubry-Mather theory of invariant sets of a dynamical system described by a Lagrangian  $L(t, x, v) = L_0(v) - U(t, x)$  with periodic potential  $U(t, x)$ , on the one hand, and idempotent spectral theory of the Bellman operator of the corresponding optimization problem, on the other hand. This connection is applied to obtain a uniqueness result for an eigenfunction of the Bellman operator in the case of irrational rotation number.

## 1 Introduction

Consider a Lagrangian

$$L(t, x, v) = L_0(v) - U(t, x),$$

where  $t, x \in \mathbb{R}$  and the functions  $L_0(v)$ ,  $v \in \mathbb{R}$ , and  $U(t, x)$  satisfy the following conditions:

- (L<sub>1</sub>)  $L_0(p)$  is of class  $C^1$ ; its derivative  $L'_0(p)$  is strictly growing:  $L'_0(p_2) > L'_0(p_1)$  iff  $p_2 > p_1$  and has a locally Lipschitzian inverse.
- (L<sub>2</sub>) For any  $N > 0$  there exists  $P > 0$  such that if  $p \in \mathbb{R}$ ,  $|p| > P$ , then  $L_0(p) > N|p|$ .
- (L<sub>3</sub>)  $U(t, x)$  is of class  $C^1$  and its derivative  $\partial U/\partial x$  is Lipschitzian.
- (L<sub>4</sub>)  $U(t, x)$  is periodic:  $U(t+1, x) = U(t, x+1) = U(t, x)$  for all  $(t, x) \in \mathbb{R}^2$ .

Let  $x, y, s, t \in \mathbb{R}$ ,  $s < t$ . We say that the set of all absolutely continuous functions  $\xi: [s, t] \rightarrow \mathbb{R}$  such that  $\xi(s) = y$ ,  $\xi(t) = x$ , and  $L(\cdot, \xi(\cdot), \xi'(\cdot))$  is Lebesgue integrable, is the set of *admissible trajectories* (or *trajectories* for

---

\*Comm. Contemporary Math. — 1999. — V. 1, N. 4. — P. 517–533.

short) and denote it by  $\Omega(s, y; t, x)$ . The functional defined on  $\Omega(s, y; t, x)$  by the formula

$$\mathcal{L}(s, y; t, x)[\xi] = \int_s^t L(\tau, \xi(\tau), \xi'(\tau)) d\tau = \int_s^t [L_0(\xi'(\tau)) - U(\tau, \xi(\tau))] d\tau$$

is called the *action functional* associated with the Lagrangian  $L(t, x, v)$ . It follows from conditions  $(L_1)$ – $(L_4)$  that this functional is bounded from below. The infimum of  $\mathcal{L}(s, y; t, x)$  over  $\Omega(s, y; t, x)$  is called the *value function* of the action functional and is denoted by  $L(s, y; t, x)$ . It is well-known (see, e.g., [4]) that this infimum is attained, i.e., that there exists a trajectory  $\xi_0 \in \Omega(s, y; t, x)$  such that  $\mathcal{L}(s, y; t, x)[\xi_0] = L(s, y; t, x)$ . This trajectory is called a *minimizer* of the functional  $\mathcal{L}(s, y; t, x)$ .

The Aubry-Mather theory [1, 2, 9, 10] deals with a generalization of the notion of minimizer to the case when  $s$  and  $t$  are allowed to be infinite,  $-\infty \leq s < t \leq \infty$ . Since the potential  $U(t, x)$  is periodic in  $t$ , we may assume that  $s$  and  $t$  are integer or infinite and consider any trajectory  $\xi(t)$  only at integer moments of time:  $x_n = \xi(n)$ . Any part of a minimizer is a minimizer itself; therefore the minimization problem for the action functional can be discretized:

$$\sum_{k=s}^{t-1} L(k, x_k; k+1, x_{k+1}) \rightarrow \min.$$

It follows from periodicity of  $U(t, x)$  in  $t$  that  $L(k, y; k+1, x) = L(0, y; 1, x)$  for any  $k \in \mathbb{Z}$ ; denote  $L(0, y; 1, x)$  by  $L(y, x)$ .

If  $s = -\infty$  or  $t = \infty$ , then the above sum diverges. In this case the definition of minimizer is modified in the following way: a sequence  $\{x_n\}$ ,  $n \in \mathbb{Z}$ ,  $s \leq n \leq t$ , is said to be *L-minimal* if it has the following property: for all  $n_1, n_2 \in \mathbb{Z}$ ,  $s \leq n_1 < n_2 \leq t$ , and any set  $\{y_n\}$ ,  $n = n_1, n_1 + 1, \dots, n_2$ , such that  $y_{n_1} = x_{n_1}$  and  $y_{n_2} = x_{n_2}$

$$\sum_{n=n_1}^{n_2-1} L(x_n, x_{n+1}) \leq \sum_{n=n_1}^{n_2-1} L(y_n, y_{n+1}).$$

This definition is due to Aubry [1, 2]; similar concepts are known in statistical mechanics (ground states in lattice models, see, e.g., [12]) and differential geometry (geodesics of type A [11, 6]).

It is clear that the definition of *L-minimality* does not change if  $L(y, x)$  is substituted by  $L(y, x) + a(y - x)$  for any  $a \in \mathbb{R}$ . Consider an optimization problem of Bolza type

$$s(x_{-n}) + \sum_{k=-n}^{-1} (L(x_k, x_{k+1}) + a(x_k - x_{k+1})) \rightarrow \min, \quad x_0 = x. \quad (1)$$

It follows from Bellman's principle of optimality that for any solution  $\{x_{-n}, x_{-n+1}, \dots, x_0 = x\}$  of this problem the following inclusion holds:

$$x_{-k} \in \arg \min_{y \in \mathbb{R}} ((B_a^{n-k} s)(y) + L(y, x_{-k+1}) + a(y - x_{-k+1})), \quad (2)$$

where operator  $B_a$  is the so-called Bellman operator [8, 14]:

$$(B_a s)(x) = \min_{y \in \mathbb{R}} (s(y) + L(y, x) + a(y - x)).$$

It is clear that any solution of the problem (1) satisfies the condition of  $L$ -minimality.

Now suppose that a function  $s^a(x)$  satisfies the following functional equation:

$$(B_a s^a)(x) = \min_{y \in \mathbb{R}} (s^a(y) + L(y, x) + a(y - x)) = s^a(x) + \lambda. \quad (3)$$

Then any solution of the problem (1) can be extended as  $-n \rightarrow -\infty$  using formula (2) without violating the condition of  $L$ -minimality. Thus to establish the existence of two-sided infinite minimizers (corresponding to the case when  $s = -\infty$ ,  $t = \infty$ ) it is sufficient to show that the initial point  $x_0 = x$  can be chosen in such a way that it is possible to find the reverse extension  $\{x_1, x_2, \dots\}$  satisfying the condition of  $L$ -minimality. This program is carried out in sections 3 and 4 after proving a number of technical lemmas in section 2.

We stress that the functional equation (3), like many other formulas in calculus of variations and optimization, can be treated as linear over a special algebraic structure, an *idempotent semiring*  $\mathbb{R}_{\min}$  [8]. By definition,  $\mathbb{R}_{\min} = \{\mathbb{R} \cup \{\infty\}, \oplus, \odot\}$ , where operations  $\oplus$  and  $\odot$  are defined as follows:  $a \oplus b = \min\{a, b\}$  (in particular,  $a \oplus \infty = \infty \oplus a = a$ ) and  $a \odot b = a + b$  (in particular,  $a \odot \infty = \infty \odot a = \infty$ ). It can easily be checked that this structure satisfies axioms of an idempotent semiring with neutral elements  $\infty$  and  $0$  with respect to operations  $\oplus$  and  $\odot$ , respectively [8]. In this context, the Bellman operator assumes the form of a linear integral operator:

$$(B_a s)(x) = \min_{y \in \mathbb{R}} (s(y) + L(y, x) + a(y - x)) = \int_{y \in \mathbb{R}}^{\oplus} s(y) \odot (L(y, x) + a(y - x))$$

and the equation (3) defines its eigenfunction  $s^a(x)$  and eigenvalue  $\lambda(a)$  [8, 14]:

$$(B_a s^a)(x) = s^a(x) + \lambda(a) = (\lambda(a) \odot s^a)(x).$$

This approach can be used further to re-establish the main results of Aubry-Mather theory in an independent way; however, in this paper, we assume a reverse viewpoint and apply Aubry-Mather theory to study of idempotent eigenfunctions  $s^a(x)$ . One question that can be hard to resolve in the general setting of idempotent analysis is whether an eigenfunction corresponding to a given eigenvalue is unique. In section 5 we exploit minimality of Aubry-Mather set in the sense of topological dynamics to prove that in the case of irrational rotation number the eigenfunction is determined uniquely up to an additive constant. This result is sharp in the sense that if the rotation number is rational, then the solution may not be unique; a counterexample is constructed in [7].

This work was motivated by the preprint [7], in which the problem was stated in terms of periodic solution to periodically forced Burgers equation rather than

eigenfunction of Bellman operator. After the work was finished [13], the author learned that in the context of the forced Burgers equation similar results, including uniqueness theorem, were obtained independently by Weinan E [5].

The author is deeply grateful to Ya. G. Sinai for setting of the problem and constant attention to this work and to Weinan E for the opportunity to read the unpublished version of his paper.

## 2 Properties of the value function

The following result is classical in calculus of variations (see, e.g., [4]):

**Lemma 1** *Suppose  $L_0(v)$  and  $U(t, x)$  satisfy assumptions  $(L_1)$ – $(L_4)$  and  $x, y, s, t \in \mathbb{R}$ ,  $s < t$ ; then there exists a minimizer  $\xi_0 \in \Omega(s, y; t, x)$  such that  $\mathcal{L}(s, y; t, x)[\xi_0] = L(s, y; t, x)$ . If  $\xi_0 \in \Omega(s, y; t, x)$  is a minimizer, then  $\xi_0$  is of class  $C^1$  in  $[s, t]$  and there exists a  $C^1$  function  $\lambda: [s, t] \rightarrow \mathbb{R}$  such that*

$$\lambda(\tau) = L'_0(\xi'_0(\tau)), \quad \lambda'(\tau) = -\frac{\partial U(\tau, \xi_0(\tau))}{\partial \xi_0}, \quad s < \tau < t. \quad (4)$$

*For all  $R, s, t \in \mathbb{R}$  such that  $R > 0$ ,  $s < t$  and all  $x, y \in \mathbb{R}$  such that  $|x - y| \leq R(t - s)$  there exists a positive constant  $C_1 = C_1(R, s, t)$  such that if  $\xi_0 \in \Omega(s, y; t, x)$  is a minimizer, then  $|\xi'_0(\tau)| \leq C_1$ ,  $s \leq \tau \leq t$ .*

**Lemma 2** *The function  $L(s, y; t, x)$  is diagonally periodic:*

$$L(s, y + 1; t, x + 1) = L(s + 1, y; t + 1, x) = L(s, y; t, x).$$

This lemma follows from periodicity of  $U(t, x)$  in  $t$  and  $x$ .

**Lemma 3** *For all  $r, s, t, x, y \in \mathbb{R}$ ,  $s < r < t$ ,*

$$L(s, y; t, x) = \min_{z \in \mathbb{R}} (L(s, y; r, z) + L(r, z; t, x)). \quad (5)$$

This is a variant of the well-known Bellman's principle of optimality.

**Lemma 4** *The function  $L(s, y; t, x)$  satisfies inequalities*

$$-M \leq \frac{1}{t-s} L(s, y; t, x) - L_0\left(\frac{x-y}{t-s}\right) \leq -m, \quad (6)$$

*where  $m = \min_{(t,x) \in \mathbb{R}^2} U(t, x)$ ,  $M = \max_{(t,x) \in \mathbb{R}^2} U(t, x)$ .*

PROOF. Consider the functional

$$\widehat{\mathcal{L}}(s, y; t, x)[\xi] = \int_s^t L_0(\xi'(\tau)) d\tau.$$

The trajectory  $\widehat{\xi}(\tau) = x(\tau-s)/(t-s) + y(t-\tau)/(t-s)$ ,  $s \leq \tau \leq t$ , is a minimizer of this functional in the set  $\Omega(s, y; t, x)$  and

$$\widehat{\mathcal{L}}(s, y; t, x)[\widehat{\xi}] = (t-s)L_0\left(\frac{x-y}{t-s}\right).$$

Let  $\xi_0$  be a minimizer of the functional  $\mathcal{L}(s, y; t, x)$  in the class  $\Omega(s, y; t, x)$ , i.e.,  $\mathcal{L}(s, y; t, x)[\xi_0] = L(s, y; t, x)$ . We have

$$\begin{aligned} L(s, y; t, x) - (t-s)L_0\left(\frac{x-y}{t-s}\right) &= \mathcal{L}(s, y; t, x)[\xi_0] - \widehat{\mathcal{L}}(s, y; t, x)[\widehat{\xi}] = \\ &= - \int_s^t U(\tau, \xi_0(\tau)) d\tau + (\widehat{\mathcal{L}}(s, y; t, x)[\xi_0] - \widehat{\mathcal{L}}(s, y; t, x)[\widehat{\xi}]) \geq \\ &\geq (t-s) \left( - \max_{(t,x) \in \mathbb{R}^2} U(t, x) \right) \end{aligned}$$

and

$$\begin{aligned} L(s, y; t, x) - (t-s)L_0\left(\frac{x-y}{t-s}\right) &= \mathcal{L}(s, y; t, x)[\xi_0] - \widehat{\mathcal{L}}(s, y; t, x)[\widehat{\xi}] \leq \\ &\leq \mathcal{L}(s, y; t, x)[\widehat{\xi}] - \widehat{\mathcal{L}}(s, y; t, x)[\widehat{\xi}] = \\ &= - \int_s^t U(\tau, \widehat{\xi}(\tau)) d\tau \leq (t-s) \left( - \min_{(t,x) \in \mathbb{R}^2} U(t, x) \right). \end{aligned}$$

Multiplying these inequalities by  $(t-s)^{-1}$ , we obtain (6).  $\square$

**Lemma 5** *The function  $L(s, y; t, x)$  is locally Lipschitzian: if  $R, x_1, x_2, y_1, y_2, s, t \in \mathbb{R}$ ,  $s < t$ ,  $R > 0$ ,  $|x_i - y_j| \leq R(t-s)$ ,  $i, j = 1, 2$ , then there exists  $C = C(R, s, t) \in \mathbb{R}$ ,  $C > 0$ , such that*

$$|L(s, y_1; t, x_1) - L(s, y_2; t, x_2)| \leq C(|y_1 - y_2| + |x_1 - x_2|). \quad (7)$$

PROOF. It is sufficient to prove the inequality

$$|L(s, y; t, x_1) - L(s, y; t, x_2)| \leq C|x_1 - x_2|, \quad (8)$$

where  $|x_i - y| \leq R(t-s)$ ,  $i = 1, 2$ . The inequality for the first pair of arguments of the function  $L$  is proved similarly; combining these two inequalities, we obtain (7).

Let  $\xi_0 \in \Omega(s, y; t, x_1)$  be a minimizer of the functional  $\mathcal{L}(s, y; t, x_1)$ . Consider the trajectory  $\xi(\tau) = \xi_0(\tau) + (x_2 - x_1)(\tau - s)/(t - s)$ ,  $s \leq \tau \leq t$ . It is clear that  $\xi \in \Omega(s, y; t, x_2)$ . We have

$$\begin{aligned} L(s, y; t, x_2) - L(s, y; t, x_1) &\leq \mathcal{L}(s, y; t, x_2)[\xi] - \mathcal{L}(s, y; t, x_1)[\xi_0] = \\ &= \int_s^t \left[ L_0 \left( \xi'_0(\tau) + \frac{x_2 - x_1}{t - s} \right) - L_0(\xi'_0(\tau)) \right] d\tau - \\ &\quad - \int_s^t \left[ U \left( \tau, \xi_0(\tau) + \frac{x_2 - x_1}{t - s}(\tau - s) \right) - U(\tau, \xi_0(\tau)) \right] d\tau. \end{aligned}$$

The functions  $U(t, x)$  and  $L_0(v)$  are of class  $C^1$ ; in addition, it follows from  $(L_3)$ ,  $(L_4)$  that  $|\partial U(t, x)/\partial x| \leq M_1$  for some positive  $M_1 \in \mathbb{R}$ . Hence,

$$\begin{aligned} L(s, y; t, x_2) - L(s, y; t, x_1) &\leq \\ &\leq \frac{|x_2 - x_1|}{t - s} (t - s) \max_{\tau \in [s, t], |\theta| \leq 1} \left| L'_0 \left( \xi'_0(\tau) + \theta \frac{|x_2 - x_1|}{t - s} \right) \right| + \\ &\quad + \frac{|x_2 - x_1|}{t - s} \frac{(t - s)^2}{2} M_1. \end{aligned}$$

By lemma 1, it follows that  $|\xi'_0(\tau)| \leq C_1 = C_1(R, s, t)$ ,  $s \leq \tau \leq t$ . Also, by  $(L_1)$ ,  $L'_0(v)$  is finite for all  $v$ ; thus for some  $C = C(M_1, R, s, t) \in \mathbb{R}$ ,  $C > 0$ ,

$$L(s, y; t, x_2) - L(s, y; t, x_1) \leq C|x_2 - x_1|.$$

Similarly,  $L(s, y; t, x_2) - L(s, y; t, x_1) \geq -C|x_2 - x_1|$ . This completes the proof of inequality (8).  $\square$

**Lemma 6** *Let  $x_1, x_2, y_1, y_2, s, r, t \in \mathbb{R}$ ,  $s < r < t$ ; suppose there exists  $z_0 \in \mathbb{R}$  such that for all  $z \in \mathbb{R}$*

$$L(s, y_i; r, z) + L(r, z; t, x_i) \geq L(s, y_i; r, z_0) + L(r, z_0; t, x_i), \quad i = 1, 2.$$

*Then either  $x_1 = x_2$  and  $y_1 = y_2$  or  $(x_1 - x_2)(y_1 - y_2) < 0$ .*

PROOF. For  $x_1 = x_2$  and  $y_1 = y_2$  there is nothing to prove. Assume  $x_1 \neq x_2$  or  $y_1 \neq y_2$ . Let  $\xi_i^< \in \Omega(s, y_i; r, z_0)$  and  $\xi_i^> \in \Omega(r, z_0; t, x_i)$  be minimizers of functionals  $\mathcal{L}(s, y_i; r, z_0)$  and  $\mathcal{L}(r, z_0; t, x_i)$ ,  $i = 1, 2$ . It follows from the condition of the lemma and Bellman's principle of optimality (lemma 3) that for  $i = 1, 2$

$$L(s, y_i; r, z_0) + L(r, z_0; t, x_i) = L(s, y_i; t, x_i).$$

Hence for  $i = 1, 2$  the trajectories

$$\xi_i(\tau) = \begin{cases} \xi_i^<(\tau), & s \leq \tau \leq r \\ \xi_i^>(\tau), & r \leq \tau \leq t \end{cases} \quad (\xi_i \in \Omega(s, y_i; t, x_i))$$

are minimizers of  $\mathcal{L}(s, y_i; t, x_i)$ . Using lemma 1, we see that the trajectories  $\xi_i$ ,  $i = 1, 2$ , are of class  $C^1$ . This means that  $(\xi_i^<)'(r - 0) = (\xi_i^>)'(r + 0) = \xi'_i(r)$ ,  $i = 1, 2$ .

We claim that  $\xi'_1(r) \neq \xi'_2(r)$ . Indeed, let  $\lambda_i$ ,  $i = 1, 2$ , be the  $C^1$  functions corresponding to trajectories  $\xi_i$  by lemma 1. Assume that  $\xi'_1(r) = \xi'_2(r)$ ; then, by continuity of  $L'_0(\cdot)$ , it follows from (4) that  $\lambda_1(r) = \lambda_2(r)$ . Consider the Cauchy problem for the system of ordinary differential equations (4) on the interval  $[s, t]$  with initial data  $\xi_0(r) = \xi_1(r) = \xi_2(r)$  and  $\lambda(r) = \lambda_1(r) = \lambda_2(r)$ . It follows from condition  $(L_3)$  that the function  $\partial U(t, x)/\partial x$  is Lipschitzian and its absolute value is bounded by the constant  $M_1 > 0$ ; thus it follows from the second equation (4) that  $|\lambda(\tau) - \lambda(r)| \leq M_1(t - s)$  for all  $\tau \in [s, t]$ . Hence it follows from condition  $(L_1)$  that the system

$$\xi'_0(\tau) = (L'_0)^{-1}(\lambda(\tau)), \quad \lambda'(\tau) = -\frac{\partial U(\tau, \xi_0(\tau))}{\partial \xi_0}, \quad s < \tau < t,$$

which is equivalent to (4), satisfies the conditions of the uniqueness theorem. Thus  $\xi_1(\tau) = \xi_2(\tau)$ ,  $s \leq \tau \leq t$ , and in particular  $x_1 = x_2$  and  $y_1 = y_2$ . This contradiction proves that  $\xi'_1(r) \neq \xi'_2(r)$ .

Now we prove that there exists  $\delta > 0$  such that  $s < r - \delta$ ,  $r + \delta < t$ , and if  $\tau \in (r - \delta, r + \delta)$ ,  $\tau \neq r$ , then  $\xi_1(\tau) \neq \xi_2(\tau)$ . Indeed, let a sequence  $\{t_n\}$  be such that  $s < t_n < t$ ,  $t_n \neq r$ ,  $\xi_1(t_n) = \xi_2(t_n)$  for all  $n$ , and  $\lim_{n \rightarrow \infty} t_n = r$ . This implies that

$$\xi'_1(r) = \lim_{n \rightarrow \infty} \frac{\xi_1(t_n) - \xi_1(r)}{t_n - r} = \lim_{n \rightarrow \infty} \frac{\xi_2(t_n) - \xi_2(r)}{t_n - r} = \xi'_2(r),$$

a contradiction.

Now assume that  $(x_1 - x_2)(y_1 - y_2) \geq 0$ ; to be precise, let  $x_1 \leq x_2$  and  $y_1 \leq y_2$ . For  $i = 1, 2$  define trajectories  $\bar{\xi}_i \in \Omega(s, y_i; t, x_i)$  by  $\bar{\xi}_1(\tau) = \min\{\xi_1(\tau), \xi_2(\tau)\}$  and  $\bar{\xi}_2(\tau) = \max\{\xi_1(\tau), \xi_2(\tau)\}$ . By the above each of  $\bar{\xi}_1$  and  $\bar{\xi}_2$  coincides with  $\xi_1$  and  $\xi_2$  on a finite number of intervals of finite length. Let us check that the trajectories  $\bar{\xi}_i$ ,  $i = 1, 2$ , are minimizers. Indeed, we have

$$\mathcal{L}(s, y_i; t, x_i)[\bar{\xi}_i] \geq \mathcal{L}(s, y_i; t, x_i)[\xi_i] = L(s, y_i; t, x_i), \quad i = 1, 2,$$

and

$$\begin{aligned} \mathcal{L}(s, y_1; t, x_1)[\bar{\xi}_1] + \mathcal{L}(s, y_2; t, x_2)[\bar{\xi}_2] &= \\ &= \mathcal{L}(s, y_1; t, x_1)[\xi_1] + \mathcal{L}(s, y_2; t, x_2)[\xi_2] = \\ &= L(s, y_1; t, x_1) + L(s, y_2; t, x_2). \end{aligned}$$

Therefore  $\mathcal{L}(s, y_i; t, x_i)[\bar{\xi}_i] = L(s, y_i; t, x_i)$ ,  $i = 1, 2$ .

By lemma 1, the trajectories  $\bar{\xi}_i$ ,  $i = 1, 2$ , are of class  $C^1$ . This implies that  $\bar{\xi}'_i(r - 0) = \bar{\xi}'_i(r + 0)$ ,  $i = 1, 2$ , where all one-sided derivatives exist. But this means that  $\xi'_1(r - 0) = \xi'_2(r + 0)$  and  $\xi'_2(r - 0) = \xi'_1(r + 0)$ , since  $r$  is the only intersection point of  $\xi_1$  and  $\xi_2$  in the interval  $(r - \delta, r + \delta)$ . Thus  $\xi'_1(r) = \xi'_2(r)$ . This contradiction concludes the proof.  $\square$

**Corollary** *Let  $x_1, x_2, y_1, y_2, s, t \in \mathbb{R}$ ,  $x_1 < x_2$ ,  $y_1 < y_2$ , and  $s < t$ . Then*

$$L(s, y_1; t, x_1) + L(s, y_2; t, x_2) < L(s, y_2; t, x_1) + L(s, y_1; t, x_2). \quad (9)$$

PROOF. Let  $\xi_1 \in \Omega(s, y_1; t, x_2)$  and  $\xi_2 \in \Omega(s, y_2; t, x_1)$  be minimizers of functionals  $\mathcal{L}(s, y_1; t, x_2)$  and  $\mathcal{L}(s, y_2; t, x_1)$ , respectively. Note that  $\xi_1(s) < \xi_2(s)$  and  $\xi_1(t) > \xi_2(t)$ ; hence there exist  $r$  and  $z_0$  such that  $s < r < t$ ,  $\xi_1(r) = \xi_2(r) = z_0$ . Therefore  $L(s, y_1; r, z_0) + L(r, z_0; t, x_2) = L(s, y_1; t, x_2)$ ,  $L(s, y_2; r, z_0) + L(r, z_0; t, x_1) = L(s, y_2; t, x_1)$  and by lemma 3  $L(s, y_1; r, z_0) + L(r, z_0; t, x_1) \geq L(s, y_1; t, x_1)$ ,  $L(s, y_2; r, z_0) + L(r, z_0; t, x_2) \geq L(s, y_2; t, x_2)$ . This implies that  $L(s, y_1; t, x_1) + L(s, y_2; t, x_2) \leq L(s, y_1; t, x_2) + L(s, y_2; t, x_1)$ . Assume that this inequality is actually an equality. This assumption means that for all  $z \in \mathbb{R}$

$$\begin{aligned} L(s, y_1; t, x_1) &= L(s, y_1; r, z_0) + L(r, z_0; t, x_1) \leq L(s, y_1; r, z) + L(r, z; t, x_1), \\ L(s, y_2; t, x_2) &= L(s, y_2; r, z_0) + L(r, z_0; t, x_2) \leq L(s, y_2; r, z) + L(r, z; t, x_2). \end{aligned}$$

It follows from the previous lemma that  $(x_1 - x_2)(y_1 - y_2) < 0$ . This contradiction concludes the proof.  $\square$

### 3 Existence of an eigenfunction of the Bellman operator

Let  $a \in \mathbb{R}$ ,  $s_0(x)$  be a continuous periodic function. We define

$$s_n(x; a, s_0) = (B_a^n s_0)(x) = \min_{y \in \mathbb{R}} (s_0(y) + L(0, y; n, x) + a(y - x)).$$

As a function of  $x$ ,  $s_n(x; a, s_0)$  is bounded from below. Further, it follows from diagonal periodicity of  $L(s, y; t, x)$  (lemma 2) that it is periodic; thus  $s_n(x; a, s_0) = \min_{k \in \mathbb{Z}} s_n(x - k; a, s_0)$ . Using this formula, lemma 2, and periodicity of  $s_0(y)$ , we get

$$s_n(x; a, s_0) = \min_{k \in \mathbb{Z}} \min_{y \in \mathbb{R}} (s_0(y) + L(0, y + k; n, x) + a(y + k - x)).$$

Thus for any  $n > 0$  and  $y, x \in \mathbb{R}$  the quantity  $L(0, y + k; n, x) + a(y + k - x)$  is bounded from below as a function of  $k \in \mathbb{Z}$ . We denote

$$L_n^a(y, x) = \min_{k \in \mathbb{Z}} (L(0, y + k; n, x) + a(y + k - x)), \quad (10)$$

where minimum is attained since  $L(s, y; t, x)$  is continuous and grows arbitrarily large as  $|x - y| \rightarrow \infty$ ; hence,

$$s_n(x; a, s_0) = \min_{y \in \mathbb{R}} (s_0(y) + L_n^a(y, x)). \quad (11)$$

**Lemma 7** *For any  $n = 1, 2, \dots$ , the function  $L_n^a(y, x)$  is periodic in both variables and Lipschitzian in  $x$ ; the Lipschitz constant  $C^*(a)$  does not depend on  $n$  and  $y \in \mathbb{R}$ . For any  $n_1 > 0$ ,  $n_2 > 0$*

$$L_{n_1+n_2}^a(y, x) = \min_{z \in \mathbb{R}} (L_{n_1}^a(y, z) + L_{n_2}^a(z, x)). \quad (12)$$

PROOF. Using the definition (10) and lemma 2, we obtain  $L_n^a(y+l, x+m) = L_n^a(y, x)$  for any integer  $l, m$ ; thus  $L_n^a(y, x)$  is periodic. Combining (10) for  $n = n_1 + n_2$  with Bellman's principle of optimality (5) for  $s = 0, r = n_1$ , and  $t = n_1 + n_2$  and using lemma 2, we get

$$L_{n_1+n_2}^a(y, x) = \min_{z \in \mathbb{R}} (L_{n_1}^a(y, z) + L(0, z; n_2, x) + a(z - x)).$$

Using periodicity of  $L_{n_1}^a(y, z)$  in  $z$ , we obtain (12).

Let us prove that  $L_1^a(y, x)$  is Lipschitzian. Since  $L_1^a(y, x)$  is a pointwise minimum of a family of continuous functions, it is upper semicontinuous. Denote by  $N$  the maximum value of  $L_1^a(y, x)$  on the set  $Q = \{(y, x) \mid 0 \leq y \leq 1, 0 \leq x \leq 1\}$ . Clearly, there exists  $R > 0$  such that  $L_0(x - y - k) + a(y + k - x) > N + \max_{(t,x) \in \mathbb{R}^2} U(t, x)$  for all  $(y, x) \in Q$  whenever  $|k| > R$ . Thus it follows from lemma 4 that for any  $(y, x) \in Q$

$$L_1^a(y, x) = \min_{k \in \mathbb{Z}, |k| \leq R} (L(0, y + k; 1, x) + a(y + k - x)).$$

On the other hand, using lemma 5, we see that all functions  $L(0, y + k; 1, x) + a(y + k - x)$ ,  $|k| < R$ , are Lipschitzian on  $Q$  with the constant  $C^*(a) = C(R + 1, 0, 1) + |a|$ . Thus  $L_1^a(y, x)$  is Lipschitzian on  $Q$  (and, by periodicity, on the whole  $\mathbb{R}^2$ ) with the same constant.

Now it follows from (12) that for any  $n \geq 2$

$$L_n^a(y, x) = \min_{z \in \mathbb{R}} (L_{n-1}^a(y, z) + L_1^a(z, x)),$$

that is for any  $y \in \mathbb{R}$   $L_n^a(y, x)$  as a function of  $x$  is a pointwise minimum of a family of functions having the same Lipschitz constant  $C^*(a)$ . Thus  $L_n^a(y, x)$  is Lipschitz continuous in  $x$  with the constant  $C^*(a)$  for all  $n = 1, 2, \dots$   $\square$

**Corollary** For any  $n = 1, 2, \dots$  the function  $s_n(x; a, s_0)$  defined in (11) is Lipschitzian with the constant  $C^*(a)$ ; it satisfies

$$\max_{x \in \mathbb{R}} s_n(x; a, s_0) - \min_{x \in \mathbb{R}} s_n(x; a, s_0) \leq C^*(a) \quad (13)$$

and for all  $m \in \mathbb{Z}, m \geq n$ ,

$$s_m(x; a, s_0) = \min_{y \in \mathbb{R}} (s_{m-n}(y; a, s_0) + L_n^a(y, x)). \quad (14)$$

PROOF. From (11) it follows that  $s_n(x; a, s_0)$  is a pointwise minimum of a family of Lipschitzian functions having the Lipschitz constant  $C^*(a)$ ; hence it is Lipschitzian with the same constant. Taking into account periodicity of  $s_n(x; a, s_0)$ , we obtain (13). Equation (14) follows from (12) for  $n_1 = m - n, n_2 = n$ .  $\square$

Denote

$$\lambda_n(a) = \min_{(y,x) \in \mathbb{R}^2} \frac{1}{n} L_n^a(y, x) = \min_{(y,x) \in \mathbb{R}^2} \left( \frac{1}{n} L(0, y; n, x) + a \frac{y - x}{n} \right). \quad (15)$$

**Lemma 8** For any  $a \in \mathbb{R}$  there exists  $\lambda(a) \in \mathbb{R}$  such that

$$|\lambda_n(a) - \lambda(a)| \leq \frac{C^*(a)}{n}, \quad (16)$$

where  $C^*(a)$  is the Lipschitz constant of the function  $L_n^a(y, x)$ . The function  $a \mapsto \lambda(a)$  is concave and satisfies inequalities

$$H_0(a) + m \leq -\lambda(a) \leq H_0(a) + M, \quad (17)$$

where  $m$  and  $M$  are defined in lemma 4 and  $H_0(a)$  is the Legendre transform of  $L_0(v)$ .

PROOF. Let  $s_0(x) = 0$ ,  $s_n(x) = s_n(x; a, s_0)$ . From (14), for any integer  $n$ ,  $n_0$ ,  $0 < n_0 < n$ ,

$$s_n(x) = \min_{y \in \mathbb{R}} (s_{n-n_0}(y) + L_{n_0}^a(y, x)). \quad (18)$$

We see that  $\min_{x \in \mathbb{R}} s_n(x) = \min_{(y,x) \in \mathbb{R}^2} L_n^a(y, x) = n\lambda_n(a)$  for any  $n = 1, 2, \dots$ . It follows from the corollary to lemma 7 that  $n\lambda_n(a) \leq s_n(x) \leq n\lambda_n(a) + C^*(a)$ . Combining these inequalities with (18) and (15), we obtain

$$\begin{aligned} n\lambda_n(a) &\leq s_n(x) \leq (n - n_0)\lambda_{n-n_0}(a) + C^*(a) + n_0\lambda_{n_0}(a), \\ (n - n_0)\lambda_{n-n_0}(a) + n_0\lambda_{n_0}(a) &\leq s_n(x) \leq n\lambda_n(a) + C^*(a). \end{aligned}$$

Hence,

$$|n\lambda_n(a) - n_0\lambda_{n_0}(a) - (n - n_0)\lambda_{n-n_0}(a)| \leq C^*(a).$$

There exist integer  $p \geq 1$  and  $0 \leq q < n$  such that  $n = pn_0 + q$ . By induction over  $p$  it is easily checked that  $|n\lambda_n(a) - pn_0\lambda_{n_0}(a) - q\lambda_q(a)| \leq pC^*(a)$ , that is

$$\left| \lambda_n(a) - \lambda_{n_0}(a) + \frac{q}{n}(\lambda_{n_0}(a) - \lambda_q(a)) \right| \leq \frac{p}{n}C^*(a) \leq \frac{1}{n_0}C^*(a). \quad (19)$$

Let  $\varepsilon > 0$  be arbitrarily small,  $n_0$  be so large that the right-hand side of this inequality is less than  $\varepsilon$ , and  $N$  be such that  $(1/N) \max_{1 \leq q \leq n_0} |q(\lambda_{n_0}(a) - \lambda_q(a))| < \varepsilon$ ; then  $|\lambda_{n_1}(a) - \lambda_{n_2}(a)| < 4\varepsilon$  for all  $n_1, n_2 \geq N$ . Thus  $\{\lambda_n(a)\}$ ,  $n = 1, 2, \dots$ , is a Cauchy sequence. Denote its limit by  $\lambda(a)$ ; then we get (16) from (19) in the limit  $n \rightarrow \infty$ .

Suppose  $a_1, a_2 \in \mathbb{R}$ ,  $0 < \alpha, \beta < 1$ ,  $\alpha + \beta = 1$ . Using definition of  $\lambda_n(a)$  (15), we obtain

$$\begin{aligned} \lambda_n(\alpha a_1 + \beta a_2) &= \frac{1}{n} \min_{(y,x) \in \mathbb{R}^2} (L(0, y; n, x) + (\alpha a_1 + \beta a_2)(y - x)) = \\ &= \frac{1}{n} \min_{(y,x) \in \mathbb{R}^2} [\alpha(L(0, y; n, x) + a_1(y - x)) + \beta(L(0, y; n, x) + a_2(y - x))] \geq \\ &\geq \alpha\lambda_n(a_1) + \beta\lambda_n(a_2). \end{aligned}$$

In the limit  $n \rightarrow \infty$  this implies that the function  $\lambda(a)$  is concave.

Finally, lemma 4 implies that  $-M \leq (1/n)L(0, y; n, x) - L_0((x-y)/n) \leq -m$ . Hence,

$$m + a \frac{x-y}{n} - L_0\left(\frac{x-y}{n}\right) \leq a \frac{x-y}{n} - \frac{1}{n}L(0, y; n, x) \leq M + a \frac{x-y}{n} - L_0\left(\frac{x-y}{n}\right).$$

Taking max over  $(y, x) \in \mathbb{R}^2$ , using (15), and denoting  $(x-y)/n$  by  $v$ , we obtain

$$m + \max_{v \in \mathbb{R}} (av - L_0(v)) \leq -\lambda_n(a) \leq M + \max_{v \in \mathbb{R}} (av - L_0(v)).$$

But by a well-known formula for the Legendre transform  $\max_{v \in \mathbb{R}} (av - L_0(v)) = H_0(a)$ . Thus we obtain (17) in the limit  $n \rightarrow \infty$ .  $\square$

**Theorem 1** *Suppose  $a \in \mathbb{R}$  and  $s_0(x)$  is a continuous and periodic function; then there exists*

$$\liminf_{n \rightarrow \infty} (s_n(x; a, s_0) - n\lambda(a)) = s^a(x) \quad (20)$$

and  $s^a(x)$  satisfies functional equation (3).

**Proof.** Using definitions of  $s_n(x; a, s_0)$  and  $\lambda_n(a)$ , we get

$$\min_{y \in \mathbb{R}} s_0(y) + n\lambda_n(a) \leq s_n(x; a, s_0) \leq \max_{y \in \mathbb{R}} s_0(y) + n\lambda_n(a).$$

Subtracting  $n\lambda(a)$  and using (16), we obtain

$$\min_{y \in \mathbb{R}} s_0(y) - C^*(a) \leq s_n(x; a, s_0) - n\lambda(a) \leq \max_{y \in \mathbb{R}} s_0(y) + C^*(a).$$

Let

$$\bar{s}_n(x) = \inf_{m \geq n} (s_m(x; a, s_0) - m\lambda(a)), \quad n = 1, 2, \dots$$

For any  $x \in \mathbb{R}$  the sequence  $\{\bar{s}_n(x)\}$  is nondecreasing; we see that it is bounded. Further, the corollary to lemma 7 implies that all functions  $\bar{s}_n(x)$  are Lipschitzian with the constant  $C^*(a)$ ; hence, this sequence is equicontinuous. It follows that there exists

$$\lim_{n \rightarrow \infty} \bar{s}_n(x) = \liminf_{n \rightarrow \infty} (s_n(x; a, s_0) - n\lambda(a)) = s^a(x)$$

and  $s^a(x)$  is periodic and continuous.

Let us check that  $s^a(x)$  satisfies functional equation (3). We have

$$\begin{aligned} \min_{y \in \mathbb{R}} (\bar{s}_n(y) + L_1^a(y, x)) - \lambda(a) &= \\ &= \min_{y \in \mathbb{R}} (\inf_{m \geq n} (\min_{z \in \mathbb{R}} (s_0(z) + L_m^a(z, y) + L_1^a(y, x) - (m+1)\lambda(a)))). \end{aligned}$$

But it follows from the definition of  $\lambda_n(a)$  and lemma 8 that  $L_m^a(z, y) - m\lambda(a) \geq -C^*(a)$  for all  $m = 1, 2, \dots$ . Therefore we can take minimum over  $y \in \mathbb{R}$  before infimum over  $m \geq n$  and obtain

$$\min_{y \in \mathbb{R}} (\bar{s}_n(y) + L_1^a(y, x)) - \lambda(a) = \bar{s}_{n+1}(x).$$

Taking into account (10) and passing to the limit  $n \rightarrow \infty$ , we see that  $s^a(x)$  satisfies functional equation (3).  $\square$

## 4 Existence of two-sided minimizers

Suppose  $a \in \mathbb{R}$ ,  $s^a(x)$  is a continuous periodic function satisfying functional equation (3). Denote  $L_a(y, x) = L(0, y; 1, x) + a(y - x) - \lambda(a)$ . Let  $Y^a(x)$  be the many-valued map of  $\mathbb{R}$  to  $\mathbb{R}$  defined as

$$Y^a(x) = \arg \min_{y \in \mathbb{R}} (s^a(y) + L_a(y, x)). \quad (21)$$

It is readily seen that if  $x, y \in \mathbb{R}$ , then  $y \in Y^a(x)$  iff

$$s^a(y) + L_a(y, x) = s^a(x) = \min_{z \in \mathbb{R}} (s^a(z) + L_a(z, x)). \quad (22)$$

For any two sets  $X \subset \mathbb{R}$ ,  $Y \subset \mathbb{R}$ , denote by  $X + Y$  the set  $\{z \in \mathbb{R} \mid z = x + y, x \in X, y \in Y\}$ . If  $Y = \{y\}$ , we denote  $X + \{y\}$  by  $X + y$ .

**Lemma 9** *For all  $x \in \mathbb{R}$*

$$Y^a(x + 1) = Y^a(x) + 1. \quad (23)$$

*If there exists  $x_0 \in \mathbb{R}$  such that  $x \in Y^a(x_0)$ , then the set  $Y^a(x)$  contains only one point. If  $x_1, x_2, y_1, y_2 \in \mathbb{R}$ ,  $x_1 < x_2$ ,  $y_1 \in Y^a(x_1)$ , and  $y_2 \in Y^a(x_2)$ , then  $y_1 < y_2$ .*

PROOF. Equation (23) follows from periodicity of  $s^a(x)$  and diagonal periodicity of  $L_a(y, x)$  (lemma 2).

Suppose  $x_1 < x_2$  and  $y_i \in Y^a(x_i)$ ,  $i = 1, 2$ . By (22), it follows that

$$\begin{aligned} s^a(y_1) + L(0, y_1; 1, x_1) + ay_1 &\leq s^a(y_2) + L(0, y_2; 1, x_1) + ay_2, \\ s^a(y_2) + L(0, y_2; 1, x_2) + ay_2 &\leq s^a(y_1) + L(0, y_1; 1, x_2) + ay_1, \end{aligned}$$

that is

$$L(0, y_1; 1, x_1) + L(0, y_2; 1, x_2) \leq L(0, y_2; 1, x_1) + L(0, y_1; 1, x_2).$$

Thus it follows from the corollary to lemma 6 that  $y_1 \leq y_2$ .

Let us check that  $y_1 < y_2$ . Assume the converse; let  $y = y_1 = y_2$ . Take any  $y_0 \in Y^a(y)$ . By (22), it follows that

$$s^a(x_i) = s^a(y_0) + L(0, y_0; 1, y) + L(1, y; 2, x_i) + a(y_0 - x_i) - 2\lambda(a), \quad i = 1, 2.$$

On the other hand, using (3), we get

$$\begin{aligned} s^a(x_i) &= \min_{z \in \mathbb{R}} \min_{t \in \mathbb{R}} (s^a(t) + L(0, t; 1, z) + L(1, z; 2, x_i) + a(t - x_i) - 2\lambda(a)) \leq \\ &\leq \min_{z \in \mathbb{R}} (s^a(y_0) + L(0, y_0; 1, z) + L(1, z; 2, x_i) + a(y_0 - x_i) - 2\lambda(a)), \\ &i = 1, 2. \end{aligned}$$

Hence it follows from lemma 6 that  $x_1 = x_2$ ; this contradiction proves that  $y_1 < y_2$ .

Finally, suppose  $x \in Y^a(x_0)$  for some  $x_0 \in \mathbb{R}$  and  $y_1, y_2 \in Y^a(x)$ . By a similar argument, we see that  $y_1 = y_2$ . Thus  $Y^a(x)$  consists of a single point.  $\square$

**Lemma 10** *Suppose  $M_0 = \mathbb{R}$ ,  $M_n = Y^a(M_{n-1})$ ,  $n = 1, 2, \dots$ ; then  $M_0 \supset M_1 \supset M_2 \supset \dots$  and there exists a nonempty closed set  $M^a$  such that (i)  $M^a = \bigcap_{n=0}^{\infty} M_n$ ; (ii) for any open  $U \subset \mathbb{R}$  such that  $M^a \subset U + \mathbb{Z}$  there exists  $N > 0$  with the following property: if the sequence  $\{y_n\}$  is such that  $y_{n+1} \in Y^a(y_n)$ ,  $n = 0, 1, \dots$ , then  $y_n \in U + \mathbb{Z}$  for all  $n \geq N$ ; (iii) the restriction  $Y^a|_{M^a}$  of  $Y^a$  to  $M^a$  is a one-to-one continuous mapping with continuous inverse.*

PROOF. Evidently, all  $M_n \neq \emptyset$  and  $M_0 = \mathbb{R}$  contains  $M_1$ . Suppose  $M_n \subset M_{n-1}$  for some  $n \geq 1$ ,  $x \in M_n$ ; then  $Y^a(x) \subset Y^a(M_n) \subset Y^a(M_{n-1}) = M_n$ . Hence  $M_{n+1} = Y^a(M_n) \subset M_n$ .

Let us show that all  $M_n$ ,  $n = 1, 2, \dots$ , are closed. Note that  $M_0 = \mathbb{R}$  is closed. Suppose  $M_n$  is closed and the sequence  $y_k \in M_{n+1}$ ,  $k = 1, 2, \dots$ , converges to a limit  $\bar{y} \in \mathbb{R}$ . It follows from lemma 9 that for any  $k$  there exists a unique  $x_k = (Y^a)^{-1}(y_k) \in M_n$  and all  $x_k$  are contained in a bounded interval. Thus there exists a subsequence  $\{x_{k_l}\}$  that converges to a limit  $\bar{x} \in M_n$ . Using (22), we get

$$s^a(y_{k_l}) + L_a(y_{k_l}, x_{k_l}) = s^a(x_{k_l}).$$

Since the functions  $s^a(x)$  and  $L_a(y, x)$  are continuous, we obtain

$$s^a(\bar{y}) + L_a(\bar{y}, \bar{x}) = s^a(\bar{x}).$$

Now it follows from (22) that  $\bar{y} \in Y^a(\bar{x})$ . Hence  $\bar{y} \in Y^a(M_n) = M_{n+1}$ , that is  $M_{n+1}$  is closed.

Consider the topology  $\mathcal{T}$  on  $\mathbb{R}$  such that  $V \in \mathcal{T}$  iff  $V = U + \mathbb{Z}$ , where  $U$  is open in the usual sense. Note that  $\mathbb{R}$  is compact in this topology. It follows from (23) that complements  $U_n$  of sets  $M_n$  are open in the topology  $\mathcal{T}$ . We see that  $U_n \subset U_{n+1}$ .

Clearly, the set  $M^a = \bigcap_{n=0}^{\infty} M_n$  is closed. Let us check that it is nonempty. Assume the converse; hence sets  $U_n$ ,  $n = 1, 2, \dots$  cover all  $\mathbb{R}$ . Since  $\mathbb{R}$  is compact in the topology  $\mathcal{T}$ , we see that there exists  $N > 0$  such that  $\mathbb{R} = \bigcup_{n=0}^N U_n = U_N$ . Thus  $M_N = \emptyset$ ; this contradiction proves that  $M^a$  is nonempty.

Now let  $V \in \mathcal{T}$  be such that  $M^a \subset V$ . Arguing as above, we see that there exists  $N > 0$  such that  $\mathbb{R} = V \cup U_N$ . On the other hand, for any sequence  $\{y_n\}$  such that  $y_{n+1} \in Y^a(y_n)$ ,  $n = 0, 1, \dots$ , it follows that  $y_n \in M_n$ . Thus for all  $n \geq N$  we obtain  $y_n \in V$ .

Denote by  $Y^a|_{M_n}: M_n \rightarrow M_{n+1}$  the restriction of  $Y^a$  to  $M_n$ ,  $n = 1, 2, \dots$ . It follows from lemma 9 that  $Y^a|_{M_n}$  is single-valued, strictly increasing as a function  $\mathbb{R} \rightarrow \mathbb{R}$ , and bijective. We claim that it is a homeomorphism. Indeed, let sequences  $\{x_k\} \subset M_n$  and  $\{y_k\} \subset M_{n+1}$  be such that  $Y^a(x_k) = y_k$  for all  $k$ . If  $y_k$  converge to  $\bar{y}$ , then it follows from strict monotonicity of  $Y^a|_{M_n}$  that  $x_k$  converge to a unique  $\bar{x}$ ; hence the map  $Y^a|_{M_n}$  has a continuous inverse. Further, if  $x_k$  converge to  $\bar{x}$ , then the set  $\{y_k\}$  is bounded. Let  $\bar{y}_1, \bar{y}_2$ ,  $y_1 \neq y_2$ , be two limit points of  $\{y_k\}$ ; then we see that  $\{\bar{y}_1, \bar{y}_2\} \subset Y^a(\bar{x})$ , since  $s^a(x)$  and  $L_a(y, x)$  are continuous. But the map  $Y^a|_{M_n}$  is single-valued; this contradiction proves that it is continuous.

By the above  $M^a$  is a closed subset of  $M_n$  for all  $n = 1, 2, \dots$ ; in addition,  $Y^a(M^a) = M^a$ . It follows that the restriction of  $Y^a|_{M_n}$  to  $M^a$  is a homeomorphism.  $\square$

We say that  $M^a$  is the *invariant set* of the many-valued map  $Y^a(x)$ .

Combining the previous lemma with the observation that sequence  $\{x_n\}$  that satisfies (2) is  $L$ -minimal, we obtain the following

**Theorem 2** *Suppose  $a \in \mathbb{R}$ ,  $s^a(x)$  is a continuous periodic function satisfying functional equation (3),  $Y^a(x)$  is the corresponding many-valued map,  $M^a$  is its invariant set, and  $x \in M^a$ . Then the two-sided sequence  $x_k = (Y^a)^{-k}(x)$ ,  $k \in \mathbb{Z}$ , is  $L$ -minimal.*

## 5 Uniqueness of an eigenfunction

To give the uniqueness condition for eigenfunction of the Bellman operator we need the following results of Aubry-Mather theory:

**Lemma 11** *Suppose a sequence  $\{x_n\}$ ,  $n \in \mathbb{Z}$ , is  $L$ -minimal; then there exists*

$$\lim_{n \rightarrow \pm\infty} \frac{x_n - x_0}{n} = \omega(\{x_n\}) \in \mathbb{R}. \quad (24)$$

*If  $\omega = \omega(\{x_n\})$  is irrational, then there exists a function  $\phi_\omega(t): \mathbb{R} \rightarrow \mathbb{R}$  with the following properties: (i) it is continuous on the right and  $\phi_\omega(t_1) < \phi_\omega(t_2)$  iff  $t_1 < t_2$ ; (ii)  $\phi_\omega(t+1) = \phi_\omega(t) + 1$  for all  $t \in \mathbb{R}$ ; (iii) if  $x_0 = \phi_\omega(t_0 \pm 0)$  then the sequence  $\{x_n\}$  defined by  $x_n = \phi_\omega(t_0 + n\omega \pm 0)$  is  $L$ -minimal; (iv) for any neighborhood  $V \in \mathcal{T}$  of the closure of the image  $\phi_\omega(\mathbb{R})$  and any  $L$ -minimal sequence  $\{x_n\}$ ,  $x_n \in V$  as soon as  $|n|$  is large enough.*

For the proof see, e.g., [10, sections 9, 11, and 12]. The number  $\omega(\{x_k\})$  is called a *rotation number* of the  $L$ -minimal sequence  $\{x_k\}$ .

**Theorem 3** *Let  $s^a(x)$  be the continuous periodic function satisfying functional equation (3),  $Y^a(x)$  be the corresponding many-valued map and  $M^a$  be its invariant set. Then the value of  $\omega$  for any  $L$ -minimal sequence of the form  $x_n = (Y^a)^{-k}(x)$  depends only on the parameter  $a$  but not on the choice of  $s^a(x)$ ; furthermore, if  $\omega(a)$  is irrational, then  $s^a(x)$  is determined uniquely up to an additive constant.*

**PROOF.** Let  $Y^a(x)$  be a many-valued map corresponding to a continuous periodic function  $s^a(x)$  satisfying functional equation (3) and  $M^a$  be its invariant set. The corresponding rotation number  $\omega$  is determined uniquely. Indeed, let  $x_k = (Y^a)^{-k}(x)$  and  $y_k = (Y^a)^{-k}(y)$ ,  $k \in \mathbb{Z}$ , for some  $x, y \in M^a$ ; we are going to demonstrate that  $\omega(\{x_k\}) = \omega(\{y_k\})$ . Take  $p \in \mathbb{Z}$  such that  $x \leq y+p \leq x+1$ . By lemma 9, it follows that  $(Y^a)^{-k}(x) \leq (Y^a)^{-k}(y+p) \leq (Y^a)^{-k}(x+1)$  for all  $k \in \mathbb{Z}$ . Using (24), we obtain  $\omega(\{x_k\}) = \omega(\{y_k\})$ .

A subset  $M \subset M^a$  is called a *minimal set* of the (single-valued) map  $Y^a|_{M^a}$  if  $M$  is closed,  $Y^a(M) = M$ , and  $M$  contains no other nonempty subset with

the same properties. Let  $x \in M^a$ ; the sequence  $\{(Y^a)^k(x)\}$ ,  $k = 1, 2, \dots$ , is called *the orbit* of the point  $x$ . It can be proved that a set  $M \subset M^a$  is minimal if and only if the orbit of any point  $x \in M$  is dense in  $M$  in topology  $\mathcal{T}$  defined in the proof of lemma 10.

The set  $M^a$  is compact in topology  $\mathcal{T}$ ; using (23), we see that  $Y^a$  is a homeomorphism of  $M^a$  in this topology. By the well-known theorem of Birkhoff [3], it follows that  $M^a$  contains at least one minimal set  $M$  of the map  $Y^a$ .

Suppose  $s_1^a(x)$  is another continuous periodic function satisfying functional equation (3); then for all  $x, y \in \mathbb{R}$  such that  $y \in Y^a(x)$

$$s^a(x) - s_1^a(x) \geq s^a(y) - s_1^a(y).$$

Indeed, it follows from (22) that  $s^a(x) = s^a(y) + L_a(y, x)$ . Using (11), we obtain

$$\begin{aligned} s^a(x) - s_1^a(x) &= s^a(y) + L_a(y, x) - s_1^a(x) = \\ &= s_1^a(y) + L_a(y, x) - s_1^a(x) + s^a(y) - s_1^a(y) \geq \\ &\geq s^a(y) - s_1^a(y). \end{aligned}$$

Clearly,  $s^a(x)$  and  $s_1^a(x)$  are continuous in topology  $\mathcal{T}$ . Let  $x_0 \in M$  be a point where  $s^a(x) - s_1^a(x)$  attains its minimal value in  $M$  and  $\{x_k\}$ ,  $k = 1, 2, \dots$ , be its orbit. By the above,

$$s^a(x_0) - s_1^a(x_0) \leq s^a(x_k) - s_1^a(x_k) \leq s^a(x_0) - s_1^a(x_0).$$

Since the closure of  $\{x_k\}$  is the set  $M$ , we see that on  $M$   $s^a(x) - s_1^a(x)$  is constant. Thus it follows from (22) that on the set  $M$   $Y^a(x)$  coincides with the map  $Y_1^a(x)$  corresponding to  $s_1^a(x)$ . In particular, the rotation number  $\omega$  depends only on  $a$ . We see also that if  $M$  is a minimal set of the map corresponding to some  $s^a(x)$  satisfying functional equation (3), then it is a minimal set of any other continuous periodic function satisfying functional equation (3) with the same  $a$ .

Now suppose  $\omega = \omega(a)$  is irrational. It follows from lemma 11 (iv) that the closure  $\bar{M}$  of the set  $\phi_\omega(\mathbb{R})$  contains limit points with respect to topology  $\mathcal{T}$  of the closed set  $M^a$ , so  $\bar{M} \cap M^a$  is nonempty. Now if  $x_0 \in M^a$  and there exists such  $t_0$  that  $x_0 = \phi_\omega(t_0 \pm 0) \in \bar{M}$ , then by lemma 11 (iii) the closure of the orbit of the point  $x_0$  with respect to topology  $\mathcal{T}$  coincides with  $\bar{M}$ , so  $\bar{M} \subseteq M^a$ . Thus statement (iv) of lemma 11 means that  $\bar{M}$  is the only minimal invariant subset of  $M^a$  not depending on the particular choice of an eigenfunction  $s^a(x)$ .

Finally let us prove that if  $\omega = \omega(a)$  is irrational, then  $s^a(x) - s_1^a(x)$  is constant for all  $x \in \mathbb{R}$ . Without loss of generality it can be assumed that  $s^a(x) - s_1^a(x) = 0$  on  $\bar{M}$ . Let  $x_0 \in \mathbb{R}$  be a point where  $s^a(x) - s_1^a(x)$  attains its minimum in  $M^a$  and  $\{x_k\}$ ,  $k = 1, 2, \dots$ , be its orbit. Arguing as above, we see that  $s^a(x_k) - s_1^a(x_k) = s^a(x_0) - s_1^a(x_0)$  for all  $k \geq 1$ . On the other hand, it follows from lemma 11 (iv) that for any neighborhood  $V \in \mathcal{T}$  of the set  $\bar{M}$  there exists  $N > 0$  such that  $x_k \in V$  for all  $k > N$ . Using continuity of the functions  $s^a(x)$  and  $s_1^a(x)$ , we see that  $s^a(x_0) - s_1^a(x_0) = 0$ . By definition of  $x_0$ , it follows that  $s^a(x) \geq s_1^a(x)$  for all  $x \in M^a$ . Likewise,  $s^a(x) \leq s_1^a(x)$ ; thus  $s^a(x) = s_1^a(x)$  on  $M^a$ . A similar argument based on lemma 10 (ii) shows that  $s^a(x) = s_1^a(x)$  for all  $x \in \mathbb{R}$ .  $\square$

## References

- [1] Aubry S., *The twist map, the extended Frenkel-Kontorova model and the devil's staircase*, Physica D **7** (1983), 240–258.
- [2] Aubry S., LeDaeron P. Y., *The discrete Frenkel-Kontorova model and its extensions I: Exact results for the ground states*, Physica D **8** (1983), 381–422.
- [3] Birkhoff G. D., *Quelques théorèmes sur les mouvements des systèmes dynamiques*, Bull. Soc. Math. de France, **40** (1912).
- [4] Cesari L., *Optimization—theory and applications: Problems with ordinary differential equations*, Springer-Verlag, 1983.
- [5] E Weinan, *Aubry-Mather theory and periodic solutions of the forced Burgers equation*, preprint (1998).
- [6] Hedlund G. A., *Geodesics on two-dimensional Riemann manifold with periodic coefficients*, Ann. of Math. **33** (1932), 719–739.
- [7] Jauslin H. R., Kreiss H. O., Moser J., *On the forced Burgers equation with periodic boundary conditions*, preprint (1997).
- [8] Kolokol'tsov V. N., Maslov V. P., *Idempotent analysis and applications*, Kluwer Academic Publishers, 1997.
- [9] Mather J. N., *Existence of quasi-periodic orbits for twist homeomorphism of the annulus*, Topology **21** (1982), 457–467.
- [10] Mather J. N., Forni G. *Action minimizing orbits in Hamiltonian systems*, Lecture Notes in Math. **1589**, Springer-Verlag, 1994.
- [11] Morse M., *A fundamental class of geodesics on any closed surface of genus greater than one*, Trans. Amer. Math. Soc. **26** (1924), 25–60.
- [12] Sinai Ya. G., *Mathematical theory of phase transitions*, Nauka, 1981 (in Russian).
- [13] Sobolevskii A. N., *On periodic solutions of Hamilton-Jacobi equation with periodic force*, Uspekhi Mat. Nauk **53** (1998) N. 6, 265–266 (in Russian).
- [14] Yakovenko S. Yu., *Fixed points of the Bellman operator semi-groups and invariant manifolds of subdifferential Hamiltonian equations*, Avtomatika i Telemekhanika (1989) N. 6, 43–52 (in Russian).