

Ultrametric space cut instead of 2d one in 2d quantum field models

D.B. Saakian

Yerevan Physics Institute, Alikhanian Brothers St. 2,
Yerevan 375036, Armenia

It is possible to formulate 2d field theory on the ultrametric space with the free correlators identical to 2d correlators and the same potential. Such model should carry some features of original model for scale invariant theories. For the case of strings and 2d conformal models it is possible to derive exact results. It is possible to investigate not only bulk structure (phase transition points) of theory, but sometimes also correlators. Such ultrametric models could be naturally expressed via random energy model and directed polymer on Cayley tree.

REM [1-8] is one of fundamental models of modern physics. It was initially introduced as the simplest model of spin glass, but later on its relation to a whole series of science fields were disclosed. This model has a decisive property of operation as an optimum coding device [2-4]. In [9-11] a relation of 2d quantum Liouville model to REM and to the Directed Polymer (DP) on Cayley tree. Using similar ideas we intend to construct 2d quantum models in ultrametric space, as well as bring out some results for strings. Let us consider the ultrametric (UM) space with some measure for the (sphere) area $d\mu_s(V, X)$, volume (ball) measure $d\mu_v(V, X)$, total surface e^V and total volume $e^V - 1$. We can construct such UM space as a limit of hierarchic lattices. Consider a tree with a constant number of branchings q in each node and number N for hierarchies. The number of end points is q^N , the number of branches $\frac{q^N - 1}{q - 1}$. We consider a set of end points as a surface of sphere, the set of branches makes a volume (of the ball), each point on the surface is connected with the origin (zero level of the hierarchi) via a single path. We determine a measure $d\mu_s(V, X) = 1$ for each end point, then $d\mu_l(V, l) = (q - 1)$ for each link. Instead of integer q consider the limit $q \rightarrow 1, N \ln q \rightarrow V_0$. Now we have for the total area $\mu_s = \int d\mu_s(V_0, X) = e^{V_0}$ and for the total volume $\mu_l = \int d\mu_l(V, l) = e^{V_0} - 1$. We determine UM distance between two points x, y on this surface $V_0 - v \equiv V_0 - \frac{v}{N}$, where n is the number of the hierarchic level, on which x and y had the last common node on trajectories to their point from origin. The maximum UM distance between two points v on the surface is V_0 (an ordinary distance as a function of V will be defined lately).

We define some field $\phi(x)$ on our surface. For determination of kinetic energy (quadratic form with the Laplacian as the kernel in the conventional space) let us consider the expansion

$$\phi(X) = f_0 + \int_0^{V_0} dV f(V, l) \quad (1)$$

Here $f(l)$ is determined on the links. The integration in (1) is made along the trajectory of point X . Since the measures on both the threads of (1) coincide ($\int d\mu_l(V, l) = 1 + \int dV d\mu_l(V, l)$), we omit the Jacobian. Now determine the kinematic part of the action for the field $\phi(x)$

$$\int_0^{V_0} dV d\mu_l(V, l) 2\pi f(V, l)^2 \quad (2)$$

Then the partition under the potential UM

$$\int d f \exp\left\{-\int_0^{V_0} dV \int d\mu_l(V, l) 2\pi f(x, V)^2\right\} \exp\left\{\int d\mu_s(V_0, X) U(\phi(X))\right\} \quad (3)$$

We have for the correlator

$$\langle \phi(X)\phi(X') \rangle = \frac{V}{4\pi} \quad (4)$$

where V is the UM distance between the points X, X' . During the treatment of (4) we have omitted the zero modes f_0 in (1) for $\phi(x)$. For usual 2d models with

$$\int d\phi_0 d\phi \exp\left\{-\frac{1}{2} dx^2 \nabla \phi(x)^2\right\} \exp\left\{\int dx U(\phi(x))\right\} \quad (5)$$

we have for the total surface πR^2 and for correlators

$$\langle \phi(X)\phi(X') \rangle = \frac{\ln r}{2\pi} \quad (6)$$

We can determine the distance from the equality $V = \ln \pi r^2$. Then our correlators coincide (in any case when $r \gg 1$). What is the advantage of representation (3)? We are in a position to calculate the partition through the iterations. Let us take some large number K and determine the iterations

$$\begin{aligned} I_1(x) &= \sqrt{2K} \int_{-\infty}^{\infty} \exp\{-2\pi K y^2 + U(x+y)\} dy \\ I_{i+1}(x) &= \sqrt{2K} \int_{-\infty}^{\infty} \exp\{-2\pi K y^2\} I_i^{\exp(\frac{V}{K})}(x+y) dy \\ Z &= \lim_{K \rightarrow \infty} I_K^{\exp(\frac{V}{K})}(0) \end{aligned} \quad (7)$$

As virtually for determination of partition we need only the equation (7), our (speculation) abstraction $q \rightarrow 1$ is quite correct. In principle, in equation (2) instead of dV one can take any measure $dV u(V)$, e.g., $u(V) = e^{\alpha V}$. In this case (the area considered by us $e^V \sim R^2$ the equation is transformed to

$$\langle \phi(X)\phi(X') \rangle \sim r^{-\frac{\alpha}{2}} \quad (8)$$

So choosing $\alpha = d(d-2)$ we have d dimensional free field correlator. Now in (7) we replace $\exp(-2\pi K y^2)$ by

$$\exp(-2\pi K y^2 u((i-1)/K)) \quad (9)$$

It is held that the equation of iteration is easily solved only when the function $u(v)$ is constant (in our case (6))

or when $u(v)$ steadily falls. In this case the SG physics corresponds to the pattern of gradual freezing of the hierarchy levels [6]. In principle, one can repeat the same reasoning for fermions also. Let us introduce the fields $\psi(V, l), \bar{\psi}(V, l)$. Then

$$\int_0^{V_0} dV d\mu_i(V, l) \bar{\psi}(V, l) \psi(V, l) e^V \psi(V, X) \psi(V', X') + \bar{\psi}(V, X) \bar{\psi}(V, X) = \delta(x - x') \delta(V - V') \quad (10)$$

This will give an expression for correlators $\Psi(V_0, X) = \int_0^{V_0} dV \psi(V, l), \bar{\Psi}(V_0, X) = \int_0^{V_0} dV \bar{\psi}(V, l)$

$$\langle \bar{\Psi}(V_0, X) \Psi(X') \rangle \sim r^{-1} \quad (11)$$

To apply these ideas to the strings one is to remember that in [10] it was obtained how the correlator in directed polymer (DP) on Cayley tree is related to correlators in the Liouville model. It is seen that the strings are closely related to the UM space approach, REM and DP. For string's (corresponding to the 2d gravitation with matter field c) partition it is known that [12-15]

$$Z = \frac{1}{V_{SL(2,C)}} \int_0^\infty dA e^{-\mu A} Z(A) \\ Z(A) = \int D_g \phi \delta(\int d^2 w \sqrt{\hat{g}} e^{\alpha\phi} - A) e^{\frac{1}{8\pi} \int d^2 w \sqrt{\hat{g}} \phi \Delta \phi + QR\phi} \quad (12)$$

where $\frac{1}{V_{SL(2,C)}}$ is some factor connected with the fixing of gauging. It is known for the coefficients

$$c = 1 - 12\alpha_0^2 \\ Q = 2\sqrt{2 + \alpha_0^2} \\ \alpha = -\frac{\sqrt{25-c}}{\sqrt{12}} + \frac{\sqrt{1-c}}{\sqrt{12}} \\ \frac{Q}{\alpha} = \frac{1}{12} [c - 25 - \sqrt{(25-c)(1-c)}] \\ \frac{1}{8\pi} \int d^2 w \sqrt{\hat{g}} R = 1 \quad (13)$$

The last equation is written for the sphere where the measure Dg is linear. If we use the technique of zero mode, we can decompose $\phi(x)$ in two parts $\phi \rightarrow \phi_0 + \tilde{\phi}$, where ϕ_0 is the zero mode. Here it is easy to make integration over ϕ_0 and obtain

$$Z = \frac{1}{V_{SL(2,C)}} \mu^{-\frac{Q}{\alpha}} \Gamma\left(\frac{Q}{\alpha}\right) \int D_g \phi e^{\frac{1}{8\pi} \int d^2 w \sqrt{\hat{g}} \phi \Delta \phi + QR\phi} \left(\int d^2 w \sqrt{\hat{g}} e^{\alpha\phi} \right)^{-\frac{Q}{\alpha}} \quad (14)$$

We see that $Z = \langle z^\mu \rangle, \mu = -\frac{Q}{\alpha}$ and $z = \int d^2 w \sqrt{\hat{g}} e^{\alpha\phi}$ resembles partition in some sense. On the language of spin glasses this expression corresponds to real positive replica (for $0 < c < 1$ μ changes in interval $2 < \mu < 2.5$) In case of other topologies with $h > 0$ we have to take $\mu = -\frac{Q(1-h)}{\alpha}$.

Let us return to UM space in (14). Here instead of $d^2 w \sqrt{\hat{g}}$ the UM measure $d\mu(X, V_0)$ is taken, and instead of $\frac{1}{8\pi} \int d^2 w \sqrt{\hat{g}} \phi \Delta \phi$ quadratic form (3). Then one can

strictly prove in analogy to [7] that the thermodynamics (14) (in the UM version) is equivalent to REM for final replicas [8]. Dividing 2d space as a whole in M parts, calculating the distribution (11), and neglecting in the thermodynamic limit the term in the action Q term (this term may turn out to be crucial at the determination of correlators) we have

$$\langle \delta(E - \phi(x)) \rangle = e^{-\frac{E^2}{2G(0)}} \\ G(0) = \ln M \quad (15)$$

where $G(r)$ is a correlator of the scalar field. Instead of our system in UM space we take a common REM (the case of final replicas) with M configurations and distribution (15) for E_i . Derrida and Gardner solved such a system [8]. Here we give only a qualitative derivation of exact results.

$$\langle \left(\sum_{i=1}^M e^{-\beta E_i} \right)^\mu \rangle \quad (16)$$

(16) is calculated for positive integral values of μ , where the averaging is made over the distribution (15) for each E_i . There are only two competing terms,- the first one has a form of $e^{-\beta \mu E_i}$

$$Z = \langle \left(\sum_{i=1}^M e^{-\mu \beta E_i} \right) \rangle = M e^{\frac{G(0)(\beta \mu)^2}{2}} \\ \ln Z = \ln M + \frac{G(0)\beta^2 \mu^2}{2} \quad (17)$$

and is one of diagonal terms in (16), and the second one is from the cross terms in expansion Z

$$Z = \langle \left(\sum_{i=1}^M e^{-\beta E_i} \right) \rangle^\mu = M^\mu e^{\mu \frac{G(0)(\beta)^2}{2}} \\ \ln Z = \mu \ln M + \frac{G(0)\beta^2 \mu}{2} \quad (18)$$

We consider these expressions in the thermodynamic limit. When the entropy $\ln Z - \frac{dZ}{d\beta} \beta$ in (17) is zeroth, then below the zeroing temperature ($T = \frac{1}{\beta}$) we take for $\ln Z$ the expression that is linear in β and is continuous in the transition point. Eventually we have for the third phase

$$\ln Z = \mu \sqrt{\beta_c} \beta \\ \beta_c = \sqrt{\frac{2 \ln M}{G(0)}} \quad (19)$$

The third phase takes place for $\mu < 1$. For $0 < d < 1$ and so $2 < \mu < 2.5$ there occur only the phases (17)-(18). So, for spherical topology our string is always in phase (17) (even at $d > 1$ when our formulas (13) give complex values for β).

For other topologies the string is in the phase (18) for $d < 1$.

Let us analytically continue (13) for $c > 1$. Here both of our parameters μ, β are complex. For higher dimensions the string is first in SG phase. As is definitely known from SG physics, there should be some probabilistic description of a physical state state. When we continue

increase the dimensionality, then at $d = 19$ there is another phase transition. It is known from REM, that at complex temperatures $\beta = \beta_1 + i\beta_2$ there is some new, Lee-Yang-Fisher phase, when

$$\begin{aligned} \beta_1 &= \frac{\beta_c}{2} \\ \beta_2 &> \frac{\beta_c}{2} \end{aligned} \quad (20)$$

This expression could be derived by means of representation $\langle \ln |Z| \rangle = \langle \frac{|Z|^\mu - 1}{\mu} \rangle$. In calculations of $\langle |Z|^\mu \rangle$ in $\mu \rightarrow 0$ limit at complex β the principal terms in (16) are the crossing ones $e^{-\beta_1(E_{i_1} + \dots + E_{i_{\frac{\mu}{2}}})}$. From expression (13) for $\alpha \equiv \beta_1 + i\beta_2$ we see that this happens at $c = 19$. So at $d = 19$ there is a second phase transition for strings (after first one at $d = 1$) and the topology higher than spherical. We can apply the same ideas to other conformal theories using the Coulomb-gas formalism with the background charge α_0 [16]. To calculate a correlator $\prod_k \exp\{i\alpha_k\}$ with screening charges $Q_+^m Q_-^n$ one has consider

$$\begin{aligned} Z &= \int D_g \phi e^{\frac{1}{8\pi} \int d^2 w \sqrt{g} \phi \Delta \phi + i2\sqrt{2}\alpha_0 R \phi} \\ &\prod_k \exp\{i\alpha_k\} \left(\int d^2 w \exp(i\alpha_+)^n \int d^2 w \exp(i\alpha_-)^m \right) \end{aligned} \quad (21)$$

We see in the ultrametric space this is equivalent to the system of 2 real replica REM-s at different imaginary temperatures.

When one considers the correlators without screening charges, the scaling indices of the 2 point correlator are the same as those in the 2d case. It is possible to derive minimal models also in this scheme. We can calculate the multipoint correlators and compare with 2d ones, bearing in mind that in our UM space all triangles are isosceles. It is interesting to compare the structural couplings for 2d and ultrametric cases. It will be interesting to apply this approach to other exactly solvable models in 2d as well as to other versions of strings.

As for the construction of theories in dimensions $d > 2$ in principle there is a phenomenological criteria for establishment of relation of any theory to REM and eventually to our ultrametric space approach. It is that, if the free energy of system is a logarithm of the number of particles. In REM the efficient number of couplings is an exponent of the number of spins or free energy. The same situation is with scale invariant theories in any dimensions. Another argument is that in critical theories system can ignore some details of geometry and so, do not distinguish real space from our UM space. So there is some chance that at the critical point our ultrametric space approach could work also at $d > 2$. This hypothesis should be checked by means of Monte-Carlo simulations. The ultrametric space approach is somewhat crude (I don't see holomorphic+antiholomorphic decomposition and Majorana fermions, as in 2d space), but is a complete concept. It could will be successful, when the real space picture is too complicated (strings, $d > 2$ critical theories, turbulence) and the main problem consists in

the closure.

I am grateful to S. Apikyan, D. Karakhanyan, C. Lang, R. Mkrtchyan, V. Ohanyan, A. Sedrakyan for discussions, Amsterdam University for partial financial support.

-
- [1] B. Derrida, Phys. Rev. Lett. **45** (1980) 79.
 - [2] N. Sourlas, Nature **239** (1989) 693.
 - [3] D.B.Saakian, JETP lett. **55**(1992),n.2
 - [4] A.Allakhverdian,D.Saakian, Nucl. Phys. **B498**(1997)604
 - [5] B.Derrida,H. Spohn, J. Stat. Phys. **51** (1988) 817
 - [6] B.Derrida, J. Stat. Physique Lett. **46** (1985) 401
 - [7] J.Cook,B.Derrida,J.Stat.Phys. **63**(1991)505
 - [8] E.Gardner,B. Derrida, J.Phys.**A22** (1989)1975
 - [9] C.C.Chamon,C.Mudry,X.G.Wen,Phys.Rev.Lett. **77**(1996)4194.
 - [10] I.I.Kogan et all Phys. Rev.Lett. **77**(1996)707.
 - [11] H.E.Castillo et all,Phys.Rev. **B56**(1997)10668
 - [12] J. Distler,H.Kawai,Nucl. Phys. **B321**(1988)171
 - [13] F. David, Mod. Phys. Lett. **A3**(1988)1651
 - [14] A.Gupta,S.Trivedi,M.Wise,Nuclear Phys.**B340**(1990)475
 - [15] E.Abdalla,M.C.Abdalla,D.Dalmazzi,A.Zadra, 2d -Gravity in Non-Critical Strings, Springer-Verlag 1994
 - [16] V.S. Dotsenko,V.Fateev,Nuclear Phys. **B240**,312 (1984)