A STOCHASTIC CASCADE MODEL FOR FX DYNAMICS

WOLFGANG BREYMANN

Olsen & Associates Seefeldstrasse 233, 8008 Zurich, Switzerland

SHOALEH GHASHGHAIE

Institut für mathematische Statistik Sidlerstr. 5, 3012 Bern, Switzerland

PETER TALKNER

Paul Scherrer Institut 5232 Villigen, Switzerland

ABSTRACT

A time series model for the FX dynamics is presented which takes into account structural peculiarities of the market, namely its heterogeneity and an information flow from long to short time horizons. The model emerges from an analogy between FX dynamics and hydrodynamic turbulence. The heterogeneity of the market is modeled in form of a multiplicative cascade of time scales ranging from several minutes to a few months, analogous to the Kolmogorov cascade in turbulence.

The model reproduces well the important empirical characteristics of FX rates for major currencies, as the heavy-tailed distribution of returns, their change in shape with increasing time interval, and the persistence of volatility.

Keywords: Hydrodynamic turbulence, Kolmogorov cascade, heterogeneous market, information flow, volatility.

1. Information cascade: an analogy with hydrodynamic turbulence

The idea of a heterogeneous market that consists of traders acting on different time horizons was first advanced by Müller et al. who investigated the absolute values of FX returns on different time scales. They observed that the price changes over longer time intervals have a stronger influence on those over shorter time intervals than conversely. This has been interpreted as an information flow from long-term to short-term traders which directly influences the volatility on different time scales.

Though being very different in its physical nature, the motion of a turbulent fluid medium is also governed by a hierarchical process in which energy flows from large to small spatial scales². Interestingly enough, this formal analogy between FX dynamics and turbulence leads to similar statistical behavior of both phenomena³. For the FX dynamics, the hierarchical process can be implemented in form of a volatility cascade, which has recently been visualized by means of wavelet techniques⁴.

In the present contribution we take this hypothesis as granted and construct a stochastic multiplicative volatility cascade driving the FX price process. In the stochastic cascade model (SCM), that we present here, the volatility at an instant of time t is assumed to be created by processes on different time scales reflecting

the heterogeneity of the market dynamics. The information flow between traders on different time horizons is modeled by a directional interaction scheme between the levels of the cascade corresponding to the different time scales. More precisely, the flow of information is accounted for by an update of stochastic "transfer" factors relating the subsequent levels of the cascade.

2. Stochastic cascade model

We denote the price at time t by p_t and define the return as $r_t \equiv \log p_t - \log p_{t-1}$. In the proposed SCM the returns are described by a stochastic volatility model, i.e.

$$r_t = \sigma_t \xi_t, \tag{1}$$

where ξ_t are i.i.d. Student random numbers with 3 degrees of freedom. The volatility σ_t is governed by a hierarchical process that reflects the heterogeneity of the market on its different time horizons. Each level k of the cascade corresponds to a time horizon of duration $\tau^{(k)}$. The largest horizon $\tau^{(0)}$ representing the first level at the top of the cascade is typically of the order of a few months while the level m at the bottom of the cascade has the smallest horizon $\tau^{(m)}$ on which dealers may act and which is of the order of minutes. For the sake of simplicity we assume that the horizon ratio between neighboring levels is constant, i.e. $\tau^{(k)}/\tau^{(k-1)} = p < 1$ is independent of k.

We now assign to each level k of the hierarchy a volatility $\sigma_t^{(k)}$ which is determined by the respective volatility on the level k-1 and a time-dependent random factor $a_t^{(k)}$ that will be specified below. Hence the level volatilities are recursively defined by:

$$\sigma_t^{(k)} = a_t^{(k)} \sigma_t^{(k-1)}. (2)$$

As a toy model we assume that the market dynamics is completely determined by the behavior of the dealers and not subject to exogenous time-dependent influences. Then, the volatility $\sigma^{(0)}$ on the largest horizon is time independent. The volatility on the shortest horizon determines the returns, as given by eq. (1) with $\sigma_t = \sigma_t^{(m)}$. Using eq. (2) we find:

$$\sigma_t = \sigma^{(0)} \prod_{k=1}^m a_t^{(k)}.$$
 (3)

3. Updating rule for the volatility factors $\boldsymbol{a}_t^{(k)}$

The time-dependence of the factors $a_t^{(k)}$ results from the following renewal scheme: At the initial time t_0 the factors $a_{t_0}^{(k)}$ are drawn from independent lognormal distributions $LN(a_k, \lambda_k^2)$ with mean values a_k and variances λ_k^2 , $k=1,\ldots,m$. At a later time t_{n+1} the factor $a_{t_{n+1}}^{(1)}$ at the top of the hierarchy maintains the corresponding value $a_{t_n}^{(1)}$ at the preceding time t_n with a probability $1-w^{(1)}$ and else is drawn from $LN(a_1, \lambda_1^2)$. In the latter case all the subsequent factors $a_{t_{n+1}}^{(k)}$, k>1 are also renewed, i.e. they aquire independent random values from the respective distributions $LN(a_k, \lambda_k^2)$. If $a_{t_{n+1}}^{(1)}$ coincides with $a_{t_n}^{(1)}$ the factor $a_{t_{n+1}}^{(2)}$ will be renewed with a probability $w^{(2)}$. These rules apply through the whole cascade down to k=m: A renewal at some level k entails one at all higher levels k'>k; if no renewal has taken place up to level k, the coefficient at level k+1 will be




Fig. 1. Distribution of returns for time intervals reaching from 1 hour (top) to 4 weeks (bottom). Full lines: SCM simulation with parameters $\sigma^{(0)}=0.075,\,\tau(k)/\tau(k-1)=\sqrt{2},\,\lambda_k^2=0.0173,$ and $a_k = a$, where a is adjusted in such a way that the scaling of the second moment correspond to the observed values. Dots: USD/CHF FX spot rates from 1986 till 1996. For better visibility, the curves have been vertically shifted with respect to each other by multiples of log 10.

renewed with probability $w^{(k+1)}$ (drawn from the distribution $LN(a_{k+1}\lambda_{k+1}^2)$) or, with probability $1-w^{(k+1)}$, its value remains constant in time, $a_{t_{n+1}}^{(k+1)}=a_{t_n}^{(k+1)}$.


The renewal probabilities $w^{(k)}$ are given by:

$$w^{(k)} = 1 - (1 - p^{m-k})/(1 - p^{m-k+1}), \quad k = 1, \dots, m.$$
(4)

According to this renewal scheme the mean life-times of a factor at level k (measured in no. of time steps of length τ (m)) is given by p^{k-m} where p is the scaling factor of the time horizons as defined above.

4. Results

The time series defined by eqs. (1 - 4) have been simulated and the parameters have been adjusted in such a way that the model simultaneously reproduces the empirical return distributions (Fig. 1), the scaling laws for the moments⁵, and the autocorrelation function of the volatility (Fig. 2) of USD/CHF FX spot rates. The distributions of returns (cf. Fig. 1) exhibit the well-known heavy tails, which lose weight with increasing time interval. The autocorrelation function of the absolute returns (Fig. 2) shows the characteristic slowly decaying tail. In both cases the SCM simulation (full line) reproduces the observed data (dotted curves) very well. The



 ${
m Fig.~2.~Autocorrelation~function~of~absolute~returns~for~the~same~simulation~(full lines)~and~observed~data~(dots)~as~in~{
m Fig.~1.}$

scaling behavior of the moments and the cross correlation function of the volatility are presented in an extended version of the paper⁵.

To summarize, the SCM is a hierarchical time series model where a net information flow from long to short time horizons is implemented in terms of a random unidirectional action of the volatility at a given time horizon on that at the next shorter one. With only three adjustable parameters this model is able to reproduce different characteristic properties of intra-day FX price series.

Note that the existence of different groups of traders and the volatility updating mechanism are elements similar to those found in market microstructure models. Therefore the SCM may provide a link between market microstructure models and more conventional models.

One of the authors (S.G.) gratefully acknowledges support from the Swiss National Science Foundation.

- [1] U. Müller, M. M. Dacorogna, R. D. Davé, R. B. Olsen, O. V. Pictet and J. E. von Weizsäcker, Volatilities of different time resolutions Analyzing the dynamics of market components. *J. of Emp. Finance* 4 (1997) 211–239.
- [2] U. Frisch, Turbulence. Cambridge University Press, Cambridge (1995).
- [3] S. Ghashghaie, W. Breymann, J. Peinke, P. Talkner and Y. Dodge, Turbulent cascades in foreign exchange markets. *Nature* 381 (1996) 767–770.
- [4] A. Arneodo, J.-F. Muzy, and D. Sornette, Causal cascade in the stock market from the "infrared" to the "ultraviolet". *Preprint*, condmat/9708012 (1997).
- [5] W. Breymann, S. Ghashghaie, and P. Talkner, to be published.