April 2000

# Study of Macroscopic Coherence and Decoherence in the SQUID by Adiabatic Inversion

P. Silvestrini

Instituto di Cibernetica del CNR, via Toiano 6, I- 80072, Arco Felice, Italy and MQC group, INFN/Napoli, Italy

L. Stodolsky

Max-Planck-Institut für Physik (Werner-Heisenberg-Institut), Föhringer Ring 6, 80805 München, Germany

(April 2000)

We suggest a procedure for demonstrating quantum coherence and measuring decoherence times between different fluxoid states of a SQUID by using "adiabatic inversion", where one macroscopic fluxoid state is smoothly transferred into the other, like a spin reversing direction by following a slowly moving magnetic field. This is accomplished by sweeping an external applied flux, and depends on a well-defined quantum phase between the two macroscopic states. Varying the speed of the sweep relative to the decoherence time permits one to move from the quantum regime, where such a well-defined phase exists, to the classical regime where it is lost and the inversion is inhibited. Thus observing whether inversion has taken place or not as a function of sweep speed offers the possibility of measuring the decoherence time. The main requirement for the feasibility of the scheme appears to be that the low temperature relaxation time among the quantum levels of the SQUID be long compared to other time scales of the problem. Applications to the "quantum computer", with the level system of the SQUID playing the role of the qbit, are briefly discussed.

### I. INTRODUCTION

One of the most interesting questions of "macroscopic quantum mechanics" concerns the demonstration of quantum coherence and interference effects between apparently grossly distinguishable states of large systems. Observation of such effects would definitively lay to rest any ideas about big or "classical" systems being fundamentally different than small systems, in some way not subject to the rules of quantum mechanics. One system which has been extensively discussed in this regard is the Josephson junction [1] under current bias and the closely related rf Squid, where a superconducting ring is interrupted by a Josephson junction. In these systems the phase across the junction or the flux  $\Phi$  in the ring plays the role of a collective coordinate. In the last decade a number of beautiful junction experiments at low temperature [2] have seen effects connected with the quantized energy levels [3] expected with respect to this coordinate. A fast sweeping method [4] has also seen the effects of these quantized levels even at relatively high temperature.

An even more striking and direct manifestation of macroscopic quantum behavior would be the demonstration of quantum coherence between two apparently macroscopically different states of the rf SQUID, between states where the current goes around the ring in opposite directions. Recently, results have been reported with microwave methods showing the repulsion of energy levels expected due to quantum mechanical mixing of the different fluxoid states [5]. Here we would like to suggest a new method for demonstrating macroscopic coherence, one which further allows a measurement of the decoherence time, the time in which the coherence between the states is lost. Our basic idea is to use the process of "adiabatic inversion" where a slowly varying field is used to reverse the states of a quantum system. In its most straightforward realization, our proposal consists of starting with the system in its lowest state, making a fast but adiabatic sweep, and reading out to see if the final state is the same or the opposite fluxoid state. If the state has switched, the system has behaved quantum mechanically, with phase coherence between the two states. If not, the phase coherence between the states was lost and the system stayed in its original configuration-behaved classically.

A very interesting aspect of the present proposal, as we shall discuss below, is the possibility of passing from the quantum to the classical regime by simply varying the sweep speed. This allows us to obtain, for adiabatic conditions, the decoherence time as the longest sweep time for which the inversion is successful.

### II. ADIABATIC INVERSION IN THE RF SQUID

For the rf SQUID biased with an external flux  $\Phi_x$ , the "potential energy" of the system for which  $\Phi$  is the coordinate is shown in Fig 1. One has, near the origin, the familiar double well potential [6]

$$U = U_0 [1/2(\Phi - \Phi_x)^2 + \beta_L \cos\Phi] .$$
 (1)

Depending on whether the system is in the left or right well, the flux through the ring has a different sign and the current goes around in opposite directions. The potential can be varied by altering the external  $\Phi_x$ , becoming symmetric for  $\Phi_x = 0$ . We suppose a situation where a few states may be possible in a well, but we will be essentially concerned with the lowest level in each. In the adiabatic inversion procedure to be discussed,  $\Phi_x$  is swept from a maximum to an minimum value, passing through zero, such that the initially asymmetric left and right wells exchange roles, the originally higher well becoming the lower one and vice versa. The asymmetry of the configurations, however, is kept small so that we effectively have only a two-state system, composed of the lowest state in each well.



FIG. 1. Double potential well with harmonic level spacing  $\omega_0$  and initial spacing  $\epsilon$  between lowest levels.

As quantum states we refer to the lowest state of the left or right well as  $|L\rangle$  and  $|R\rangle$ , and the tunneling between |L> and |R> will generally lead to eigenstates which are linear combinations of the two. Any two-state system may viewed as constituting the two components of a "spin", and this provides an easy visualization which has been used in many contexts [7]. In the present problem "spin up" can be identified with, say, the state |L>(flux one way) and "spin down" with  $|R\rangle$  (flux the other way), while the spin in a general direction represents a quantum mechanical linear combination of the two states, with some definite relative magnitude and phase. Various influences like the external bias or the tunneling amplitude may be thought of as creating a kind of pseudomagnetic field  $\mathbf{V}$  about which the spin or polarization  $\mathbf{P}$ precesses (Fig. 2) according to the equation

$$\dot{\mathbf{P}} = \mathbf{V} \times \mathbf{P} - D\mathbf{P}_{tr} , \qquad (2)$$

where  $\mathbf{V}$  can be time dependent. The quantity D is the decoherence parameter, which we neglect for the moment and will deal with below.

Now it is a familiar fact under adiabatic conditions, where **V** varies slowly, that the "spin" **P** will tend to "follow" the magnetic field  $\mathbf{V}(t)$ . This is a completely familiar procedure when rotating the spin of say an atom or a neutron by a magnetic field. Here the motion of  $\mathbf{V}$ will be accomplished by sweeping  $\Phi_x$ ; it is then merely necessary to identify the components of  $\mathbf{V}$  and the meaning of "adiabatic" or "slow".



FIG. 2. Precession of the "spin" vector  $\mathbf{P}$  around the pseudo-magnetic field vector  $\mathbf{V}(t)$ . In adiabatic inversion  $\mathbf{V}$ , swings from "up" to "down", and carries  $\mathbf{P}$  with it.

The vertical component of  $\mathbf{V}$  corresponds to the difference in the two potential wells or lowest energy levels, for small asymmetry it is linear in  $\Phi_x$ . If  $\epsilon$  is the initial level splitting when  $\Phi_x = \Phi_x^{max}$ , we can write  $V_{vert}(t) = \epsilon(\Phi_x(t)/\Phi_x^{max})$ . Thus as  $\Phi_x$  sweeps from its positive maximum value to its negative minimum value  $V_{vert}$  reverses direction. The transverse component of  $\mathbf{V}_{tr}$  corresponds to the tunneling energy between the two quasi-degenerate states  $V_{tr} = \omega_{tunnel}$ . As  $V_{vert}$  passes through zero at  $\Phi_x = 0$ , the |L > and |R > states are strongly mixed and the splitting of the resulting energy eigenstates is determined by  $V_{tr}$  alone. The magnitude of V(t) at a given time,  $|\mathbf{V}| = \sqrt{V_{vert}^2 + V_{tr}^2}$  gives the instantaneous splitting of the two levels, which varies from approximately  $\epsilon$  in the vertical position of  $\mathbf{V}$  to  $\omega_{tunnel}$ in the horizontal position.

As to the meaning of "adiabatic" or "slow", the imposed time variation of  $\mathbf{V}$  should not significantly contain frequencies or fourier components corresponding to the energy splitting between levels. Or in terms of time, the rate of variation of  $\mathbf{V}$  should be slow on the time scale corresponding to the tunneling time between the two states. Thus we have the requirement on  $\mathbf{V}$  that its relative rate of variation  $\dot{V}/V$  always be small compared to V itself. Since the varying component of  $\mathbf{V}$  is  $V_{vertical}$  (neglecting the indirect effect of  $\Phi_x$  on  $\omega_{tunnel}$ ) we require  $\dot{V}_{vertical}/V \ll V$ . Thus taking the near degenerate configuration where  $V \approx \omega_{tunnel}$  we find

$$\epsilon \frac{\dot{\Phi}_x(t)}{\Phi_x^{max}} \approx \epsilon \omega_{sweep} \ll \omega_{tunnel}^2 \tag{3}$$

Ф

as the condition for adiabaticity. If the adiabatic condition is violated, then  $\mathbf{P}$  cannot follow  $\mathbf{V}$  and undergoes wide swings. We stress that for adiabatic inversion there must be a well-defined quantum phase between the two states, and that when this phase is lost the inversion is suppressed. This is the topic of the next section.

## III. DAMPING

Thus far we have not considered dissipative or damping effects tending to destroy the quantum coherence of the system; we have neglected the D term in Eq [2]. The quantity  $\mathbf{P}_{tr}$  has the interpretation of the degree of phase coherence between the two states, and D gives the rate of loss of this coherence [8]. In particular, we are interested in the effect of D on the inversion process. With the loss of phase coherence between the two states, we expect the situation to become more "classical" and for the inversion to be inhibited. Indeed, in solving Eq[2] for large D one finds that the inversion is strongly blocked and that one arrives in the "Turing-Watched Pot-Zeno" regime [7]. Eq [2] with moving  $\mathbf{V}(\mathbf{t})$  has been studied numerically [9]. One finds that while for weak damping D has little effect on the inversion, values of  $D/V_{tr} \sim 0.1$ are enough to stall the inversion and values  $D/V_{tr} >> 1$ block it altogether. Given adiabatic conditions, the important relation is that between the sweep speed and the decoherence time 1/D. This relation determines if the system has time to "decohere" during the sweep. For  $D/V_{tr} = 0.1$ , for example, one finds that a factor of ten increase in the sweep speed will change the stalled inversion into a successful one. Such behavior is of great interest to us here, since if the experimental situation is favorable, we can pass from the quantum regime (small D) to the classical regime (large D) by varying the sweep speed. In the former case the system ends up in the other potential well while in the latter case, where the phase coherence is destroyed, the system behaves classically as  $\mathbf{P}_{tr} \to 0$  and remains in the original state.

Concerning the absolute magnitude of D, a number of subtle issues are involved which cannot be treated here. However, for orientation we can take the result of calculations based on the Caldeira-Leggett approach, where we may identify the "decay rate" for weak dissipation with D. Weiss and Grabert [10] give  $T/Re^2$ for this parameter (their  $\Gamma$ ). With  $R = 5 \ M\Omega$ , this gives  $D = 0.08 \ mk = 9.6 \ MHZ$  at  $T = 100 \ mk$ , and  $D = 0.008 \ mk = 960 \ kHZ$  for  $T = 10 \ mK$ . (Units:  $1 \ K = 8 \ meV = 120 \ GHZ$ ,  $1/e^2 = 4 \ k\Omega$ ). If these estimates are correct, we are in an interesting range since it is possible to suppose sweeps both slow and fast on these scales and thus the direct measurement of D and its temperature dependence.

## **IV. TIME SCALES**

Since we envision working at time scales shorter than have been common in this field, it may be useful to give a qualitative discussion of the various time scales which arise. The highest frequency involved is the ordinary harmonic frequency  $\omega_0$  in one of the potential wells, giving the approximate level spacing in the well. For typical conditions this spacing may be on the order of several Kelvin (K). Since we envision experiments in the Kelvin to millikelvin range, this implies that for the equilibrium system at the start of the sweep only the lowest level is populated. The next parameter is the tunneling frequency through the barrier. This is a sensitive function of the SQUID parameters, with sample values  $[\beta_L = 1.8, C = .01 \ pF, L = 1 \ Nh, R = 5 \ M\Omega]$  we obtain  $\approx 20 \ GHZ$ , and  $\omega_{tunnel}/\omega_0 \approx 4 \cdot 10^{-2}$ .

The next two parameters concern the extent and speed of the sweep. The initial asymmetry  $\epsilon$  (Fig. 1) should be small compared to  $\omega_0$  in order to retain the approximate two-state character of the system, but large compared to  $\omega_{tunnel}$  to avoid initial mixing of the two states, say  $\epsilon/\omega_0 \sim 10^{-1} - 10^{-2}$ . The speed of the sweep,  $\omega_{sweep}$  will be the easiest experimental parameter to control, and the behavior of the results as  $\omega_{sweep}$  is varied will be an important check on the theory. It must not be so fast as to lose adiabaticity, but not so slow as to allow relaxation processes to mask the results. We may suppose it to be in the range  $10^{-3}\omega_0 - 10^{-4}\omega_0 = 10 - 100 MHZ$ . With these numbers, it is possible to satisfy the adiabatic condition Eq [3].

Finally there are dissipative parameters D, and  $\omega_{relax}$ , the relaxation rate for transitions among the SQUID levels. From the above estimate of 1/D in the 1-10 MHZ range we have, as mentioned, the interesting possiblity of being able to choose sweep speeds either slow or fast with respect to the decoherence time while still retaining the adiabatic condition. The resulting switches between classical and quantum behavior would provide persuasive evidence for the correctness of our general picture, and allow the measurement of this parameter and its temperature dependence. Finally, the relaxation parameter  $\omega_{relax}$  gives the rate of conventional kinetic relaxation processes induced by the environment, say thermal jumping over the barrier or direct transitions with the emission of some energy. Estimates based on the formulas of Ovichinnikov and Schmid [11] lead to  $\omega_{relax} \sim 100 \; kHZ$ which, as desired, would be much slower than the sweep time. This large difference between the tunneling time and the relaxation time is due to the many orders of magnitude suppression associated with the emission of energy in the relaxation process, particularly for underdamped SQUIDS.

In conclusion, there are five important (inverse) timescales;  $\omega_0$ ,  $\omega_{tunnel}$ ,  $\epsilon$ ,  $\omega_{sweep}$ , D, and  $\omega_{relax}$ . Naturally a wide range of behavior is available by changing the parameters and some, like the sweep time or the tem-

perature, can be adjusted online. Probably the most important condition for feasibility is the smallness of  $\omega_{relax}$ : relaxation processes should not be significant during the sweep and the subsequent readout.

Concerning the readout itself, we have examined a scheme involving a switchable flux linkage to a DC SQUID, which would be sufficiently fast for the above estimates [12]. Here one profits from the fact that observation of the system is only necessary after the procedure is completed.

## V. QBITS AND THE QUANTUM COMPUTER

The two-state system under discussion here suggests itself as a physical embodiment of the "quantum computer". The "qbit" itself is naturally represented by L and R playing the role of 0 and 1. A linear combination of L and R may be created by adiabatically rotating **P** from some starting position. Adiabatic inversion is evidently an embodiment of NOT since it will turn one linear combination into another one with the weights of L and R interchanged. As for CNOT, the other basic operation, a NOT is performed or not performed on a "target bit" according to the state of a second, "control bit", which itself does not change its state. One straightforward realization of this would be to perform the NOT operation as just described in the presence of an additional linking flux supplied by a second SQUID nearby. The magnitude and direction of this linking flux would be so arranged that the inversion of the first SQUID is or is not successful depending on the state of the second SQUID. This and many other interesting combinations of junctions and flux linkages may be contemplated and are under study [13].

SQUID systems like these would seem to be particularly well suited for the embodiment of the quantum computer, where we wish to generate a series of unitary transformations for the various steps of computation. This may be done by creating a "moving landscape" of potential maxima and minima, as in our simplest one-dimensional example of the adiabatic inversion. This imaginary landscape can be produced and manipulated by controlling various external parameters (as with our sweeping flux), performing the various operations in one physical device. Naturally, practicality will depend very much on the relation between the speed of these operations and the decoherence/ relaxation times which we hope to determine by the present methods.

Although our interest here has been the SQUID, it will be evident that the principle of determining decoherence times through the inhibition of adiabatic inversion could be applied to many other types of systems as well.

Preliminary reports on this work were presented by one of us (P.S.) at ICSSUR VI (Napoli, Italy, 24-29 May 1999), at NATO ASI (Ankara, 13-25 June 1999), and at MICRO AND NANO CRYOCENICS (Jyvaskyla, Finland, August 1-3 1999).

- See A. J. Leggett, Les Houches, session XLVI (1986) Le Hasard et La Matiere; North -Holland (1987), and references cited therein.
- [2] R. Rouse, S. Y. Han, and J. E. Lukens, Phys. Rev. Let.75, 1615 (1995).
- [3] J. Clarke, A. N. Cleland, M. H. Devoret, D. Esteve, J. M. Martinis, Science 239, 992 (1988).
- [4] P. Silvestrini, V.G. Palmieri, B. Ruggiero, and M. Russo Phys. Rev. Let. **79**, 3046 (1997).
- [5] J. Friedman, V. Patel, W.Chen, S.K. Tolpygo and J.E. Lukens, cond-mat/0004293, submitted to Nature. Also see C.S. van der Wal, A.C. ter Haar, F. K. Wilhelm, R. N. Schouten, C.J.P.M. Harmans, T.P. Orlando, Seth Lloyd, and J.E. Mooij, submitted to MQC2 (Conference on Macroscopic Quantum Coherence and Computing, Naples, June 2000.) Proceedings to be published by Academic-Plenum.
- [6] A. Barone and G. Paternó, Physics and Applications of the Josephson Effect, Wiley, N.Y., 1982.
- [7] R. A. Harris and L. Stodolsky, Phys. Lett. 116B, 464 (1982). For a general introduction and review of these concepts see L. Stodolsky, "Quantum Damping and Its Paradoxes" in *Quantum Coherence*, J. S. Anandan ed. World Scientific, Singapore (1990). For recent work involving mesoscopic devices see L. Stodolsky, Phys. Lett. B459, 193 (1999) and for the "fluctuation- decoherence relation", Physics Reports 320, 51 (1999).
- [8] Note that in the absence of D, according to Eq [2], **P** cannot change its length, so the density matrix, which **P** parameterizes, retains its degree of purity. In using Eq [2] one assumes the temperature low enough that relaxation processes like barrier hoping may be neglected, thus damping of only the transverse components,  $\mathbf{P}_{tr}$ .
- [9] The effect of damping on adiabatic inversion has been extensively studied in neutrino physics where it is called the "MSW effect". See L. Stodolsky, Phys. Rev. D36 2273 (1987), and chapter 9 of G.G. Raffelt, *Stars as Lab*oratories for Fundamental Physics (Univ. Chicago Press, 1996). We quote the numerical results from a diploma thesis by J. Flaig, University of Munich (1989).
- [10] U. Weiss and H. Grabert, *Europhys. Lett.* 2 667 (1986). See Eq (16) of this paper. The general approach is based on Caldeira and Leggett, Phys. Rev. Lett. 46, 211 (1981).
- [11] Yu. Ovchinnikov and A. Schmid, Phys. Rev. B50 6332 (1994).
- [12] B. Ruggiero, V. Corato, E. Esposito, C. Granata, M. Russo, P. Silvestrini, and L. Stodolsky, submitted to SATT 10, to be published in Intl. Jnl. Mod. Phys.B.
- [13] D.V. Averin, Solid State Communications 105, 659 (1998), has discussed related ideas using adiabatic operations on the charge states of small Josephson junctions.