

Ising model on the square $M \times N$ lattice: exact finite-size calculations.N.Sh. Izmailian^{1,2} and Chin-Kun Hu^{1,3*}¹ Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan² Yerevan Physics Institute, Alikhanian Br. 2, 375036 Yerevan, Armenia³ Department of Physics, National Dong Hwa University, Hualien 97401, Taiwan

Let f , U and C represent, respectively, the free energy, the internal energy and the specific heat of the critical Ising model on the square $M \times N$ lattice with periodic boundary conditions, and f_∞ represents f as $M, N \rightarrow \infty$. We find that f , U and C can be written as: $N(f - f_\infty) = \sum_{i=1}^{\infty} f_i/N^{2i-1}$, $U = -\sqrt{2} + \sum_{i=1}^{\infty} u_i/N^{2i-1}$ and $C = \frac{8}{\pi} \ln N + \sum_{i=0}^{\infty} c_i/N^i$ and Nf and U are well-defined odd function of N^{-1} . We also find that the ratio u_{2i+1}/c_{2i+1} of subdominant (N^{-2i-1}) finite-size correction amplitudes for the internal energy and the specific heat are constant, namely, $u_{2i+1}/c_{2i+1} = 1/\sqrt{2}$ and $u_{2i}/c_{2i} = 0$. We obtain analytic equations for f_i , u_i and c_i up to $i = 3$.

PACS numbers: 05.50+q, 75.10-b

The concept of finite - size scaling [1] plays a fundamental role in the theory of finite-size effects near phase transitions [1–4] and is indispensable for the analysis of numerical studies of critical phenomena in small systems [5]. The theory of finite-size effects and of finite-size scaling in general has been most successful in deriving critical and noncritical properties of infinite systems from those of their finite or partially finite counterparts. Of course the main difficulty is that the finite-size scaling is valid only for large N and that the numerical calculations can be done only for small N . Therefore the corrections to finite-size scaling cannot be neglected. The dominant scaling properties of a number of two-dimensional models have been determined in recent years, but questions remain about the subdominant finite-size corrections to scaling. This paper presents new results for finite-size effects in the Ising model on a large $M \times N$ square lattice at the critical point.

The critical properties of a finite Ising model consisting of a square $M \times N$ lattice with periodic boundary condition has been investigated by A. Ferdinand and M. E. Fisher [6]. Among other results, they gave explicit expressions for the internal energy per spin U and the specific heat C at critical point ($T = T_c$), and for a large lattice they obtained the following asymptotic expansions

$$f = f_\infty + \frac{1}{\xi N^2} \left[\ln(\theta_2 + \theta_3 + \theta_4) - \frac{1}{3} \ln(4\theta_2\theta_3\theta_4) \right] + O\left(\frac{1}{N^3}\right), \quad (1)$$

$$U = -\sqrt{2} - \frac{1}{N} \frac{2\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} + O\left(\frac{1}{N^2}\right), \quad (2)$$

$$\begin{aligned} C = & \frac{8}{\pi} \ln N + \frac{8}{\pi} \left(\ln \frac{2^{5/2}}{\pi} + C_E - \frac{\pi}{4} \right) \\ & - \frac{16}{\pi} \frac{\sum_{i=2}^4 \theta_i \ln \theta_i}{\theta_2 + \theta_3 + \theta_4} - 4\xi \left(\frac{\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} \right)^2 \\ & - 2\sqrt{2} \frac{\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} \frac{1}{N} + O((\ln N)^3/N^2), \end{aligned} \quad (3)$$

where $\xi = M/N$, $\theta_i = \theta_i(0, q)$ ($i = 2, 3, 4$) is elliptic theta functions of modulus $q = e^{-\pi\xi}$, C_E is the Euler constant and f_∞ is the free energy in the thermodynamic limit $M, N \rightarrow \infty$.

In this paper we extend the calculation of the critical internal energy and specific heat up to $1/N^5$ and $1/N^3$ orders, respectively. We also calculate the critical free energy heat up to $1/N^5$ order.

Consider an Ising ferromagnet on an $M \times N$ lattice with periodic boundary conditions (*i.e.* a torus). The Hamiltonian of the system is

$$\beta H = -J \sum_{\langle ij \rangle} s_i s_j, \quad (4)$$

where $\beta = (k_B T)^{-1}$, the Ising spins $s_i = \pm 1$ are located at the sites of the lattice and the summation goes over all nearest-neighbor pairs of the lattice. The partition function $Z_{MN}(T)$ of a finite $M \times N$ square Ising lattice wrapped on a torus is

$$Z_{MN}(T) = \frac{1}{2} (2 \sinh 2J)^{\frac{1}{2}MN} \sum_{i=1}^4 Z_i. \quad (5)$$

The partial partition functions Z_i are defined by

$$Z_1 = \prod_{r=0}^{N-1} 2 \cosh \frac{M}{2} \gamma_{2r+1} = P_1 \exp \left[\frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r+1} \right], \quad (6)$$

$$Z_2 = \prod_{r=0}^{N-1} 2 \sinh \frac{M}{2} \gamma_{2r+1} = P_2 \exp \left[\frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r+1} \right], \quad (7)$$

$$Z_3 = \prod_{r=0}^{N-1} 2 \cosh \frac{M}{2} \gamma_{2r} = P_3 \exp \left[\frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r} \right] (1 + e^{-M\gamma_0}), \quad (8)$$

$$Z_4 = \prod_{r=0}^{N-1} 2 \sinh \frac{M}{2} \gamma_{2r} = P_4 \exp \left[\frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r} \right] (1 - e^{-M\gamma_0}), \quad (9)$$

where

$$\begin{aligned} P_1 &= \prod_{r=0}^{N-1} (1 + e^{-M\gamma_{2r+1}}), & P_2 &= \prod_{r=0}^{N-1} (1 - e^{-M\gamma_{2r+1}}), \\ P_3 &= \prod_{r=1}^{N-1} (1 + e^{-M\gamma_{2r}}), & P_4 &= \prod_{r=1}^{N-1} (1 - e^{-M\gamma_{2r}}), \end{aligned} \quad (10)$$

and γ_r is implicitly given by

$$\cosh \gamma_r = \frac{\cosh^2 2J}{\sinh 2J} - \cos \frac{r\pi}{N}. \quad (11)$$

At the critical point J_c of the sq lattice Ising model, where $J_c = \frac{1}{2} \ln(1 + \sqrt{2})$, one then obtains

$$\gamma_r^{(cr)} = 2\psi_{sq} \left(\frac{r\pi}{2N} \right), \quad \text{with} \quad \psi_{sq}(x) = \ln \left(\sin x + \sqrt{1 + \sin^2 x} \right). \quad (12)$$

The free energy and the internal energy per spin are given by

$$f = \frac{1}{MN} \ln Z_{MN} = \frac{1}{2} \ln 2 \sinh 2J + \frac{1}{MN} \ln \frac{1}{2} \left(\sum_{i=1}^4 Z_i \right), \quad (13)$$

$$U = -\frac{1}{MN} \frac{d}{dJ} \ln Z_{MN} = -\coth 2J - \frac{1}{MN} \left(\sum_{i=1}^4 Z'_i \right) / \left(\sum_{i=1}^4 Z_i \right), \quad (14)$$

while the specific heat per spin is

$$\begin{aligned} C &= \frac{1}{MN} \frac{d^2}{dJ^2} \ln Z_{MN} \\ &= -\frac{2}{\sinh^2 2J} + \frac{1}{MN} \left\{ \left(\sum_{i=1}^4 Z''_i \right) / \left(\sum_{i=1}^4 Z_i \right) - \left[\left(\sum_{i=1}^4 Z'_i \right) / \left(\sum_{i=1}^4 Z_i \right) \right]^2 \right\}, \end{aligned} \quad (15)$$

where the primes denote differentiation with respect to J . At the critical point ($T = T_c$) the partial partition functions Z_i and their first and second derivatives are given by

$$Z_1 = P_1 e^A, \quad Z_2 = P_2 e^A, \quad Z_3 = 2P_3 e^B, \quad Z_4 = 0; \quad (16)$$

$$Z'_1 = 0, \quad Z'_2 = 0, \quad Z'_3 = 0, \quad Z'_4 = 4MP_4 e^B; \quad (17)$$

$$\frac{Z''_1}{MN} = Q_1 Z_1, \quad \frac{Z''_2}{MN} = Q_2 Z_2, \quad \frac{Z''_3}{MN} = Q_3 Z_3, \quad Z''_4 = -\sqrt{2}Z'_4, \quad (18)$$

where

$$A = \frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r+1}^{(cr)}, \quad B = \frac{M}{2} \sum_{r=0}^{N-1} \gamma_{2r}^{(cr)}, \quad (19)$$

$$Q_1 = \frac{1}{2N} \sum_{r=0}^{N-1} \gamma_{2r+1}^{(cr)} \tanh \frac{M\gamma_{2r+1}^{(cr)}}{2}, \quad (20)$$

$$Q_2 = \frac{1}{2N} \sum_{r=0}^{N-1} \gamma_{2r+1}^{(cr)} \coth \frac{M\gamma_{2r+1}^{(cr)}}{2}, \quad (21)$$

$$Q_3 = \frac{4}{N} + \frac{1}{2N} \sum_{r=1}^{N-1} \gamma_{2r}^{(cr)} \tanh \frac{M\gamma_{2r}^{(cr)}}{2}, \quad (22)$$

and P_i are given by Eq. (10) with $\gamma_r = \gamma_r^{(cr)}$ and $\gamma_r^{(cr)}$ denote the second derivative of γ_r with respect to J at the critical point $J = J_c$.

Then the exact expression for the free energy, the internal energy and the specific heat of a finite Ising model at critical point ($T = T_c$) can be written as

$$f = \frac{1}{2} \ln 2 + \frac{1}{MN} A + \frac{1}{MN} \ln \frac{P_1 + P_2 + 2P_3 \exp(B - A)}{2}, \quad (23)$$

$$U = -\sqrt{2} - \frac{4}{N} \frac{P_4}{2P_3 + (P_1 + P_2) \exp(A - B)}, \quad (24)$$

$$C = \sqrt{2}U - \xi \left(\frac{4P_4}{2P_3 + (P_1 + P_2) \exp(A - B)} \right)^2 + \frac{2Q_3 P_3 + (Q_1 P_1 + Q_2 P_2) \exp(A - B)}{2P_3 + (P_1 + P_2) \exp(A - B)}. \quad (25)$$

We will consider only sequences of lattices in which $\xi = M/N$ remains positive and finite as the thermodynamic limit $M, N \rightarrow \infty$ is approached. Using Taylor's theorem we obtained

$$M\gamma_r^{(cr)} = \sum_{i=0}^{\infty} \frac{a_i}{N^{2i+1}} = \pi \xi r - \frac{\pi^3 \xi}{12} \frac{r^3}{N^2} + \frac{\pi^5 \xi}{96} \frac{r^5}{N^4} + \dots \quad (26)$$

Using Euler-Maclaurin summation formula

$$\begin{aligned} \sum_{r=0}^N f(r\delta) &= \frac{1}{\delta} \int_0^{\delta N} f(x) dx + \frac{1}{2} [f(0) + f(\delta N)] \\ &+ \sum_{i=1}^m \frac{B_{2i}}{(2i)!} [f^{(2i-1)}(\delta N) - f^{(2i-1)}(0)] \delta^{2i-1} \\ &- \frac{\delta^{2m+1}}{(2m+2)!} \int_0^{\delta N} \bar{B}_{2m+2}(x) f^{(2m+2)}(x) dx, \end{aligned} \quad (27)$$

where B_{2i} are the Bernoulli numbers, $B_j(x)$ are the Bernoulli polynomials and $\bar{B}_j(x)$ are the periodic extension of $B_j(x)$, we can expand A and B up to arbitrary order

$$\begin{aligned} A &= \frac{MN}{\pi} \int_0^\pi \psi_{sq}(x) dx + M \sum_{k=1}^{\infty} \frac{2B_{2k}}{(2k)!} (2^{2k-1} - 1) \psi_{sq}^{(2k-1)}(0) \left(\frac{\pi}{2N}\right)^{2k-1} \\ &= \frac{2G}{\pi} MN + \frac{\pi\xi}{12} + \frac{7\pi^3\xi}{1440} \frac{1}{N^2} + \frac{31\pi^5\xi}{24192} \frac{1}{N^4} + \frac{10033\pi^7\xi}{9676800} \frac{1}{N^6} + \dots, \end{aligned} \quad (28)$$

$$\begin{aligned} B &= \frac{MN}{\pi} \int_0^\pi \psi_{sq}(x) dx - M \sum_{k=1}^{\infty} \frac{2B_{2k}}{(2k)!} \left(\frac{\pi}{N}\right)^{2k-1} \psi_{sq}^{(2k-1)}(0) \\ &= \frac{2G}{\pi} MN - \frac{\pi\xi}{6} - \frac{\pi^3\xi}{180} \frac{1}{N^2} - \frac{\pi^5\xi}{756} \frac{1}{N^4} - \frac{79\pi^7\xi}{75600} \frac{1}{N^6} - \dots \end{aligned} \quad (29)$$

Let us now evaluate the products P_i for $i = 1, 2, 3, 4$. It is easy to see from Eqs. (10) and (26) that P_i contains only even power of $1/N$

$$P_i = \pi_i(0, \xi) \left(1 + \sum_{j=1}^{\infty} \frac{p_{ij}}{N^{2j}} \right) \quad (i = 1, 2, 3, 4), \quad (30)$$

$$\begin{aligned} P_1 &= \pi_1(0, \xi) \left(1 + \frac{p_{11}}{N^2} + \frac{p_{12}}{N^4} + \dots \right), & P_2 &= \pi_2(0, \xi) \left(1 + \frac{p_{21}}{N^2} + \frac{p_{22}}{N^4} + \dots \right), \\ P_3 &= \pi_3(0, \xi) \left(1 + \frac{p_{31}}{N^2} + \frac{p_{32}}{N^4} + \dots \right), & P_4 &= \pi_4(0, \xi) \left(1 + \frac{p_{41}}{N^2} + \frac{p_{42}}{N^4} + \dots \right), \end{aligned}$$

with

$$\pi_1(0, \xi) = \frac{\theta_3}{\theta_0}, \quad \pi_2(0, \xi) = \frac{\theta_4}{\theta_0}, \quad \pi_3(0, \xi) = \frac{\theta_0^2}{\theta_3\theta_4}, \quad \pi_4(0, \xi) = \theta_0^2, \quad e^{\frac{\pi\xi}{4}} = \frac{2\theta_0^3}{\theta_2\theta_3\theta_4},$$

where $\theta_i = \theta_i(0, q)$ is elliptic theta functions of modulus $q = e^{-\pi\xi}$. As readily sees from Eqs. (23), (24) and (28) - (30) that the finite-size estimates of the free energy (Nf) and the internal energy (U) must be well-defined odd function of N^{-1} .

$$N(f - f_\infty) = \sum_{i=1}^{\infty} \frac{f_i}{N^{2i-1}}, \quad (31)$$

$$U = -\sqrt{2} + \sum_{i=1}^{\infty} \frac{u_i}{N^{2i-1}}. \quad (32)$$

Let us now evaluate the coefficients p_{i1}, p_{i2} for $i = 1, 2, 3, 4$. After little algebra we obtained

$$\begin{aligned} p_{11} &= -\frac{1}{3}\pi^3\xi \left[\frac{1}{4} \sum_{j=1}^{\infty} (-1)^j \frac{\cosh \pi\xi j}{\sinh^2 \pi\xi j} + \frac{3}{2} \sum_{j=1}^{\infty} (-1)^j \frac{\cosh \pi\xi j}{\sinh^4 \pi\xi j} \right], \\ p_{21} &= -\frac{1}{3}\pi^3\xi \left[\frac{1}{4} \sum_{j=1}^{\infty} \frac{\cosh \pi\xi j}{\sinh^2 \pi\xi j} + \frac{3}{2} \sum_{j=1}^{\infty} \frac{\cosh \pi\xi j}{\sinh^4 \pi\xi j} \right], \\ p_{31} &= -\frac{1}{3}\pi^3\xi \left[\sum_{j=1}^{\infty} \frac{(-1)^j}{\sinh^2 \pi\xi j} + \frac{3}{2} \sum_{j=1}^{\infty} \frac{(-1)^j}{\sinh^4 \pi\xi j} \right], \\ p_{41} &= -\frac{1}{3}\pi^3\xi \left[\sum_{j=1}^{\infty} \frac{1}{\sinh^2 \pi\xi j} + \frac{3}{2} \sum_{j=1}^{\infty} \frac{1}{\sinh^4 \pi\xi j} \right], \end{aligned} \quad (33)$$

$$\begin{aligned}
p_{12} &= \frac{1}{2}p_{11}^2 + \frac{4\pi^6\xi^2}{9} \left[\frac{3}{\pi\xi}\Psi_1(\pi\xi) + \frac{d}{d(\pi\xi)}\Psi_1(\pi\xi) \right], \\
p_{22} &= \frac{1}{2}p_{21}^2 + \frac{4\pi^6\xi^2}{9} \left[\frac{3}{\pi\xi}\Psi_2(\pi\xi) + \frac{d}{d(\pi\xi)}\Psi_2(\pi\xi) \right], \\
p_{32} &= \frac{1}{2}p_{31}^2 + \frac{4\pi^6\xi^2}{9} \left[\frac{3}{\pi\xi}\Psi_3(\pi\xi) + \frac{d}{d(\pi\xi)}\Psi_3(\pi\xi) \right], \\
p_{42} &= \frac{1}{2}p_{41}^2 + \frac{4\pi^6\xi^2}{9} \left[\frac{3}{\pi\xi}\Psi(\pi\xi) + \frac{d}{d(\pi\xi)}\Psi(\pi\xi) \right],
\end{aligned} \tag{34}$$

with

$$\begin{aligned}
\Psi_1(x) &= \frac{1}{128} \sum_{j=1}^{\infty} (-1)^j \cosh xj \left(\frac{1}{\sinh^2 xj} + \frac{60}{\sinh^4 xj} + \frac{120}{\sinh^6 xj} \right), \\
\Psi_2(x) &= \frac{1}{128} \sum_{j=1}^{\infty} \cosh xj \left(\frac{1}{\sinh^2 xj} + \frac{60}{\sinh^4 xj} + \frac{120}{\sinh^6 xj} \right), \\
\Psi_3(x) &= \frac{1}{16} \sum_{j=1}^{\infty} (-1)^j \left(\frac{2}{\sinh^2 xj} + \frac{15}{\sinh^4 xj} + \frac{15}{\sinh^6 xj} \right), \\
\Psi(x) &= \frac{1}{16} \sum_{j=1}^{\infty} \left(\frac{2}{\sinh^2 xj} + \frac{15}{\sinh^4 xj} + \frac{15}{\sinh^6 xj} \right).
\end{aligned} \tag{35}$$

The coefficients p_{i1} and p_{i2} can be rewritten in a more symmetrical way

$$\begin{aligned}
p_{11} &= \frac{1}{3}\pi^3\xi \left[\frac{1}{8}R(\xi/2) - \frac{5}{4}R(\xi) + 2R(2\xi) \right], \\
p_{21} &= \frac{1}{3}\pi^3\xi \left[R(\xi) - \frac{1}{8}R(\xi/2) \right], \\
p_{31} &= \frac{1}{3}\pi^3\xi [R(\xi) - 2R(2\xi)], \\
p_{41} &= -\frac{1}{3}\pi^3\xi R(\xi),
\end{aligned} \tag{36}$$

and

$$\begin{aligned}
\Psi_1(\xi) &= -2\Psi(2\xi) + \frac{17}{16}\Psi(\xi) - \frac{1}{32}\Psi(\xi/2), \\
\Psi_2(\xi) &= -\Psi(\xi) + \frac{1}{32}\Psi(\xi/2), \\
\Psi_3(\xi) &= 2\Psi(2\xi) - \Psi(\xi),
\end{aligned} \tag{37}$$

with

$$R(x) = S_2(x) + \frac{3}{2}S_4(x), \tag{38}$$

$$\Psi(x) = \frac{1}{16} (2S_2(x) + 15S_4(x) + 15S_6(x)). \tag{39}$$

Here for convenience we use the following notation:

$$S_n(x) = \sum_{j=1}^{\infty} \frac{1}{\sinh^n x j} \quad \text{for} \quad n = 2, 4, 6, \quad (40)$$

and $S_2(x)$ is given by (see [7] p. 721)

$$S_2(x) = \frac{1}{6} + \frac{2(2-k^2)}{3\pi^2} K^2(k) - \frac{2}{\pi^2} K(k)E(k), \quad (41)$$

where $x = \pi K'(k)/K(k)$ with $K(k)$ and $E(k)$ the elliptic integrals of the first and second kind, respectively. The $S_4(x)$ and $S_6(x)$ are given by

$$S_4(x) = -\frac{11}{90} - \frac{4(2-k^2)}{9\pi^2} K^2(k) + \frac{4}{3\pi^2} K(k)E(k) + \frac{8(1-k^2+k^4)}{45\pi^4} K^4(k), \quad (42)$$

$$S_6(x) = \frac{191}{1890} + \frac{32(2-k^2)}{45\pi^2} K^2(k) - \frac{16}{15\pi^2} K(k)E(k) - \frac{8(1-k^2+k^4)}{45\pi^4} K^4(k) - \frac{32(2-3k^2-3k^4+2k^6)}{945\pi^6} K^6(k). \quad (43)$$

Thus we are now in a position to evaluate $R(x)$ and $\Psi(x)$ given by Eqs. (38) and (39) respectively. The result is

$$R(x) = -\frac{1}{60} + \frac{4(1-k^2+k^4)}{15\pi^4} K^4(k), \quad (44)$$

$$\Psi(x) = \frac{1}{1008} - \frac{2(2-3k^2-3k^4+2k^6)}{63\pi^6} K^6(k). \quad (45)$$

The expressions for $R(2x)$, $R(x/2)$ and $\Psi(2x)$, $\Psi(x/2)$ can be written as a function of the modulus k by using properties of the elliptic functions.

Thus for the coefficients p_{i1} (for $i = 1, 2, 3, 4$) we finally obtain

$$\begin{aligned} p_{11} &= -\frac{7\pi^3\xi}{1440} + \frac{(7+8k^2-8k^4)\xi}{90\pi} K^4(k), \\ p_{21} &= -\frac{7\pi^3\xi}{1440} + \frac{(7-22k^2+7k^4)\xi}{90\pi} K^4(k), \\ p_{31} &= \frac{\pi^3\xi}{180} + \frac{(-8+8k^2+7k^4)\xi}{90\pi} K^4(k), \\ p_{41} &= \frac{\pi^3\xi}{180} - \frac{4(1-k^2+k^4)\xi}{45\pi} K^4(k). \end{aligned} \quad (46)$$

After little algebra the expressions for p_{i2} (for $i = 1, 2, 3, 4$) can be written as

$$\begin{aligned} p_{12} &= \frac{\xi^2 K^8}{189\pi^2} \left[31 - 88k^2 + 88k^4 + (31 - 78k^2 + 48k^4 - 32k^6) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \\ &\quad - \frac{31\pi^5\xi}{24192} + \frac{p_{11}^2}{2}, \end{aligned}$$

$$\begin{aligned}
p_{22} &= \frac{\xi^2 K^8}{189\pi^2} \left[31 - 67k^2 + 25k^4 - 21k^6 + (31 - 15k^2 - 15k^4 + 31k^6) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \\
&\quad - \frac{31\pi^5 \xi}{24192} + \frac{p_{21}^2}{2}, \\
p_{32} &= -\frac{\xi^2 K^8}{189\pi^2} \left[32 - 80k^2 + 38k^4 - 21k^6 + (32 - 48k^2 + 78k^4 - 31k^6) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \\
&\quad + \frac{\pi^5 \xi}{756} + \frac{p_{31}^2}{2}, \\
p_{42} &= -\frac{16\xi^2 K^8}{189\pi^2} \left[2 - 5k^2 + 5k^4 + (2 - 3k^2 - 3k^4 + 2k^6) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \\
&\quad + \frac{\pi^5 \xi}{756} + \frac{p_{41}^2}{2},
\end{aligned} \tag{47}$$

where for simplicity we denote $K \equiv K(k)$, $K' \equiv K'(k)$, $E \equiv E(k)$ and $E' \equiv E'(k)$.

Substituting Eqs. (28) - (30), (46) and (47) in Eqs. (23) and (24) we can write the expansions of the free energy (Nf) and the internal energy (U) at the critical point ($T = T_c$) up to $1/N^5$ order. The final result is

$$\begin{aligned}
N(f - f_\infty) &= \frac{\xi^{-1}}{N} \left[\ln(\theta_2 + \theta_3 + \theta_4) - \frac{1}{3} \ln(4\theta_2\theta_3\theta_4) \right] \\
&\quad - \frac{\pi^3}{N^3} \frac{8\theta_2\theta_3\theta_4 [\theta_3^3(\theta_2^3 + \theta_4^3) - \theta_2^3\theta_4^3] - 7(\theta_2^9 + \theta_3^9 + \theta_4^9)}{1440(\theta_2 + \theta_3 + \theta_4)} \\
&\quad + \frac{1}{N^5} \frac{\xi K^8}{189\pi^2} \frac{f_{51} + \left(\frac{E'}{K'} - \frac{E}{K} \right) f_{52}}{\theta_2 + \theta_3 + \theta_4} + O\left(\frac{1}{N^7}\right),
\end{aligned} \tag{48}$$

with

$$\begin{aligned}
f_{52} &= \theta_2(-32 + 48k^2 - 78k^4 + 31k^6) + \theta_3(31 - 78k^2 + 48k^4 - 32k^6) \\
&\quad + \theta_4(31 - 15k^2 - 15k^4 + 31k^6)
\end{aligned} \tag{49}$$

$$\begin{aligned}
f_{51} &= \theta_2(-32 + 80k^2 - 38k^4 + 21k^6) + \theta_3(31 - 88k^2 + 88k^4) \\
&\quad + \theta_4(31 - 67k^2 + 25k^4 - 21k^6) \\
&\quad + \frac{21}{8} \frac{\theta_3\theta_4 k^4 (1 + k'^2)^2 + \theta_2\theta_3 k'^4 (1 + k^2)^2 + \theta_2\theta_4 (k'^2 - k^2)^2}{\theta_2 + \theta_3 + \theta_4},
\end{aligned} \tag{50}$$

and

$$\begin{aligned}
U &= -\sqrt{2} - \frac{2}{N} \frac{\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} + \frac{2}{N^3} \frac{\theta_2\theta_3\theta_4(\theta_2^9 + \theta_3^9 + \theta_4^9)}{(\theta_2 + \theta_3 + \theta_4)^2} \frac{\pi^3 \xi}{96} \\
&\quad + \frac{2}{N^5} \frac{\theta_2\theta_3\theta_4}{(\theta_2 + \theta_3 + \theta_4)^2} \frac{\xi^2 K^8}{3\pi^2} \left[U_{51} + \left(\frac{E'}{K'} - \frac{E}{K} \right) U_{52} \right] + O\left(\frac{1}{N^7}\right)
\end{aligned} \tag{51}$$

with

$$U_{52} = \theta_3(k'^2 - k^2) - \theta_2 k^4 (1 + k'^2) + \theta_4 k'^4 (1 + k^2), \tag{52}$$

$$\begin{aligned}
U_{51} &= \theta_3 \frac{23 - 64k^2 + 64k^4}{24} + \theta_2 k^4 \frac{16 + 8k^2 - k^4}{24} + \theta_4 (1 - k^2)^2 \frac{23 - 6k^2 - k^4}{24} \\
&\quad + \frac{\theta_3\theta_4 k^4 (1 + k'^2)^2 + \theta_2\theta_3 k'^4 (1 + k^2)^2 + \theta_2\theta_4 (k'^2 - k^2)^2}{12(\theta_2 + \theta_3 + \theta_4)},
\end{aligned} \tag{53}$$

where $f_\infty = -0.5 \ln 2 - 2G/\pi$ and $\theta_2, \theta_3, \theta_4$ are elliptic function

$$\theta_2 = \sqrt{\frac{2kK(k)}{\pi}}, \quad \theta_3 = \sqrt{\frac{2K(k)}{\pi}}, \quad \theta_4 = \sqrt{\frac{2k'K(k)}{\pi}}. \quad (54)$$

To calculate the specific heat we must evaluate asymptotically the sums appearing in the expression (25) for C , namely Q_1, Q_2 and Q_3 . Since the analysis follows the same general lines as in the case of the free energy and internal energy, we will not present the details of calculations and we quote here only results, namely, at the critical point $T = T_c$ the asymptotic expansion of the sums Q_1, Q_2 and Q_3 can be written as

$$Q_i = \frac{8}{\pi} \ln N + \sum_{j=0}^{\infty} \frac{q_{ij}}{N^{2j}} \quad \text{for} \quad (i = 1, 2, 3) \quad (55)$$

It is easy to see from Eqs. (24), (25), (30) and (55) that the asymptotical expansion of the specific heat, can be written as

$$C = \frac{8}{\pi} \ln N + \sum_{i=0}^{\infty} c_i / N^i. \quad (56)$$

Surprisingly, the next terms in the the asymptotical expansion of the specific heat are N^i , without multiplicative logarithms.

It is also clear that the contribution to odd (N^{-2i-1}) order in the specific heat expansion give only first term in right-hand side of the Eq. (25). Thus, we can obtained immediately that the ratio u_{2i+1}/c_{2i+1} of subdominant (N^{-2i-1}) finite-size corrections term in the internal energy and the specific heat expansions are constant, namely, $u_{2i+1}/c_{2i+1} = 1/\sqrt{2}$ as well that $u_{2i}/c_{2i} = 0$.

Several questions arise: (i) can one prove that these ratios are universal? (ii) If, so, how do such amplitudes behave in other models, for example in the three-state Potts model?

Let us now evaluate the first few terms in the specific heat expansion. After tedious calculation we obtain

$$q_{10} = \frac{8}{\pi} (C_E + \ln \frac{2^{5/2}}{\pi} - 2 \ln \theta_3) \quad (57)$$

$$q_{20} = \frac{8}{\pi} (C_E + \ln \frac{2^{5/2}}{\pi} - 2 \ln \theta_4) \quad (58)$$

$$q_{30} = \frac{8}{\pi} (C_E + \ln \frac{2^{5/2}}{\pi} - 2 \ln \theta_2) \quad (59)$$

$$q_{11} = -\frac{8K^4\xi}{9\pi^2} \left[1 + (1 - 2k^2) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \quad (60)$$

$$q_{21} = -\frac{8K^4\xi}{9\pi^2} \left[1 - 3k^2 + (1 + k^2) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \quad (61)$$

$$q_{31} = \frac{8K^4\xi}{9\pi^2} \left[2 - 3k^2 + (2 - k^2) \left(\frac{E'}{K'} - \frac{E}{K} \right) \right] \quad (62)$$

with $D_1 = E'/K' - E/K$ and

$$D_2 = \frac{9\pi^2 + 50(E'K - EK' - KK')^2 - 10\pi(1 - k^2)KK'}{20(1 - k^2)K^2K'^2} \quad (63)$$

Substituting Eqs. (28) - (30), (46), (47), (55), (57) - (62) in Eq. (25) we have finally obtained the expansions of the specific heat (C) at the critical point ($T = T_c$) up to $1/N^3$ order

$$\begin{aligned} C = & \frac{8}{\pi} \ln N + \frac{8}{\pi} \left(\ln \frac{2^{5/2}}{\pi} + C_E - \frac{\pi}{4} \right) - 4\zeta \left(\frac{\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} \right)^2 \\ & - \frac{16}{\pi} \frac{\sum_{i=2}^4 \theta_i \ln \theta_i}{\theta_2 + \theta_3 + \theta_4} - 2\sqrt{2} \frac{\theta_2\theta_3\theta_4}{\theta_2 + \theta_3 + \theta_4} \frac{1}{N} + \frac{c_2}{N^2} \\ & + \frac{1}{N^3} \frac{\pi^3 \xi}{24\sqrt{2}} \frac{\theta_2\theta_3\theta_4(\theta_2^9 + \theta_3^9 + \theta_4^9)}{(\theta_2 + \theta_3 + \theta_4)^2} + O\left(\frac{1}{N^4}\right) \end{aligned} \quad (64)$$

with

$$\begin{aligned} c_2 = & \frac{\pi^2 \xi}{6} \frac{(\theta_2^8 - \theta_3^8)\theta_2\theta_3 \ln \frac{\theta_3}{\theta_2} + (\theta_4^8 - \theta_3^8)\theta_4\theta_3 \ln \frac{\theta_3}{\theta_4} + (\theta_4^8 - \theta_2^8)\theta_4\theta_2 \ln \frac{\theta_4}{\theta_2}}{(\theta_2 + \theta_3 + \theta_4)^2} \\ & + \frac{\pi^3 \xi^2}{12} \frac{\theta_2^2\theta_3^2\theta_4^2(\theta_2^9 + \theta_3^9 + \theta_4^9)}{(\theta_2 + \theta_3 + \theta_4)^3} \\ & + \frac{2\theta_2\theta_4}{3(\theta_2 + \theta_3 + \theta_4)} \left[(\theta_2^3 - \theta_4^3)(EK_1 - E_1K) + (\theta_2^3 + \theta_4^3)KK_1 \right] \\ & - \frac{2}{9} \left[(\theta_2^4 - \theta_4^4)(EK_1 - E_1K) + (\theta_2^4 + \theta_4^4)KK_1 \right]. \end{aligned} \quad (65)$$

It is of interest to compare this finding with other results. The exact and numerical estimates [8] of the subdominant correction amplitudes for the internal energy for $\xi = 1$ are presented in Table I, which shows that the numerical values obtained by J. Salas and A. Sokal [8] are very close to our exact results.

Let us now briefly summarize our results. In this paper we derive exact finite-size corrections for several thermodynamic quantities of the ferromagnetic Ising model on the square $M \times N$ lattice with periodic boundary conditions. We find that the asymptotical expansions of f , U and C can be written as: $N(f - f_\infty) = \sum_{i=1}^{\infty} f_i/N^{2i-1}$, $U = -\sqrt{2} + \sum_{i=1}^{\infty} u_i/N^{2i-1}$ and $C = \frac{8}{\pi} \ln N + \sum_{i=0}^{\infty} c_i/N^i$. Surprisingly, the finite-size estimates of the free energy (Nf) and the internal energy (U) are well-defined odd function of N^{-1} . We find also that the ratio u_{2i+1}/c_{2i+1} of subdominant (N^{-2i-1}) finite-size corrections term in the internal energy and the specific heat expansions are constant, namely, $u_{2i+1}/c_{2i+1} = 1/\sqrt{2}$ as well that $u_{2i}/c_{2i} = 0$. The free energy and the internal energy at the critical point is calculated asymptotically for fixed $\xi = M/N$ up to N^{-5} order, and the specific heat up to N^{-3} order.

	TABLES internal energy	
	exact	numerical
N^{-3}	0.206683145...	0.206683133
N^{-5}	0.730182312347...	0.73018231235

TABLE I. Comparison exact (Eqs. (51)-(53) and numerical [8] values for subdominant finite-size corrections terms in internal energy asymptotic expansion for $\xi = 1$.

The calculation of finite-size corrections is of interest for a number of reasons. For critical models the energy and the correlation length corrections are closely related to important parameters like central charge c and the magnetic scaling dimension x_N and, besides its intrinsic interest, the treatment of lattice models for finite but large N can give important insights into finite-size studies. The large- N limit is one of the very few opportunities of having analytical results, and therefore a good quantitative understanding of finite-size effects.

The results of this paper inspire several problems for further studies: (i) can one prove that the ratio u_{2i+1}/c_{2i+1} of subdominant (N^{-2i-1}) finite-size corrections term in the internal energy and the specific heat expansions are universal? (ii) If, so, how do such amplitudes behave in other models, for example in the three-state Potts model?

This work was supported in part by the National Science Council of the Republic of China (Taiwan) under Contract No. NSC 89-2112-M-001-005.

REFERENCES

- * Electronic address: huck@phys.sinica.edu.tw.
- [1] M. E. Fisher, in *Critical Phenomena, Proceedings of the 1970 International School of Physics "Enrico Fermi"*, Course 51, edited by M. S. Green (Academic, New York, 1971).
 - [2] M. N. Barber, in *Phase Transition and Critical Phenomena*, edited by C. Domb, L. J. lebowits (Academic Press, New York, 1983), Vol. 8, p. 145.
 - [3] *Finite Size Scaling and Numerical Simulation of Statistical Systems*, edited by V. Privman (World Scientific, Singapore, 1990)
 - [4] V. Privman, P. C. Hohenberg, A. Aharony, in *Phase Transition and Critical Phenomena*, edited by C. Domb, L. J. lebowits (Academic Press, New York, 1991), Vol. 14, p. 1.
 - [5] K. Binder, Ann. Rev. Phys. Chem. **43**, 33 (1992); in *Computational Methods in Field Theory*, edited by H. Gausterer, C. P. Lang (Springer, berlin, 1992), p. 59.
 - [6] A. Ferdinand and M. E. Fisher, Phys. Rev. **185**, 832, 1969.
 - [7] A.P.Prudnikov, Yu. A Brychkov, O.I. Marichev *Integrals and series* New York, 1992.
 - [8] J. Salas and A. D. Sokal, cond-mat/9904038 and private communication.