## Charge superselection rule does not rule out pure states of subsystems to be coherent superpositions of states with different charges

Akira Shimizu\* and Takayuki Miyadera

Department of Basic Science, University of Tokyo, Komaba, Meguro-ku, Tokyo 153-8902, Japan

(Received 23 February 2001)

We consider a huge quantum system that is subject to the charge superselection rule, which requires that any pure state must be an eigenstate of the total charge. We regard some parts of the system as "subsystems"  $S_1, S_2, \dots, S_M$ , and the rest as an environment E. We assume that one does not measure anything of E, i.e., one is only interested in observables of the joint subsystem  $S \equiv S_1 + S_2 + \dots + S_M$ . We show that there exist states  $|\Phi\rangle_{tot}$  with the following properties: (i) The reduced density operator  $\text{Tr}_E (|\Phi\rangle_{tot tot} \langle \Phi|)$  is completely equivalent to a vector state  $|\varphi\rangle_S \ _S \langle \varphi|$  of S, for any gauge-invariant observable of S. (ii)  $|\varphi\rangle_S$  is a simple product of vector states of individual subsystems;  $|\varphi\rangle_S = |C^{(1)}\rangle_1 |C^{(2)}\rangle_2 \cdots$ , where  $|C^{(k)}\rangle_k$  is a vector state in  $S_k$  which is *not* an eigenstate of the charge in  $S_k$ . Furthermore, one can associate to each subsystem  $S_k$  the vector state  $|C^{(k)}\rangle_k$ and observables which are *not* necessarily gauge invariant in each subsystem, and  $|C^{(k)}\rangle_k$  is then a pure state. These results justify taking (a) superpositions of states with different charges, and (b) non-gauge-invariant operators, such as the order parameter of the breaking of the gauge symmetry, as observables, for subsystems.

PACS numbers: 03.65.Bz, 11.15.Ex, 03.75.Fi, 74.20.-z

In quantum theory, some superpositions of states are not permitted as pure states [1]. In particular, the charge superselection rule (CSSR) forbids coherent superpositions of states with different charges [1]. Namely, any pure state must be an eigenstate of the total charge  $\hat{N}_{tot}$ . However, it is customary to take such superpositions when one discusses the breaking of a gauge symmetry. Superconductors and superfluids are typical examples. If the system size V is infinite, this does not conflict with the CSSR because  $\hat{N}_{tot}$  becomes ill-defined as  $V \to \infty$ , and the CSSR becomes inapplicable. In real physical systems, however, phase transitions practically occur for finite ( $V < +\infty$ ) systems as well. In particular, phase transitions have been observed in relatively small systems, including small superconductors [2], Helium atoms in a micro bubble [3], and laser-trapped atoms [4]. The meaning of the symmetry breaking in such systems has been a subject of active research [5–8]. The purpose of this paper is to present a general discussion which justifies taking coherent superpositions of states with different charges for finite quantum systems subject to the CSSR, such as charged particles and massive bosons. Furthermore, we also justify taking non-gauge-invariant operators such as the order parameter of the breaking of the gauge symmetry, as observables of subsystems.

We consider a huge quantum system that is subject to the CSSR. We regard some parts of the system as "subsystems"  $S_1, S_2, \dots, S_M$ , and the rest as the environment E. We assume the usual situation where (i) E is much larger than the joint subsystem  $S \equiv S_1 + S_2 + \dots + S_M$ , and (ii) one is not interested in (thus, one will not measure) degrees of freedom of E, i.e, one is only interested in S (or some parts of S). The Hilbert space  $\mathcal{H}_{tot}$  of the total system is the tensor product of the Hilbert spaces of  $S_1, S_2, \dots, S_M$  and  $E; \mathcal{H}_{tot} = \mathcal{H}_S \otimes \mathcal{H}_E$ , where  $\mathcal{H}_S \equiv \mathcal{H}_1 \otimes \mathcal{H}_2 \otimes \cdots \otimes \mathcal{H}_M$ . The total charge (in some unit)  $\hat{N}_{tot}$  is the sum of the charges of  $S_1, S_2, \dots, S_M$  and E;

$$\hat{N}_{\text{tot}} = \sum_{k=1}^{M} \hat{N}_k + \hat{N}_{\text{E}}.$$
 (1)

Products of eigenfunctions  $|N_1n_1\rangle_1, \dots, |N_Mn_M\rangle_M$ , and  $|N_{\rm E}\ell\rangle_{\rm E}$  of  $\hat{N}_1, \dots, \hat{N}_M$ , and  $\hat{N}_{\rm E}$ , respectively, form complete basis sets of  $\mathcal{H}_{\rm tot}$ . Here,  $N_k$  ( $k = 1, 2, \dots, M$ ) and  $N_{\rm E}$  are eigenvalues of  $\hat{N}_k$  and  $\hat{N}_{\rm E}$ , respectively, and  $n_k$  and  $\ell$  denote additional quantum numbers.

The CSSR requires that any pure state of the total system must be an eigenstate of  $\hat{N}_{tot}$ , i.e., superposition is allowed only among states with a fixed value of the eigenvalue  $N_{tot}$  of  $\hat{N}_{tot}$ . Consider the following state that satisfies this requirement:

$$|\Phi\rangle_{\rm tot} = \sum_{N_1, n_1} \cdots \sum_{N_M, n_M} \sum_{\ell} C_{N_1 n_1}^{(1)} \cdots C_{N_M n_M}^{(M)} C_{N_{\rm S}\ell}^{({\rm E})} |N_1 n_1\rangle_1 \cdots |N_M n_M\rangle_M |N_{\rm tot} - N_{\rm S}, \ell\rangle_{\rm E},$$
(2)

where  $N_{\rm S} = \sum_k N_k$ , and the superposition coefficients are normalized as

$$\sum_{N,n} |C_{Nn}^{(k)}|^2 = \sum_{\ell} |C_{N_{\mathcal{S}\ell}}^{(\mathcal{E})}|^2 = 1.$$
(3)

For this state, the probability of finding  $N_{\text{tot}} - N_{\text{S}}$  bosons in E takes almost the same values for all  $N_{\text{S}}$  such that  $|N_{\text{S}} - \langle N_{\text{S}} \rangle| < \langle \delta N_{\text{S}}^2 \rangle^{1/2}$  [9]. This property seems natural for a huge environment. Since we assume that one will not measure degrees of freedom of E, we are interested in the reduced density operator  $\hat{\rho}_{\text{S}}$  of S, which is evaluated as

$$\hat{\rho}_{\rm S} = \operatorname{Tr}_{\rm E} \left( |\Phi\rangle_{\rm tot \ tot} \langle \Phi| \right) \\
= \sum_{N'_1, n'_1} \cdots \sum_{N'_M, n'_M} \sum_{N_1, n_1} \cdots \sum_{N_M, n_M} \delta_{N_1 + \dots + N_M, \ N'_1 + \dots + N'_M} C^{(1)}_{N'_1 n'_1} \cdots C^{(M)}_{N'_M n'_M} C^{(M)*}_{N_M n_M} \cdots C^{(1)*}_{N_1 n_1} \\
\times |N'_1 n'_1\rangle_1 \cdots |N'_M n'_M\rangle_{MM} \langle N_M n_M | \cdots |\langle N_1 n_1 |.$$
(4)

We can easily show that  $(\hat{\rho}_{\rm S})^2 \neq \hat{\rho}_{\rm S}$  unless

$$\sum_{n} |C_{Nn}^{(k)}|^2 = \delta_{N,N_0^{(k)}} \text{ for all } k$$
(5)

for some set of numbers  $N_0^{(1)}, \dots, N_0^{(M)}$ . We exclude this trivial case (where the CSSR is satisfied in each subsystem) from our consideration. Then,  $(\hat{\rho}_S)^2 \neq \hat{\rho}_S$ , and one may say that  $\hat{\rho}_S$  represents a mixed state. However, the relation  $(\hat{\rho}_S)^2 \neq \hat{\rho}_S$  only ensures that for any vector state  $|\varphi\rangle_S (\in \mathcal{H}_S)$  there exists some *operator*  $\hat{\Xi}_S$  (on  $\mathcal{H}_S$ ) for which

$$\operatorname{Tr}_{S}\left(\hat{\rho}_{S}\hat{\Xi}_{S}\right) \neq {}_{S}\langle\varphi|\hat{\Xi}_{S}|\varphi\rangle_{S}.$$
(6)

Note that such a general operator  $\hat{\Xi}_{\rm S}$  is not necessarily gauge-invariant. Hence,  $\hat{\Xi}_{\rm S}$  might not be an *observable*, which must be gauge-invariant. In fact, we first show that  $\hat{\rho}_{\rm S}$  is equivalent to a vector state  $|\varphi\rangle_{\rm S}$  for all gauge-invariant (thus physical) observables on  $\mathcal{H}_{\rm S}$ . This statement might not sound surprising because a vector state is not necessarily a pure state [1,10]. [Here, we use the precise definition of pure and mixed states [11], rather than misleading definitions such as  $\hat{\rho}^2 = \hat{\rho}$  and  $\hat{\rho}^2 \neq \hat{\rho}$ .] In fact, the equivalence of  $|\varphi\rangle_{\rm S}$  to  $\hat{\rho}_{\rm S}$  means that the vector state  $|\varphi\rangle_{\rm S}$  is a mixed state. In other words,  $|\varphi\rangle_{\rm S}$  is a vector state in a *reducible* representation of the algebra of gauge-invariant observables [10].

Actually, we first show a stronger statement:  $|\varphi\rangle_{\rm S}$  is a simple product of vector states of individual subsystems;

$$\hat{\rho}_{\rm S}$$
 is equivalent to  $|\varphi\rangle_{\rm S} = |C^{(1)}\rangle_1 |C^{(2)}\rangle_2 \cdots |C^{(M)}\rangle_M$  for any gauge-invariant observables in  $\mathcal{H}_{\rm S}$ , (7)

where  $|C^{(k)}\rangle_k$  is a coherent superposition of states with different charges;

$$|C^{(k)}\rangle_k = \sum_{N,n} C^{(k)}_{Nn} |Nn\rangle_k.$$
(8)

To see this, we recall that one will not measure degrees of freedom of E. This means that one measures only observables which take the following form;

$$\hat{A}_{\rm S} \otimes \hat{1}_{\rm E},$$
 (9)

where  $\hat{A}_{\rm S}$  is an operator on  $\mathcal{H}_{\rm S}$ , and  $\hat{1}_{\rm E}$  denotes the unity operator on  $\mathcal{H}_{\rm E}$ . Note that  $\hat{A}_{\rm S} \otimes \hat{1}_{\rm E}$  must be gauge-invariant because of the gauge invariance of the total system, hence  $\hat{A}_{\rm S}$  must also be gauge-invariant. This requires that  $N_{\rm S}$  $(=\sum_k N_k)$  should be conserved by the operation of  $\hat{A}_{\rm S}$ . Hence, the matrix elements of  $\hat{A}_{\rm S}$  should take the following form;

$${}_{1}\langle N_{1}n_{1}|\cdots_{M}\langle N_{M}n_{M}| \ \hat{A}_{S} \ |N'_{M}n'_{M}\rangle_{M}\cdots|N'_{1}n'_{1}\rangle_{1} = \delta_{N_{1}+\cdots+N_{M},N'_{1}+\cdots+N'_{M}}A^{N_{1}n_{1}\cdots\cdot N_{M}n_{M}}_{N'_{1}n'_{1}\cdots\cdot N'_{M}n'_{M}}.$$
(10)

Hence, the expectation value of  $\hat{A}_{\rm S}$  for  $\hat{\rho}_{\rm S}$  is evaluated as

$$\langle A_{\rm S} \rangle = \operatorname{Tr}_{\rm S} \left( \hat{\rho}_{\rm S} \hat{A}_{\rm S} \right)$$

$$= \sum_{N'_1, n'_1} \cdots \sum_{N'_M, n'_M} \sum_{N_1, n_1} \cdots \sum_{N_M, n_M} C^{(1)}_{N'_1 n'_1} \cdots C^{(M)}_{N'_M n'_M} C^{(M)*}_{N_M n_M} \cdots C^{(1)*}_{N_1 n_1} \delta_{N_1 + \dots + N_M, N'_1 + \dots + N'_M} A^{N_1 n_1 \dots N_M n_M}_{N'_1 n'_1 \dots N'_M n'_M}$$

$$= {}_{\rm S} \langle \varphi | \hat{A}_{\rm S} | \varphi \rangle_{\rm S},$$

$$(11)$$

and Eq. (7) is proved. The point is that Eqs. (4) and (10) contain the same Kronecker's delta,  $\delta_{N_1+\dots+N_M,N'_1+\dots+N'_M}$ . Although this factor in Eq. (4) makes  $\hat{\rho}_S$  different from  $|\varphi\rangle_S {}_S\langle\varphi|$ , the difference becomes totally irrelevant to  $\langle A_S\rangle$  because of the same factor in Eq. (10). (Recall that  $(\delta_{N_1+\dots+N_M,N'_1+\dots+N'_M})^2 = \delta_{N_1+\dots+N_M,N'_1+\dots+N'_M}$ .) Moreover, we can easily show that  $\hat{\rho}_{S-S_M} \equiv \text{Tr}_M[\hat{\rho}_S]$  is also equivalent to a vector state,  $|C^{(1)}\rangle_1|C^{(2)}\rangle_2 \cdots |C^{(M)}\rangle_{M-1}$ , where  $\text{Tr}_M$  denotes the trace operation over  $\mathcal{H}_M$ . This result is easily generalized: For states of the form of (2), its reduced density operator for any set of subsystems is completely equivalent to a vector state, which is a simple product of vector states of the individual subsystems, if one measures only gauge-invariant observables of the subsystems. Note that there is no 'entanglement' between any subsystems [12]. In general, the absence of entanglement among subsystems means the 'separability', i.e., it is possible to control quantum states of individual subsystems independently by local operations [13]. Hence, one can control the state  $|C^{(k)}\rangle_k$  of each subsystem by local operations, without perturbing the other subsystems [14].

For example, for interacting many bosons, which are confined in a large but *finite* box, one may take  $|C^{(k)}\rangle_k$  to be the 'coherent state of interacting bosons' (CSIB), which is defined by [8,15,16]

$$|\alpha_k, \mathbf{G}\rangle_k = e^{-|\alpha_k|^2/2} \sum_{N=0}^{\infty} \frac{\alpha_k^N}{\sqrt{N!}} |N, \mathbf{G}\rangle_k,$$
(12)

where  $\alpha_k = |\alpha_k| e^{i\phi_k}$  is a complex amplitude, and  $|N, G\rangle_k$  denotes the ground state that has *exactly* N bosons, which we call the 'number state of interacting bosons' (NSIB) [8,15]. One may also take  $|C^{(l)}\rangle_l$  as  $|\alpha_l, G\rangle_l$ , where  $\alpha_l = |\alpha_l| e^{i\phi_l}$ . Then,  $\hat{\rho}_s$  is equivalent to

$$|\varphi\rangle_{\rm S} = \cdots |\alpha_k, {\rm G}\rangle_k \cdots |\alpha_l, {\rm G}\rangle_l \cdots$$
(13)

This state has an almost definite value of the relative phase  $\phi_{kl}$  between two condensates in  $S_k$  and  $S_l$ . To see this, we may define the operator (acting on  $\mathcal{H}_k \otimes \mathcal{H}_l$ ) corresponding to  $e^{i\phi_{kl}}$  by

$$\widehat{e^{i\phi_{kl}}} \equiv \widehat{e^{i\phi_k}} \ (\widehat{e^{i\phi_l}})^{\dagger}. \tag{14}$$

Here,  $\hat{e^{i\phi_k}}$ , acting on  $\mathcal{H}_k$ , is not an exponential of some phase operator  $\hat{\phi}_k$ , but is defined by

$$\widehat{e^{i\phi_k}} \equiv (\hat{b}_{0k}^{\dagger}\hat{b}_{0k} + 1)^{-1/2}b_{0k}, \tag{15}$$

where  $\hat{b}_{0k}$  denotes the operator  $\hat{b}_0$ , which is defined in Ref. [15] as a nonlinear function of free operators, for  $S_k$ . This operator is a linear combination of the cosine and sine operators of Ref. [15]. In the same way,  $\hat{e^{i\phi_l}}$ , acting on  $\mathcal{H}_l$ , is defined using  $\hat{b}_{0l}$ , which is  $\hat{b}_0$  for  $S_l$ . By similar calculations as in Ref. [15], we can show that  $\hat{e^{i\phi_l}}$ ,  $(\hat{e^{i\phi_l}})^{\dagger}$ , and  $\hat{e^{i\phi_{kl}}}$  have almost definite values, and we can regard  $\phi_{kl} \simeq \phi_k - \phi_l$ . Hence, the state (13) has the almost definite value of the relative phase. It should not be confused with states of the following form, which have been frequently discussed in the literature [17];

$$|\varphi_{\text{ent}}\rangle_{\text{S}} = \sum_{N} C_{N} \sum_{N_{k}} \cdots e^{iN_{k}\phi_{k}} R_{N_{k}} |N_{k}, \text{G}\rangle_{k} \cdots e^{i(N-N_{k})\phi_{l}} R_{N-N_{k}} |N-N_{k}, \text{G}\rangle_{l} \cdots,$$
(16)

where  $C_N$  is a complex coefficient and R's are real ones. This state also has an almost definite relative phase if R's are appropriately taken. However, since  $S_k$  and  $S_l$  are strongly entangled in this state [12], one cannot control the state of  $S_k$  by local operations on  $S_k$ , without perturbing the state of  $S_l$ . In contrast, this is possible for the simple product state  $|\varphi\rangle_S$ . Moreover, we now show that  $|\varphi\rangle_S$  allows us to treat observables of individual subsystems separately.

As exemplified by  $e^{i\phi_{kl}}$  above,  $\hat{A}_{S}$  is generally a sum of products of operators of subsystems [18];

$$\hat{A}_{\rm S} = \sum \hat{A}_k \hat{A}_l \cdots, \tag{17}$$

where  $\hat{A}_k$  ( $\hat{A}_l$ ) is an operator on  $\mathcal{H}_k$  ( $\mathcal{H}_l$ ), and the sum is over combinations of  $\hat{A}_k \hat{A}_l \cdots$ . For the product state  $|\varphi\rangle_{\rm S}$ , the expectation value of  $\hat{A}_{\rm S}$  is simply evaluated from the expectation values of individual subsystems as

$${}_{\mathrm{S}}\langle\varphi|\hat{A}_{\mathrm{S}}|\varphi\rangle_{\mathrm{S}} = \sum_{k}\langle C^{(k)}|\hat{A}_{k}|C^{(k)}\rangle_{k} \ _{l}\langle C^{(l)}|\hat{A}_{l}|C^{(l)}\rangle_{l}\cdots.$$
(18)

From Eqs. (7) and (18), we can consider each subsystem separately from the other subsystems. Namely, for subsystem  $S_k$ , we can consider that its state is the vector state  $|C^{(k)}\rangle_k$ , and that  $\hat{A}_k$  is one of its observables. When the expectation value of  $\hat{A}_S$  for the set of subsystems is necessary, it can be evaluated from the expectation values of individual subsystems, as Eq. (18). Note that the gauge invariance of  $\hat{A}_S$  does not require the gauge invariance of  $\hat{A}_k$ 

of each subsystem: It rather requires that only gauge-invariant combinations of  $\hat{A}_k \hat{A}_l \cdots$  should appear in the sum. [Namely, each  $N_k$  can vary by the operation of  $\hat{A}_k$ , whereas  $N_S (= \sum_k N_k)$  is conserved by the operation of  $\hat{A}_k \hat{A}_l \cdots$ .] For example,  $e^{i\phi_{kl}}$  is gauge invariant, whereas neither  $e^{i\phi_k}$  nor  $e^{i\phi_l}$  is. Hence, by considering  $\hat{A}_k$  as an observable of  $S_k$ , we can include non-gauge invariant operators into the set of observables in  $\mathcal{H}_k$ , with the restriction that among various combinations of  $\hat{A}_k \hat{A}_l \cdots$  we must take only gauge-invariant ones as physical combinations: Results for other combinations should be discarded. This formulation gives correct results for all gauge-invariant (thus physical) combinations. The vector state  $|C^{(k)}\rangle_k$  then becomes a pure state of an irreducible representation of the algebra of observables of  $S_k$ , because non-gauge invariant observables are now included in the algebra [10]. We have thus arrived at non-gauge invariant observables and a pure state which is a superposition of states with different charges, for each subsystem.

The restriction that we must take only gauge-invariant ones among various combinations of  $\hat{A}_k \hat{A}_l \cdots$  has the following physical meaning: If  $\hat{A}_k$  is a non-gauge invariant observable of  $S_k$ , then its value can only be defined relative to some reference observable  $\hat{A}_l$  of some reference system  $S_l$ . When  $|\varphi\rangle_S$  takes the form of Eq. (13), for example, we can consider that  $S_k$  is in a pure state  $|\alpha_k, G\rangle_k$ , and that the phase factor  $e^{i\phi_k}$  is one of observables on  $\mathcal{H}_k$ . However, like the classical phase factor, the quantum phase factor  $\widehat{e^{i\phi_k}}$  can only be defined relative to some reference (such as  $(\widehat{e^{i\phi_l}})^{\dagger}$  of  $S_l$ ). In contrast to the position observable, which also requires a reference system but it can be any system composed of any material, the reference system of the phase factor should contain particles of the same kind as the particles in  $S_k$ . Hence, the system of interest  $(S_k)$  and the reference system  $(S_l)$  should be subsystems of a larger system of the same kind of particles. When the larger system contains a huge system(s), of which we are not interested in, we may call it an environment E, and the present model is applicable. When one is only interested in  $S_k$ , the reference system  $S_l$  may be considered as a part of the external systems, which include an apparatus with which one measures  $\phi_{kl}$ . In this case, the reference system S<sub>l</sub> can be regarded as a part of the measuring apparatus. (In the case of measurement of the position, for example, a material which defines the origin of the position coordinate can be considered as a part of the measuring apparatus of the position.) Although  $e^{i\phi_k}$  is not gauge invariant, results of measurement is gauge invariant because  $e^{i\phi_k}$   $(e^{i\phi_l})^{\dagger}$  is gauge invariant. In general terms, although results of any physical measurements must be gauge invariant, it does not necessarily mean that any observables of a subsystem must be gauge-invariant, because the gauge invariance of  $\hat{A}_k \hat{A}_l$  (which ensures the gauge invariance of results of measurements) does not necessarily mean the gauge invariance of  $\hat{A}_k$ .

Finally, we present some significant applications of the present results. First, Our results give a natural interpretation of the order parameter O of a finite system, which can exhibit the breaking of the gauge symmetry in the infinite-volume limit. Although one might suspect that  $\hat{O}$  must be excluded from observables since it is not gauge invariant, our results allow one to include it among observables as one of  $\hat{A}_k$ 's. Second, one might also suspect that the CSSR would forbid a pure state which has a finite expectation value of  $\hat{O}$  because its expectation value vanishes for any eigenstate of  $\hat{N}$ . However, our results show that such a pure state is allowed as a state of a subsystem,  $|C^{(k)}\rangle_k$ , which is not entangled with states of the other subsystems The absence of entanglement allows the preparation of such a pure state of a subsystem without perturbing the other subsystems (with perturbing the environment only). For interacting many bosons, for example, the order parameter of the condensation is usually taken as the field operator  $\dot{\psi}$  of bosons. The present work justifies taking such a non-gauge-invariant operator as an observable of a subsystem. The expectation value of  $\hat{\psi}$  is finite only for superpositions of states with different numbers of bosons. The present work justifies taking such a superposition as a pure state of a subsystem. More concretely, the expectation value of  $\hat{\psi}$  is finite for the CSIB,  $\langle \alpha, G | \hat{\psi} | \alpha, G \rangle = \mathcal{O}(1)$ , whereas it vanishes for the NSIB,  $\langle N, G | \hat{\psi} | N, G \rangle = 0$  [8,15]. Although both the NSIB and CSIB have the off-diagonal long-range order [8,15], the gauge symmetry is broken, in the sense that  $\langle \hat{\psi} \rangle = \mathcal{O}(1)$ , only for the CSIB [19]. Since the state vectors of the NSIB and CSIB are quite different, they have different properties. The most striking difference is that the NSIB decoheres much faster than the CSIB when bosons have a finite probability of leakage into a huge environment, whose boson density is zero [8]. Moreover, it was shown that the CSIB has the 'cluster property,' which ensures that fluctuations of any intensive variables are negligible, in consistency with thermodynamics [20]. Although the CSIB may look against the CSSR, the present work justifies taking it as a pure state of a subsystem, which, unlike the NSIB, is robust, symmetry breaking, and consistent with thermodynamics. In particular, generalizing Eq. (13), one can take CSIB's for all subsystems. In this case, the joint subsystem S is in a pure state in which each subsystem is in a CSIB. Namely, the state vector of S behaves locally as a CSIB, which is robust, symmetry breaking, and consistent with thermodynamics. Moreover, unlike Eq. (16), one can controle states of individual subsystems independently by local operations. These justify macroscopic theories such as the Ginzburg-Landau theory and the Gross-Pitaevskii theory, which assume, sometimes implicitly, that the order parameters can be defined locally, and that their fluctuations are negligible, and that the state vectors are robust against weak perturbation from environments, and that local operations do not cause global changes. We consider

that such a locally-CSIB state should be (a good approximation to) a quantum state of real physical systems at low temperature, except for extreme cases such as bosons in a perfectly-closed box at a ultra-low temperature. Finally, we remark that we have not assumed that the particles in the environment are in (or, not in) a condensed phase.

We thank H. Tasaki and M. Ueda for discussions.

- Electronic address: shmz@ASone.c.u-tokyo.ac.jp
- [1] See, e.g., R. Haag, Local Quantum Physics (Springer, Berlin, 1992).
- [2] See, e.g., M. Tinkham, Superconductivity 2nd ed. (McGraw-Hill, New York, 1996).
- [3] E. G. Syskakis, F. Pobell and H. Ullmaier, Phys. Rev. Lett. 55 (1985) 2964.
- M. H. Anderson et al., Science 269 (1995) 198; C. C. Bradley et al., Phys. Rev. Lett. 75 (1995) 1687; K. B. Davis et al., Phys. Rev. Lett. 75 (1995) 3969.
- [5] M. R. Andrews et al., Science 275 (1997) 637.
- [6] J. Javanainen and S. M. Yoo, Phys. Rev. Lett. 76 (1996) 161.
- [7] S. M. Barnett, K. Burnett and J. A. Vaccaro, J. Res. Natl. Inst. Stand. Technol. 101, 593 (1996).
- [8] A. Shimizu and T. Miyadera, Phys. Rev. Lett. 85, 688 (2000).
- [9] We assume that the product  $\prod_k C_{N_k n_k}^{(k)}$  is finite only when  $N_S \ll N_{tot}$ . Since E is assumed to be much larger than S, states with low energies would satisfy this condition.
- [10] A vector state is a state represented by a vector (more precisely, ray) in a Hilbert space, which is not necessarily an irreducible representation: Since the Hilbert space consists of many charge sectors, it becomes an irreducible representation of the algebra of observables only when the observables include non-gauge invariant ones (such as a field operator), which connect different sectors [1].
- [11] Let us denote a quantum state symbolically by  $\omega$  and the expectation value of an observable A by  $\omega(A)$ . A state  $\omega$  is called *mixed* iff there exist states  $\omega_1$  and  $\omega_2 \ (\neq \omega_1)$ , and a positive number  $\lambda \ (0 < \lambda < 1)$ , such that  $\omega(A) = \lambda \omega_1(A) + (1 \lambda) \omega_2(A)$  for any *local* observable A. Otherwise,  $\omega$  is called a pure state. See, e.g., Ref. [1].
- [12] Here, the entanglement is defined by the increase of the von Neumann entropy for the reduced density operator which is obtained by tracing over a part of subsystems.
- [13] See, e.g., A. Peres, Phys. Scripta T76, 52 (1998) (quant-ph/9707026).
- [14] On the other hand, the environment E might be perturbed by the control of a subsystem because  $|\Phi\rangle_{tot}$  is not a simple product of  $\hat{\rho}_{\rm S}$  and a state vector of E. However, we do not measure effects of such perturbations on E because we assume that we do not measure anything of E.
- [15] A. Shimizu and J. Inoue, Phys. Rev. A 60, 3204 (1999).
- [16] Although this relation is the same as the relation between the corresponding states of free bosons,  $|\alpha, G\rangle$  is a complicated wave function because  $|N, G\rangle$  is complicated [8].
- [17] For example, the state (16) appears for two condensates, each of which initially has a definite number of bosons, when an interference pattern is developed and observed. See, e.g., J. I. Cirac et al., Phys. Rev. A54, R3714 (1996).
- [18] If necessary, one can make  $\hat{A}_{\rm S}$  self-adjoint by adding its conjugate operator.
- [19] This is a generic property of quantum phase transitions. For example, if we treat a ferromagnetic Ising spin system as a quantum spin system, then two ferromagnetic states,  $|\uparrow\uparrow\cdots\rangle$  and  $|\downarrow\downarrow\cdots\rangle$ , correspond to CSIB's, whereas their superpositions  $(|\uparrow\uparrow\cdots\rangle\pm|\downarrow\downarrow\cdots\rangle)/\sqrt{2}$  correspond to NSIB's: Although all these states have a long-range order, the up-down symmetry is broken only for the former states, in the sense that  $\langle M_z \rangle \neq 0$ .
- [20] A. Shimizu and T. Miyadera, cond-mat/0009258.