

# Kronecker's Double Series and Exact Asymptotic Expansion for Free Models of Statistical Mechanics on Torus

E.V. Ivashkevich<sup>1,\*</sup>, N.Sh. Izmailian<sup>2,3</sup> and Chin-Kun Hu<sup>2,4</sup>

<sup>1</sup>*Bogoliubov Laboratory of Theoretical Physics, J.I.N.R., Dubna 141980, Russia*

<sup>2</sup>*Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan*

<sup>3</sup>*Yerevan Physics Institute, Alikhanian Br. 2, Yerevan 375036, Armenia*

<sup>4</sup>*Department of Physics, National Dong Hwa University, Hualien 97401, Taiwan*

(February 14, 2019)

## Abstract

For the free models of statistical mechanics on torus, exact asymptotic expansion of the free energy in the vicinity of the critical point is found. It is shown that there is direct relation between the terms of the expansion and the Kronecker's double series. The latter can be expressed in terms of the elliptic  $\theta$ -functions in all orders of the asymptotic expansion.

05.50.+q, 05.70.Jk, 64.60.Cn.

## I. INTRODUCTION

It is well known that the singularities in thermodynamic functions associated with a critical point occur only in the thermodynamic limit when dimension  $L$  of the system under consideration tends to infinity. In such a limit, the critical fluctuations are correlated over a distance of the order of correlation length  $\xi_{\text{corr}}$  which may be defined as the length scale governing the exponential decay of correlation functions. Besides these two fundamental lengths,  $L$  and  $\xi_{\text{corr}}$ , there is also the microscopic length of interactions  $a$ . Thermodynamic quantities thus may in principle depend on the dimensionless ratios  $\xi_{\text{corr}}/L$  and  $a/L$ . The Finite-Size Scaling (FSS) hypothesis [1] assumes that in the scaling interval, for temperatures so close to the critical point that  $a \ll \xi_{\text{corr}} \sim L$ , the microscopic length drops out and the behaviour of any thermodynamic quantity can be described in terms of the universal scaling function of the scaling variable  $t = \xi_{\text{corr}}/L$ . However, non-universal corrections to FSS do exist. These sometimes can be viewed as asymptotic series in powers of  $a/L$ .

Two-dimensional models have long served as a proving ground in attempts to understand critical behavior and to test the general ideas of FSS. There are very few two-dimensional models that have been solved exactly [2] notably the so-called free models of statistical mechanics which can be treated in terms of non-interacting quasi-particles on the lattice. These are best represented by the Ising model which is known to be equivalent to the model of free lattice fermions [2,3] and its boson analog often referred to as Gaussian model.

For all free models exact asymptotic expansion of the free energy on an infinite cylinder of circumference  $L$  can easily be obtained by direct application of Euler-Maclaurin summation formula [4,5]. However, derivation of such an expansion on a torus of area  $S$  and aspect ratio  $\xi$  is much more difficult problem. In the case of Ising model such an expansion has been studied by Ferdinand and Fisher [6]. Starting with Onsager's explicit expression for the partition function of Ising model [7] they have calculated first two leading terms  $f_{\text{bulk}}$  and  $f_0(\xi)$  of the expansion

$$F_{T=T_c}(\xi, S) = f_{\text{bulk}} S + f_0(\xi) + \sum_{p=1}^{\infty} f_p(\xi) S^{-p} \quad (1)$$

Their calculations have been pushed forward to get two next sub-leading terms of the expansion [8,9].

It is the purpose of this paper to derive *all* terms of the exact asymptotic expansion of the logarithm of the partition function on torus for a class of free exactly solvable models of statistical mechanics. Our approach is based on an intimate relation between the terms of the asymptotic expansion and the so-called Kronecker's double series. Besides the aesthetic appeal of the exact expansion there is also physical motivation to study non-universal corrections to FSS. The problem is that in numerical simulations of lattice models one usually studies relatively small lattices. Therefore, to compare the results of high precision numerical simulations with the theoretical predictions one cannot neglect sub-leading corrections to FSS [10]. Non-universal terms in the asymptotic expansion also provide an important information about the structure of irrelevant operators in conformal field theory [11].

## II. FREE MODELS OF STATISTICAL MECHANICS

In this section we formulate three basic models of statistical mechanics: Ising model, dimer model and Gaussian model. These models are often referred to as "free" since they were shown to be equivalent to free fermions or free bosons on the lattice. Partition functions of all these models can be written in terms of the only object — the partition function with twisted boundary conditions  $Z_{\alpha,\beta}(\mu)$ . Exact asymptotic expansion of the later will be our main objective in the subsequent section.

### A. Ising model

Ising model is usually formulated as follows. Consider a planar square lattice of size  $M \times N$  with periodic boundary conditions, i.e. torus. To each site  $(m, n)$  of the torus a spin variable is ascribed,  $s_{mn}$ , with two possible values:  $+1$  or  $-1$ . Two nearest neighbor spins,

say  $s_{mn}$  and  $s_{m n+1}$  contribute a term  $-J s_{mn} s_{m n+1}$  to the Hamiltonian, where  $J$  is some fixed energy. So, the Hamiltonian is simply the sum of all such terms, one for each edge of the lattice

$$H(s) = -J \sum_{n=0}^{N-1} \sum_{m=0}^{M-1} (s_{mn} s_{m+1 n} + s_{mn} s_{m n+1}) \quad (2)$$

The partition function of the Ising model is given by the sum over all spin configurations on the lattice

$$Z_{\text{Ising}}(J) = \sum_{\{s\}} e^{-H(s)}$$

It is convenient to set up another parameterizations of the interaction constant  $J$  in terms of the mass variable  $\mu = \ln \sqrt{\text{sh } 2J}$ . Critical point corresponds to the massless case  $\mu = 0$ .

Onsager's explicit expression for the partition function of the Ising model can be written as [7]

$$Z_{\text{Ising}}(\mu) = \frac{1}{2} (\sqrt{2}e^\mu)^{MN} \left\{ Z_{\frac{1}{2}, \frac{1}{2}}(\mu) + Z_{0, \frac{1}{2}}(\mu) + Z_{\frac{1}{2}, 0}(\mu) + Z_{0, 0}(\mu) \right\} \quad (3)$$

where we have introduced the partition function with twisted boundary conditions

$$Z_{\alpha, \beta}^2(\mu) = \prod_{n=0}^{N-1} \prod_{m=0}^{M-1} 4 \left[ \sin^2 \left( \frac{\pi(n+\alpha)}{N} \right) + \sin^2 \left( \frac{\pi(m+\beta)}{M} \right) + 2 \text{sh}^2 \mu \right] \quad (4)$$

Here  $\alpha = 0$  corresponds to the periodic boundary conditions for the underlying free fermion in the  $N$ -direction while  $\alpha = \frac{1}{2}$  stands for anti-periodic boundary conditions. Similarly  $\beta$  controls boundary conditions in  $M$ -direction. With the help of the identity [12]

$$4 |\text{sh}(M\omega + i\pi\beta)|^2 = 4 \left[ \text{sh}^2 M\omega + \sin^2 \pi\beta \right] = \prod_{m=0}^{M-1} 4 \left[ \text{sh}^2 \omega + \sin^2 \left( \frac{\pi(m+\beta)}{M} \right) \right]$$

the partition function with twisted boundary conditions  $Z_{\alpha, \beta}$  can be transformed into simpler form

$$Z_{\alpha, \beta}(\mu) = \prod_{n=0}^{N-1} 2 \left| \text{sh} \left[ M\omega_\mu \left( \frac{\pi(n+\alpha)}{N} \right) + i\pi\beta \right] \right| \quad (5)$$

where lattice dispersion relation has appeared

$$\omega_\mu(k) = \text{Arsh} \sqrt{\sin^2 k + 2 \text{sh}^2 \mu} \quad (6)$$

This is the functional relation between energy  $\omega_\mu$  and momentum  $k$  of a free quasi-particle on the planar square lattice.

## B. Dimer model

A "dimer" is a two-atom molecule. When drawn on a lattice it covers two adjacent sites of the lattice and the bond that joins them. The "dimer problem" is to determine the number of ways of covering of a given lattice with dimers, so that all sites are occupied and no two dimers overlap. If we consider planar square lattice of size  $2M \times 2N$  wrapped on a torus then the number of dimers must be  $2MN$  and the number of distinct dense coverings of the lattice (the partition function) has been calculated by Kasteleyn and Fisher [13,14]. This can be expressed in terms of the same partition function with twisted boundary conditions Eq.(5) as

$$Z_{\text{Dimer}} = \frac{1}{2} \left\{ Z_{\frac{1}{2}, \frac{1}{2}}^2(0) + Z_{0, \frac{1}{2}}^2(0) + Z_{\frac{1}{2}, 0}^2(0) - Z_{0, 0}^2(0) \right\} \quad (7)$$

First two leading terms of the asymptotic expansion of this partition function has been obtained by Ferdinand [15]. Let us also mention that dimers are always in the critical point and has no phase transition.

## C. Gaussian model

Let us turn now to a boson analog of Ising model which is often referred to as Gaussian model. Again, we consider planar square lattice of size  $N \times M$  wrapped on a torus. To each site  $(m, n)$  of the lattice we assign a continuous variable  $x_{mn}$ . The Hamiltonian of the model is

$$H(x) = -J \sum_{n=0}^{N-1} \sum_{m=0}^{M-1} \left( x_{mn} x_{m+1n} + x_{mn} x_{mn+1} - 2x_{mn}^2 \right) \quad (8)$$

The partition function of the model can be written as

$$Z(J) = \int_{\mathbf{R}^{MN}} e^{-H(x)} d\sigma(x)$$

If the measure  $d\sigma(x)$  in the phase space  $\mathbf{R}^{MN}$  is chosen to be Gaussian

$$d\sigma_{\text{Gauss}}(x) = \pi^{-MN/2} \prod_{n=0}^{N-1} \prod_{m=0}^{M-1} e^{-x_{mn}^2} dx_{mn}$$

the integration can be done explicitly and partition function of the free boson model can be written in terms of the partition function with twisted boundary conditions Eq.(5) and parameterization  $J^{-1} = 4 \operatorname{ch}^2 \mu$  as

$$Z_{\text{Gauss}}(\mu) = \left(\sqrt{2} \operatorname{ch} \mu\right)^{MN} \left[ Z_{0,0}(\mu) \right]^{-1} \quad (9)$$

This model exhibit phase transition at the point  $\mu_c = 0$  where the partition function is divergent. This is due to the presence of zero mode. In other words, due to the symmetry transformation  $x_{mn} \rightarrow x_{mn} + \text{const}$  which leave the Hamiltonian (8) invariant. Correlation functions of disorder operator in this model have been studied by Sato, Miwa and Jimbo [16].

The reason why this model is often considered as boson analog of Ising model is that one can choose another measure in the phase space which makes this model equivalent to the Ising model considered above, namely

$$d\sigma_{\text{Ising}}(x) = 2^{-MN} \prod_{n=0}^{N-1} \prod_{m=0}^{M-1} [\delta(x_{mn} - 1) + \delta(x_{mn} + 1)] dx_{mn}$$

where  $\delta$ 's are Dirac  $\delta$ -functions. With such a definition the variables  $x_{mn}$  can actually take only two values:  $+1$  or  $-1$ , so that  $x_{mn}^2 = 1$ . In this case integration can be replaces by summation over discrete values of  $x_{mn} = \pm 1$  and the Hamiltonian (8) coincides with the Hamiltonian of the Ising model (2) up to a constant.

### III. ASYMPTOTIC EXPANSION IN TERMS OF KRONECKER'S DOUBLE SERIES

In the previous section it was shown that partition functions of three basic free models of statistical mechanics can be expressed in terms of the only object: the partition function with twisted boundary condition  $Z_{\alpha,\beta}(\mu)$ . In this section we shall obtain exact asymptotic  $1/S$ -expansion of the partition function in the vicinity of the critical point. For reader's convenience, all the technical details of our calculations and the definitions of the special functions are summarized in the appendices attached to the paper.

First of all, let us mention the symmetry properties of the partition function  $Z_{\alpha,\beta}(\mu)$ . From its definition (5) one can easily verify that it is even and periodic with respect to its arguments  $\alpha$  and  $\beta$

$$\begin{aligned} Z_{\alpha,\beta}(\mu) &= Z_{\alpha,-\beta}(\mu) = Z_{-\alpha,\beta}(\mu) \\ Z_{\alpha,\beta}(\mu) &= Z_{1+\alpha,\beta}(\mu) = Z_{\alpha,1+\beta}(\mu) \end{aligned}$$

These imply that twist angles  $\alpha$  and  $\beta$  can be taken from the interval  $[0, 1]$ . Then, one can note that for all twists  $(\alpha, \beta) \neq (0, 0)$  the partition function  $Z_{\alpha,\beta}(\mu)$  is even with respect to its mass argument  $\mu$ . Hence, in the vicinity of the critical point ( $\mu = 0$ ) we have

$$Z_{\alpha,\beta}(\mu) = Z_{\alpha,\beta}(0) + \frac{\mu^2}{2!} Z''_{\alpha,\beta}(0) + \dots \quad (\alpha, \beta) \neq (0, 0) \quad (10)$$

The only exception is the point where both  $\alpha$  and  $\beta$  are equal to zero. This case has to be treated separately since at this point the partition function turns to zero. As a result we have

$$Z_{0,0}(\mu) = \mu Z'_{0,0}(0) + \frac{\mu^3}{3!} Z'''_{0,0}(0) + \dots \quad (\alpha, \beta) = (0, 0) \quad (11)$$

In what follows notation  $Z_{\alpha,\beta}(\mu)$  will imply  $(\alpha, \beta) \neq (0, 0)$ .

### A. Asymptotic Expansion of $Z_{\alpha,\beta}(0)$

Considering logarithm of the partition function with twisted boundary conditions Eq.(5) we note that it can be transformed as

$$\ln Z_{\alpha,\beta}(0) = M \sum_{n=0}^{N-1} \omega_0 \left( \frac{\pi(n+\alpha)}{N} \right) + \sum_{n=0}^{N-1} \ln \left| 1 - e^{-2[M\omega_0 \left( \frac{\pi(n+\alpha)}{N} \right) + i\pi\beta]} \right| \quad (12)$$

Note, that the second sum here vanishes in the formal limit  $M \rightarrow \infty$  when the torus turns into infinitely long cylinder of circumference  $N$ . The first sum gives the logarithm of the partition function with twisted angle  $\alpha$  on that cylinder. Its asymptotic expansion can be found with the help of the Euler-Maclaurin summation formula (Appendix A)

$$M \sum_{n=0}^{N-1} \omega_0\left(\frac{\pi(n+\alpha)}{N}\right) = \frac{S}{\pi} \int_0^\pi \omega_0(x) dx - \pi \lambda \xi B_2^\alpha - 2\pi \xi \sum_{p=1}^{\infty} \left(\frac{\pi^2 \xi}{S}\right)^p \frac{\lambda_{2p}}{(2p)!} \frac{B_{2p+2}^\alpha}{2p+2} \quad (13)$$

Here  $B_p^\alpha$  are so called Bernoulli polynomials. We have also used the symmetry property,  $\omega_0(k) = \omega_0(\pi - k)$ , of the lattice dispersion relation (6) and its Taylor expansion

$$\omega_0(k) = k \left( \lambda + \sum_{p=1}^{\infty} \frac{\lambda_{2p}}{(2p)!} k^{2p} \right) \quad (14)$$

where  $\lambda = 1$ ,  $\lambda_2 = -2/3$ ,  $\lambda_4 = 4$ , etc. In what follows we shall not use the special values of these coefficients assuming the possibility for generalizations.

The second term in Eq.(12) we may transform as

$$\begin{aligned} & \sum_{n=0}^{N-1} \ln \left| 1 - e^{-2[M\omega_0(\frac{\pi(n+\alpha)}{N}) + i\pi\beta]} \right| = \\ & = - \operatorname{Re} \sum_{m=1}^{\infty} \frac{1}{m} \left\{ \sum_{n=0}^{[N/2]-1} e^{-2m[M\omega_0(\frac{\pi(n+\alpha)}{N}) + i\pi\beta]} \right. \\ & \quad \left. + \sum_{n=0}^{N-[N/2]-1} e^{-2m[M\omega_0(\frac{\pi(n+1-\alpha)}{N}) + i\pi\beta]} \right\} \end{aligned} \quad (15)$$

The argument of the first exponent can be expanded in powers of  $1/S$  if we replace the lattice dispersion relation  $\omega_0(x)$  by its expansion (14)

$$\exp \left\{ -2\pi m [\lambda \xi (n + \alpha) + i\beta] - 2\pi m \xi \sum_{p=1}^{\infty} \frac{\lambda_{2p}}{(2p)!} \left(\frac{\pi^2 \xi}{S}\right)^p (n + \alpha)^{2p+1} \right\}$$

Taking into account the relation between moments and cumulants (Appendix B) we obtain asymptotic expansion of the first exponent itself in powers of  $1/S$

$$\begin{aligned} e^{-2m[M\omega_0(\frac{\pi(n+\alpha)}{N}) + i\pi\beta]} &= e^{-2\pi m [\lambda \xi (n + \alpha) + i\beta]} \\ & - 2\pi m \xi \sum_{p=1}^{\infty} \left(\frac{\pi^2 \xi}{S}\right)^p \frac{\Lambda_{2p}}{(2p)!} (n + \alpha)^{2p+1} e^{-2\pi m [\lambda \xi (n + \alpha) + i\beta]} \end{aligned}$$

where we have also introduced differential operators  $\Lambda_{2p}$  which can be expressed via coefficients  $\lambda_{2p}$  of the expansion of the lattice dispersion relation (14)

$$\begin{aligned} \Lambda_2 &= \lambda_2 \\ \Lambda_4 &= \lambda_4 + 3\lambda_2^2 \frac{\partial}{\partial \lambda} \end{aligned}$$

$$\begin{aligned}
\Lambda_6 &= \lambda_6 + 15\lambda_4\lambda_2 \frac{\partial}{\partial\lambda} + 15\lambda_2^3 \frac{\partial^2}{\partial\lambda^2} \\
&\vdots \\
\Lambda_p &= \sum_{r=1}^p \sum \left( \frac{\lambda_{p_1}}{p_1!} \right)^{k_1} \cdots \left( \frac{\lambda_{p_r}}{p_r!} \right)^{k_r} \frac{p!}{k_1! \cdots k_r!} \frac{\partial^k}{\partial\lambda^k}
\end{aligned}$$

Here summation is over all positive numbers  $\{k_1 \dots k_r\}$  and different positive numbers  $\{p_1, \dots, p_r\}$  such that  $p_1 k_1 + \dots + p_r k_r = p$  and  $k = k_1 + \dots + k_r - 1$ .

The expansion for the second exponent in Eq.(15) can be obtained along the same lines by substitution:  $\alpha \rightarrow 1 - \alpha$ . Plugging the expansion of both of the exponents back into Eq.(15) we obtain

$$\begin{aligned}
&\sum_{n=0}^{N-1} \ln \left| 1 - e^{-2[M\omega_0(\frac{\pi(n+\alpha)}{N})+i\pi\beta]} \right| = \\
&= -\operatorname{Re} \sum_{m=1}^{\infty} \frac{1}{m} \left\{ \sum_{n=0}^{[N/2]-1} e^{-2\pi m[\lambda\xi(n+\alpha)+i\beta]} + \sum_{n=0}^{N-[N/2]-1} e^{-2\pi[\lambda\xi(n+1-\alpha)+i\beta]} \right\} \\
&+ 2\pi\xi \sum_{p=1}^{\infty} \left( \frac{\pi^2\xi}{S} \right)^p \frac{\Lambda_{2p}}{(2p)!} \operatorname{Re} \sum_{m=1}^{\infty} \left\{ \sum_{n=0}^{[N/2]-1} (n+\alpha)^{2p+1} e^{-2\pi m[\lambda\xi(n+\alpha)+i\beta]} \right. \\
&\quad \left. + \sum_{n=0}^{N-[N/2]-1} (n+1-\alpha)^{2p+1} e^{-2\pi m[\lambda\xi(n+1-\alpha)+i\beta]} \right\}
\end{aligned}$$

In all these series summation over  $n$  can be extended to infinity. The resulting errors are exponentially small and does not affect our asymptotic expansion in any finite power of  $1/S$ . The key point of our analysis is the observation that all the series that appear here can be obtained by resummation of either elliptic theta function,  $\theta_{\alpha,\beta}(\tau)$  (Appendix C), or Kronecker's double series,  $K_p^{\alpha,\beta}(\tau)$  (Appendix D). Namely, with the help of the identities (C4) and (D6) we obtain

$$\begin{aligned}
\sum_{n=0}^{N-1} \ln \left| 1 - e^{-2[M\omega_0(\frac{\pi(n+\alpha)}{N})+i\pi\beta]} \right| &= \ln \left| \frac{\theta_{\alpha,\beta}(i\lambda\xi)}{\eta(i\lambda\xi)} \right| + \pi\lambda\xi B_2^\alpha \\
&- 2\pi\xi \sum_{p=1}^{\infty} \left( \frac{\pi^2\xi}{S} \right)^p \frac{\Lambda_{2p}}{(2p)!} \frac{\operatorname{Re} K_{2p+2}^{\alpha,\beta}(i\lambda\xi) - B_{2p+2}^\alpha}{2p+2}
\end{aligned} \tag{16}$$

Substituting Eqs.(13) and (16) into Eq.(12) we finally obtain exact asymptotic expansion of the logarithm of the partition function with twisted boundary conditions in terms of the Kronecker's double series

$$\begin{aligned} \ln Z_{\alpha,\beta}(0) &= \frac{S}{\pi} \int_0^\pi \omega_0(x) dx + \ln \left| \frac{\theta_{\alpha,\beta}(i\lambda\xi)}{\eta(i\lambda\xi)} \right| \\ &\quad - 2\pi\xi \sum_{p=1}^{\infty} \left( \frac{\pi^2\xi}{S} \right)^p \frac{\Lambda_{2p}}{(2p)!} \frac{\operatorname{Re} K_{2p+2}^{\alpha,\beta}(i\lambda\xi)}{2p+2} \end{aligned} \quad (17)$$

Note, that Bernoulli polynomials  $B_p^\alpha$  have finally dropped out from the asymptotic expansion on torus. This actually means that Kronecker's double series can be considered as elliptic generalizations of Bernoulli polynomials.

### B. Asymptotic Expansion of $Z'_{0,0}(0)$

As it has already been mentioned, we have to treat the case  $(\alpha, \beta) = (0, 0)$  separately. Taking the derivative of Eq.(5) with respect to mass variable  $\mu$  and then considering limit  $\mu \rightarrow 0$  we obtain

$$Z'_{0,0}(0) = 2\sqrt{2}M \prod_{n=1}^{N-1} 2 \left| \operatorname{sh} \left[ M\omega_0 \left( \frac{\pi n}{N} \right) \right] \right| \quad (18)$$

Asymptotic expansion of this expression can be found along the same lines as above. The final result for the asymptotic expansion in terms of the Kronecker's double series states

$$\begin{aligned} \ln Z'_{0,0}(0) &= \frac{S}{\pi} \int_0^\pi \omega_0(x) dx + \frac{1}{2} \ln 8\xi S + 2 \ln |\eta(i\lambda\xi)| \\ &\quad - 2\pi\xi \sum_{p=1}^{\infty} \left( \frac{\pi^2\xi}{S} \right)^p \frac{\Lambda_{2p}}{(2p)!} \frac{\operatorname{Re} K_{2p+2}^{0,0}(i\lambda\xi)}{2p+2} \end{aligned} \quad (19)$$

Expansions (17) and (19) are the main results of the paper. Kronecker's double series  $K_p^{\alpha,\beta}$  with  $\alpha$  and  $\beta$  taking values 0 and 1/2 can all be expressed in terms of the elliptic  $\theta$ -functions only (Appendix E).

### C. Asymptotic Expansion of the Free Energy

After reaching this point, one can easily write down exact asymptotic expansion of the free energy (1) for all three models under consideration. For Ising model first few terms of the expansion are

$$f_{\text{bulk}} = -\ln \sqrt{2} - \frac{2\gamma}{\pi} \quad (20)$$

$$f_0(\xi) = -\ln \frac{\theta_2 + \theta_3 + \theta_4}{2\eta} \quad (21)$$

$$f_1(\xi) = -\frac{\pi^3 \xi^2 \frac{7}{8} (\theta_2^9 + \theta_3^9 + \theta_4^9) + \theta_2 \theta_3 \theta_4 [\theta_2^3 \theta_4^3 - \theta_3^3 \theta_2^3 - \theta_3^3 \theta_4^3]}{180 (\theta_2 + \theta_3 + \theta_4)} \quad (22)$$

⋮

For dimer model the expansion is somewhat different

$$f_{\text{bulk}} = -\frac{\gamma}{\pi} \quad (23)$$

$$f_0(\xi) = -\ln \frac{\theta_2^2 + \theta_3^2 + \theta_4^2}{2\eta^2} \quad (24)$$

$$f_1(\xi) = -\frac{\pi^3 \xi^2 \frac{7}{8} (\theta_2^{10} + \theta_3^{10} + \theta_4^{10}) + \theta_2^2 \theta_3^2 \theta_4^2 [\theta_2^2 \theta_4^2 - \theta_2^2 \theta_3^2 - \theta_3^2 \theta_4^2]}{90 (\theta_2^2 + \theta_3^2 + \theta_4^2)} \quad (25)$$

⋮

Finally, the asymptotic expansion of the free energy of the Gaussian model after subtraction of the zero mode,  $\ln \mu \sqrt{8S}$ , can be written as

$$f_{\text{bulk}} = \frac{2\gamma}{\pi} \quad (26)$$

$$f_0(\xi) = \ln \sqrt{\xi} \eta^2 \quad (27)$$

$$f_1(\xi) = \frac{\pi^3 \xi^2}{180} (\theta_2^4 \theta_4^4 - \theta_2^4 \theta_3^4 - \theta_3^4 \theta_4^4) \quad (28)$$

⋮

#### IV. SUMMARY

In this paper we have derived exact asymptotic expansion of the partition function with twisted boundary conditions in the vicinity of the critical point, Eqs. (17) and (19). As the application of this result, we have obtained exact asymptotic expansion of the free energy for a class of free exactly solvable models of statistical mechanics on torus. Generalization of our results to the case of internal energy and specific heat is straightforward. Moreover, the partition function of dimer model on the Klein bottle can also be expressed via the

partition function with twisted boundary conditions, namely  $Z_{\text{Klein}} = Z_{\frac{1}{4}, \frac{1}{2}}(0)$  [17]. Exact asymptotic expansion of the latter can immediately be written down with the help of our general formula (17). An interesting question is: whether this is also the case for other free models. This, however, is the problem for the future.

## APPENDIX A: EULER-MACLAURIN SUMMATION FORMULA.

Euler-Maclaurin summation formula states [18]

$$\sum_{n=0}^{N-1} f(n + \alpha) = \int_0^N f(x) dx + \sum_{p=1}^{\infty} \frac{B_p^\alpha}{p!} \left( f^{(p-1)}(N) - f^{(p-1)}(0) \right) \quad (\text{A1})$$

Here  $B_p^\alpha$  are so called Bernoulli functions

$$B_p^\alpha = -\frac{p!}{(-2\pi i)^p} \sum_{n \neq 0} \frac{e^{-2\pi i n \alpha}}{n^p} \quad (\text{A2})$$

These are actually polynomials defined on the interval  $\alpha \in [0, 1]$

$$\begin{aligned} B_1^\alpha &= \alpha - \frac{1}{2} \\ B_2^\alpha &= \alpha^2 - \alpha + \frac{1}{6} \\ B_3^\alpha &= \alpha^3 - \frac{3}{2}\alpha^2 + \frac{1}{2}\alpha \\ B_4^\alpha &= \alpha^4 - 2\alpha^3 + \alpha^2 - \frac{1}{30} \\ &\vdots \end{aligned}$$

The generating function for the polynomials is

$$\mathcal{B}^\alpha(z) = \frac{e^{z\alpha}}{e^z - 1} = \frac{1}{z} \left( 1 + \sum_{p=1}^{\infty} \frac{z^p}{p!} B_p^\alpha \right) \quad (\text{A3})$$

Fourier transform of the generating function gives the following important identity

$$\frac{e^{2\pi i z \alpha}}{e^{2\pi i z} - 1} = - \sum_{n=0}^{\infty} e^{2\pi i z(n+\alpha)} = \frac{1}{2\pi i} \sum_{n=-\infty}^{+\infty} \frac{e^{-2\pi i n \alpha}}{z + n} \quad (\text{A4})$$

## APPENDIX B: RELATION BETWEEN MOMENTS AND CUMULANTS

Moments  $Z_k$  and cumulants  $F_k$  which enters the expansion of exponent

$$\exp \left\{ \sum_{k=1}^{\infty} \frac{x^k}{k!} F_k \right\} = 1 + \sum_{k=1}^{\infty} \frac{x^k}{k!} Z_k \quad (\text{B1})$$

are related to each other as [19]

$$\begin{aligned} Z_1 &= F_1 \\ Z_2 &= F_2 + F_1^2 \\ Z_3 &= F_3 + 3F_1F_2 + F_1^3 \\ Z_4 &= F_4 + 4F_1F_3 + 3F_2^2 + 6F_1^2F_2 + F_1^4 \\ &\vdots \\ Z_k &= \sum_{r=1}^k \sum \left( \frac{F_{k_1}}{k_1!} \right)^{i_1} \cdots \left( \frac{F_{k_r}}{k_r!} \right)^{i_r} \frac{k!}{i_1! \cdots i_r!} \end{aligned} \quad (\text{B2})$$

where summation is over all positive numbers  $\{i_1 \dots i_r\}$  and different positive numbers  $\{k_1, \dots, k_r\}$  such that  $k_1 i_1 + \dots + k_r i_r = k$ .

## APPENDIX C: ELLIPTIC THETA FUNCTIONS

We use the following definition of Jacobi  $\theta$ -functions (compare with the notations of Mumford [20])

$$\theta_{\alpha, \beta}(z, \tau) = \sum_{n \in \mathbb{Z}} \exp \left\{ \pi i \tau \left( n + \frac{1}{2} - \alpha \right)^2 + 2\pi i \left( n + \frac{1}{2} - \alpha \right) \left( z + \frac{1}{2} - \beta \right) \right\} \quad (\text{C1})$$

$$\begin{aligned} &= \eta(\tau) \exp \left\{ \pi i \tau \left( \alpha^2 - \alpha + \frac{1}{6} \right) + 2\pi i \left( \frac{1}{2} - \alpha \right) \left( z + \frac{1}{2} - \beta \right) \right\} \\ &\times \prod_{n=0}^{\infty} \left[ 1 - e^{2\pi i \tau (n+\alpha) - 2\pi i (z-\beta)} \right] \left[ 1 - e^{2\pi i \tau (n+1-\alpha) + 2\pi i (z-\beta)} \right] \end{aligned} \quad (\text{C2})$$

The relation to standard notations is follows

$$\theta_{0,0}(z, \tau) = \theta_1(z, \tau)$$

$$\theta_{0, \frac{1}{2}}(z, \tau) = \theta_2(z, \tau)$$

$$\theta_{\frac{1}{2}, 0}(z, \tau) = \theta_4(z, \tau)$$

$$\theta_{\frac{1}{2}, \frac{1}{2}}(z, \tau) = \theta_3(z, \tau)$$

The Dedekind  $\eta$ -function is usually defined as

$$\eta(\tau) = e^{\pi i \tau / 12} \prod_{n=1}^{\infty} [1 - e^{2\pi i \tau n}] \quad (\text{C3})$$

Taking into account Eq.(C2) and considering the functions  $\theta_{\alpha,\beta} = \theta_{\alpha,\beta}(0, i\xi)$  and  $\eta = \eta(i\xi)$  with argument  $z = 0$  and pure imaginary aspect ratio  $\tau = i\xi$  we obtain

$$\begin{aligned} \ln \left| \frac{\theta_{\alpha,\beta}}{\eta} \right| + \pi \xi B_2^\alpha &= \sum_{n=0}^{\infty} \ln \left| 1 - e^{-2\pi \xi (n+\alpha) - 2\pi i \beta} \right| \\ &+ \sum_{n=0}^{\infty} \ln \left| 1 - e^{-2\pi \xi (n+1-\alpha) - 2\pi i \beta} \right| \end{aligned} \quad (\text{C4})$$

This is the identity that have appeared in our derivation of the asymptotic expansions.

#### APPENDIX D: KRONECKER'S DOUBLE SERIES

The building blocks of our construction are so-called Kronecker's double series [21]

$$K_p^{\alpha,\beta}(\tau) = -\frac{p!}{(-2\pi i)^p} \sum_{\substack{m,n \in \mathbb{Z} \\ (m,n) \neq (0,0)}} \frac{e^{-2\pi i(n\alpha+m\beta)}}{(n+\tau m)^p} \quad (\text{D1})$$

These can be obtained from the generating function

$$\mathcal{K}^{\alpha,\beta}(z|\tau) = \sum_{m,n \in \mathbb{Z}} \frac{e^{-2\pi i(n\alpha+m\beta)}}{z+n+\tau m} = 2\pi \eta^3(\tau) e^{2\pi i \beta} \frac{\theta(z+w)}{\theta(z)\theta(w)} \quad (\text{D2})$$

where  $w = \alpha\tau + \beta$ . However, in this form Kronecker's double series cannot be directly applied to our analysis. First we need to cast them in a different form. Let us separate from the double sum a subseries with  $m = 0$

$$K_p^{\alpha,\beta}(\tau) = -\frac{p!}{(-2\pi i)^p} \sum_{n \neq 0} \frac{e^{-2\pi i n \alpha}}{n^p} - \frac{p!}{(-2\pi i)^p} \sum_{m \neq 0} \sum_{n \in \mathbb{Z}} \frac{e^{-2\pi i(n\alpha+m\beta)}}{(n+\tau m)^p} \quad (\text{D3})$$

Here the first sum gives nothing but Fourier representation of Bernoulli polynomials Eq.(A2).

The second sum can be rearranged in a different way with the help of the identity

$$\frac{p!}{(-2\pi i)^p} \sum_{n \in \mathbb{Z}} \frac{e^{-2\pi i n \alpha}}{(z+n)^p} = p \sum_{n=0}^{\infty} (n+\alpha)^{p-1} e^{2\pi i z(n+\alpha)} \quad (\text{D4})$$

which can easily be derived from Eq.(A4) differentiating it  $p$  times. The final result of our resummation of double Kronecker sum is

$$K_p^{\alpha,\beta}(\tau) = B_p^\alpha - p \sum_{m \neq 0} \sum_{n=0}^{\infty} (n + \alpha)^{p-1} e^{2\pi i m(\tau(n+\alpha)-\beta)} \quad (\text{D5})$$

Considering real part of the Kronecker sums with pure imaginary aspect ratio  $\tau = i\xi$  we can further rearrange this expression to get summation only over positive  $m \geq 1$

$$\begin{aligned} B_{2p}^\alpha - \text{Re} K_{2p}^{\alpha,\beta}(i\xi) &= \frac{(2p)!}{(-4\pi^2)^p} \text{Re} \sum_{m=1}^{\infty} \sum_{n \in \mathbb{Z}} \frac{e^{-2\pi i(n\alpha+m\beta)} + e^{-2\pi i(n(1-\alpha)-m\beta)}}{(n + i\xi m)^{2p}} \\ &= 2p \text{Re} \left\{ \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} (n + \alpha)^{2p-1} e^{-2\pi m(\xi(n+\alpha)+i\beta)} \right. \\ &\quad \left. + \sum_{m=1}^{\infty} \sum_{n=0}^{\infty} (n + 1 - \alpha)^{2p-1} e^{-2\pi m(\xi(n+1-\alpha)+i\beta)} \right\} \quad (\text{D6}) \end{aligned}$$

This last expression has appeared in our calculation of asymptotic expansion of partition function with twisted boundary conditions.

## APPENDIX E: REDUCTION OF KRONECKER'S DOUBLE SERIES TO THETA FUNCTIONS

Let us consider the Weierstrass function [22]

$$\wp(z) = \frac{1}{z^2} + \sum_{(n,m) \neq (0,0)} \left[ \frac{1}{(z - n - \tau m)^2} - \frac{1}{(n + \tau m)^2} \right] \quad (\text{E1})$$

and its Laurent expansion in powers of  $z$

$$\wp(z) = \frac{1}{z^2} + \sum_{p=2}^{\infty} a_p(\tau) z^{2p-2} \quad (\text{E2})$$

The coefficients  $a_p(\tau)$  of the expansion can all be written in terms of the elliptic  $\theta$ -functions with the help of the recursion relation

$$a_p = \frac{3}{(p-3)(2p+1)} (a_2 a_{p-2} + a_3 a_{p-3} + \dots + a_{p-2} a_2) \quad (\text{E3})$$

where first terms of the sequence are

$$\begin{aligned} a_2 &= \frac{\pi^4}{15} (\theta_2^4 \theta_3^4 - \theta_2^4 \theta_4^4 + \theta_3^4 \theta_4^4) \\ a_3 &= \frac{\pi^6}{189} (\theta_2^4 + \theta_3^4) (\theta_4^4 - \theta_2^4) (\theta_3^4 + \theta_4^4) \end{aligned}$$

$$\begin{aligned}
a_4 &= \frac{1}{3}a_2^2 \\
a_5 &= \frac{3}{11}(a_2a_3) \\
a_6 &= \frac{1}{39}(2a_2^3 + 3a_3^2) \\
&\vdots
\end{aligned}$$

Kronecker functions  $K_{2p}^{0,0}(\tau)$  are related directly to the coefficients  $a_p(\tau)$ , namely we have

$$K_{2p}^{0,0}(\tau) = -\frac{(2p)!}{(-4\pi^2)^p} \frac{a_p(\tau)}{(2p-1)} \quad (\text{E4})$$

Kronecker functions  $K_{2p}^{\alpha,\beta}(\tau)$  with  $\alpha$  and  $\beta$  taking values 0 or 1/2 can in their turn be related to the function  $K_{2p}^{0,0}(\tau)$  by means of simple resummation of Kronecker's double series

$$K_p^{0,\frac{1}{2}}(\tau) = 2K_p^{0,0}(2\tau) - K_p^{0,0}(\tau) \quad (\text{E5})$$

$$K_p^{\frac{1}{2},0}(\tau) = 2^{1-p}K_p^{0,0}(\tau/2) - K_p^{0,0}(\tau) \quad (\text{E6})$$

$$K_p^{\frac{1}{2},\frac{1}{2}}(\tau) = (1 + 2^{2-p})K_p^{0,0}(\tau) - 2^{1-p}K_p^{0,0}(\tau/2) - 2K_p^{0,0}(2\tau) \quad (\text{E7})$$

Thus, Kronecker functions  $K_{2p}^{\alpha,\beta}(\tau)$  with  $\alpha$  and  $\beta$  taking values 0 or 1/2 can all be expressed in terms of the elliptic  $\theta$ -functions only. For practical calculations the following identities are also helpful

$$\begin{aligned}
2\theta_2^2(2\tau) &= \theta_3^2 - \theta_4^2 & \theta_2^2(\tau/2) &= 2\theta_2\theta_3 \\
2\theta_3^2(2\tau) &= \theta_3^2 + \theta_4^2 & \theta_3^2(\tau/2) &= \theta_2^2 + \theta_3^2 \\
2\theta_4^2(2\tau) &= 2\theta_3\theta_4 & \theta_4^2(\tau/2) &= \theta_3^2 - \theta_2^2
\end{aligned}$$

From the general formulas above we can easily write down all the Kronecker functions that have appeared in our asymptotic expansions

$$\begin{aligned}
K_4^{0,0}(\tau) &= \frac{1}{30}(\theta_2^4\theta_4^4 - \theta_2^4\theta_3^4 - \theta_3^4\theta_4^4) \\
K_4^{0,\frac{1}{2}}(\tau) &= \frac{1}{30}(\frac{7}{8}\theta_2^8 - \theta_3^4\theta_4^4) \\
K_4^{\frac{1}{2},0}(\tau) &= \frac{1}{30}(\frac{7}{8}\theta_4^8 - \theta_2^4\theta_3^4) \\
K_4^{\frac{1}{2},\frac{1}{2}}(\tau) &= \frac{1}{30}(\frac{7}{8}\theta_3^8 + \theta_2^4\theta_4^4)
\end{aligned}$$

$$\begin{aligned}
K_6^{0,0}(\tau) &= \frac{1}{84}(\theta_2^4 + \theta_3^4)(\theta_4^4 - \theta_2^4)(\theta_3^4 + \theta_4^4) \\
K_6^{0,\frac{1}{2}}(\tau) &= \frac{1}{84}(\theta_3^4 + \theta_4^4)\left(\frac{31}{16}\theta_2^8 + \theta_3^4\theta_4^4\right) \\
K_6^{\frac{1}{2},0}(\tau) &= -\frac{1}{84}(\theta_2^4 + \theta_3^4)\left(\frac{31}{16}\theta_4^8 + \theta_2^4\theta_3^4\right) \\
K_6^{\frac{1}{2},\frac{1}{2}}(\tau) &= \frac{1}{84}(\theta_2^4 - \theta_4^4)\left(\frac{31}{16}\theta_3^8 - \theta_4^4\theta_2^4\right) \\
&\vdots
\end{aligned}$$

We would like to thank V.B. Priezhev for critical reading of the manuscript. This work was supported by the Russian Foundation for Basic Research through grant No. 99-01-00882, by the Swiss National Science Foundation through grant No. 7-SUPJ-62295 and by the National Science Council of the Republic of China (Taiwan) through grant No. NSC89-2112-M-001-084.

## REFERENCES

- \* Electronic address: ivashkev@thsun1.jinr.ru
- [1] Privman, V. (Ed.): *Finite Size Scaling and Numerical Simulation of Statistical Systems*. Singapore: World Scientific, 1990 (see also references therein)
- [2] Baxter, R.J.: *Exactly Solved Models in Statistical Mechanics*. London: Academic Press, 1982
- [3] McCoy, B. and Wu, T.T.: *The Two-Dimensional Ising Model*. Cambridge: Harward University Press, 1973
- [4] Cardy, J.L.: *Conformal Invariance and Statistical Mechanics in Les Houches Session XLIX: Fields, Strings and Critical Phenomena*. Eds. Brezin, E. and Zinn-Justin, J., Amsterdam: North-Holland, 1990
- [5] Izmailian, N.Sh. and Hu, C.K.: Exact Universal Amplitude Ratios for the Planar Ising Model and a Quantum Spin Chain. cond-mat/0009102 (2000)
- [6] Ferdinand, A.E. and Fisher, M.E.: Bounded and Inhomogeneous Ising Models. I. Specific-Heat Anomaly of a Finite Lattice. Phys. Rev. **185**, 832 (1969)
- [7] Onsager, L.: Crystal Statistics. I. A Two-Dimensional Model with an Order-Disorder Transition. Phys. Rev. **65**, 117 (1944)
- [8] Izmailian, N.Sh. and Hu, C.K.: Ising Model on The Square  $M \times N$  Lattice: Exact Finite-Size Calculations. cond-mat/0009024 (2000)
- [9] Salas, J.: Exact Finite-Size Scaling Corrections to the Critical Two-Dimensional Ising Model on Torus. cond-mat/0009054 (2000)
- [10] Salas, J. and Sokal, A.D.: Universal Amplitude Ratios in the Critical Two-Dimensional Ising Model on a Torus. J. Stat. Phys. **88**, 567 (1997)

- [11] Christe, P. and Henkel, M.: *Introduction to Conformal Invariance and its Application to Critical Phenomena*. Berlin: Springer-Verlag, 1993
- [12] Gradshteyn, I.S. and Ryzhik, I.M.: *Table of Integrals, Series and Products*. New York: Academic Press, 1965
- [13] Kasteleyn, P.W.: The Statistics of Dimers on a Lattice. *Physica* **27**, 1209 (1961)
- [14] Fisher, M.E.: Statistical Mechanics of Dimers on a Plane Lattice. *Phys. Rev.* **124**, 1664 (1961)
- [15] Ferdinand, A.E.: Statistical Mechanics of Dimers on a Quadratic Lattice. *J. Math. Phys.* **8**, 2332 (1967)
- [16] Sato, M., Miwa, T. and Jimbo, M.: *Holonomic Quantum Fields V*, Publ. RIMS, Kyoto Univ. **16**, 531 (1980).
- [17] Lu, W.T and Wu, F.Y.: Dimer Statistics on the Möbius strip and the Klein bottle. *Phys. Lett.* **A259**, 108 (1999)
- [18] Hardy, G.H.: *Divergent Series*. Oxford: Clarendon Press, 1949
- [19] Prohorov, Yu.V. and Rozanov, Yu.A.: *Probability Theory*. New York: Springer-Verlag, 1969
- [20] Mumford, D.: *Tata lectures on Theta*. Boston-Basel-Stuttgart: Birkhäuser, 1983
- [21] Weil, A.: *Elliptic functions according to Eisenshtein and Kronecker*. Berlin-Heidelberg-New York: Springer-Verlag, 1976
- [22] Korn, G.A. and Korn, T.M.: *Mathematical Handbook*. New-York: McGraw-Hill, 1968