Thermal magnetization fluctuations in thin films and a new physical form for magnetization damping

Vladimir L. Safonov and H. Neal Bertram

Center for Magnetic Recording Research, University of California - San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0401

(November 1, 2018)

The effect of thermal fluctuations on a thin film magnetoresistive element has been calculated. The technique involves adding to the basic spin dynamics a general form of interaction with a thermal bath. For a general anisotropic magnetic system the resulting equation can be written as a Langevin equation for a harmonic oscillator. Our approach predicts two times smaller noise power at low frequencies than the conventional stochastic Landau-Lifshitz-Gilbert equation. It is shown that equivalent results can be obtained by introducing a tensor phenomenological damping term to the gyromagnetic dynamics driven by a thermal fluctuating field.

Recently Smith and Arnett [1] have reported that thermal magnetization noise may be a serious problem for giant magnetoresistive spin-valve heads designed for high areal storage density. Using the phenomenological Landau-Lifshitz-Gilbert (LLG) equation with random fields and the fluctuation-dissipation theorem, they obtained an estimate for the low-frequency noise intensity. They considered the GMR head as a single-domain anisotropic magnetic particle.

The goal of this Note is to show that the analysis in the framework of conventional stochastic LLG equation gives an overstated noise level due to inconsistency of isotropic phenomenological damping with basic dynamic equations for anisotropic magnetic system. Using normal mode approach, we derive a formula for the noise power spectrum, which at low frequencies is two times smaller than theoretical estimate of Ref. [1]. We also show that this result may be obtained in the framework of stochastic gyromagnetic equation with tensor phenomenological damping.

Thermal noise is a phenomenon that is a result of microscopic spin fluctuations due to interactions with a thermal bath. It has been customary to include thermal fluctuations by simply adding random fields to dynamic equations with phenomenological damping, such as the LLG equation [2]. However the gyromagnetic term describes an averaged precession of a large number of spins. A direct connection with microscopic physics must be found for the phenomenological damping term.

Lax [3] has shown that independent of the specific coupling to the heat bath, the resulting dynamic equation in the vicinity of equilibrium has the form of a damped harmonic oscillator driven by a fluctuating force. Here we show that for small fluctuations, an anisotropic magnetic dynamic system can be transformed into the general form analyzed by Lax. We do this by a series of coordinate transformations from the original magnetic variables to a harmonic oscillator. By back transformation we have found a tensor form of the phenomenological damping for the magnetization dynamic equation. With this new form random fields may now be added to this dynamic equation, while properly characterizing the physics of interaction with a thermal bath.

Consider thermal fluctuations in a single-domain magnetic particle with the following energy density:

$$\mathcal{E}/V_0 = -K_{\parallel} (M_z/M_s)^2 + K_{\perp} (M_x/M_s)^2 - M_z H_0.$$
(1)

Here V_0 is the particle volume, K_{\parallel} and K_{\perp} are the easy (z) and hard (x) axis anisotropy constants, respectively, $\mathbf{M} = (M_x, M_y, M_z)$ is the magnetization $(|\mathbf{M}| = M_s)$ and H_0 is the external magnetic field oriented along the easy axis. For a thin film y and z are in the plane.

For convenience of calculations we introduce the classical spin $\mathbf{S} = -\mathbf{M}V_0/\hbar\gamma$, where γ is the gyromagnetic ratio. Planck's constant \hbar is used as a dimensional parameter (it could be another parameter of the same dimensionality, e.g., $M_{\rm s}V_0/\gamma$). All our analysis is pure classical and final result does not contain \hbar . We shall describe small oscillations of the magnetization in terms of complex variables a^* , a, which are classical analogs of creation and annihilation operators of a harmonic oscillator. This technique is used for a description of spin waves (see, e.g., [4]). Complex variables can be introduced by an expanded Holstein-Primakoff transformation [5]:

$$\frac{S_x}{\sqrt{S/2}} \simeq \frac{a-a^*}{i}, \ \frac{S_y}{\sqrt{S/2}} \simeq a+a^*, \ S_z = a^*a-S.$$
 (2)

The quadratic part of the energy (1) can be written as

$$\mathcal{E}^{(2)}/\hbar = Aa^*a + (B/2)(aa + a^*a^*), \qquad (3)$$

where $A = \gamma (H_{\rm K}^{(\parallel)} + H_0 + H_{\rm K}^{(\perp)}/2)$, $B = -\gamma H_{\rm K}^{(\perp)}/2$, $H_{\rm K}^{(\parallel)} \equiv 2K_{\parallel}/M_{\rm s}$ and $H_{\rm K}^{(\perp)} \equiv 2K_{\perp}/M_{\rm s}$. To eliminate the nondiagonal terms from (3) we shall use the linear canonical transformation:

$$a = uc + vc^{*}, \quad a^{*} = uc^{*} + vc,$$
(4)
$$u = \sqrt{\frac{A + \omega_{0}}{2\omega_{0}}}, \quad v = -\frac{B}{|B|}\sqrt{\frac{A - \omega_{0}}{2\omega_{0}}},$$

where $\omega_0 = \sqrt{A^2 - B^2}$. Thus, we find the normal mode of the system and the energy of an harmonic oscillator

$$\mathcal{E}^{(2)}/\hbar = \omega_0 c^* c, \tag{5}$$

$$\omega_0 = \gamma \sqrt{(H_{\rm K}^{(\parallel)} + H_0 + H_{\rm K}^{(\perp)})(H_{\rm K}^{(\parallel)} + H_0)}, \qquad (6)$$

where ω_0 is the ferromagnetic resonance frequency.

Using the energy (5), the dynamic equation for the complex amplitude c(t) is:

$$dc/dt = -i\omega_0 c \tag{7}$$

and is of a form that corresponds exactly to small amplitude gyromagnetic rotation. Now we consider the interaction of this harmonic oscillator with a reservoir (thermal bath)

$$\mathcal{E}^{(int)}/\hbar = c \sum_{k} g_k^* + c^* \sum_{k} g_k, \qquad (8)$$

which will introduce two related effects: damping and fluctuations. Here g_k , g_k^* are the reservoir variables and incorporate specific physical processes. Such interaction could have been added to the energy in the a, a^* variables. However the linear transformation (4) does not alter this general form. Following [3], [6], the oscillator dynamics for this general form (8) can be represented by the following Langevin equation:

$$dc/dt = -(i\omega_0 + \eta)c + f(t), \qquad (9)$$

where η is defined by reservoir parameters and f(t) is the random force related to η . It should be noted that Eq.(9) is an exact classical analog of a corresponding quantum mechanical equation [3]. Note also that phenomenological introduction of dissipation by assigning an imaginary part to the resonant frequency $\omega_0 \pm i\eta$ is a common procedure in the theory of spin-wave instabilities [7].

We can start with Eq.(9) and rewrite it as a stochastic differential equation with a damping parameter η and a thermodynamically consistent random field with diffusion coefficient $D = (\eta/\omega_0)k_{\rm B}T/\hbar$ [8], [9]:

$$dc = -(i\omega_0 + \eta)cdt + \sqrt{D}[dW_1(t) + idW_2(t)].$$
 (10)

Here T is the temperature and $W_1(t)$ and $W_2(t)$ are two independent Wiener processes. The solution of (10) is

$$\frac{c(t)}{\sqrt{D}} = \int_{-\infty}^{t} e^{-(i\omega_0 + \eta)(t - t_1)} [dW_1(t_1) + idW_2(t_1)].$$
(11)

Taking into account the rule of Ito's calculus (see, e.g., [9]) $dW_m(t_1)dW_n(t_2) = \delta_{mn}\Delta(t_1 - t_2)dt_1$, where $\Delta(t)$ is the Kronecker delta function, we can obtain the correlation function with thermal averaging $\langle ... \rangle$:

$$\langle c^*(t)c(0)\rangle = (D/\eta)\exp(i\omega_0 t - \eta t), \qquad (12)$$

where $c^*(t)$ is the complex conjugate of c(t).

The power spectral density of the complex amplitude fluctuations is defined by

$$S_{c^*c}(\omega) = \int_{-\infty}^{\infty} \langle c^*(t)c(0) \rangle e^{-i\omega t} dt.$$
 (13)

Substituting (12) into (13) and including the correct meaning of a damping process for both negative and positive time, one obtains

$$S_{c^*c}(\omega) = \frac{k_{\rm B}T}{\hbar\omega_0} \int_{-\infty}^{\infty} e^{i(\omega_0 - \omega)t - \eta t \cdot \operatorname{sign}(t)} dt$$
$$= \frac{2(\eta/\omega_0)k_{\rm B}T/\hbar}{(\omega_0 - \omega)^2 + \eta^2}.$$
(14)

Let us consider now an application of our analysis to a magnetoresistive head. We shall follow the model [1] of a GMR head as a thin film of the volume V_0 , uniformly magnetized along the z axis of uniaxial anisotropy. The transverse anisotropy is defined by shape demagnetization: $K_{\perp} \simeq 2\pi M_{\rm s}^2$. The power spectral density of random fluctuations in the voltage $\delta V(t)$ is defined by

$$S_{VV}(\omega) = \int_{-\infty}^{\infty} \langle \delta V(t) \delta V(0) \rangle e^{i\omega t} dt.$$
 (15)

The voltage deviation for a GMR head is $\delta V(t) = I_{\text{bias}} \delta R(t)$, which can be expressed in terms of the inplane transverse magnetization fluctuation

$$\delta V(t) = C_V \delta m_x(t), \quad C_V = I_{\text{bias}} \frac{\partial R}{\partial H_{\text{ext}}} \frac{\partial H_{\text{ext}}}{\partial m_x}.$$
 (16)

Thus, Eq.(15) can be rewritten as

$$S_{VV}(\omega) = C_V^2 S_{m_y m_y}(\omega), \qquad (17)$$

where

$$S_{m_y m_y}(\omega) = \int_{-\infty}^{\infty} \langle \delta m_y(t) \delta m_y \rangle(0) e^{i\omega t} dt \qquad (18)$$

defines the magnetization noise.

Since $\langle m_y(t) \rangle = 0$, the deviation $\delta m_y(t) = m_y(t)$. Taking into account Eqs. (2) and (4), one obtains

$$m_y = -S_y/S \simeq -(u+v)(c^*+c)/\sqrt{2S}.$$
 (19)

Thus

$$S_{m_y m_y}(\omega) = \frac{(u+v)^2}{2S} \Big[\int_{-\infty}^{\infty} \langle c(t)c^*(0) \rangle e^{i\omega t} dt \\ + \int_{-\infty}^{\infty} \langle c^*(t)c(0) \rangle e^{i\omega t} dt \Big].$$

Utilizing (13) and (14), we have

$$S_{m_y m_y}(\omega) = \frac{\gamma k_{\rm B} T}{M_{\rm s} V_0} \left(\frac{H_{\rm K} + 4\pi M_{\rm s} + H_0}{H_{\rm K} + H_0} \right)^{1/2} \frac{\eta}{\omega_0} F(\omega), \quad (20)$$
$$F(\omega) = \frac{1}{(\omega_0 - \omega)^2 + \eta^2} + \frac{1}{(\omega_0 + \omega)^2 + \eta^2}.$$

Examination of (20) indicates that the damping parameter η affects linearly the low frequency noise ($\omega \ll \omega_0$) and broadens the resonance peak. However the total power is independent of η .

In the case of small frequencies $\omega \ll \omega_0 \simeq \gamma \sqrt{4\pi M_{\rm s} H_{\rm K}}$ for $H_0 = 0$ and $4\pi M_{\rm s} \gg H_{\rm K}$. From (20) we obtain the noise power (as measured for positive frequencies):

$$NP \simeq 2S_{m_y m_y}(0) \simeq \eta k_{\rm B} T / [\pi (\gamma H_{\rm K} M_{\rm s})^2 V_0].$$
 (21)

The corresponding noise power of [1] (in our notation) is

$$NP^{(\mathrm{SA})} \simeq 4\alpha k_{\mathrm{B}} T / [\gamma (H_{\mathrm{K}})^2 M_{\mathrm{s}} V_0], \qquad (22)$$

where α is the Gilbert damping parameter. To compare (21) and (22) we need to find a relation between η and α . This can be done by comparing the dynamic equation for the transverse magnetization given in Ref. [1] and the corresponding transverse magnetization equation in our approach. From [1] we have

$$d^{2}m_{y}/dt^{2} + \alpha 4\pi\gamma M_{s}dm_{y}/dt + \omega_{0}^{2}m_{y} = h(t), \qquad (23)$$

where h(t) is a random "force". Corresponding to (10) dynamic equation is:

$$d^{2}m_{y}/dt^{2} + 2\eta dm_{y}/dt + \omega_{0}^{2}m_{y} = h(t).$$
(24)

From (23) and (24), we obtain the following relation:

$$\alpha = \eta / 2\pi \gamma M_{\rm s}.\tag{25}$$

Substituting (25) into (22), one can see that at low frequencies our theory (21) predicts 2 times smaller (3 dB smaller) noise power than the result of [1]: $NP/NP^{(SA)} = 1/2$. It should be also noted that the conventional approach with stochastic LLG equation gives different noise frequency dependence than our result (20).

Now we demonstrate the inconsistency of the conventional phenomenological damping term with basic equations for a harmonic oscillator and show how this isotropic damping can be replaced by a tensor damping term to obtain the above results.

From the LLG equation

$$\frac{d\mathbf{M}}{dt} = -\gamma \mathbf{M} \times \mathbf{H}_{\text{eff}} - \frac{\alpha \gamma}{M_{\text{s}}} \mathbf{M} \times (\mathbf{M} \times \mathbf{H}_{\text{eff}}) \qquad (26)$$

one can derive the following linearized equations:

$$\frac{d}{dt} \begin{pmatrix} m_x \\ m_y \end{pmatrix}$$

$$= \begin{pmatrix} -\alpha\gamma(H_{\rm K} + 4\pi M_{\rm s} + H_0) & -\gamma(H_{\rm K} + H_0) \\ \gamma(H_{\rm K} + 4\pi M_{\rm s} + H_0) & -\alpha\gamma(H_{\rm K} + H_0) \end{pmatrix} \begin{pmatrix} m_x \\ m_y \end{pmatrix}$$
(27)

On the other hand, we can obtain the linearized equation for m_x and m_y from the normal mode equations (Eq.(10) without the thermal term)

$$(d/dt + \eta)c = -i\omega_0 c, \quad (d/dt + \eta)c^* = i\omega_0 c^*.$$
 (28)

Substituting $c = ua - va^*$, $c^* = ua^* - va$ into (28) and taking into account (2) and $\mathbf{m} = -\mathbf{S}/S$, we obtain

$$\frac{d}{dt} \begin{pmatrix} m_x \\ m_y \end{pmatrix}$$

$$= \begin{pmatrix} -\eta & -\gamma(H_{\rm K} + H_0) \\ \gamma(H_{\rm K} + 4\pi M_{\rm s} + H_0) & -\eta \end{pmatrix} \begin{pmatrix} m_x \\ m_y \end{pmatrix}.$$
(29)

By comparison with (27) we see that the nondiagonal terms coincide. The diagonal terms in (27) and (29), responsible for relaxation, differ from each other.

In order to have the magnetization equations (29), one should start with the magnetization equation containing a tensor damping of the form:

$$\frac{d\mathbf{M}}{dt} = -\gamma \mathbf{M} \times \mathbf{H}_{\text{eff}} - \gamma \frac{\mathbf{M}}{M_{\text{s}}} \times [\stackrel{\leftrightarrow}{\alpha} \cdot (\mathbf{M} \times \mathbf{H}_{\text{eff}})], \qquad (30)$$

$$\overset{\leftrightarrow}{\alpha} = \frac{\eta \gamma}{\omega_0^2} \begin{pmatrix} H_{\rm K}^{(||)} + H_0 + H_{\rm K}^{(\perp)} & 0 & 0\\ 0 & H_{\rm K}^{(||)} + H_0 & 0\\ 0 & 0 & H_0 \end{pmatrix}.$$
(31)

For small oscillations about equilibrium, the α_{zz} component does not enter into the equations. The conventional LLG equation can be used only for a uniaxial system, where $H_{\rm K}^{(\perp)} = 0$ and the tensor is effectively a scalar.

We thank H. Suhl, C. E. Patton and N. Smith for valuable discussions, and X. Wang for his help with the analysis of paper [1]. This work was partly supported by matching funds from the Center for Magnetic Recording Research at the University of California - San Diego and CMRR incorporated sponsor accounts.

- [1] N. Smith and P. Arnett, Appl. Phys. Lett. 78, 1448 (2001).
- [2] W. F. Brown, Jr., Phys. Rev. **130**, 1677 (1963).
- [3] M. Lax, Phys. Rev. 145, 110 (1966).
- [4] V. S. L'vov, Wave Turbulence Under Parametric Excitation (Springer, Berlin, 1994).
- [5] T. Holstein and H. Primakoff: Phys. Rev. 58, 1098 (1940).
- [6] M. O. Scully and M. S. Zubairy, *Quantum Optics* (Cambridge University Press, 1997), Chapter 9.
- [7] E. Schlömann, R. I. Joseph, and I. Bady, J. Appl. Phys. 34, 672 (1963).
- [8] V. L. Safonov, J. Magn. Magn. Mater. 195, 526 (1999).
- [9] C. W. Gardiner: Handbook of Stochastic Methods (Springer, Berlin, 1985).