

# Lattice animals on a staircase and generalized Fibonacci numbers

Loïc Turban

*Laboratoire de Physique des Matériaux, UMR CNRS 7556,  
Université Henri Poincaré, BP 239,  
F-54506 Vandœuvre lès Nancy Cedex, France  
e-mail: turban@lpm.u-nancy.fr*

---

## Abstract

We study the statistics of column-convex lattice animals generated by the stacking of squares on a staircase with step height  $p$ . We calculate the number of animals with area  $k$  living on  $l$  stairs. The total number of animals with area  $k$  is given by the generalized Fibonacci number  $F_p(k)$ . Exact results for the mean length and mean height of animals with area  $k$  are also obtained and we examine their asymptotic behaviour.

*Key words:* lattice animals, polyominoes, generalized Fibonacci numbers

---

## 1 Introduction

A lattice animal is a connected cluster of occupied sites on a lattice or the corresponding cluster of occupied cells on the dual lattice, also called a polyomino. In two dimensions, animals can be enumerated either according to their area, the number of sites belonging to a cluster, or according to their perimeter, the number of vacant sites which are first neighbours of occupied sites.

The enumeration problem is quite difficult in general. Only some bounds on the asymptotic behaviour are known [10]. For restricted classes of animals such as Ferrers graphs, convex and/or directed animals, some exact results have been obtained during the last two decades [1,3,4,5,7,8,11,12]. A review of the state of the problem before 1996 can be found in [2].

In the present work, we consider lattice animals resulting from the stacking of squares on a staircase with step height  $p$ . Two square faces are connected




Fig. 1. The 6 different animals obtained by stacking 7 squares on  $l = 1, 2, 3$  stairs in a staircase with step height 2. The  $3l - 2$  dark squares build the minimal cluster necessary to cover  $l$  stairs.

when they share an edge as shown in figure 1. Such animals are called column-convex or vertically convex because the intersection of a vertical line with the perimeter generates at most two connected components. The enumeration problem and the morphology of the animals involve the generalized Fibonacci numbers  $F_p(k)$ . The case  $p = 0$ , which corresponds to the stacking of squares on a line, is well known (see [13] for example). With  $p = 1$ , a case recently considered in [14], one obtains a relation with the ordinary Fibonacci numbers  $F_k = F_1(k)$ . A similar connection has been noticed a long time ago between column-convex directed animals and ordinary Fibonacci numbers with odd indices [9]

In section 2, the number of animals with area  $k$  living on  $l$  stairs is deduced from their generating function. Their total number is given by the generalized Fibonacci numbers  $F_p(k)$ . In sections 3 and 4, we obtain exact expressions for the mean length and mean height of an animal with area  $k$ . The asymptotic behaviour is discussed in section 5.

## 2 Number of animals

We consider animals with area  $k$  living on  $l$  stairs in a staircase with step height  $p$  as shown in figure 1. In order to calculate their number  $F_p(k, l)$ , we introduce the generating function

$$F_p(z, t) = \sum_{k, l=1}^{\infty} F_p(k, l) z^k t^l \quad (1)$$

which satisfies the following relation:

$$\begin{aligned} F_p(z, t) &= t(z + z^2 + \cdots + z^p) + t \frac{z^{p+1}}{1-z} [1 + F_p(z, t)] \\ &= tz \frac{1-z^p}{1-z} + t \frac{z^{p+1}}{1-z} [1 + F_p(z, t)]. \end{aligned} \quad (2)$$

The first term corresponds to animals with up to  $p$  squares stacked on a single stair. In the second term, the first factor corresponds to a column with more than  $p$  squares on the lowest stair, such that the animal may or not continue to grow on the next stair, as indicated in the second factor.

Equation (2) leads to the generating function

$$F_p(z, t) = \frac{tz}{1-z-tz^{p+1}} \quad (3)$$

which can be expanded to give:

$$\begin{aligned} F_p(z, t) &= tz \sum_{m=0}^{\infty} z^m (1 + tz^p)^m = \sum_{m=0}^{\infty} \sum_n \binom{m}{n} z^{m+pn+1} t^{n+1} \\ &= \sum_{l \geq 1} \sum_{k=(p+1)(l-1)+1}^{\infty} \binom{k-p(l-1)-1}{l-1} z^k t^l. \end{aligned} \quad (4)$$

Comparing equations (1) and (4), the number of animals with area  $k$  living on  $l$  stairs is given by

$$F_p(k, l) = \binom{k-p(l-1)-1}{l-1} \quad (5)$$

when  $k \geq (p+1)(l-1)+1$  and vanishes otherwise. This expression can be rewritten as

$$F_p(k, l) = \binom{l+m-1}{l-1}, \quad m = k - k_{min} = k - [(l-1)(p+1)+1], \quad (6)$$

where  $k_{min}$  is the number of squares needed to cover  $l$  successive stairs with a minimal connected cluster forming the basis of an animal, as shown in figure 1. Thus  $F_p(k, l)$  is also the number of unconstrained configurations of the remaining  $m = k - k_{min}$  squares, placed on top of this basis, which is given by the binomial coefficient in (6).

From the addition/induction relation for the binomial coefficients,

$$\binom{m+1}{n+1} = \binom{m}{n+1} + \binom{m}{n}$$

we deduce the following recursion relation:

$$F_p(k+1, l+1) = F_p(k, l+1) + F_p(k-p, l), \quad k \geq pl+1. \quad (7)$$

The total number of animals with area  $k$  is given by:

$$F_p(k) = \sum_{l=1}^{\infty} F_p(k, l) = \sum_{0 \leq m \leq \frac{k-1}{p+1}} \binom{k-pm-1}{m}. \quad (8)$$

Its generating function is obtained by setting  $t = 1$  in (3) and reads:

$$F_p(z) = \frac{z}{1 - z - z^{p+1}}. \quad (9)$$

According to (7) and (8),  $F_p(k)$  satisfies the recursion relation for the generalized Fibonacci numbers:

$$F_p(k+1) = F_p(k) + F_p(k-p), \quad F_p(1) = \dots = F_p(p+1) = 1.$$

### 3 Mean length

We define the mean length of an animal with area  $k$  as the mean number of successive stairs it occupies:

$$\bar{l}_p(k) = \frac{A_p(k)}{F_p(k)}, \quad A_p(k) = \sum_{l=1}^{\infty} l F_p(k, l). \quad (10)$$

Introducing the generating function

$$A_p(z) = \sum_{k=1}^{\infty} A_p(k) z^k$$

and making use of equations (1), (3) and (9), we obtain:

$$\begin{aligned}
A_p(z) &= \left. \frac{\partial F_p(z, t)}{\partial t} \right|_{t=1} = \frac{z}{1 - z - z^{p+1}} + \frac{z^{p+2}}{(1 - z - z^{p+1})^2} \\
&= \frac{1 - z}{z} \left( \frac{z}{1 - z - z^{p+1}} \right)^2 = (z^{-1} - 1) F_p^2(z).
\end{aligned}$$

With  $F_p^2(z) = \sum_{k=2}^{\infty} F_p^{(2)}(k) z^k$ , the last equation leads to:

$$A_p(k) = [z^k] (z^{-1} - 1) F_p^2(z) = F_p^{(2)}(k+1) - F_p^{(2)}(k) \quad (11)$$

where  $[z^k] f(z)$  is the coefficient of  $z^k$  in  $f(z)$ . According to (9) we have

$$(1 - z - z^{p+1}) F_p^2(z) = \frac{z^2}{1 - z - z^{p+1}} = z F_p(z)$$

so that:

$$\begin{aligned}
F_p(k) &= [z^k] F_p(z) = [z^k] (z^{-1} - 1 - z^p) F_p^2(z) \\
&= F_p^{(2)}(k+1) - F_p^{(2)}(k) - F_p^{(2)}(k-p).
\end{aligned} \quad (12)$$

Hence, combining (10), (11) and (12), the mean length can be rewritten as:

$$\bar{l}_p(k) = 1 + \frac{F_p^{(2)}(k-p)}{F_p(k)}. \quad (13)$$

$F_p^{(2)}(k)$  can be expressed in terms of  $F_p(k)$  using the relation

$$z^2 \frac{d}{dz} \left[ \frac{F_p(z)}{z} \right] = [1 + (p+1) z^p] F_p^2(z)$$

which follows from (9) and leads to:

$$(k-1) F_p(k) = F_p^{(2)}(k) + (p+1) F_p^{(2)}(k-p).$$

Replacing  $F_p^{(2)}(k-p)$  by its value taken from (12), one obtains:

$$(p+1) F_p^{(2)}(k+1) - p F_p^{(2)}(k) = (k+p) F_p(k).$$

Thus we have the recursion relation

$$F_p^{(2)}(k) = \frac{k+p-1}{p+1} F_p(k-1) + \frac{p}{p+1} F_p^{(2)}(k-1), \quad F_p^{(2)}(1) = 0,$$

which can be iterated to give:

$$\begin{aligned} F_p^{(2)}(k) &= \sum_{n=1}^{k-1} \left( \frac{p}{p+1} \right)^n \frac{k+p-n}{p} F_p(k-n) \\ &= \left( \frac{p}{p+1} \right)^k \sum_{n=1}^{k-1} \left( \frac{p+1}{p} \right)^n \frac{p+n}{p} F_p(n), \quad k \geq 2. \end{aligned} \quad (14)$$

Together with equation (13), this leads to the mean length in terms of generalized Fibonacci numbers.

When  $p = 1$  the coefficients  $F_1^{(2)}(k)$  can be obtained more directly than above for a general value of  $p$  (see reference [6], p. 353). They are given by

$$F_1^{(2)}(k) = \frac{2k F_1(k+1) - (k+1) F_1(k)}{5}$$

which, together with (14) for  $p = 1$ , leads to the following identity for ordinary Fibonacci numbers:

$$\sum_{n=1}^{k-1} 2^n (n+1) F_1(n) = 2^k \frac{2k F_1(k+1) - (k+1) F_1(k)}{5}.$$

#### 4 Mean height

The mean height of an animal with area  $k$ , measured from the staircase, is defined as:

$$\overline{h}_p(k) = \frac{B_p(k)}{F_p(k)}, \quad B_p(k) = \sum_{l=1}^{\infty} \frac{k}{l} F_p(k, l) = \sum_{l=1}^{\infty} \frac{k}{l} \binom{k-p(l-1)-1}{l-1}.$$

The absorption/extraction identity gives:

$$\binom{k-p(l-1)}{l} = \frac{k-p(l-1)}{l} \binom{k-p(l-1)-1}{l-1}, \quad l \geq 1.$$

Hence we have

$$B_p(k) = \frac{kp}{k+p} \sum_l \binom{k-p(l-1)-1}{l-1} + \frac{k}{k+p} \sum_{l \geq 1} \binom{k-p(l-1)}{l} \quad (15)$$

where the last sum can be rewritten as:

$$\begin{aligned}
\sum_{l \geq 1} \binom{k - p(l-1)}{l} &= \sum_l \binom{k - p(l-1)}{l} - 1 \\
&= \sum_l \binom{k + p + 1 - p(l-1) - 1}{l-1} - 1.
\end{aligned} \tag{16}$$

Taking into account the combinatorial definition of the generalized Fibonacci numbers in (8), equations (15) and (16) yield

$$B_p(k) = \frac{k}{k+p} [F_p(k+p+1) + p F_p(k) - 1]$$

so that the mean height is finally given by:

$$\overline{h_p}(k) = \frac{k}{k+p} \frac{F_p(k+p+1) + p F_p(k) - 1}{F_p(k)}. \tag{17}$$

## 5 Asymptotic behaviour

In order to obtain the asymptotic behaviour of the generalized Fibonacci numbers, the generating function  $F_p(z)$  in equation (9) can be written as the partial fraction expansion

$$F_p(z) = \frac{C_p}{1 - \phi_p z} + r(z) = \sum_{k=1}^{\infty} C_p (\phi_p z)^k + r(z) \tag{18}$$

where  $\phi_p$  is the reciprocal of the smallest root in modulus of the denominator of  $F_p(z)$ , which is assumed to be a simple root. It satisfies the relations:

$$\phi_p^{p+1} = 1 + \phi_p^p = \frac{\phi_p}{\phi_p - 1}. \tag{19}$$

In equation (18),  $r(z)$  gives the contribution of the  $p-1$  other roots to the partial fraction expansion. It remains finite at  $z = \phi_p^{-1}$ . At large  $k$ -values, the leading contribution to  $F_p(k)$  comes from the first term so that

$$F_p(k) \simeq C_p \phi_p^k, \quad k \gg 1. \tag{20}$$

The calculation of the amplitude  $C_p$  proceeds as follows. Taking the derivative of the inverse of  $F_p(z)$ , we have:

$$\left. \frac{dF_p^{-1}(z)}{dz} \right|_{z=\phi_p^{-1}} = \lim_{z \rightarrow \phi_p^{-1}} \frac{F_p^{-1}(z)}{z - \phi_p^{-1}} = -\frac{\phi_p}{C_p}. \tag{21}$$

Thus, from (9), we deduce

$$C_p = \frac{1}{1 + (p+1)\phi_p^{-p}} = \frac{1}{(p+1)\phi_p - p}$$

where the last expression is obtained using (19). It is easy to verify that a multiple smallest root would lead to a vanishing derivative in (21), which justifies our previous assumption.

Let us now study the behaviour of the mean length at large  $k$ -values. For this we need an asymptotic expression for  $F_p^{(2)}(k)$ . Equation (14) leads to:

$$F_p^{(2)}(k) \simeq C_p \left( \frac{p}{p+1} \right)^k \sum_{n=1}^{k-1} \frac{p+n}{p} \left( \frac{p+1}{p} \phi_p \right)^n.$$

Summing the series, to leading order, one obtains:

$$F_p^{(2)}(k) \simeq \frac{C_p}{(p+1)\phi_p - p} k \phi_p^k. \quad (22)$$

According to equations (13), (20) and (22), the mean length behaves asymptotically as

$$\overline{l_p}(k) = \frac{k}{p + \phi_p^{p+1}} + O(1) = \frac{\phi_p - 1}{(p+1)\phi_p - p} k + O(1)$$

where (19) has been used to reduce the denominator.

For the mean height in equation (17), we have the following asymptotic form:

$$\overline{h_p}(k) = p + \phi_p^{p+1} + O(k^{-1}) = p + \frac{\phi_p}{\phi_p - 1} + O(k^{-1}).$$

These results can be recovered and extended by considering the behaviour when  $k \gg 1$  of  $F_p(k, l)$  in equation (5). It is obtained by expanding  $\ln F_p(k, l)$  to second order near its maximum, making use of the Stirling approximation. The asymptotic distribution is Gaussian and reads:

$$F_p(k, l) \simeq \frac{C_p \phi_p^k}{\sqrt{2\pi \Delta l_p^2}} \exp \left[ -\frac{(l - \overline{l_p}(k))^2}{2\Delta l_p^2} \right].$$



Here  $\overline{l}_p(k)$  and

$$\overline{\Delta l_p^2} = \frac{\phi_p(\phi_p - 1)}{[(p+1)\phi_p - p]^3} k$$

are the leading contributions to the mean value of  $l_p(k)$  and to its mean-square deviation. The prefactor follows from the normalization to  $F_p(k)$ .

Finally let us examine some special values of  $p$ .

The case  $p = 0$  corresponds to the staking of squares on a flat surface [13] for which

$$F_0(k) = 2^{k-1}, \quad F_0^{(2)}(k) = (k-1)2^{k-2}, \quad \phi_0 = 2.$$

The mean length in (13) and the mean height in (17) behave as:

$$\overline{l}_0(k) = \frac{k+1}{2} = \frac{k}{2} + O(1), \quad \overline{h}_0(k) = \frac{2^k - 1}{2^{k-1}} = 2 + O(2^{-k}).$$

The statistics of animals on a staircase with  $p = 1$  is connected to the usual Fibonacci numbers,  $F_1(k) = 1, 1, 2, 3, 5, 8, \dots$ . Then, according to (19),  $\phi_1 = (1 + \sqrt{5})/2$  is the golden mean  $\phi$  and

$$\overline{l}_1(k) = \frac{k}{1 + \phi^2} + O(1), \quad \overline{h}_1(k) = 1 + \phi^2 + O(k^{-1}),$$

in agreement with the results of reference [14].

## References

- [1] M. Bousquet-Mélou, A method for the enumeration of various classes of column-convex polygons, *Discrete Math.* **154** (1996) 1–25.
- [2] M. Bousquet-Mélou, Combinatoire Énumérative, *Université de Bordeaux I Habilitation Report* (1996) p. 37.
- [3] R. Brak and A.J. Guttmann, Exact solution of of the staircase and row-convex polygon perimeter and area generating function, *J. Phys. A* **23** (1990) 4581–4588.
- [4] D. Dhar, Equivalence of the two-dimensional directed-site animal problem to Baxter’s hard square lattice gas model, *Phys. Rev. Lett.* **49** (1982) 959–962.
- [5] D. Dhar, M.K. Phani and M. Barma, Enumeration of directed site animals on two-dimensional lattices, *J. Phys. A* **15** (1982) L279–L284.
- [6] R.L. Graham, D.E. Knuth and O. Patashnik, *Concrete Mathematics: a Foundation for Computer Science*, 2nd edition (Addison-Wesley, Readings, 1994).
- [7] V. Hakim and J.P. Nadal, Exact results for 2D directed animals on a strip of finite width, *J. Phys. A* **16** (1983) L213–L218.
- [8] G.S. Joyce and A.J. Guttmann, Exact results for the generating function of directed column-convex animals on the square lattice, *J. Phys. A* **27** (1994) 4359–4367.
- [9] D.A. Klarner, Some results concerning polyominoes, *Fibonacci Quart.* **3** (1965) 9–20.
- [10] D.A. Klarner and R.L. Rivest, A procedure for improving the upper bound for the number of  $n$ -ominoes, *Can. J. Math.* **13** (1973) 585–602.
- [11] K.Y. Lin, Exact solution of the convex polygon perimeter and area generating function, *J. Phys. A* **24** (1991) 2411–2417.
- [12] J.P. Nadal, B. Derrida and J. Vannimenus, Directed animals in two dimensions: numerical and exact results, *J. Phys.* **43** (1982) 1561–1574.
- [13] V. Privman and N.M. Švrakić, *Directed Models of Polymers, Interfaces, and Clusters: Scaling and Finite-Size Properties* (Springer, New-York, 1989).
- [14] L. Turban, Lattice animals on a staircase and Fibonacci numbers, *J. Phys. A* **33** (2000) 2587–2595.