## Conformal off-diagonal boundary density profiles on a semi-infinite strip

L Turban

Laboratoire de Physique des Matériaux, Université Henri Poincaré (Nancy I), BP 239, F–54506 Vandœuvre lès Nancy Cedex, France

Abstract. The off-diagonal profile  $\phi^{\rm b}_{\rm od}(v)$  associated with a local operator  $\hat{\phi}(v)$  (order parameter or energy density) close to the boundary of a semi-infinite strip with width L is obtained at criticality using conformal methods. It involves the surface exponent  $x^s_{\phi}$  and displays a simple universal behaviour which crosses over from surface finite-size scaling when v/L is held constant to corner finite-size scaling when  $v/L \to 0$ .

The finite-size behaviour of order parameter or energy density profiles has been the subject of much interest during the last two decades following the work of Fisher and de Gennes [1]. These profiles have been studied in the vivinity of the critical point in the mean-field approximation [2], using field-theoretical methods [3] and through exact solutions [4, 5]. Such profiles display universal behaviour at criticality and in two-dimensional systems they can be deduced from ordinary scaling and covariance under conformal transformation [6–17]. A short review can be found in reference [18].

With symmetry-breaking boundary conditions, one may consider diagonal order parameter profiles [6], i.e., ground-state expectation values. Otherwise, off-diagonal profiles can be used with any type of boundary conditions [13]. For the order parameter with Dirichlet boundary conditions, off-diagonal matrix elements must be considered since a diagonal order parameter profile then vanishes for symmetry reasons.

On a strip with fixed boundary conditions at v = 0 and v = L the diagonal order-parameter profile  $\phi(v)$  associated with an operator  $\hat{\phi}$  takes the following form at criticality [6]

$$\phi(v) = \langle 0 | \hat{\phi}(v) | 0 \rangle = \mathcal{A} \left[ \frac{L}{\pi} \sin\left(\frac{\pi v}{L}\right) \right]^{-x_{\phi}} \qquad 0 < v < L \,. \tag{1}$$

The exponent  $x_{\phi}$  is the bulk scaling dimension of  $\hat{\phi}$ ,  $|0\rangle$  is the ground state of the Hamiltonian  $\mathcal{H} = -\ln \mathcal{T}$  where  $\mathcal{T}$  denotes the row-to-row transfer operator on the strip. When  $L \to \infty$  with a fixed value of the ratio v/L, one obtains the bulk finite-size scaling behaviour  $\phi(L) \sim L^{-x_{\phi}}$ . When  $L \to \infty$  while keeping v fixed, one obtains the profile  $\phi(v) \sim v^{-x_{\phi}}$  on the half-plane with fixed boundary conditions which is a consequence of ordinary scaling. Actually the profile on the strip in (1) follows from the profile on the half-plane through the logarithmic conformal transformation  $w = (L/\pi) \ln z$  [6].



Figure 1. Conformal mapping of the half-plane y > 0 on the semi-infinite strip u > 0, 0 < v < L.

The off-diagional critical profile with general symmetric boundary conditions at v = 0 and L is obtained as [13]

$$\phi_{\rm od}(v) = \langle \phi | \hat{\phi}(v) | 0 \rangle \sim \left(\frac{L}{\pi}\right)^{-x_{\phi}} \left[ \sin\left(\frac{\pi v}{L}\right) \right]^{x_{\phi}^{\rm s} - x_{\phi}} \qquad 0 < v < L \,, \qquad (2)$$

where  $|\phi\rangle$  is the lowest excited state of  $\mathcal{H}$  leading to a non-vanishing matrix element. Besides the bulk exponent  $x_{\phi}$  the off-diagonal profile involves the surface scaling dimension  $x_{\phi}^{\rm s}$  of the operator  $\hat{\phi}$ . It can be identified by considering the transformation of the connected two-point correlation function  $\mathcal{G}_{\phi\phi}^{\rm con}(z_1, z_2)$  from the half-plane to the strip under the logarithmic conformal mapping. For the order parameter with fixed boundary conditions,  $x_{\phi}^{\rm s} = 0$ , and (2) gives an off-diagonal profile in agreement with (1). When  $L \to \infty$ , equation (2) shows the crossover from bulk finite-size scaling  $\phi_{\rm od}(L) \sim L^{-x_{\phi}}$  when the ratio v/L is constant, to surface finite-size scaling  $\phi_{\rm od}(L) \sim L^{-x_{\phi}}$  when v is constant, i.e., when  $v/L \to 0$ .

Let us now consider a half-strip in the (u, v)-plane with  $0 < u < \infty$ , 0 < v < Land uniform boundary conditions. If one crosses the semi-infinite strip at  $u \gg L$ the behaviour of the off-diagonal profile will be the same as for the infinite strip in equation (2). A different behaviour is expected close to the boundary of the semiinfinite strip at  $u \ll L$ . The profile should then involve the surface exponent  $x_{\phi}^{s}$  and, when  $v \ll L$  or  $L - v \ll L$ , the corner exponent  $x_{\phi}^{c}$ .

In order to obtain the profiles on the semi-infinite strip, we consider the transformation of the connected two-point correlation function  $\mathcal{G}_{\phi\phi}^{\mathrm{con}}(z_1, z_2)$  from the half-plane  $z = x + \mathrm{i}y$ , y > 0 to the half-strip  $w = u + \mathrm{i}v$ ,  $0 < u < \infty$ , 0 < v < L as shown in figure 1. The two geometries are related by the conformal transformation [6]

$$z = \cosh\left(\frac{\pi w}{L}\right) \tag{3}$$

or

$$x = \cosh\left(\frac{\pi u}{L}\right)\cos\left(\frac{\pi v}{L}\right) \qquad y = \sinh\left(\frac{\pi u}{L}\right)\sin\left(\frac{\pi v}{L}\right) \,.$$
(4)

Going from the half-plane to the half-strip, the dilatation factor is given by:

$$b(z) = \left|\frac{\mathrm{d}z}{\mathrm{d}w}\right| = \left|\frac{\pi}{L}\sinh\left(\frac{\pi w}{L}\right)\right| = \frac{\pi}{L}\left[\sinh^2\left(\frac{\pi u}{L}\right) + \sin^2\left(\frac{\pi v}{L}\right)\right]^{1/2}.$$
 (5)

At criticality, the form of  $\mathcal{G}_{\phi\phi}^{\mathrm{con}}(z_1, z_2)$  is strongly constrained by conformal invariance. Using an infinitesimal special conformal transformation which preserves the surface geometry, one obtains a system of partial differential equation for the connected two-point correlation function on the half-plane, from which the following scaling form is deduced [19]:

$$\mathcal{G}_{\phi\phi}^{\rm con}(z_1, z_2) = (y_1 y_2)^{-x_{\phi}} g(\omega) \qquad \omega = \frac{(x_2 - x_1)^2 + (y_2 - y_1)^2}{y_1 y_2} \,. \tag{6}$$

We are mainly interested in the behaviour of the profile close to the boundary of the half-strip. Thus we may consider the correlations between two points, the first close to the boundary located at u = 0 and the second far from it:

$$\frac{u_1}{L} \ll 1 \qquad 0 < v_1 < L \qquad \frac{u_2}{L} \gg 1 \qquad 0 < v_2 < L \,. \tag{7}$$

Considering (4) and (5) in the limits of equation (7) we have:

$$\begin{aligned} x_2 - x_1 &\simeq \frac{1}{2} \exp\left(\frac{\pi u_2}{L}\right) \cos\left(\frac{\pi v_2}{L}\right) \quad y_2 - y_1 \simeq \frac{1}{2} \exp\left(\frac{\pi u_2}{L}\right) \sin\left(\frac{\pi v_2}{L}\right) \\ y_1 y_2 &\simeq \frac{\pi u_1}{2L} \sin\left(\frac{\pi v_1}{L}\right) \sin\left(\frac{\pi v_2}{L}\right) \exp\left(\frac{\pi u_2}{L}\right) \\ b(z_1) &\simeq \frac{\pi}{L} \sin\left(\frac{\pi v_1}{L}\right) \quad b(z_2) \simeq \frac{\pi}{L} \exp\left(\frac{\pi u_2}{L}\right) \end{aligned}$$
(8)

Thus the crossover variable  $\omega$  defined in (6) takes the form

$$\omega \simeq \frac{L \exp(\pi u_2/L)}{2\pi u_1 \sin(\pi v_1/L) \sin(\pi v_2/L)} \gg 1.$$
(9)

In this limit, ordinary scaling leads to  $g(\omega) \sim \omega^{-x_{\phi}^{s}}$  so that, in the half-plane geometry:

$$\mathcal{G}_{\phi\phi}^{\rm con}(z_1, z_2) \sim \frac{(y_1 y_2)^{x_{\phi}^{\rm s} - x_{\phi}}}{\left[(x_2 - x_1)^2 + (y_2 - y_1)^2\right]^{x_{\phi}^{\rm s}}}.$$
 (10)

The conformal mapping (3) leads to the correlation function in the half-strip geometry [20]

$$\mathcal{G}_{\phi\phi}^{\rm con}(w_1, w_2) \sim b(z_1)^{x_{\phi}} b(z_2)^{x_{\phi}} \mathcal{G}_{\phi\phi}^{\rm con}(z_1, z_2) \,. \tag{11}$$

Making use of (8) in equations (10) and (11), we finally obtain:

$$\mathcal{G}_{\phi\phi}^{\mathrm{con}}(w_1, w_2) \sim u_1^{x_{\phi}^{\mathrm{s}} - x_{\phi}} \left[ \left( \frac{2\pi}{L} \right) \sin\left( \frac{\pi v_1}{L} \right) \right]^{x_{\phi}^{\mathrm{s}}} \left( \frac{L}{\pi} \right)^{-x_{\phi}} \left[ \sin\left( \frac{\pi v_2}{L} \right) \right]^{x_{\phi}^{\mathrm{s}} - x_{\phi}} \exp\left( -\frac{\pi x_{\phi}^{\mathrm{s}} u_2}{L} \right).$$
(12)

In order to identify the different contributions to the two-point correlation function in (12), we can rewrite it using the row-to-row transfer operator  $\mathcal{T}$  on the strip with width L. The two-point correlation function on the semi-infinite strip reads

$$\mathcal{G}_{\phi\phi}(w_1, w_2) = \frac{\langle B|\hat{\phi}(v_1)\mathcal{T}^{u_2}\hat{\phi}(v_2)|0\rangle}{\langle B|\mathcal{T}^{u_2}|0\rangle} = \frac{\sum_n \langle B|\hat{\phi}(v_1)|n\rangle \exp(-u_2 E_n)\langle n|\hat{\phi}(v_2)|0\rangle}{\langle B|0\rangle \exp(-u_2 E_0)}$$
(13)

where  $|0\rangle$  is the ground state of  $\mathcal{H}$  which is selected by the transfer from  $u = u_2$  to  $u = \infty$  and  $|B\rangle$  is a state vector appropriate for the boundary conditions at u = 0. In the case of free boundary conditions, it describes the free summation over the boundary states. In the last expression the summation is over the complete set of eigenstates  $|n\rangle$  of  $\mathcal{H}$  with eigenvalues  $E_n$ .

The connected two-point correlation function is then obtained by substracting the ground-state contribution to the eigenstate expansion:

$$\mathcal{G}_{\phi\phi}^{\mathrm{con}}(w_1, w_2) = \mathcal{G}_{\phi\phi}(w_1, w_2) - \frac{\langle B | \hat{\phi}(v_1) | 0 \rangle}{\langle B | 0 \rangle} \langle 0 | \hat{\phi}(v_2) | 0 \rangle$$

$$\simeq \frac{\langle B | \hat{\phi}(v_1) | \phi \rangle}{\langle B | 0 \rangle} \langle \phi | \hat{\phi}(v_2) | 0 \rangle \exp[-u_2(E_{\phi} - E_0)]$$
(14)

In the last expression we took into account the condition  $u_2 \gg L$ . In this limit, the eigenstate expansion is dominated by the contribution of the lowest excited state  $|\phi\rangle$  of  $\mathcal{H}$  for which the matrix elements are non-vanishing.

Comparing with the conformal expression in (12), we can read the gap-exponent relation [20] in the exponential factor,  $E_{\phi} - E_0 = \pi x_{\phi}^s/L$ , and the off-diagonal profile at  $u_2 \gg L$  in agreement with equation (2). The remaining part can be then identified as the boundary profile on the half-strip at  $u_1 \ll L$  and we obtain

$$\phi_{\rm od}^{\rm b}(v) = \frac{\langle B|\hat{\phi}(v)|\phi\rangle}{\langle B|0\rangle} \sim \left[\frac{\pi}{L}\sin\left(\frac{\pi v}{L}\right)\right]^{x_{\phi}^{\rm s}} \qquad 0 < v < L\,. \tag{15}$$

In the case of the two-dimensional Ising model with free boundary conditions  $\mathcal{H}$  is the Hamiltonian of the Ising model in a transverse field; if one associates the Pauli spin operator  $\sigma_l^x$  (l = 1, L) with the order parameter, then the boundary state vector  $|B\rangle$  is explicitly given by [21]

$$|B\rangle = \prod_{l=1,L} \frac{1}{\sqrt{2}} (|\sigma_l^x = +1\rangle + |\sigma_l^x = -1\rangle) = \prod_{l=1,L} |\sigma_l^z = +1\rangle.$$
(16)

Both  $|0\rangle$  and  $|B\rangle$  are even under the operator  $P = \prod_{l=1,L} \sigma_l^z$  [21]. In the expression of the order parameter profile the state  $|\phi\rangle = |\sigma\rangle$ , which contains a single fermionic excitation, is odd under P. At l = 1, the order parameter profile coincides with the corner magnetization obtained in reference [21]. The surface magnetic exponent is  $x_{\sigma}^s = 1/2$  [22]. For the energy density profile the state  $|\phi\rangle = |\epsilon\rangle$  contains two fermionic excitations and is even under P. The surface energy exponent is then  $x_{\epsilon}^s = 2$  [23, 24].

When  $L \to \infty$  with a fixed v/L value, one obtains the surface finite-size scaling behaviour  $\phi^{\rm b}_{\rm od}(L) \sim L^{-x^{\rm s}_{\phi}}$  while keeping v constant leads to the corner finite-size scaling behaviour

$$\phi_{\rm od}^{\rm b}(L) \sim v^{x^{\rm s}_{\phi}} L^{-2x^{\rm s}_{\phi}} \qquad v \ll L.$$
 (17)

Thus the corner exponent  $x_{\phi}^{c}(\pi/2)$  is given by  $2x_{\phi}^{s}$ . This result is in agreement with the general expression  $x_{\phi}^{c}(\theta) = \pi x_{\phi}^{s}/\theta$  for a corner with opening angle  $\theta$ , which also follows from conformal invariance [19, 21].

The Laboratoire de Physique des Matériaux is Unité Mixte de Recherche CNRS No 7556.

## References

- [1] Fisher M E and de Gennes P G 1978 C. R. Acad. Sci. Paris B 287 207
- Binder K 1983 Phase Transitions and Critical Phenomena vol 8 ed C Domb and J L Lebowitz (London: Academic Press) p 1
- Diehl H W 1986 Phase Transitions and Critical Phenomena vol 10 ed C Domb and J L Lebowitz (London: Academic Press) p 75
- [4] Au-Yang H and Fisher M E 1980 Phys. Rev. B 21 3956
- [5] Bariev R Z 1988 Theor. Math. Phys. 77 1090
- [6] Burkhardt T W and Eisenriegler E 1985 J. Phys. A: Math. Gen. 18 L83
- [7] Burkhardt T W and Cardy J L 1987 J. Phys. A: Math. Gen. 20 L233
- [8] Burkhardt T W and Guim I 1987 Phys. Rev. B 36 2080
- [9] Cardy J L 1990 Phys. Rev. Lett. 65 1443
- [10] Burkhardt T W and Xue T 1991 Phys. Rev. Lett. 66 895
- [11] Burkhardt T W and Xue T 1991 Nucl. Phys. B **354** 653
- [12] Burkhardt T W and Eisenriegler E 1994 Nucl. Phys. B **424** 487
- [13] Turban L and Iglói F 1997 J. Phys. A: Math. Gen. **30** L105
- [14] Iglói F and Rieger H 1997 Phys. Rev. Lett. 78 2473
   [15] Karevski D and Henkel M 1997 Phys. Rev. B 55 6429
- [16] Carlon E and Iglói F 1998 Phys. Rev. B 57 7877
- [10] Carlon E and Igiol F 1998 *Phys. Rev.* B **51** 1811
- [17] Karevski D, Turban L and Iglói F 2000 J. Phys. A: Math. Gen. 33 2663
- [18] Henkel M 1999 Conformal Invariance and Critical Phenomena (Berlin: Springer) p 346
- [19] Cardy J L 1984 Nucl. Phys. B 240 514
- [20] Cardy J L 1984 J. Phys. A: Math. Gen. 17 L385
- [21] Barber M N, Peschel I and Pearce PA 1984 J. Stat. Phys. 37 497
- [22] McCoy B M and Wu T T 1973 The Two-Dimensional Ising Model (Cambridge: Harvard University Press) p 132
- [23] von Gehlen G and Rittenberg V 1986 J. Phys. A: Math. Gen. 19 L631
- [24] Cardy J L 1986 Nucl. Phys. B 275 200