## Ground states of a system of interacting particles in a parabolic trap

M. S. Hussein and O.K. Vorov

Instituto de Fisica, Universidade de Sao Paulo Caixa Postal 66318, 05315-970, Sao Paulo, SP, Brasil

(13 June 2001)

## Abstract

Within the "lowest Landau level approximation", we develop a method to find the ground state of a 2d system of interacting particles confined by a parabolic potential.

Keywords: Ground-state, 2d-systems, BEC

The problem of finding exact eigenstates of a 2d system of particles interacting by the two-body central forces V(r) and confined by an external parabolic potential arises in many subfields, including cooled and trapped fermionic atoms, Bose-Einstein condensates in traps, quantum dots and electrons in magnetic field. Assuming spherical symmetry, we are interested here in the lowest energy states of such system as a function of exact quantum numbers. The lowest Landau level approximation is equivalent to the weak interaction limit

$$V(r) \ll \hbar \omega$$

where  $\omega$  is the oscillator frequency of the external trapping potential. The effective Hamiltonian of the N-body system is the sum of the energy of degenerate level [1] with total angular momentum L ( $\hbar=m=1$ )

$$H = \omega(L+N) + \tilde{V} = P(L) \left[ \sum_{i}^{N} \left( \vec{p}_{i}^{2} + \omega^{2} \vec{r}_{i}^{2} \right) / 2 + \sum_{i>j} V(r_{ij}) \right] P(L)$$
(1)

which comes from the noninteracting Hamiltonian in the 2d parabolic trap, and the projected interaction  $\tilde{V}$ . Here, P(L) is the projector onto the space of lowest Landau level, S. The space S is formed by the homogeneous polynomials in  $z_i=x_i+iy_i$  of degree L times the Gaussian factor  $|0\rangle=e^{-\frac{1}{2}\sum |z_k|^2}$ . Hereafter, we set  $\omega=1$  for simplicity. The polynomials are antisymmetric (symmetric) for identical fermions (bosons). The eigenstates of the H (1) are found from diagonalization of  $\tilde{V}$  in the space S.

The effective interaction  $\tilde{V}$  can be found in the form [2]

$$\tilde{V} = \sum_{k=0}^{[L/2]} V_k, \qquad V_k \equiv s_{2k} (B^{2k} - B^{2k-1}) + (s_{2k+2} - s_{2k+1}) B^{2k+1}, \tag{2}$$

where  $B^k = \sum_{j>i}^N (a_i^{\dagger} - a_j^{\dagger})^k (a_i - a_j)^k$  and  $a_i^{\dagger} = z_i/2 - \partial/\partial z_i^*$  and  $a_i = z_i^*/2 + \partial/\partial z_i$  are the standard ladder operators. Here,  $B^0 \equiv N(N-1)/2$  and  $B^{-1} \equiv 0$ ; [k] denotes the integer part of k. In (2),  $s_k = \int_0^\infty \frac{dt}{k!} M(k+1,1,-t) V(\sqrt{2t})$  with M the Kummer function.

In order to find the ground state without solving the whole spectrum, we use the following idea [2]. Suppose that the Hamiltonian can be written as a sum

$$\tilde{V} = V_0 + V_S,$$
 (i)  $V_S|0) = 0,$  (ii)  $V_0|0) = E_{min}|0),$  (3)

such that the second term (i) annihilates a state  $|0\rangle$ , and (ii)  $|0\rangle$  is the ground state for the first term (this ground state may be degenerate). If (iii) the second term is non-negative definite (i.e., it has no negative eigenvalues)

$$V_S \ge 0,\tag{4}$$

then the state  $|0\rangle$  will still be the ground state of the full Hamiltonian  $V_0 + V_S$ , with the same eigenvalue  $E_{min}$  [2].

In the case of Bose system, the polynomials of the basis states are  $S_L = P_S m_L P_S$ , where  $m_L = z_1^{l_1} z_2^{l_2} \dots z_N^{l_N}$  is the monomial with a given partition of the integer  $L = \sum_n l_n$ , and  $P_S$  is the operator of symmetrization over N variables. The above decomposition (3,4) with the properties (i) and (ii) can be found by inspecting action of (2) on the basis states. The simplest basis state

$$|0_L\rangle = P_S z_1 z_2 \dots z_L |0\rangle, \qquad (\tilde{V} - V_0)|0_L\rangle = 0,$$
(5)

is annihilated by any term in (2), except for  $V_0$ . The term  $V_0$  can be written as

$$V_0 = (N/2)[(N-1)s_0 - (L-v)(s_1 - s_2)].$$
(6)

Here,

$$v = \frac{1}{N} \sum_{j,k}^{N} a_j^{\dagger} a_k$$

is the collective contribution to the angular momentum, which is also a conserved quantum number,  $[v, \tilde{V}] = 0$ . The allowed values of v are 0, 1, 2, ..., L - 2, L [2]. The state (5) can be represented as a sum

$$|0_L) = \sum_{v=0}^N |0_{L,v}\rangle, \qquad |0_{L,v}\rangle = Z^v P_S \prod_{k=1}^{L-v} (z_k - Z)|0\rangle, \tag{7}$$

where  $Z = \sum_{k=1}^{N} z_k / N$  is the collective variable. Each term  $|0_{L,v}\rangle$  is the eigenvector of vand therefore of  $V_0$ . Each v-sector is represented by a single state  $|0_{L,v}\rangle$  in expansion (7). The spectrum of  $V_0$  consists of L equidistant degenerate levels. Conditions (i) and (ii) are therefore satisfied for the states  $|0_{L,v}\rangle$ . In particular, this means that those are eigenstates of  $\tilde{V}$  with eigenvalues given by (6) [2], see also [3].

If the term  $V_S \equiv \tilde{V} - V_0$  is non-negative definite, the condition (iii) is satisfied and each state  $|0_{L,v}\rangle$  will be the ground state in corresponding sector L, v. In fact, it is possible to find a broad universality class of the potentials V(r) which give  $V_S \equiv \tilde{V} - V_0 \ge 0$ . We work in the space S and use permutation symmetry to write  $P_S V_S P_S$  in the form

$$P_S \sum_{i>j} V_{S,ij} P_S = \frac{N(N-1)}{2} P_S V_{S,12} P_S = \frac{N(N-1)}{2} P_S p_{12} V_{S,12} p_{12} P_S$$
(8)

where  $V_{S,ij}$  is the contribution from pair of particles i, j to  $V_S$  (2,3), and  $p_{ij}$  is the operator of symmetrization over this pair. The condition  $p_{12}V_{S,12}p_{12} \ge 0$  is a sufficient condition for  $P_SV_SP_S \ge 0$  [2]. The eigenvalues of  $p_{12}V_{S,12}p_{12}$  can be easily evaluated, using the full space of monomials  $m_L$ . As a result, the condition  $p_{12}V_{S,12}p_{12} \ge 0$  reduces to the following set of integral inequalities

$$\int_{0}^{\infty} dt V(r) e^{-r^{2}/2} \left[ \frac{r^{4n}}{2^{2n}(2n)!} - 1 + n \left( 1 - \frac{r^{4}}{8} \right) \right] \ge 0, \tag{9}$$

which must be satisfied for integer  $n, 0 \leq 2n \leq L$ . These conditions describe the broad universality class of interaction potentials V(r), for which the lowest energy states in sectors with quantum numbers v and  $L \leq N$  are given by  $|0_{L,v}\rangle$  [see Eq.(7)], and their energies are equal to the eigenvalues of  $V_0$ , Eq.(6), with  $s_0 = \int_0^\infty dt e^{-t} V(\sqrt{2t})$ ,  $s_1 - s_2 = \int_0^\infty dt e^{-t}$  $\left(\frac{1}{2} - \frac{t^2}{4}\right) V(\sqrt{2t})$ . Conditions (9) are valid, in particular, for the Gaussian interaction  $V(r) = U_0 \frac{e^{-r^2/R^2}}{\pi R^2}, U_0 \geq 0$ , for the  $\delta$ -function interaction  $V = U_0 \delta(\vec{r}), U_0 \geq 0$  and for the 2d log-Coulomb interaction  $V = U_0 log(1/r), U_0 \geq 0$  and for many others potentials, which are predominantly repulsive. These results are obtained for the Bose case. Generalization of the method for the Fermi case is possible.

The work was supported by FAPESP.

## REFERENCES

- R.B. Laughlin, Phys. Rev. Lett. **50** (1983) 1395; N. K. Wilkin, J. M. Gunn, and R. A.
   Smith, Phys. Rev. Lett. **80** (1998) 2265; G. F. Bertsch and T. Papenbrock, Phys. Rev.
   Lett. **83** (1999) 5412; B. Mottelson, Phys. Rev. Lett. **83** (1999) 2695.
- [2] M.S. Hussein and O.K. Vorov, subm. to Phys. Rev. Lett., cond-mat/0102505.
- [3] T. Papenbrock and G.F. Bertsch, J. Phys. A34 (2001) 603; R.A. Smith and N.K. Wilkin, Phys. Rev. A62 (2000) 061602.