Large-N expansion based on the Hubbard-operator path integral representation and its application to the t-J model

Adriana Foussats and Andrés Greco

Facultad de Ciencias Exactas Ingeniería y Agrimensura and Instituto de Física Rosario (UNR-CONICET). Av. Pellegrini 250-2000 Rosario-Argentina.

(October 29, 2018)

In the present work we have developed a large-N expansion for the t-J model based on the path integral formulation for Hubbard-operators. Our large-N expansion formulation contains diagrammatic rules, in which the propagators and vertex are written in term of Hubbard operators. Using our large-N formulation we have calculated, for J=0, the renormalized O(1/N) boson propagator. We also have calculated the spin-spin and charge-charge correlation functions to leading order 1/N. We have compared our diagram technique and results with the existing ones in the literature.

PACS: 71.10.-w,71.27.+a

I. INTRODUCTION

The role of electronic correlations is an important and open problem in solid state physics. Its close connection with the phenomena of high- T_c superconductivity¹ makes this problem relevant in present days.

One of the most popular models in the context of high- T_c superconductivity is the t-J model. A natural representation of the t-J model is in terms of Hubbard operators^{2,3}. Although the t-J model is quadratic, when is written using Hubbard-operators³, there are several difficulties in the calculation of physical quantities. These difficulties are mainly: a)the complicate commutation rules of the Hubbard operators² (X-operators) and, b) any of the model parameters can be taken to be small in a perturbative evaluation of the observables.

To handle the point a), a popular method is to use slave particles (slave boson and fermion) to decouple the original X-operator in usual bosons and fermions⁴. For point b), one of the nonperturbative techniques (which will be relevant in the present paper) is the large-N expansion. N is the number of electronic degrees of freedom per site (see below) and 1/N can be considered as a small parameter.

The large-N expansion was developed in the framework of slave boson representation^{5,6} and, using Baym-Kadanoff⁷ functional theory in terms of the X-operators^{8,9}.

In this paper we develop a large-N expansion for the t-J model based on the path integral representation for X-operators.

In a series of papers^{10,11} we have studied the path integral formulation for Hubbard-operators. Our point of view is to write a path integral for t-J model by using directly the Hubbard X-operators as fields variables, without any decoupling. Our starting point was the construction of a particular family of first-order constrained Lagrangians by using the Faddeev-Jackiw and Dirac methods¹². We showed that the constrains of the model are second class. The canonical quantization of

this constrained theory leads to the commutation rules of the Hubbard-operators. Next, by using path-integral techniques, the correlation functional and effective Lagrangian were constructed.

In Ref. [11], we found a particular family of constrained Lagrangians and showed that the corresponding path-integral can be mapped to that of the slave-boson representation in the radial gauge 13,5. The path-integral formulation defined in [11] is the starting point for our large-N expansion.

In section II we give a summary of the formalism which will be useful for understanding our large-N expansion. In section III, we develop our large-N method. We present explicit calculations for the case J=0 of the t-J model (U-infinite Hubbard model). In section IV, we calculate spin-spin and charge-charge correlation functions to leading order of large-N expansion. Finally, in section V we give our main discussions and conclusions.

II. PRELIMINARIES AND DEFINITIONS

In this section we give a summary of the main results obtained in [11], which are the starting point for our large-N expansion.

As we mentioned in the introduction, although the $t\!-\!J$ model is quadratic when is written using X-operators, the complicate commutation rules between them make this model highly nontrivial. To attack the problem, our method works as follows:

a) We construct a Lagrangian written in terms of Hubbard operators

$$L = \sum_{i} a_{i\alpha\beta}(X) \dot{X}_{i}^{\alpha\beta} - H(X)$$
 (1)

where the t-J model Hamiltonian H(X), with the addition of a chemical potential μ , reads as:

$$H(X) = \sum_{i,j,\sigma} t_{ij} X_i^{\sigma 0} X_j^{0\sigma} + \frac{1}{2} \sum_{ij,\sigma} J_{ij} (X_i^{\sigma \bar{\sigma}} X_j^{\bar{\sigma}\sigma} - X_i^{\sigma \sigma} X_j^{\bar{\sigma}\bar{\sigma}}) - \mu \sum_{i,\sigma} X_i^{\sigma \sigma}.$$

$$(2)$$

In (1) the coefficients $a_{i\alpha\beta}(X)$ are unknown and must be determined.

The indices α and β can take both 0 value (empty state) or spin index $\sigma = \pm$ (up and down state, respectively). The five Hubbard \hat{X} -operators $X^{\sigma\sigma'}$ and X^{00} are boson-like and the four Hubbard \hat{X} -operators $X^{\sigma 0}$ and $X^{0\sigma}$ are fermion-like². In (2), t_{ij} and J_{ij} are hoping and exchange parameters, respectively, between sites i and j.

- b) We introduce (via Lagrangian multipliers $\lambda^{\alpha\beta}$) a set of bosonic constrains $\Omega^{\alpha\beta}$ which must be also determined.
- c) After these definitions and proposals, we impose that the Dirac brackets¹² between X-variables obtained using our constrained theory verify the correct commutation rules for Hubbard operators. With this prescription, we obtain a set of differential equations for the coefficients $a_{i\alpha\beta}(X)$ and constrains Ω_a . A particular solution for the coefficients is

$$a_{i0\sigma} = \frac{i}{2X_i^{00}} X_i^{\sigma 0} , \qquad a_{i\sigma 0} = \frac{i}{2X_i^{00}} X_i^{0\sigma}$$
 (3)

and for the constrains,

$$\Omega_i^{00} = X_i^{00} + \sum_{\sigma} X_i^{\sigma\sigma} - 1 = 0 , \qquad (4)$$

$$\Omega_i^{\sigma\sigma'} = X_i^{\sigma\sigma'} - \frac{X_i^{\sigma 0} X_i^{0\sigma'}}{X_i^{00}} = 0.$$
(5)

The boson-like Lagrangian coefficients are all zero. The constrains are second class because none of them commutes with all the other ones.

Consequently, the dynamics is given by the Lagrangian

$$L(X, \dot{X}) = -\frac{i}{2} \sum_{i,\sigma} \frac{(\dot{X_i}^{0\sigma} X_i^{\sigma 0} + \dot{X_i}^{\sigma 0} X_i^{0\sigma})}{X_i^{00}} - H(X) \quad (6)$$

and the constrains (4) and (5).

Now, in order to write the path integral representation we use the method of Faddeev-Senjanovich¹⁴. This method is currently used in quantum field theory to obtain the path integral of constrained systems.

The partition function reads as:

$$Z = \int \mathcal{D}X_i^{\alpha\beta} \, \delta[X_i^{00} + \sum_{\sigma} X_i^{\sigma\sigma} - 1] \, \delta[X_i^{\sigma\sigma'} - \frac{X_i^{\sigma 0} X_i^{0\sigma'}}{X_i^{00}}]$$

$$\times \left(sdet M_{AB} \right)_i^{\frac{1}{2}} exp \left(i \int dt \, L(X, \dot{X}) \right). \tag{7}$$

where the superdeterminant

$$(sdet M_{AB})^{\frac{1}{2}} = 1/\frac{1}{(-X_{\cdot}^{00})^2}$$
 (8)

is just the superdeterminant of the Dirac matrix¹² (or equivalently the sympletic matrix of the Faddeev-Jackiw method) formed with the set of all the second class constrains of the theory.

The power two on X^{00} in $(sdet M_{AB})^{\frac{1}{2}}$ appears because we are working with two spin projections. This point will be very important for our large-N extension in the next section.

Note the first constrain (4) (the first δ -function in (7)) is the completeness condition which means that "double occupancy" at each site is forbidden. The constrains (5) (the second δ -function in (7)) is obtained in our formalism as a consequence of imposing that the commutation rules of X-operators must be fulfilled.

From the above equations our theory corresponds to a configuration in which the bosons are totally constrained and the dynamics is carried out only by the fermions. As it was shown in Ref[11], the path integral formalism corresponding to this dynamical situation is mapped in the slave boson representation.

Equation (7) is our main formula to develop the large-N expansion.

III. THE LARGE-N EXPANSION

In this section, we present the large-N expansion in the framework of the path integral representation for Hubbard operators. We present our explicit results for the J=0 case (U-infinite Hubbard model) which will be very useful to show how our method works and to compare our results with previously ones found in the literature.

Our starting point is the partition function (7) written in the Euclidean form:

$$Z = \int \mathcal{D}X_{i}^{\alpha\beta} \, \delta[X_{i}^{00} + \sum_{\sigma} X_{i}^{\sigma\sigma} - 1] \, \delta[X_{i}^{\sigma\sigma'} - \frac{X_{i}^{\sigma0} X_{i}^{0\sigma'}}{X_{i}^{00}}]$$

$$\times \left(sdet M_{AB} \right)_{i}^{\frac{1}{2}} exp \left(- \int d\tau \, L_{E}(X, \dot{X}) \right) . \tag{9}$$

The Euclidean Lagrangian $L_E(X, X)$ in (9) is:

$$L_E(X, \dot{X}) = \frac{1}{2} \sum_{i,\sigma} \frac{(\dot{X}_i^{0\sigma} X_i^{\sigma 0} + \dot{X}_i^{\sigma 0} X_i^{0\sigma})}{X_i^{00}} + H(X) . \quad (10)$$

First, we make the following steps in the path integral (9):

- a) we integrate over the boson variables $X^{\sigma\sigma'}$ using the second δ -function in (9),
- b) the spin index $\sigma = \pm$, is extended to a new index p running from 1 to N. In order to get a finite theory in the N-infinite limit, we re-scale the hoping t_{ij} to t_{ij}/N in the Hamiltonian,

- c) the completeness condition $(X_i^{00} + \sum_p X_i^{pp} = N/2)$ can be exponentiated, as usual, by using the Lagrangian multipliers λ_i ,
- d) in order to prepare the path integral for the large-N expansion, we write the boson fields in terms of static mean-field values and dynamic fluctuations

$$X_i^{00} = Nr_0(1 + \delta R_i) , \qquad (11)$$

$$\lambda_i = \lambda_0 + \delta \lambda_i \,\,, \tag{12}$$

e) finally, we make the following change of variables

$$f_{ip}^{+} = \frac{1}{\sqrt{Nr_o}} X_i^{p0}, \tag{13}$$

$$f_{ip} = \frac{1}{\sqrt{Nr_o}} X_i^{0p},\tag{14}$$

By following the steps a-e, we find the effective Lagrangian for the J=0 case

$$L_{eff} = -\frac{1}{2} \sum_{i,p} \left(\dot{f}_{ip} f_{ip}^{+} + \dot{f}_{ip}^{+} f_{ip} \right) \frac{1}{(1 + \delta R_i)}$$

$$+ \sum_{i,j,p}^{N} t_{ij} \ r_o f_{ip}^{+} f_{jp} - (\mu - \lambda_0) \sum_{i,p} f_{ip}^{+} f_{ip} \frac{1}{(1 + \delta R_i)}$$

$$+ N \ r_0 \sum_{i} \delta \lambda_i \ \delta R_i + \sum_{i,p} f_{ip}^{+} f_{ip} \frac{1}{(1 + \delta R_i)} \ \delta \lambda_i$$

$$+ L_{ghost}. \tag{15}$$

In (15), L_{ghost} is defined via the exponentiation of the superdenterminant using ghost fields. After the extension to large-N, the superdeterminant becomes:

$$(sdet M_{AB})_i^{\frac{1}{2}} = \frac{(Nr_0)^N}{\left(\frac{-1}{1+\delta R_i}\right)^N}.$$
 (16)

The numerator of (16) together with the Jacobian of the transformations (11-14) contribute to the path-integral normalization factor. The denominator of (16) can be seen as a $N \times N$ diagonal matrix and the integral representation is given in terms of complex boson ghost fields \mathcal{Z}_p . Therefore,

$$L_{ghost}(\mathcal{Z}) = -\sum_{ip} \mathcal{Z}_{ip}^{\dagger} \left(\frac{1}{1 + \delta R_i} \right) \mathcal{Z}_{ip} . \tag{17}$$

To implement the 1/N expansion, the nonpolynomial L_{eff} should be developed in powers of δR . Up to order 1/N, it is sufficient to retain terms up to δR^2 . Then,

$$L_{eff} = -\frac{1}{2} \sum_{i,p}^{N} \left(\dot{f}_{ip} f_{ip}^{+} + \dot{f}_{ip}^{+} f_{ip} \right) (1 - \delta R_{i} + \delta R_{i}^{2})$$

$$+ \sum_{i,j,p}^{N} t_{ij} r_{o} f_{ip}^{+} f_{jp} - \mu \sum_{i,p} f_{ip}^{+} f_{ip} (1 - \delta R_{i} + \delta R_{i}^{2})$$

$$+ N r_{0} \sum_{i} \delta \lambda_{i} \delta R_{i} + \sum_{i,p} f_{ip}^{+} f_{ip} (1 - \delta R_{i} + \delta R_{i}^{2}) \delta \lambda_{i}$$

$$- \sum_{ip} \mathcal{Z}_{ip}^{\dagger} \left(1 - \delta R_{i} + \delta R_{i}^{2} \right) \mathcal{Z}_{ip}, \tag{18}$$

where we have changed μ to $\mu - \lambda_0$ and dropped constant and linear terms in the fields.

Looking at the effective Lagrangian (18), the Feynman rules can be obtained as usual. The bilinear parts give rise to the propagators and the remaining pieces are represented by vertices. Besides, we assume the equation (18) written in the momentum space once the Fourier transformation was performed.

In summary, the Feynman rules are:

i) Propagators: We associate with the two component $\delta X^a=(\delta R\ ,\ \delta \lambda)$ boson field, the propagator

$$D_{(0)ab}(q,\omega_n) = \begin{pmatrix} 0 & \frac{1}{Nr_0} \\ \frac{1}{Nr_0} & 0 \end{pmatrix}$$
 (19)

which is represented by a dashed line in Fig.1 connecting two generic components a and b.

The quantities q and ω_n are the momentum and the Bose Matsubara frequency of the bosonic field, respectively.

We associate with the N-component fermion field f_p , the propagator

$$G_{(0)pp'}(k,\nu_n) = -\frac{\delta_{pp'}}{i\nu_n - (\varepsilon_k - \mu)}$$
 (20)

which is represented by a solid line in Fig.1 connecting two generic components p and p'.

In (20) the electron bare dispersion was defined as $\varepsilon_k = -2tr_o(\cos k_x + \cos k_y)$, where t is the hopping between nearest neighbors sites on the square lattice.

The quantities k and ν_n are the momentum and the Fermionic Matsubara frequency of the fermionic field, respectively.

We associate with the N-components ghost field $\mathcal{Z}_{p'}$, the propagator

$$\mathcal{D}_{pp'} = -\delta_{pp'} \tag{21}$$

which is represented by a dotted line in Fig.1 connecting two generic components p and p'.

ii) Vertices: The expressions of the different three-leg and four-leg vertices are

$$\Lambda_a^{pp'} = (-1) \left(\frac{i}{2} (\nu + \nu') + \mu ; 1 \right) \delta^{pp'}$$
(22)

representing the interaction between two fermions and one boson (Fig.1a);

$$\Lambda_{ab}^{pp'} = (-1) \begin{pmatrix} -\frac{i}{2} (\nu + \nu') - \mu & -\frac{1}{2} \\ -\frac{1}{2} & 0 \end{pmatrix} \delta^{pp'}, \qquad (23)$$

representing the interaction between two fermions and two bosons (Fig.1b);

$$\Gamma_{nn'}^a = (-1)(\delta_{pp'}, 0),$$
 (24)

representing the interaction between two ghosts and one boson (Fig.1c); and

$$\Gamma_{pp'}^{ab} = (-1) \begin{pmatrix} -1 & 0 \\ 0 & 0 \end{pmatrix} \delta_{pp'}$$
(25)

representing the interaction between two bosons and two ghosts (Fig.1d). Each vertex conserves the momentum and energy.

In the N-infinite limit (order O(1)), we only have free fermions with renormalized band due to correlations, $\varepsilon_k = -2tr_0(\cos k_x + \cos k_y)$. For a given value of μ , r_0 (which from (11) and the completeness condition is equal to $\delta/2$, where δ is the hole doping away from half-filling) must be determined self-consistently. This result is in agreement with previous calculation 13,8.

The bare boson propagator (19), which is of order O(1/N), is renormalized by a series of electronic bubbles which contribute also in O(1/N). Then, in O(1/N), the boson propagator is dressed by electronic interactions.

By looking at the diagrammatic, it can be seen that the irreducible boson self-energy Π_{ab} is given by the sum of the contributions corresponding to the one-loop diagrams (in order 1/N) seen in Fig.1e.




FIG. 1. Types of vertex, up to order O(1/N), between two fermions and one boson (a), two fermions and two bosons (b), two ghosts and one boson (c) and, two bosons and two ghosts (d). (e) represents the sum of all one-loop diagrams contributing to the irreducible boson-self energy.

The presence of the last two diagrams in Fig.1e involving ghost fields is very important. It is possible to show that these two diagrams exactly cancel the infinities, due to the frequency dependence of our vertices, of the two first diagrams in Fig.1e. Therefore, using our Feynman rules, the final (and finite) result for each component of the boson-self energy Π_{ab} is:

$$\Pi_{RR} (q, \omega_n) = -\frac{N}{N_s} \frac{1}{4} \sum_k \left[2 n_F(\varepsilon_k - \mu)(\varepsilon_{k+q} - \varepsilon_k) + (\varepsilon_{k+q} + \varepsilon_k)^2 \frac{\left[n_F(\varepsilon_{k+q} - \mu) - n_F(\varepsilon_k - \mu) \right]}{-i\omega_n + \varepsilon_{k+q} - \varepsilon_k} \right], (26)$$

$$\Pi_{\lambda R}(q,\omega_n) = -\frac{N}{N_s} \frac{1}{2} \sum_k \left(\varepsilon_{k+q} + \varepsilon_k \right) \\
\times \frac{\left[n_F(\varepsilon_{k+q} - \mu) - n_F(\varepsilon_k - \mu) \right]}{-i\omega_n + \varepsilon_{k+q} - \varepsilon_k}$$
(27)

and.

$$\Pi_{\lambda\lambda}(q,\omega_n) = -\frac{N}{N_s} \sum_{k} \frac{\left[n_F(\varepsilon_{k+q} - \mu) - n_F(\varepsilon_k - \mu) \right]}{-i\omega_n + \varepsilon_{k+q} - \varepsilon_k} . (28)$$

From the Dyson equation $(D_{ab})^{-1} = (D_{(0)ab})^{-1} - \Pi_{ab}$ the dressed components D_{ab} of the boson propagator can be found.

Our expressions (26)-(28) agree with the previously irreducible boson self energies calculated in the context of the large-N slave boson approach⁵.

The ghost fields interact only with the boson fields (Fig.1c and Fig.1d). The only role of the ghost fields, in order 1/N, is to cancel infinities in the boson self energy Π_{ab} due to the frequency dependence of our vertices (22) and (23). For the evaluation of quantities to higher orders than 1/N, we must include and check the contribution of the ghost fields in each particular calculation. Moreover it should be noted that in this case we must consider terms beyond δR^2 in developing the nonpolinomial parts of L_{eff} . Although this seems to be complicated, all the contributions can be controlled by the small number 1/N via the vertices and propagators. At this point we want to remark that so far not many calculations go beyond the order 1/N. In order 1/N, diagramatic rules are well defined in the present paper.

In summary, we have developed a diagrammatic technique for a large-N expansion in the same spirit of the large-N expansion in quantum field theory. It means, via the order of the propagators and vertices, we can determine the order of the diagram contribution. Note that our Green's functions are calculated in terms of the original Hubbard operators. For example, from (13) and (14) we can see that our fermions f_{ip} are always proportional to the Fermi-like X-operator X^{op} and not only in the leading order like in the slave-boson formulation¹³.

IV. CHARGE-CHARGE AND SPIN-SPIN CORRELATION FUNCTION

In this section we calculate numerically the chargecharge and spin-spin correlation functions on the square lattice for the nearest neighbor hopping t. We choose t = 1.0 as the energy unit and the temperature T = 0.0K. We can define retarded density-density \tilde{D} and spin-spin \tilde{S} Green's functions as 15

$$\tilde{D}_{ij} = \frac{1}{N} \sum_{pq} \langle T_{\tau} X_i^{pp} X_j^{qq} \rangle \tag{29}$$

and

$$\tilde{S}_{ij} = \langle T_{\tau} X_i^{pq} X_i^{qp} \rangle, \tag{30}$$

respectively.

Using $\sum_q X_i^{qq} = N/2 - X_i^{00}$ and (11) we find for \tilde{D} , in O(1), in the Fourier space

$$\tilde{D}(Q,\omega_n) = -N(\frac{\delta}{2})^2 D_{RR}(Q,\omega_n)$$
(31)

Using the constrains (5), it is easy to prove that \tilde{S} , in O(1), is the electron bubble formed with two bare electron Green's functions (20). The analytical expression for \tilde{S} agrees with $\Pi_{\lambda\lambda}$.

In Fig.2, we show $D(Q,\omega) = -Im[\tilde{D}(Q,\omega)]$ and $S(Q,\omega) = -Im[\tilde{S}(Q,\omega)]$ using $\eta = 0.1t$ in the analytical continuation $i\omega_n = \omega + i\eta$.

To make explicit comparison with exact diagonalization 16 and analytical 15 calculation we choose the same densities and momenta Q of these two references.

In Fig.2a, we plot $D(Q,\omega)$ (solid line) and $S(Q,\omega)$ (dashed line) for doping $\delta=0.12$. In Fig.2b, we present similar results for doping $\delta=0.78$. In agreement with analytical calculation based on X-operator canonical approach¹⁵ (see also slave boson calculation¹³), there is no collective excitations (like magnons) in the spin-spin correlation function to leading order of the 1/N expansion. The spin-spin correlation functions is determined by the particle-hole bubble with the renormalized electron dispersion by correlations. In contrast to $S(Q,\omega)$, $D(Q,\omega)$ contains collective excitations (zero sound) that can live inside or outside of the spectrum¹⁵.

For low doping (Fig.2a) the collective excitations lie outside the particle-hole spectrum. Therefore, they show up as sharp peaks. In contrast, for high doping (Fig. 2b) the collective excitations lie inside the particle-hole spectrum and they show up as broad peaks. The dominance of collective excitations in D, for low doping, can be clearly seen for $Q = (\pi, \pi)$.

It is also very important to point out that spin and charge fluctuations occur at low doping (Fig.2a), near half filling, on different energy scales. This picture disappears at high doping as it can be seen in Fig.2b. This behavior is not present in the uncorrelated case where spin and charge excitations are essentially degenerate.

The agreement between our results and previous ones (see Fig.1 of Ref[15]) is excelent, proving the consistency of our large-N method.




FIG. 2. Spin-spin (dashed lines) and charge-charge (solid lines) correlation functions as a function of ω for several momenta for dopings $\delta = 0.12$ (a) and $\delta = 0.78$ (b).

V. CONCLUSIONS AND DISCUSSIONS

Based on our Hubbard operator path-integral representation, defined for the first time in Ref.[11], we have developed a large-N expansion for the t-J model.

Because our path integral is written in terms of X-operators, we do not need any decoupling scheme for the Hubbard operators. Then, we do not invoke a gauge invariance or Bose condensation, like in the slave boson formulation¹³. Our dynamical variables are the original Hubbard operators and, there is no need to introduce spinons and holons excitations.

The use of a path integral allows us to define a Feynman diagram technique which is a very important and interesting problem in electron correlated systems¹⁷.

As we mentioned previously, the large-N expansion was formulated using canonical quantization in terms of

X-operator^{8,9} or slave boson path integral approach¹³. The first one cannot be defined in a diagrammatic form while the second one can be developed in terms of Feynman diagrams but needs the introduction of spinons and holons. In the formulation presented in this paper we have diagrammatic rules: propagators and vertices in terms of the original X-operators.

One important point is that in our diagram technique the fermion operator f_{σ} is proportional to the fermion X- operator $X^{0\sigma}$ for all orders in the large N expansion. This is very important for the interpretation of the experimental spectral susceptibilities. In the slave boson approach the spinon is only proportional to the fermion $X^{0\sigma}$ operator in the leading order of 1/N expansion¹³.

It is also important to note that our diagram technique contains ghost fields that cancel infinities to order O(1/N) in the boson propagator. In the slave boson approach in the radial gauge, the ghost fields necessary to treat the Jacobian, cancel infinities in $O(1/N^2)^{18}$.

Using our large-N expansion, we have calculated the boson propagator and the charge-charge and spin-spin correlation functions in O(1). In sections III and IV we have shown that these quantities agree with those calculated by slave boson¹³ and X-operator canonical approach¹⁵. These results, which are far nontrivial, show the consistency of our formulation.

We showed that in the N-infinite limit we have well defined electronic excitations with renormalized band due to correlations. This means (and we think) that the large-N expansion, in the form presented in this paper, is a better approximation to describe high-Tc cuprates in optimal doping and overdoped than in underdoped close to antiferromagnetism.

We believe that our formulation will be useful (and simpler than previous ones) to calculate quantities in the next order in O(1/N), such as superconductivity and electronic self-energy corrections. We are also planning to extend the method to the case of finite J.

Acknowledgments

The authors thank to O. Zandron for valuable discussions. A.G. wishes to acknowledge many interesting discussions with A. Dobry, O. Fojón and R.Zeyher.

- ⁸ R. Zevher and M. Kulić, Phys.Reb. B **53**, 2850(1996).
- ⁹ R. Zeyher and A. Greco, Eur. Phys. J.B **6**, 473 (1998).
- A. Foussats, A. Greco and O. S. Zandron, Annals of Physics (NY) 275, 238(1999); A. Foussats, A. Greco and O. S. Zandron, Annals of Physics(NY) 279, 263(2000).
- ¹¹ A. Foussats, A. Greco, C. Repetto, O. P. Zandron and O. S. Zandron, Journal of Physics A 33, 5849(2000).
- ¹² L. Faddeev and R. Jackiw, Phys. Rev. Lett **60**, 1692 (1988); P. Dirac, Lectures on Quantum Mechanics (New York: Yeshiva University Pressi, 1964); K. Sundermayer, Constrained Dynamics (Berlin: Spinger, 1982); J. Govaerts, Int. J. Mod. Physics A **5**, 3625(1990).
- ¹³ Z. Wang, Int. Journal of Modern Physiscs B **6**, 155(1992).
- ¹⁴ L. Faddeev, Theoret. Mat. Phys. 1, 1(1970); P. Senjanovich, Ann. Phys. (NY) 100, 227(1976).
- ¹⁵ L. Gehlhoff and R. Zeyher; Phys. Rev. B **52**, 4635(1995).
- ¹⁶ T. Tohyama, P. Horsch and S. Maekawa, Phys. Rev. Lett. 74, 980(1995).
- ¹⁷ Y. Izyumov and Y. Skrabin; Statistical Mechanics of Magnetically Ordered Systems, Consultansts Bureau, New York (1988).
- ¹⁸ N. Read and D.Newns, J. Phys. C **16**, 3273(1983).

 $^{^1}$ P.W. Anderson, The Theory of superconductivity in High- Tc Cuprates (Princeton University Press, Princeton, New Jersey, USA,1997).

² J. Hubbard, Proc. R. Soc. London Ser. A **276**, 238 (1963).

³ A. Izyumov, Physics-Uspekhi **40**, 445 (1997).

⁴ J. C. Le Guillou and E. Ragoucy, Phys. Rev. B **52** , 2403 (1995).

⁵ G. Kotliar and J. Liu, Phys. Rev. Lett. **61**,1784 (1988).

⁶ M. Grilli and G. Kotliar, Phys.Rev. lett. **64**, 1170 (1990).

⁷ G. Baym and L. Kadanoff, Phys. Rev. B **124**, 287 (1961).