Effect of the additional second neighbor hopping on the charge dynamics in the t-J model

Feng Yuan¹ and Shiping Feng^{1,2,3}

¹Department of Physics, Beijing Normal University, Beijing 100875, China

²The Key Laboratory of Beam Technology and Material Modification of Ministry of Education,

Beijing Normal University, Beijing 100875, China

³National Laboratory of Superconductivity, Academia Sinica, Beijing 100080, China

Abstract

The effect of the additional second neighbor hopping t' on the charge dynamics of the t-J model in the underdoped regime is studied within the fermion-spin theory. The conductivity spectrum of the t-t'-J model shows the low-energy peak and unusual midinfrared band, while the resistivity exhibits a nearly temperature linear dependence with deviation at low temperature in the underdoped regime. Although the qualitative feature of the charge dynamics in the t-t'-J model is the same as in the case of the t-J model, the additional second neighbor hopping t' leads to a clear shift of the position of the midinfrared band to the higher energies in the conductivity spectrum, and suppress the range of the deviation from the temperature linear dependence in the resistivity.

71.27.+a, 74.72.-h, 72.10.-d

It has become clear in the past several years that copper oxide materials show many unusual normal-state properties [1]. The normal-state properties exhibit a number of anomalous properties in the sense that they do not fit in the conventional Fermi-liquid theory, and some properties mainly depend on the extent of dopings [1]. Among the striking features of the anomalous normal-state properties stands out the extraordinary charge dynamics, which is manifested by the optical conductivity and resistivity [2]. It has been shown from the experiments [1–5] that the optical conductivity spectrum of copper oxide materials shows the non-Drude behavior at low energies and exotic midinfrared band in the charge-transfer gap, while the resistivity exhibits a linear behavior in the temperature in the optimally doped regime and a nearly temperature linear dependence with deviations at low temperatures in the underdoped regime. The single common feature of copper oxide materials is the two-dimensional (2D) CuO₂ plane [1,2], and it seems evident that the anomalous behaviors are governed by this plane. Since the copper oxide superconductors are doped Mott insulators, many authors [6,7] have suggested that the essential physics of these materials can be effectively described by the 2D t-J model acting on the space with no doubly occupied sites, where t is the nearest neighbor hopping matrix element, and J is the nearest neighbor magnetic exchange interaction. This model has been used to study the charge dynamics of copper oxide materials in the underdoped regime, and the results obtained [8–11] from the analytical methods and numerical simulations are in qualitative agreement with the experiments [1–5].

However, the recent angle-resolved photoemission spectroscopy measurements [12] on copper oxide materials show that although the highest energy filled electron band is well described by the t-J model in the direction between the (0,0) point and the (π,π) point in the momentum space, but both the experimental data near $(\pi,0)$ point and overall dispersion may be properly accounted by generalizing the t-J model to include the second- and third-nearest neighbors hopping terms t' and t''. These photoemission results also show that the electron band width is reduced from the several eV expected from the band theory to of order J, which indicates that the coupling of the electron to the antiferromagnetic background

plays an important role in the electronic structure [12]. On the other hand, the charge response is a powerful probe for the systems of interacting electrons, and provides very detailed informations of the excitations, which interact with carriers in the normal-state [1–5]. It is believed that both experiments from the angle-resolved photoemission spectroscopy measurements and charge response produce interesting data that introduce important constraints on the microscopic models and theories. In this case, a natural question is what is the effect of these additional hoppings on the charge dynamics of the t-J model. In this paper, we study this issue within t-t'-J model. Our results indicate that the conductivity spectrum of the t-t'-J model shows the low-energy peak and unusual midinfrared band, while the resistivity exhibits a nearly temperature linear dependence with deviation at low temperature in the underdoped regime. Although these qualitative feature of the charge dynamics in the t-t'-J model is the same as in the case of the t-J model [8–11], the additional second neighbor hopping t' leads to a clear shift of the position of the midinfrared band to the higher energies in the conductivity spectrum, and suppress the range of the deviation from the temperature linear dependence in the resistivity.

We begin with the t-t'-J model defined on a square lattice,

$$H = -t \sum_{i\hat{\eta}\sigma} C_{i\sigma}^{\dagger} C_{i+\hat{\eta}\sigma} + t' \sum_{i\hat{\tau}\sigma} C_{i\sigma}^{\dagger} C_{i+\hat{\tau}\sigma} + \mu \sum_{i\sigma} C_{i\sigma}^{\dagger} C_{i\sigma} + J \sum_{i\hat{\eta}} \mathbf{S}_{i} \cdot \mathbf{S}_{i+\hat{\eta}}, \tag{1}$$

where $\hat{\eta} = \pm \hat{x}, \pm \hat{y}, \ \hat{\tau} = \pm \hat{x} \pm \hat{y}, \ C_{i\sigma}^{\dagger} \ (C_{i\sigma})$ is the electron creation (annihilation) operator, $\mathbf{S}_i = C_i^{\dagger} \sigma C_i / 2$ are spin operators with $\sigma = (\sigma_x, \sigma_y, \sigma_z)$ as the Pauli matrices, and μ is the chemical potential. The strong electron correlation in the t-J and t-t'-J model manifests itself by this electron single occupancy on-site local constraint [6,7], and then the on-site local constraint should be treated properly. Recently a fermion-spin theory based on the charge-spin separation [13,14], $C_{i\uparrow} = h_i^{\dagger} S_i^{-}$ and $C_{i\downarrow} = h_i^{\dagger} S_i^{+}$, has been proposed to incorprate this on-site local constraint, where the spinless fermion operator h_i describes the charge (holon) degrees of freedom, while the pseudospin operator S_i describes the spin (spinon) degrees of freedom. The main advantage of this approach is that the on-site local constraint can be treated exactly in analytical calculations. In the fermion-spin representation, the

t-t'-J model can be expressed [13,14] as,

$$H = t \sum_{i\hat{\eta}} h_{i+\hat{\eta}}^{\dagger} h_i (S_i^+ S_{i+\hat{\eta}}^- + S_i^- S_{i+\hat{\eta}}^+) - t' \sum_{i\hat{\tau}} h_{i+\hat{\tau}}^{\dagger} h_i (S_i^+ S_{i+\hat{\tau}}^- + S_i^- S_{i+\hat{\tau}}^+)$$

$$- \mu \sum_{i} h_i^{\dagger} h_i + J_{eff} \sum_{i\hat{\eta}} (\mathbf{S}_i \cdot \mathbf{S}_{i+\hat{\eta}}),$$
(2)

where $J_{eff} = J[(1 - \delta)^2 - \phi_1^2]$, the holon particle-hole order parameter $\phi_1 = \langle h_i^{\dagger} h_{i+\hat{\eta}} \rangle$, and S_i^+ and S_i^- are the pseudospin raising and lowering operators, respectively.

Since the on-site electron local constraint has been treated exactly within the framework of the fermion-spin theory, then the extra gauge degree of freedom related with the electron on-site local constraint under the charge-spin separation does not appear in the fermion-spin theory. In this case, the spin fluctuation couples only to spinons [15], while the charge fluctuation couples only to holons [11], but the strong correlation between holons and spinons still is considered through the holon's order parameters entering in the spinon's propagator and the spinon's order parameters entering in the holon's propagator, therefore both holons and spinons contribute to the charge and spin dynamics. Within the fermion-spin theory, the charge dynamics of the t-J model in the underdoped regime has been discussed [11] by considering the holon fluctuation around the mean-field solution, where the holon part is treated by the loop expansion to the second-order. Following their discussions, we can obtain the optical conductivity in the present t-t-t-J model as,

$$\sigma(\omega) = \frac{1}{2} (2Ze)^2 \frac{1}{N} \sum_{k} \gamma_{sk}^2 \int_{-\infty}^{\infty} \frac{d\omega'}{2\pi} A_h(k, \omega' + \omega) A_h(k, \omega') \frac{n_F(\omega' + \omega) - n_F(\omega')}{\omega}, \tag{3}$$

where Z is the number of the nearest neighbor or second-nearest neighbor sites, $\gamma_{sk} = t\chi_1(\sin k_x + \sin k_y)/2 - t'\chi_2(\sin k_x \cos k_y + \cos k_x \sin k_y)$, the spinon correlation functions $\chi_1 = \langle S_i^+ S_{i+\hat{\eta}}^- \rangle$, $\chi_2 = \langle S_i^+ S_{i+\hat{\tau}}^- \rangle$, $n_F(\omega)$ is the fermion distribution function, and the holon spectral function $A_h(k,\omega)$ is obtained as $A_h(k,\omega) = -2\operatorname{Im} g(k,\omega)$. The full holon Green's function $g^{-1}(k,\omega) = g^{(0)-1}(k,\omega) - \Sigma_h^{(2)}(k,\omega)$ with the mean-field holon Green's function $g^{(0)-1}(k,\omega) = \omega - \xi_k$ and the second-order holon self-energy from the spinon pair bubble [11],

$$\Sigma_h^{(2)}(k,\omega) = \left(\frac{Z}{N}\right)^2 \sum_{pp'} \gamma_{12}^2(k,p,p') \frac{B_{p'}B_{p+p'}}{4\omega_{p'}\omega_{p+p'}} \times \left(2\frac{F_1(k,p,p')}{\omega + \omega_{p+p'} - \omega_{p'} - \xi_{p+k}}\right)$$

$$+ \frac{F_2(k, p, p')}{\omega + \omega_{p'} + \omega_{p+p'} - \xi_{p+k}} - \frac{F_3(k, p, p')}{\omega - \omega_{p+p'} - \omega_{p'} - \xi_{p+k}} \right), \tag{4}$$

where $\gamma_{12}(k,p,p') = t(\gamma_{\mathbf{p'}-\mathbf{k}} + \gamma_{\mathbf{p'}+\mathbf{p}+\mathbf{k}}) - t'(\gamma'_{\mathbf{p'}-\mathbf{k}} + \gamma'_{\mathbf{p'}+\mathbf{p}+\mathbf{k}}), \quad \gamma_{\mathbf{k}} = (1/Z) \sum_{\hat{\eta}} e^{i\mathbf{k}\cdot\hat{\eta}},$ $\gamma'_{\mathbf{k}} = (1/Z) \sum_{\hat{\tau}} e^{i\mathbf{k}\cdot\hat{\tau}}, \quad F_1(k,p,p') = n_F(\xi_{p+k})[n_B(\omega_{p'}) - n_B(\omega_{p+p'})] + n_B(\omega_{p+p'})[1 + n_B(\omega_{p'})],$ $F_2(k,p,p') = n_F(\xi_{p+k})[1 + n_B(\omega_{p+p'}) + n_B(\omega_{p'})] + n_B(\omega_{p'})n_B(\omega_{p+p'}), \quad F_3(k,p,p') = n_F(\xi_{p+k})[1 + n_B(\omega_{p+p'}) + n_B(\omega_{p'})] - [1 + n_B(\omega_{p'})][1 + n_B(\omega_{p+p'})], \quad n_B(\omega_p) \text{ is the boson distribution functions,}$ $B_k = \Delta_1[2\chi_1^z(\epsilon\gamma_{\mathbf{k}} - 1) + \chi_1(\gamma_{\mathbf{k}} - \epsilon)] - \Delta_2(2\chi_2^z\gamma'_{\mathbf{k}} - \chi_2), \quad \Delta_1 = 2ZJ_{eff}, \quad \Delta_2 = 4Z\phi_2t',$ $\epsilon = 1 + 2t\phi_1/J_{eff}, \text{ the mean-field holon spectrum } \xi_k = 2Zt\chi_1\gamma_{\mathbf{k}} - 2Zt'\chi_2\gamma'_{\mathbf{k}} - \mu, \text{ and the mean-field spinon spectrum,}$

$$\omega_{k}^{2} = \Delta_{1}^{2} \left(\left[\alpha C_{1}^{z} + \frac{1}{4Z} (1 - \alpha) - \alpha \epsilon \chi_{1}^{z} \gamma_{k} - \frac{1}{2Z} \alpha \epsilon \chi_{1} \right] (1 - \epsilon \gamma_{k}) \right. \\
+ \frac{1}{2} \epsilon \left[\alpha C_{1} + \frac{1}{2Z} (1 - \alpha) - \alpha \chi_{1} \gamma_{k} - \frac{1}{2} \alpha \chi_{1}^{z} \right] (\epsilon - \gamma_{k}) \right) \\
+ \Delta_{2}^{2} \left(\left[\alpha \chi_{2}^{z} \gamma_{k}' - \frac{3}{2Z} \alpha \chi_{2} \right] \gamma_{k}' + \frac{1}{2} \left[\alpha C_{2} + \frac{1}{2Z} (1 - \alpha) - \frac{1}{2} \alpha \chi_{2}^{z} \right] \right) \\
+ \Delta_{1} \Delta_{2} \left(\alpha \chi_{1}^{z} \gamma_{k}' (1 - \epsilon \gamma_{k}) + \frac{\alpha}{2} (\chi_{1} \gamma_{k}' - C_{3}) (\epsilon - \gamma_{k}) + \alpha \gamma_{k}' (C_{3}^{z} - \epsilon \chi_{2}^{z} \gamma_{k}) \right. \\
- \frac{1}{2} \alpha \epsilon (C_{3} - \chi_{2} \gamma_{k}) \right), \tag{5}$$

with the spinon correlation functions $\chi_1^z = \langle S_i^z S_{i+\hat{\eta}}^z \rangle$, $\chi_2^z = \langle S_i^z S_{i+\hat{\tau}}^z \rangle$, $C_1 = (1/Z) \sum_{\hat{\eta}'} \langle S_{i+\hat{\eta}}^+ S_{i+\hat{\eta}'}^- \rangle$, $C_1 = (1/Z) \sum_{\hat{\eta}'} \langle S_{i+\hat{\eta}}^z S_{i+\hat{\eta}'}^z \rangle$, $C_2 = (1/Z) \sum_{\hat{\tau}'} \langle S_{i+\hat{\tau}}^+ S_{i+\hat{\tau}'}^- \rangle$, $C_3 = (1/Z) \sum_{\hat{\tau}} \langle S_{i+\hat{\eta}}^z S_{i+\hat{\tau}}^z \rangle$, and the holon particle-hole order parameter $\phi_2 = \langle h_i^{\dagger} h_{i+\hat{\tau}} \rangle$. In order not to violate the sum rule of the correlation function $\langle S_i^+ S_i^- \rangle = 1/2$ in the case without antiferromagnetic long-range-order, the important decoupling parameter α has been introduced in the mean-field calculation, which can be regarded as the vertex correction [14,16]. All the above mean-field order parameters are determined by the self-consistent calculation [14].

Although the charge dynamics of the copper oxide materials is very complicated, the transport studies of electron excitations have revealed much about the nature of the charge carriers [1–5]. In this paper, we are interested in the effect of the additional second-neighbor hopping on the charge dynamics of the t-J model. To make the discussion simpler, we only study the charge dynamics of the t-t'-J model in the underdoped regime. We have

performed the numerical calculation for the optical conductivity of the t-t'-J model, and the zero temperature results with doping concentration (a) $\delta = 0.06$ and (b) $\delta = 0.12$ for parameter t/J = 2.5, t'/J = 0.1 (dashed line), t'/J = 0.3 (dotted line), and t'/J = 0.5(dash-dotted line) are plotted in Fig. 1, where we have taken charge e as the unit. For comparison, the corresponding result |11| of the t-J model (solid line) is also shown in Fig. 1. The conductivity spectrum of the t-t'-J model shows the low-energy peak at $\omega < 0.5t$ and midinfrared band appearing inside the charge-transfer gap of the undoped systems. Although the qualitative feature of the conductivity spectrum in the t-t'-J model is the same as in the case of the t-J model [8,11], the additional second neighbor hopping t' leads to a clear shift of the position of the midinfrared band to the higher energies in the conductivity spectrum. Moreover, the charge-transfer gap in the conductivity is doping dependent, decreases with increasing dopings. Since the spectral weight from both the low-energy peak and midinfrared sideband represents the actual free-carrier density, the suppression of the midinfrared band with increasing dopings means that only few amount of the free-carrier are taken from the Drude absorption to the midinfrared band, which leads to the unusual decay of conductivity at low energies as $\sigma(\omega) \propto 1/\omega$ in the underdoped regime, which are consistent with the experiments [1–5].

 linear dependence in the resistivity is also suppressed with increasing the values of t'.

Since the scattering of holons dominates the charge dynamics in the fermion-spin theory, therefore the present study indicates that in the underdoped regime the observed crossover from the temperature linear to the nonlinear range in the resistivity in low temperatures is closely related to the pseudogap in holon excitations. This can be understood from the physical property of the holon density of states (DOS) $\Omega(\omega)=1/N\sum_k A_h(k,\omega)$. The numerical result of the holon DOS $\Omega(\omega)$ at doping $\delta = 0.06$ for parameters t/J = 2.5 and t'=0 in temperature T=0 is shown in Fig. 3. For comparison, the corresponding mean-field result (dashed line) is also shown in Fig. 3. From Fig. 3, it is found that the holon DOS in the mean-field approximation consists of the central part only, which comes from the noninteracting particles. After including fluctuations the central part is renormalized and a V-shape holon pseudogap near the chemical potential μ in the underdoped regime appears. For the further understanding the physical property of this holon pseudogap, we plot the phase diagram $T^* \sim \delta$ at parameters t/J=2.5 and t'=0 in Fig. 4, where T^* marks the development of the holon pseudogap in the holon density of states. As seen from Fig. 4, this holon pseudogap is doping and temperature dependent, and grows monotonously as the doping δ decreases, and disappear in higher doping. Moreover, this holon pseudogap decreases with increasing temperatures, and vanishes at higher temperatures. Now we discuss the effect of t' on the pseudogap. In Fig. 5, we plot the phase diagram $T^* \sim t'$ at doping $\delta = 0.06$ with parameters t/J = 2.5, which shows that the holon pseudogap also decreases with increasing the values of t', and vanishes for t'/J > 0.1. The t-J model is characterized by a competition between the kinetic energy (t) and magnetic energy (J). The magnetic energy J favors the magnetic order for spins, while the kinetic energy t favors delocalization of holes and tends to destroy the magnetic order. Our results show that the additional second neighbor hopping t' in the t-J model may be equivalent to increase the kinetic energy, and its influence on the charge dynamics of the t-J model is similar to the effect of dopings. This is why at least for small values of t' the qualitative behavior of the charge dynamics in the t-t'-J model is the same as these obtained [8–11] from the t-J model. Since the full holon Green's function (then the holon spectral function and DOS) is obtained by considering the second-order correction due to the spinon pair bubble, therefore the holon pseudogap is closely related to the spinon fluctuation. For small dopings and lower temperatures, the holon kinetic energy is much smaller than the magnetic energy, in this case the magnetic fluctuation is strong enough to lead to the holon pseudogap. This holon pseudogap would reduce the holon scattering and thus is responsible for the deviation from the temperature linear behavior in the resistivity. With increasing temperatures, dopings, or values of t', the holon kinetic energy is increased, while the spinon magnetic energy is decreased. In the region where the holon pseudogap closes at high temperatures, higher doping levels, or for large values of t', the charged holon scattering would give rise to the temperature linear resistivity.

In summary, we have discussed the effect of the additional second neighbor hopping t' on the charge dynamics of the t-J model in the underdoped regime within the fermionspin theory. Our results indicate that the conductivity spectrum of the t-t'-J model shows the low-energy peak and unusual midinfrared band, while the resistivity exhibits a nearly temperature linear dependence with deviation at low temperature in the underdoped regime. Although the qualitative feature of the charge dynamics in the t-t'-J model is the same as in the case of the t-J model, the additional second neighbor hopping t' leads to a clear shift of the position of the midinfrared band to the higher energies in the conductivity spectrum, and suppress the range of the deviation from the temperature linear dependence in the resistivity.

Finally, we also note that some physical properties of the t-t'-J model have been discussed by Martins et al. [17], they found that there is a tendency of holes to generate nontrivial spin environments, this effect leads to decouple charge from spin. Moreover, they [17] have shown that introducing t' in the t-J model is equivalent to effectively renormalizing J (decreasing the magnetic energy). Therefore, within the charge-spin separation of the t-t'-J model, the scattering of spinons dominates the spin dynamics [15], while the scattering of holons dominates the charge dynamics, the two rates observed in the experiments are attributed to

the scattering of two distinct excitations, spinons and holons.

ACKNOWLEDGMENTS

This work was supported by the National Natural Science Foundation of China under Grant No. 10074007, 90103024, and 10125415, and the Grant from Ministry of Education of China.

REFERENCES

- See, e.g., Proceedings of Los Alamos Symposium, edited by K.S. Bedell, D. Coffey, D.E. Meltzer, D. Pines, and J.R. Schrieffer (Addison-Wesley, Redwood city, California, 1990);
 A.P. Kampf, Phys. Rep. 249, 219 (1994); M.A.Kastner, R.J. Birgeneau, G. Shiran, Y. Endoh, Rev. Mod. Phys. 70, 897 (1998).
- [2] D.B. Tanner and T. Timusk, in *Physical Properties of High Temperature Superconductors* III, edited by D.M. Ginsberg (World Scientific, Singapore, 1992), p. 363.
- [3] H. Takagi, B. Batlogg, H.L. Kao, J. Kwo, R.J. Cava, J.J. Krajewski, and W.F. Peck, Phys. Rev. Lett. 69, 2975 (1992); T. Ito, K. Takenaka, and S. Uchida, Phys. Rev. Lett. 70, 3995 (1993).
- [4] S. Uchida, Mod. Phys. Lett. B4, 513 (1990); T. Timusk, M. Reedyk, R. Hughes, D.A. Bonn, J.D. Garret, J.E. Greedan, C.V. Stager, D.B. Tanner, Feng Gao, S.L. Herr, K. Kamaras, G.A. Thomas, S.L. Cooper, J. Orenstein, L.F. Schneemeyer, and A.J. Millis, Physica C162-164, 841 (1989); S. Uchida, T. Ido, H. Takagi, T. Arima, Y. Tokura, and S. Taima, Phys. Rev. B43, 7942 (1991); S.L. Cooper, D. Reznik, A. Kotz, M.A. Karlow, R. Liu, M. V. Klein, B.W. Veal, and P. Paulikas, Phys. Rev. B47, 8233 (1993); A.V. Puchkov, P. Fournier, T. Timusk, and N.N. Kolesnikov, Phys. Rev. Lett. 77, 1853 (1996).
- [5] N.W. Preyer, R.J. Birgeneau, C.Y. Chen, D.R. Gabbe, H.P. Jenssen, M.A. Kastner, P.J. Picone, and T. Thio, Phys. Rev. B 39, 11563 (1989); J.P. Falck, A.Levy, M.A. Kastner, and R.J. Birgeneau, Phys. Rev. B48, 4043 (1993).
- [6] P. W. Anderson, in Frontiers and Borderlines in Many Particle Physics, edited by R.A. Broglia and J.R. Schrieffer (North-Holland, Amsterdam, 1987), p. 1; Science 235, 1196 (1987).
- [7] F. C. Zhang and T. M. Rice, Phys. Rev. B37, 3759 (1988); T. M. Rice, in *Proceedings*

- of the International Conference on Materials and Mechanisms of Superconductivity and High Temperature Superconductors V, Beijing, China, 1997, edited by Y.S. He, P.H. Wu, L.F. Xu, and Z.X. Zhao [Physica C282-287, xix (1997)].
- [8] W. Stephan and P. Horsch, Phys. Rev. B42, 8736 (1990).
- [9] A. Moreo and E. Dogotto, Phys. Rev. B42, 4786 (1990); E. Dagotto, Rev. Mod. Phys.66, 763 (1994), and references therein.
- [10] J. Jaklič and P. Prelovšek, Phys. Rev B **50**, 7129 (1994).
- [11] Shiping Feng and Zhongbing Huang, Phys. Lett. A232, 293 (1997).
- [12] B.O. Wells, Z.X. Shen, A. Matsuura, D.M. King, M.A. Kastner, M. Greven, and R.J. Birgeneau, Phys. Rev. Lett. 74 (1995) 964; C.P. Kim, P.J. White, Z.X. Shen, T. Tohyama, Y. Shibata, S. Maekawa, B.O. Wells, Y.J. Kim, R.J. Birgeneau, and M.A. Kastner, Phys. Rev. Lett. 80, 4245 (1998).
- [13] Shiping Feng, Z.B. Su, and L. Yu, Phys. Rev. B 49, 2368 (1994); Mod. Phys. Lett. B7, 1013 (1993).
- [14] Shiping Feng and Y. Song, Phys. Rev. B55, 642 (1997).
- [15] Shiping Feng and Zhongbing Huang, Phys. Rev. B 57, 10328 (1998).
- [16] J. Kondo and K. Yamaji, Prog. Theor. Phys. 47, 807 (1972).
- [17] G.B. Martins, R. Eder, and E. Dagotto, Phys. Rev. B60, R3716 (1999); G.B. Martins, J.C. Xavier, C. Gazza, M. Vojta, and E. Dagotto, Phys. Rev. B63, 014414 (2000); G.B. Martins, C. Gazza, J.C. Xavier, A. Feiguin, and E. Dagotto, Phys. Rev. Lett. 84, 5844 (2000); G.B. Martins, J.C. Xavier, L. Arrachea, and E. Dagotto, cond-mat/0109405.

FIGURES

- FIG. 1. The conductivity of the t-t'-J model with doping concentration (a) $\delta = 0.06$ and (b) $\delta = 0.12$ for parameters t/J = 2.5, t'/J = 0.1 (dashed line), t'/J = 0.3 (dotted line), and t'/J = 0.5 (dash-dotted line) in temperature T = 0. The solid line is the corresponding result of the t-J model.
- FIG. 2. The resistivity at doping concentration $\delta = 0.06$ for parameters t/J = 2.5, (a) t' = 0 and (b) t'/J = 0.1.
- FIG. 3. The holon density of states for parameters t/J=2.5 and t'=0 at doping $\delta=0.06$. The dashed line is the result at the mean-field level.
- FIG. 4. The normal-state phase diagram $T^* \sim \delta$ for parameters t/J = 2.5 and t' = 0. T^* marks the development of the holon pseudogap in the holon density of states
 - FIG. 5. The normal-state phase diagram $T^* \sim t'$ at doping $\delta = 0.06$ for parameter t/J = 2.5.









