

WEAKLY NONLOCAL CONTINUUM PHYSICS - THE GINZBURG-LANDAU EQUATION

PETER VÁN

ABSTRACT. In this paper we investigate and refine Liu's procedure for the exploitation of the entropy inequality. We give a new-old proof of Liu's theorem. We shortly compare the Coleman-Noll and Liu techniques and call the attention to the proper application of the constraints in the procedure. As an example the Ginzburg-Landau equation and some variants are derived.

1. INTRODUCTION

In 1972 Liu introduced a method of the exploitation of the entropy principle [1]. The method become a basic tool to exploit the entropy inequality, one of the fundamental ingredients of the Second Law. The method is based on a linear algebraic theorem, called Liu's theorem in the thermodynamic literature [2, 3, 4] and an interpretation of role of the entropy inequality, one of the fundamental ingredients of the Second Law.

Recently Hauser and Kirchner recognized that Liu's theorem is a consequence of a famous statement of optimization theory and linear programming, the so called Farkas's lemma [5]. That theorem was proved first by Farkas in 1894 and independently by Minkowski in 1896 [6]. In the first part of this paper we formulate Liu's theorem in a way that is mostly adapted for our purposes and shows the whole train of thought from Farkas's lemma to Liu's theorem giving a simple proof to every statement in question.

At the second part we call the attention to some fine points in the application of the Liu procedure, namely, it requires a careful choice of the independent variables and an investigation of the constraints from the point of view of independency of these variables. This is important in weakly nonlocal continuum theories, as it is demonstrated in the last section, where the application of the treated ideas lead to a straightforward derivation of the Ginzburg-Landau equation. Some other derivations and variants and generalizations of the Ginzburg-Landau equation are compared and discussed.

2. FARKAS'S LEMMA AND SOME OF ITS CONSEQUENCES

Lemma 2.1. (*Farkas*) Let $\mathbf{a}_i \neq \mathbf{0}$ be vectors in a finite dimensional vector space \mathbb{V} , $i = 1 \dots n$, and $S = \{\mathbf{p} \in \mathbb{V}^* | \mathbf{p} \cdot \mathbf{a}_i \geq 0, i = 1 \dots n\}$. The following statements are equivalent for all $\mathbf{b} \in \mathbb{V}$:

- (i) $\mathbf{p} \cdot \mathbf{b} \geq 0$, for all $\mathbf{p} \in S$.
- (ii) There are nonnegative real numbers $\lambda_1, \dots, \lambda_n$ such that $\mathbf{b} = \sum_{i=1}^n \lambda_i \mathbf{a}_i$.

Proof:

- (ii) \Rightarrow (i) $\mathbf{p} \cdot \sum_{i=1}^n \lambda_i \mathbf{a}_i = \sum_{i=1}^n \lambda_i \mathbf{p} \cdot \mathbf{a}_i \geq 0$ if $\mathbf{p} \in S$.

(i) \Rightarrow (ii) It is sufficient to prove the statement for a maximal, linearly independent subset of $\mathbf{a}_1, \dots, \mathbf{a}_l$, because that subset generates S and we can choose zero multipliers for the vectors that are not independent.

Let $S_0 = \{\mathbf{y} \in \mathbb{V}^* | \mathbf{y} \cdot \mathbf{a}_i = 0, i = 1 \dots l\}$ Clearly $\emptyset \neq S_0 \subset S$.

If $\mathbf{y} \in S_0$ then $-\mathbf{y}$ is also in S_0 , therefore $\mathbf{y} \cdot \mathbf{b} \geq 0$ and $-\mathbf{y} \cdot \mathbf{b} \geq 0$ together. Therefore for all $\mathbf{y} \in S_0$ it is true that $\mathbf{y} \cdot \mathbf{b} = 0$.

As a consequence \mathbf{b} is in the set generated by $\{\mathbf{a}_i\}$, that is there are real numbers $\lambda_1, \dots, \lambda_l$ such that $\mathbf{b} = \sum_{i=1}^l \lambda_i \mathbf{a}_i$.

Moreover, for all $k \in \{1, \dots, l\}$ there is a $\mathbf{p}_k \in \mathbb{V}$ such that $\mathbf{p}_k \cdot \mathbf{a}_k = 1$ and $\mathbf{p}_k \cdot \mathbf{a}_i = 0$ if $i \neq k$. As a consequence for all $\mathbf{p} \in S$ $0 \leq \mathbf{p} \cdot \mathbf{b} = \mathbf{p} \cdot \sum_{i=1}^l \lambda_i \mathbf{a}_i = \sum_{i=1}^l \lambda_i \mathbf{p} \cdot \mathbf{a}_i = \lambda_k$ is valid for all k . ■

Remark 2.1. In the following the elements of \mathbb{V}^* are called independent variables and \mathbb{V}^* itself is called the space of independent variables. The inequality in the first statement of the lemma is called aim inequality and the nonnegative numbers in the second statement are called Lagrange-Farkas multipliers. The inequalities determining S are the constraints.

In the calculations an excellent reminder is to use Lagrange- Farkas multipliers similarly to the Lagrange multipliers in case of conditional extremum problems:

$$\mathbf{p} \cdot \mathbf{b} - \sum_{i=1}^n \lambda_i \mathbf{p} \cdot \mathbf{a}_i = \mathbf{p} \cdot (\mathbf{b} - \sum_{i=1}^n \lambda_i \cdot \mathbf{a}_i) \geq 0, \quad \forall \mathbf{p} \in \mathbb{V}^*$$

From this form we can read out the second statement of the lemma.

Remark 2.2. The geometric interpretation of the theorem is important and graphic: if the vector \mathbf{b} does not belong to the cone generated by the vectors \mathbf{a}_i , there exists a hyperplane separating \mathbf{b} from the cone.

2.1. Affine Farkas's lemma. This generalization of the previous lemma was first published simultaneously by A. Haar and J. Farkas as subsequent papers in the same journal, with different proofs [7, 8]. Lately it was reproved independently by others several times (e.g. [9, 10]).

Theorem 2.2. (Affine Farkas) Let $\mathbf{a}_i \neq \mathbf{0}$ be vectors in a finite dimensional vector space \mathbb{V} and α_i real numbers, $i = 1 \dots n$ and $S_A = \{\mathbf{p} \in \mathbb{V}^* | \mathbf{p} \cdot \mathbf{a}_i \geq \alpha_i, i = 1 \dots n\}$. The following statements are equivalent for a $\mathbf{b} \in \mathbb{V}$ and a real number β :

(i) $\mathbf{p} \cdot \mathbf{b} \geq \beta$, for all $\mathbf{p} \in S_A$.

(ii) There are nonnegative real numbers $\lambda_1, \dots, \lambda_n$ such that $\mathbf{b} = \sum_{i=1}^n \lambda_i \mathbf{a}_i$ and $\beta \leq \sum_{i=1}^n \lambda_i \alpha_i$.

Proof:

(ii) \Rightarrow (i) $\mathbf{p} \cdot \mathbf{b} = \mathbf{p} \cdot \sum_{i=1}^n \lambda_i \mathbf{a}_i = \sum_{i=1}^n \lambda_i \mathbf{p} \cdot \mathbf{a}_i \geq \sum_{i=1}^n \lambda_i \alpha_i \geq \beta$.

(i) \Rightarrow (ii) First we will show indirectly that the first condition of Farkas's lemma is a consequence of the first condition here, that is if (i) is true then $\mathbf{p} \cdot \mathbf{b} \geq 0$, for all $\mathbf{p} \in S$.

Thus let us assume the contrary, hence there is $\mathbf{p}' \in S$, for which $\mathbf{p}' \cdot \mathbf{b} < 0$. Take an arbitrary $\mathbf{p} \in S_A$, then $\mathbf{p} + k\mathbf{p}' \in S_A$ for all real numbers k . But now $(\mathbf{p} + k\mathbf{p}') \cdot \mathbf{b} = \mathbf{p} \cdot \mathbf{b} + k\mathbf{p}' \cdot \mathbf{b} < \beta$, if $k \frac{\mathbf{p}' \cdot \mathbf{b}}{\mathbf{p} \cdot \mathbf{b}} < -1$. That is a contradiction.

Therefore, according to Farkas's lemma exist $\lambda \in \mathbb{R}^{n+}$ Lagrange-Farkas multipliers $\lambda_1, \dots, \lambda_n$ such that $\mathbf{b} = \sum_{i=1}^n \lambda_i \mathbf{a}_i$. Hence $\beta \leq \inf_{\mathbf{p} \in S_A} \{\mathbf{p} \cdot \sum_{i=1}^n \lambda_i \mathbf{a}_i\} = \inf_{\mathbf{p} \in S_A} \{\sum_{i=1}^n \lambda_i \mathbf{p} \cdot \mathbf{a}_i\} = \sum_{i=1}^n \lambda_i \alpha_i$. ■

Remark 2.3. The multiplier form is a good reminder again

$$(\mathbf{p} \cdot \mathbf{b} - \beta) - \sum_{i=1}^m \lambda_i (\mathbf{p} \cdot \mathbf{a}_i - \alpha_i) = \mathbf{p} \cdot (\mathbf{b} - \sum_{i=1}^m \lambda_i \cdot \mathbf{a}_i) - \beta + \sum_{i=1}^m \lambda_i \alpha_i \geq 0, \quad \forall \mathbf{p} \in \mathbb{V}^*$$

Remark 2.4. The geometric interpretation is similar to the previous one, but everything is affine now. If the line (one dimensional affine hyperplane) (\mathbf{b}, β) does not belong to the (affine) cone generated by (\mathbf{a}_i, α_i) there exists an affine hyperplane separating \mathbf{b} from the cone.

2.2. Liu's theorem. Here the constraints are equalities instead of inequalities, therefore the multipliers are not necessarily positive.

Theorem 2.3. (Liu) Let $\mathbf{a}_i \neq \mathbf{0}$ be vectors in a finite dimensional vector space \mathbb{V} and α_i real numbers, $i = 1 \dots n$ and $S_L = \{\mathbf{p} \in \mathbb{V}^* | \mathbf{p} \cdot \mathbf{a}_i = \alpha_i, i = 1 \dots n\}$. The following statements are equivalent for a $\mathbf{b} \in \mathbb{V}$ and a real number β :

- (i) $\mathbf{p} \cdot \mathbf{b} \geq \beta$, for all $\mathbf{p} \in S_L$,
- (ii) There are real numbers $\lambda_1, \dots, \lambda_n$ such that

$$(1) \quad \mathbf{b} = \sum_{i=1}^n \lambda_i \mathbf{a}_i$$

and

$$(2) \quad \beta \leq \sum_{i=1}^n \lambda_i \alpha_i.$$

Proof:

A straightforward consequence of the previous affine form of Farkas's lemma because S_L can be given in a form S_A with the vectors \mathbf{a}_i and $-\mathbf{a}_i$, $i = 1, \dots, n$: $S_L = \{\mathbf{p} \in \mathbb{V}^* | \mathbf{p} \cdot \mathbf{a}_i \geq \alpha_i, \mathbf{p} \cdot (-\mathbf{a}_i) \geq \alpha_i, i = 1 \dots n\}$.

Therefore there are nonnegative real numbers $\lambda_1^+, \dots, \lambda_n^+$ and $\lambda_1^-, \dots, \lambda_n^-$ such, that $\mathbf{b} = \sum_{i=1}^n (\lambda_i^+ \mathbf{a}_i - \lambda_i^- \mathbf{a}_i) = \sum_{i=1}^n (\lambda_i^+ - \lambda_i^-) \mathbf{a}_i = \sum_{i=1}^n \lambda_i \mathbf{a}_i$ and $\beta \leq \sum_{i=1}^n (\lambda_i^+ \alpha_i - \lambda_i^- \alpha_i)$. ■

Remark 2.5. The multiplier form is a help in the applications again

$$0 \leq (\mathbf{p} \cdot \mathbf{b} - \beta) - \sum_{i=1}^n \lambda_i (\mathbf{p} \cdot \mathbf{a}_i - \alpha_i) = \mathbf{p} \cdot (\mathbf{b} - \sum_{i=1}^n \lambda_i \mathbf{a}_i) - \beta + \sum_{i=1}^n \lambda_i \alpha_i, \quad \forall \mathbf{p} \in \mathbb{V}^*.$$

Remark 2.6. In the theorem with Lagrange multipliers for local conditional extremum of differentiable function we apply exactly the above theorem of linear algebra after a linearization of the corresponding functions at the extremum point.

Considering the requirements of the applications we generalize Liu's theorem to take into account vectorial constraints:

Theorem 2.4. (vector Liu) Let $\mathbf{A} \neq \mathbf{0}$ in a tensor product $\mathbb{V} \otimes \mathbb{U}$ of finite dimensional vector spaces \mathbb{V} and \mathbb{U} . Let $\alpha \in \mathbb{U}$ and $S_L = \{\mathbf{p} \in \mathbb{V}^* | \mathbf{p} \cdot \mathbf{A} = \alpha\}$. The following statements are equivalent for a $\mathbf{b} \in \mathbb{V}$ and a real number β :

- (i) $\mathbf{p} \cdot \mathbf{b} \geq \beta$, for all $\mathbf{p} \in S_L$.
- (ii) There is a vector λ in the dual of \mathbb{U} such that

$$(3) \quad \mathbf{b} = \mathbf{A} \cdot \lambda,$$

and

$$(4) \quad \beta \leq \lambda \cdot \alpha.$$

Proof:

Let us observe that we can get back the previous form of the theorem by introducing a linear bijection $\mathbf{K} : \mathbb{U} \rightarrow \mathbb{R}^n$, a *coordinatization* in \mathbb{U} . Therefore, applying it for $\mathbf{K} \cdot \mathbf{A} = (\mathbf{A})_i = \mathbf{a}_i$, $\mathbf{K} \cdot \alpha = (\alpha)_i = \alpha_i$ and $\mathbf{K}' \cdot \mathbf{A} = \mathbf{a}'_i$, $\mathbf{K}' \cdot \alpha = \alpha'_i$ we get that $\mathbf{b} = \sum_{i=1}^n \lambda_i \mathbf{a}_i = \sum_{i=1}^n \lambda'_i \mathbf{a}'_i$. Thus $\lambda'_i = \mathbf{K}'^{*-1} \cdot \mathbf{K}^* \cdot \lambda_i$. Therefore there is

a $\lambda \in \mathbb{U}$, independently of the coordinatization, with the components λ_i and λ'_i in the coordinatizations \mathbf{K} and \mathbf{K}' . ■

The previously excluded degenerate case of $\mathbf{A} = \mathbf{0}$ deserves a special attention. Now we require the validity of the aim inequality for all $\mathbf{p} \in \mathbb{V}^*$ without any constraint. The consequences can be formulated as previously and the proof is trivial.

Theorem 2.5. (*degenerate Liu*) *The following statements are equivalent for a $\mathbf{b} \in \mathbb{V}$ and a real number β :*

- (i) $\mathbf{p} \cdot \mathbf{b} \geq \beta$ for all $\mathbf{p} \in \mathbb{V}^*$.
- (ii) *There is a vector λ in the dual of \mathbb{U} such that $\mathbf{b} = \mathbf{0}$ and $\beta \leq 0$.*

Remark 2.7. *The practical application rule is that if $\mathbf{A} = \mathbf{0}$ then the multiplier is zero.*

Remark 2.8. *In continuum physics and thermodynamics the corresponding form of (3) and (4) are called Liu equation(s) and the dissipation inequality, respectively. We apply the same names for the degenerate case, too. The Lagrange-Farkas multipliers are called simply Lagrange multipliers. Our nomination takes into account the rightful priority of Farkas, and emphasizes the difference between the two kind of multipliers. It can be important also to make a clear distinction of a similar but different nomination and method in variational principle construction in continuum physics [11].*

3. APPLICATION IN THERMODYNAMICS - METHODS OF LIU AND COLEMAN-NOLL

Originally Farkas developed his lemma to formulate correctly the Fourier's principle of mechanics of mass-points, which is the generalization of d'Alembert's principle in case of inequality constraints [12]. The role of Liu's theorem in continuum physics is similar in some sense: we are to formulate the correct form of the evolution equations (restricting the constitutive, material functions) taking into account the requirement of the entropy inequality.

The most predictive formulation of the problem is far from being trivial and originated from Coleman and Noll (see [3] and the references therein) and essentially reverses the way of thinking: we are looking for the solution of the entropy inequality taking into account the evolution equations as constraints. In continuum physics the dynamic equations are given in a determined form (as a balance for extensives), except some constitutive, material functions. The task is to ensure a nonnegative entropy production with appropriate constitutive assumptions. Therefore one should determine the undetermined material functions, that in case of all possible solutions of the dynamic equations the form of the constitutive functions, the material properties, ensure the nonnegativity of the entropy production. The entropy and the entropy current are also constitutive and are to be determined according to the above requirement. The entropy possibly should preserve its potential character in a general sense, therefore solving the above problem (e.g. Liu equations) the practical aim is to simplify the problem in such a way that every constitutive quantity (in the dynamic equations) and also the entropy current should be calculated from the entropy function.

There are two methods to exploit the entropy inequality.

- *Coleman-Noll procedure.* In the Coleman-Noll procedure one exploits the dynamic equations directly, substituting them into the entropy production. The constraints are substituted into the inequality and the degenerate form of Liu's theorem is applied. In this case one should assume a specific form of the entropy current. There are essentially two choices here. The entropy current can be the classical one ($\mathbf{j}_s = \mathbf{q}/T$), this is the usual assumption in also in weakly nonlocal considerations (see e.g. [13, 14]). The other choice

is to consider generalized entropy currents (or currents of other thermodynamic potentials). That can be done on different grounds and the method can be combined with the procedure.

- *Liu procedure.* With Liu procedure one applies Liu's theorem with Lagrange-Farkas multipliers. At the first glance, from the mathematical point of view, the application of this method seems to have only practical advantages. As the Lagrange multiplier method preserves the symmetry of functions in question in case of extremum problems, with Lagrange-Farkas multipliers we can preserve and exploit the structure of the entropy inequality. However, the question is not purely mathematical. First of all there can be cases when the multipliers could not be eliminated and they can get physical significance. Moreover, an inevitable advantage of Liu's method is that the structure of entropy inequality makes possible to solve completely the physical problem. With a proper choice of the state space we can determine the entropy current (solving the Liu equations). The entropy current should be considered as an independent constitutive quantity as it was suggested in extended rational thermodynamics [15]. Moreover, the entropy inequality can be solved, too, determining all constitutive quantities. This will ensure, that our theory is independent of further artificial constraints, the entropy inequality becomes a consequence of material properties. A basic philosophical requirement is, that the acceptable theories are those, where the entropy inequality is the property of the material and not our model (e.g. initial conditions).

There are some additional practical rules that we should take into account in this particular application of Farkas lemma. In Liu procedure the condition is a differential equation (a partial one in continuum physics). The functions in the differential equation form the *basic state space*. The constitutive quantities depend on this functions, on the *basic state* and some of its derivatives. To make the problem algebraically manageable the basic state variables and their derivatives are considered as independent quantities. Some of them can be included into the *constitutive state space* (or large state space [4]), into the domain of the constitutive functions. The aim inequality has a special balance form, and determines the independent variables of the algebraic problem: those are the derivatives of the constitutive state, the so called *process directions*. The choice of the constitutive state space is crucial and can result in different kind of restricted constitutive functions, after applying Liu procedure [16]. In the following we will show some examples.

Remark 3.1. *The mentioned exploitation methods of the second law, being algebraic, are essentially independent of the solvability of the dynamic equations, whether the associated problems are well or ill posed, on the applied function spaces, etc... For example the number of the equations can be less than the number of variables in the basic state space.*

Let us see a simple example to clarify the most important practical rules in the application of the formalism.

Example 1. *In this example the basic state space is formed by two times differentiable real functions $x : \mathbb{R} \mapsto \mathbb{R}$. Our constitutive space let be spanned by the basic state and its derivative (x, Dx) . We are looking for scalar valued differentiable functions F and S interpreted on the state space that*

$$D(S(x, Dx)) \geq 0$$

for all (x, Dx) satisfying

$$(5) \quad F(x, Dx) = 0.$$

Evidently $DS(x, Dx) = \partial_1 SDx + \partial_2 SD^2x$, where ∂_n denotes the partial derivative according to the n th variable. Therefore, the space of the process directions (the space of independent variables) is spanned by D^2x . We are looking for conditions on S and F that the above inequality should be true for all (x, Dx) solving (5), but independently of the values of D^2x . Thus the degenerate case of Liu's theorem gives some conditions. The Liu equation

$$\partial_2 S = 0.$$

Therefore S is independent on Dx . The dissipation inequality can be written in the following simple form

$$(6) \quad \partial_1 SDx = \frac{dS}{dx}(x)Dx \geq 0$$

The above inequality does not give any condition for F from S .

e. with $\lambda_1 \equiv 0$ $\lambda_1 F \geq 0$ is satisfied. It does not mean that the inequality (6) is meaningless or is always fulfilled independently of F . With a given F one can determine λ_1 and give conditions for the form of S . But one cannot make reversely, given an S it is easy to find a λ_1 so that the inequality is fulfilled for any F . In the following λ_1 will be zero.

However, let us observe, that one of our previous assumptions was false. The independent variable D^2x is not really independent on the state space, (5) gives a restriction

$$\partial_1 FDx + \partial_2 FD^2x = 0.$$

Considering this condition we apply Liu's theorem with the multiplier method

$$\begin{aligned} \partial_1 SDx + \partial_2 SD^2x - \lambda_2(\partial_1 FDx + \partial_2 FD^2x) - \lambda_1 F \\ = (\partial_1 S - \lambda_2(\partial_1 F)Dx + (\partial_2 S - \lambda_2\partial_2 F)D^2x - \lambda_1 F \geq 0 \end{aligned}$$

Therefore we can read the Liu equation as follows

$$\lambda_2\partial_2 F + \partial_2 S = 0$$

Expressing the multiplier and substituting into the dissipation inequality we get

$$\partial_1 SDx - \lambda_2\partial_1 FDx - \lambda_1 F = (\partial_1 S - \partial_2 S(\partial_2 F)^{-1}\partial_1 F)Dx \geq 0$$

With $\lambda_1 = 0$ we can give the general solution of the above inequality, as

$$\partial_1 S - \partial_2 S(\partial_2 F)^{-1}\partial_1 F = L(x, Dx)Dx,$$

where L is a nonnegative. Given an S function we can calculate F , with appropriate conditions on L . For example if $S(x, Dx) = x \cdot Dx$ and $L = \text{const.}$ then $F(x, Dx) = f(x^{L-1}Dx)$ is a solution of the above equation for any f .

4. WEAKLY NONLOCAL THERMODYNAMICS – GINZBURG-LANDAU AND THERMODYNAMIC GINZBURG-LANDAU EQUATIONS

Ginzburg-Landau equation and variants appear in different fields of physics and are applied to several phenomena. Its physical content and the way how we get it is sound and transparent [17]. On the other hand Gurtin recognized that its derivation has an ad-hoc character and a natural expectation, that the Ginzburg-Landau equation should be compatible with the general balance and constitutive structure of continuum physics. He also gave a method to get it based on the concept of microforce balance [13]. There are some other attempts of the derivation also on different other grounds [18, 16, 19]. Here we will show that the physical assumptions to get the Ginzburg-Landau equation are even more moderate, it is a straightforward consequence of the entropy inequality.

Let us denote the internal variable (e.g. it can be an order parameter of a second order phase transition) characterizing the microstructure of the material by ξ . In this case the Ginzburg-Landau equation can be written as

$$(7) \quad \dot{\xi} = -\gamma_1 \Gamma_\xi + \gamma_2 \Delta \xi,$$

where Γ_ξ is partial derivative of the appropriate thermodynamic potential (e.g. $\partial_\xi f$, where f is the free energy), and the point denotes a substantial derivative. The usual form of the Ginzburg-Landau free energy density is

$$(8) \quad f(\xi, \nabla \xi) = f_0(\xi) - \gamma(\nabla \xi)^2/2.$$

Ván argued that the Ginzburg-Landau equation is the first nonlocal extension of any kind of equation for an internal variable. Its special form is can be derived only from the requirement of compatibility with the Second Law, without any variational principle. Hence, the reason of its wide-range applicability is well founded, because any internal variable that can characterize an material structure and is independent of other requirements is described by that kind of equation in the first nonlocal approximation. The arguments were supported by calculations based on Liu's theorem. However, Ván did not receive the Ginzburg-Landau equation, but only a very similar one, that he called thermodynamic Ginzburg-Landau equation

$$(9) \quad \dot{\xi} = \gamma_1 \Gamma_\xi + \gamma_2 \Delta \Gamma_\xi.$$

Moreover, the entropy (free energy) was proved to be gradient independent. One can see, that the equations (7) and (9) are similar but not the same at all. The essential qualitative difference is that the equilibrium solutions of the thermodynamic Ginzburg-Landau equation are homogeneous but equilibrium solutions of the original Ginzburg-Landau equation are not, they can form structures. The situation is well known and understood in superconductors. The London equation does not determine the structure of the penetration length of the magnetic field but the Ginzburg-Landau equation gives that [20]. It is remarkable that contrary to the quite general assumptions, the derivation given in [16] prefers a London like equation and forbid the Ginzburg-Landau like structure. In the following we will derive the Ginzburg-Landau equations from the same general assumptions and show how to get the Gurtin terms from a thermodynamic derivation.

Therefore, we are looking for a dynamic equation of ξ in the following general form

$$(10) \quad \dot{\xi} - \mathcal{F} = 0,$$

where \mathcal{F} is a constitutive function, which form is to be restricted by the Second Law. The basic state space is spanned by ξ . Let us assume that the state space is spanned by ξ , $\nabla \xi$ and $\nabla^2 \xi$. In this case the entropy inequality will be

$$\begin{aligned} \dot{s} + \nabla \cdot \mathbf{j}_s = \\ \partial_1 s \dot{\xi} + \partial_2 s \cdot (\nabla \dot{\xi}) + \partial_3 s : (\nabla^2 \dot{\xi}) + \partial_1 \mathbf{j}_s \cdot \nabla \dot{\xi} + \partial_2 \mathbf{j}_s : \nabla^2 \dot{\xi} + \partial_3 \mathbf{j}_s : \nabla^3 \dot{\xi} \geq 0. \end{aligned}$$

One can see, that the space of the independent variables is spanned by $\dot{\xi}$, $\nabla \dot{\xi}$, $\nabla^2 \dot{\xi}$ and $\nabla^3 \dot{\xi}$. However, let us observe that these variables are not really independent, the derivative of (10) connect them. Therefore, in addition to (10) one should consider the following constraint, too

$$(11) \quad \nabla \dot{\xi} + \nabla \mathcal{F} = \nabla \dot{\xi} + \partial_1 \mathcal{F} \nabla \xi + \partial_2 \mathcal{F} \cdot \nabla^2 \xi + \partial_3 \mathcal{F} : \nabla^3 \xi = 0.$$

Introducing the Γ_1 and Γ_2 Farkas multipliers for the constraints (10) and (11) respectively, one can get the following Liu equations

$$\begin{aligned}\partial_1 s &= \Gamma_1, \\ \partial_2 s &= \Gamma_2, \\ \partial_3 s &= \mathbf{0}, \\ \partial_3 \mathbf{j}_s - \Gamma_2 \partial_3 \mathcal{F} &= \mathbf{0}.\end{aligned}$$

The first two equations determine the multipliers. From the third follows, that the entropy does not depend on the second derivative of ξ . Taking into account these requirements one can solve the fourth equation and determine the entropy current in the following form

$$(12) \quad \mathbf{j}_s(\xi, \nabla \xi, \nabla^2 \xi) = \partial_2 s(\xi, \nabla \xi) \mathcal{F}(\xi, \nabla \xi, \nabla^2 \xi) + \mathbf{j}_0(\xi, \nabla \xi).$$

For the sake of brevity we explicitly denoted the variables of the corresponding functions. With the above solution of the Liu equations the dissipation inequality can be considerably simplified

$$(13) \quad \nabla \cdot \mathbf{j}_0 + (\nabla \cdot \partial_2 s - \partial_1 s) \cdot \mathcal{F} \geq 0.$$

Assuming, that $\mathbf{j}_0 \equiv 0$ one can give the general solution of the above inequality. That solution can be interpreted as the well known traditional method of irreversible thermodynamics, choosing appropriate forces and currents so, that the constitutive quantity to be determined should be proportional to the given one

$$(14) \quad \dot{\xi} = \mathcal{F} = L(\nabla \cdot \partial_2 s - \partial_1 s)$$

with an L nonnegative state dependent constitutive function. $s(\xi, \nabla \xi)$ is the given (determined from static measurements) entropy function. (14) is the Ginzburg-Landau equation, and one can get back the very traditional (7) form if uses the specific entropy functional (8) and deals with a strictly linear theory in a thermodynamic sense, hence L is a constant function. The choice of the right thermodynamic potential (entropy or free energy) depends on the other thermodynamic boundary conditions.

If we do not restrict the space of the independent variables by (11), by the derivative of the original constraint, then according to [16] one can get the following form of the dissipation inequality

$$\nabla \cdot \mathbf{j}_0(\xi, \nabla \xi) - \partial_1 s(\xi) \mathcal{F}(\xi, \nabla \xi) \geq 0$$

It is important here, that the entropy must not depend on the gradients of the basic state. According to [16] one can solve the above inequality, too, considering two additional physical requirements. First that ξ is a dynamic variable in a thermodynamic sense therefore ξ is zero in equilibrium. On the other hand there is no entropy flow connected to the dynamic variable if its value is zero. Ván argued that these requirements are very weak and cover all possibilities that appeared up to now in the practice. With these assumptions one can specify \mathbf{j}_0 with the Nyíri-form in the entropy current [21] as $\mathbf{j}_0(\xi, \nabla \xi) = \mathbf{A}(\xi, \nabla \xi) \xi$, or equivalently $\mathbf{j}_0(\xi, \nabla \xi) = \hat{\mathbf{A}}(\xi, \nabla \xi) \partial_\xi s$. Here \mathbf{A} and $\hat{\mathbf{A}}$ are constitutive functions to be determined. With the second form of \mathbf{j}_0 the solution of the dissipation inequality gives the thermodynamic Ginzburg-Landau equation (9). Introduction of another new dynamic variable gives the additional Gurtin-term in the thermodynamic Ginzburg-Landau equation. Let us observe, that the key assumption determining the form of the additional entropy current was that the entropy current should not affect the equilibrium solutions. The most direct application of that physical requirement is to assume \mathbf{j}_0 in the following form

$$\mathbf{j}_0(\xi, \nabla \xi) = \hat{\mathbf{B}}(\xi, \nabla \xi) \mathcal{F}.$$

In this case the entropy production (13) is

$$(15) \quad \hat{\mathbf{B}} \nabla \mathcal{F} + (\nabla \cdot \hat{\mathbf{B}} + \nabla \cdot \partial_2 s - \partial_1 s) \cdot \mathcal{F} \geq 0.$$

The inequality has an Onsagerian general solution, the currents and forces are determined by the constitutive dependencies

$$\begin{aligned} \hat{\mathbf{B}} &= L_1 \nabla \mathcal{F} \\ \mathcal{F} &= L_2 (\nabla \cdot \hat{\mathbf{B}} + \nabla \cdot \partial_2 s - \partial_1 s). \end{aligned}$$

Here L_1 and L_2 are nonnegative constitutive functions. \mathbf{B} can be eliminated from the above equations and we get

$$(16) \quad \dot{\xi} = \mathcal{F} = L_2 (\nabla \cdot \partial_2 s - \partial_1 s) + L_2 \nabla \cdot (L_1 \nabla \dot{F}).$$

This is the Ginzburg-Landau equation with the characteristic additional Gurtin-term.

5. CONCLUSIONS AND DISCUSSION

The requirement a nonnegative entropy production is a relatively strong and not a complete form of the Second Law. Strong form because it is a local requirement and other weaker formulations require only the validity of integral inequalities. Not a complete one because an increasing entropy is only a part of the physical content of the Second Law. Material stability, the physical content of the Second law incorporates some other conditions (concave entropy function), too.

The traditional derivation of the Ginzburg-Landau equation has two main ingredients

- The static, equilibrium part is derived from a variational principle.
- The dynamic part is added by obscure stability arguments (relaxational form).

In this paper we unified the two parts in a thermodynamic derivation, where we did not refer to any kind of variational principle, however, the derived static part had the form that one could get from variational thinking, it has a complete Euler-Lagrange form. The dynamic part contains a first order time derivative therefore one cannot hope to derive it from a variational principle of Hamiltonian type [22]. In our approach we get the "reversible" part as a specific case of the thermodynamic, irreversible thinking, but one cannot get the irreversible part from a variational, reversible thinking.

These critical remarks were clearly formulated by Gurtin [13], who investigated that kind of additional terms in the Ginzburg-Landau and Cahn-Hilliard equations for mechanical systems, with the concept of microforce balance. However, that term appears in very different fields of physics starting from such classical equations as the Guyer-Krumhansl equation of weakly nonlocal heat conduction [23, 24] up to recent studies of the motion of granular materials [25], among others. Our last example with Ginzburg-Landau equation can be considered as a refinement of the investigations of [16] regarding the thermodynamic origin of that term. On the other hand, from a general point of view, the mathematical background and the key ingredients of an efficient formalism to exploit the second law inequality in weakly nonlocal continuum theories is given.

REFERENCES

- [1] I-Shih Liu. Method of Lagrange multipliers for exploitation of the entropy principle. *Archive of Rational Mechanics and Analysis*, 46:131–148, 1972.
- [2] I. Müller. On the frame dependence of stress and heat flux. *Archive of Rational Mechanics and Analysis*, 45:241–250, 1972.

- [3] W. Muschik and H. Ehrentraut. An amendment to the Second Law. *Journal of Non-Equilibrium Thermodynamics*, 21:175–192, 1996.
- [4] W. Muschik, C. Papenfuss, and H. Ehrentraut. A sketch of continuum thermodynamics. *Journal of Non-Newtonian Fluid Mechanics*, 96:255–290, 2001.
- [5] R. A. Hauser and N. P. Kirchner. A historical note on the entropy principle of Müller and Liu. *Continuum Mechanics and Thermodynamics*, 14:223–226, 2002. Lecture held on CIMRF’01, Berlin, 3-6 September, 2001.
- [6] H. Minkowski. *Geometrie der Zahlen*. Teubner, Leipzig und Berlin, 1896.
- [7] A. Haar. A lineáris egyenlőtlenségekről. *Mathematikai és Természettudományi Értesítő*, 36:279–296, 1918. in Hungarian.
- [8] Gy. Farkas. Egyenlőtlenség alkalmazásának új módjai. *Mathematikai és Természettudományi Értesítő*, 36:297–308, 1918. in Hungarian.
- [9] J. von Neuman. Discussion of a maximum principle. In A. H. Taub, editor, *Collected Works*, volume VI, pages 89–95, New York and Oxford, 1961. Pergamon Press.
- [10] A. Schriver. *Theory of linear and integer programing*. John Wiley and Sons, Chicester-etc., 1998.
- [11] S. Sieniutycz and R. S. Berry. Variational theory for thermodynamics of thermal waves. *Physical Review E*, 65(046132), 2002.
- [12] Gy. Farkas. A Fourier-féle mechanikai elv alkalmazásai. *Mathematikai és Természettudományi Értesítő*, 12:457–472, 1894. in Hungarian.
- [13] M. G. Gurtin. Generalized Ginzburg-Landau and Cahn-Hilliard equations based on a micro-force balance. *Physica D*, 92:178–192, 1996.
- [14] V. A. Cimmelli and Kosiński W. Gradient generalization to internal state variables and a theory of superfluidity. *Journal of Theoretical and Applied Mechanics*, 35(4):763–779, 1997.
- [15] I. Müller and T. Ruggeri. *Rational Extended Thermodynamics*, volume 37 of *Springer Tracts in Natural Philosophy*. Springer Verlag, New York-etc., 2nd edition, 1998.
- [16] P. Ván. Weakly nonlocal irreversible thermodynamics. cond-mat/0112214, submitted to *Annalen der Physik*, 2001.
- [17] P. C. Hohenberg and B. I. Halperin. Theory of dynamic critical phenomena. *Reviews of Modern Physics*, 49(3):435–479, 1977.
- [18] P. M. Mariano. Multifield theories in mechanics of solids. *Advances in Applied Mechanics*, 38:1–94, 2002.
- [19] P. Ván. Weakly nonlocal irreversible thermodynamics - the Ginzburg -Landau equation. *Technische Mechanik*, 22(2):104–110, 2002. cond-mat/0111307.
- [20] L. D. Landau and L. P. Pitajevskij. *Statistical Physics II*, volume 9 of *Course of Theoretical Physics*. Pergamon Press, Oxford-etc., 1982.
- [21] B. Nyíri. On the entropy current. *Journal of Non-Equilibrium Thermodynamics*, 16:179–186, 1991.
- [22] P. Ván and W Muschik. Structure of variational principles in nonequilibrium thermodynamics. *Physical Review E*, 5(4):3584–3590, 1995.
- [23] R. A. Guyer and J. A. Krumhansl. Solution of the linearized phonon Boltzmann equation. *Physical Review*, 148(2):766–778, 1966.
- [24] R. A. Guyer and J. A. Krumhansl. Thermal conductivity, second sound and phonon hydrodynamic phenomena in nonmetallic crystals. *Physical Review*, 148(2):778–788, 1966.
- [25] T. Unger, Brendel L., D. E. Wolf, and Kertész J. Elastic behaviour in contact dynamics of rigid particles. *Physical Review E*, 65, 2002. 061305.

BUDAPEST UNIVERSITY OF TECHNOLOGY AND ECONOMICS, DEPARTMENT OF CHEMICAL PHYSICS,
1521 BUDAPEST, BUDAFOKI ÚT 8.

E-mail address: vpet@phyndi.fke.bme.hu