

# A possible mechanism for cold denaturation of proteins at high pressure

Manuel I. Marqués<sup>1,\*</sup>, Jose M. Borreguero<sup>1</sup>, H. Eugene Stanley<sup>1</sup>, and Nikolay V. Dokholyan<sup>2</sup>

<sup>1</sup> *Center for Polymer Studies and Department of Physics, Boston University, Boston, MA 02215, USA*

<sup>2</sup> *Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, School of Medicine, Chapel Hill, NC 27599, USA*

We study the cold denaturation of a model protein at high pressure. Using multicanonical Monte Carlo simulations of a model protein in a water bath, we study the effect of water density fluctuations on the stability of a protein under these conditions. We find that a possible mechanism for the cold denaturation of proteins is the loss of local water structure due to the disruption of hydrogen bonds between water molecules at high pressures.

Some proteins become thermodynamically unstable at low temperatures, a phenomenon termed cold denaturation [1, 2, 3]. This phenomenon has been mainly observed at high pressures, ranging from approximately 200 MPa up to 700 MPa [4, 5]. An explanation of the  $P-T$  phase diagram of a protein with cold denaturation has been proposed [6, 7, 8], but a microscopic understanding of the mechanisms leading to cold denaturation has yet to be developed, due in part to the complexity of protein-solvent interactions.

Existing theories of folding and unfolding of diluted proteins consider amino acid solvation (hydrophobicity) as the driving force of protein stability [9, 10, 11, 12, 13, 14]. The origin of hydrophobicity has been identified with the partial ordering of water around nonpolar molecules. Water molecules are highly organized, forming hydrogen bonded cages around the nonpolar molecules (clathrate structures) [15]. The simplest hydrophobic model is an effective interaction between hydrophobic molecules [16], but this simple models does not predict cold denaturation. An explicit account of water around the hydrophobic molecules is required in order to understand the cold denaturation process. Theoretical attempts modeled the water-polymer interactions with a bimodal description of the energy of water in the shell around the hydrophobic molecules [17, 18, 19] and a simplified lattice model based on this bimodal description was able to reproduce the cold denaturation of the model [20]. Similar results were obtained using a lattice model of a random hydrophobic-hydrophilic (HP) heteropolymer interacting with the solvent [21]. Several models have also been proposed to mimic the ice-like structures formed by the water molecules around the non-polar monomers [22, 23, 24, 25, 26].

One possible reason for the inability of simple models to capture the molecular details of cold denaturation is their neglect of (i) long range correlations among water molecules near the freezing point, and (ii) density anomalies due to the tetrahedral structure of the hydrogen bonded network. Here, we implement a two-dimensional lattice model of water that captures these above-mentioned water phenomena, and that has been successfully applied to understand water anomalies [27].

In the model, the possible orientations of water molecules are set by the allowed values of a  $q$ -state Potts variable  $\sigma_i$ . Only when two neighbor molecules ( $i, j$ ) are in the correct orientation ( $\sigma_i = \sigma_j$ ) a hydrogen bond (HB) with energy  $-J$  is formed. Once a HB is formed, the volume of the system increases by  $\delta V$ .

The Hamiltonian for our model of water-water interaction may be written as [27]

$$\mathcal{H}_{HB} = -J \sum_{\langle i,j \rangle} \delta_{\sigma_i, \sigma_j}, \quad (1)$$

where  $J > 0$  is the scale of the interaction between water molecules upon hydrogen bond formation. The total volume of the system is given by  $V = V_0 + N_{HB}\delta V$ , where  $N_{HB} = \sum_{\langle i,j \rangle} \delta_{\sigma_i, \sigma_j}$  is the total number of hydrogen bonds in the system. The sum  $\sum_{\langle i,j \rangle}$  extends only to nearest neighbors. The enthalpy of the system ( $\mathcal{H}_{HB} + PV$ ) is given by

$$\mathcal{H}_{HB} + PV = -(J - P\delta V) \sum_{\langle i,j \rangle} \delta_{\sigma_i, \sigma_j}, \quad (2)$$

where  $P$  is the pressure applied to the system.

For  $P > J/\delta V$ , formation of a hydrogen bond is not favorable from the energetic point of view, and the system undergoes a transition from the disordered state to an ordered, high density state with no hydrogen bonds ( $N_{HB} = 0$ ). For  $P < J/\delta V$ , the water model exhibits a first order phase transition from a disorder state to an ordered, low density state, with all hydrogen bonds formed at the critical temperature of the Potts model  $T_c = (J - P\delta V)/(\ln(1 + \sqrt{q}))$  [27]. Thus, our model reproduces the freezing of water to low and high density ice, since for  $P < J/\delta V$  ( $\approx 200$ MPa in real water) water freezes to the low density ice Ih, and for  $P > J/\delta V$  water freezes to the high density ice II [28]. A relation between these two phases of ice and polymer folding has been already suggested from a thermodynamic point of view in Ref. [29].

We model the protein dynamics as a self-avoiding random walk embedded in the water bath. For simplicity, we consider a non-polar homopolymer that interacts with

the surrounding water molecules via the hydrophobic effect, which is the result of the partial ordering of water forming clathrate structures around the homopolymer [15]. We mimic such interaction using the following Hamiltonian,

$$\mathcal{H}_p = J_r n_{HB} (n_{max} - \sum_{\langle i,j \rangle} n_i n_j), \quad (3)$$

where  $J_r > 0$  is the strength of the repulsive interaction,  $n_{HB} = N_{HB}/N_{water}$  is the number of hydrogen bonds per water molecule ( $N_{water}$  is the number of water molecules),  $n_i = 1$  if the position  $i$  is occupied with a monomer ( $n_i = 0$  otherwise) and finally,  $n_{max}$  is the maximum number of contacts that the homopolymer is able to make. The hydrophobic repulsion driving the polymer to a compact state is equal to zero when the polymer is in the lowest energy state ( $\sum_{\langle i,j \rangle} n_i n_j \approx n_{max}$ ) or when the water network is not able to form clathrate structures ( $n_{HB} \approx 0$ ).

We hypothesize that the inability of water molecules to form clathrate structures is the principal mechanism responsible for protein cold denaturation. At low pressures ( $P < J/\delta V$ ) and low temperatures, water molecules form a hydrogen bond network, so the polymer is forced to adopt a compact state. At high pressures ( $P > J/\delta V$ ), the water is not able to form the low density ice Ih, in which case, since no clathrates are formed, the apolar monomers feel no repulsion and the cold-denaturation of the homopolymer takes place. Our hypothesis is supported by the experimental observation that cold denaturation exists mainly at high pressures (of the order of Kbar), where water only freezes in the dense ice II phase.

Next we demonstrate that our model of a polymer embedded to a water network with an enthalpy given by

$$\mathcal{W} = \mathcal{H}_p + \mathcal{H}_{HB} + PV, \quad (4)$$

gives rise to both cold and warm denaturation of the polymer and agrees with the experimentally observed protein denaturation at high pressures.

Since the energy landscape of the polymer interacting with the water network is characterized by a multitude of local minima, at low temperatures we perform multicanonical Monte Carlo simulations to avoid transient trapping of our water-protein system in local energy minima. Specifically, we use the multiple-range random walk algorithm proposed by Wang and Landau [30] to calculate the density of states. We adopt the algorithm in order to embed the self-avoiding polymer into the lattice, and to calculate the two-parameter density of states  $g(N_{HB}, N_c)$ , where  $N_c$  is the number of monomer-monomer contacts ( $N_c = \sum_{\langle i,j \rangle} n_i n_j$ ). From the density of states we calculate the temperature and pressure dependence of the average number of monomer-monomer contacts,




FIG. 1: Normalized number of monomer-monomer contacts vs. temperature for pressure values  $PV_0/J = 0, 0.25, 0.5, 0.75, 1, 1.1, 1.125, 1.15, 1.175, 1.2, 1.25, 1.4, 1.5, 2$ . A magnified region where the cold denaturation takes place is shown in the inset. The dotted line corresponds to  $\bar{N}_c/n_{max} > 0.96$ . We represent the curve corresponding to  $P = J/\delta V$  by a bold line. The values of the parameters used are:  $J = 1$ ,  $J_r = 10$ ,  $\delta V = 1$  and  $q = 10$ .

$$\bar{N}_c = \sum_{N_{HB}} \sum_{N_c} N_c g(N_{HB}, N_c) \frac{e^{-\mathcal{W}(N_{HB}, N_c)/T}}{Z} \quad (5)$$

where  $Z$  is the partition function.

We perform Monte Carlo simulations of a system of 383 water molecules and a polymer consisting of 17 non-polar monomers with periodic boundary conditions. We present the dependence of  $\bar{N}_c/n_{max}$  on temperature for different values of the pressure both above and below the critical pressure  $P = J/\delta V$  on Fig.1. The calculated density of states  $g(N_{HB}, N_c)$  converges to the true value with an accuracy of the order of  $10^{-5}$ . The value of  $\bar{N}_c/n_{max}$  ranges from  $\bar{N}_c/n_{max} = 1$  (lowest energy state) to  $\bar{N}_c/n_{max} \approx 0.71$  (which is the average number of monomer-monomer contacts obtained at high temperatures). Only when the value of the pressure is larger than  $J/\delta V$ , we observe the cold denaturation of the homopolymer.

We also reconstruct the phase diagram of the water-polymer system in the  $P - T$  plane (Fig.2). We consider a homopolymer to be in the collapsed state if 96% of all possible contacts are formed, i.e., if  $\bar{N}_c/n_{max} > 0.96$ . For each pressure value, the freezing lines of water shown on Fig.2 are given by the temperature at which we observe the maximum in the specific heat of the water bath. We compare our findings to the experimental observations of Zhang et al. [31] for ribonuclease A (RNase A) from bovine pancreas studied by  $^1\text{H}$  NMR spectroscopy. We




FIG. 2:  $P$ - $T$  phase diagram for the polymer derived from Fig.1. Dashed lines indicate the freezing lines for model water. Water freezes in hydrogen bonded ice Ih for  $PV_0/J < 1$  and in dense non-hydrogen bonded ice II for  $PV_0/J > 1$ . In the inset we present the experimental results obtained by Zhang et al. [31] for the ribonuclease A (RNase A). Two typical configurations of the homopolymer are shown, one in the compact state and the other in the denaturated state.

find remarkable qualitative agreement between the experimental and numerical  $P - T$  phase diagrams. In both experimental and numerical  $P - T$  phase diagrams, we observe that cold denaturation occurs at high pressures and low temperatures, close to the freezing line and in the region where water molecules are not capable of forming clathrate structures.

Not all proteins behave equally as we decrease temperature at high pressure. In particular, there are some proteins that do not exhibit cold denaturation. We reproduce the variability of proteins dynamics at high pressure and low temperature by varying the hydrophobic parameter  $J_r$  to a lower value, effectively impeding a stable compact state for pressures above the  $P = J/\delta V$  line. In Fig.3 we present the phase diagrams obtained for different values of the hydrophobic interaction parameter  $J_r$ , ranging from 2 to 20. The shape of the phase diagram changes as we increase the value of the repulsive interaction  $J_r$ , allowing stabilization of the compact state and cold denaturation above the  $P = J/\delta V$  line.

We conclude that the effect of pressure on the water density is a key ingredient for understanding cold denaturation of proteins. The density anomaly of water is the main driving force forming the clathrate structures responsible of the hydrophobic effect, and the resulting compactness of the protein [2, 11, 13]. At extreme pressures water is not capable of forming hydrogen bonds and there is no density anomaly. In this scenario, the hydrophobic effect disappears and cold denaturation takes place. Our model supports this scenario. Interactions




FIG. 3:  $P$ - $T$  Phase diagram for polymers with  $J_r = 2, 5, 10, 20$ . Dashed lines indicate the computed freezing lines for water. Dotted line indicates the  $P = J/\delta V$  line.

among amino acids determine the dynamics of proteins at low temperature and high pressure, thus making some proteins more stable than others at these  $P - T$  conditions.

We thank NSF Grant CHE0096892 for support. M.I. Marqués acknowledges the Postdoctoral financial support of the Spanish Ministry of Education. J.M. Borreguero acknowledges the support of the Petroleum Research Fund.

---

\* Electronic address: manuel@argento.bu.edu

- [1] C. N. Pace and Ch. Tanford, *Biochemistry* **7**, 198 (1968).
- [2] P. L. Privalov, *Crit. Rev. Biochem. Mol. Biol.* **25**, 281 (1990); in T. E. Creighton, *Protein Folding* (W.H. Freeman, New York, 1992); G. P. Privalov and P. L. Privalov, *Methods Enzymol.* **323**, 31 (2000).
- [3] J. Jonas, *ACS Symp. S* **676**, 310 (1997).
- [4] S. Kunugi and N. Tanaka, *Biochimica et Biophysica Acta* **1595**, 329 (2002).
- [5] L. Smeller, *Biochimica et Biophysica Acta* **1595**, 11 (2002).
- [6] S. A. Hawley, *Biochemistry* **10**, 2436 (1971).
- [7] S. A. Hawley, *Methods Enzymol.* **49**, 14 (1978).
- [8] J. Zipp and W. Kauzmann, *Biochemistry* **12**, 4217 (1973).
- [9] K. A. Dill, D. O. V. Alonso and K. Hutchinson, *Biochemistry* **28**, 5439 (1989).
- [10] K. F. Lau and K. A. Dill, *Macromolecules* **22**, 3986 (1989).
- [11] K. A. Dill, *Biochemistry* **29**, 7133 (1990).
- [12] K. A. Dill and D. Stigter, *Adv Prot Chem* **46**, 59 (1995).
- [13] H. S. Chan and K. A. Dill, *Phys. Today* **46**(2), 24 (1993).
- [14] W. Kauzmann, *Adv. Protein Chem.* **14**, 1 (1959).
- [15] H. S. Frank and M. W. Evans, *J. Chem. Phys.* **13**, 507

- (1945)
- [16] C. Vanderzande, *Lattice Models of Polymers* (Cambridge University Press, Cambridge, 1998).
  - [17] N. Muller, *Acc. Chem. Res.* **23**, 23 (1990).
  - [18] B. Lee and G. Graziano, *J. Am. Chem. Soc.* **22**, 5163 (1996).
  - [19] K. A. T. Silverstein, A. D. J. Haymet and K. A. Dill, *J. Chem. Phys.* **111**, 8000 (1999).
  - [20] P. De los Rios and G. Caldarelli, *Phys. Rev. E* **62**, 8449 (2000); *Phys. Rev. E* **63**, 031802 (2001); G. Caldarelli and P. De los Rios, *J. Biol. Phys.* **27**, 279 (2001).
  - [21] A. Trovato, J. van Mourik and A. Marian, *Eur. Phys. J. B* **6**, 63 (1998).
  - [22] A. Hansen, M. H. Jensen, K. Sneppen and G. Zocchi, *Eur. Phys. J. B* **6**, 157 (1998).
  - [23] A. Bakk, *Phys. Rev. E* **63**, 061906 (2001); A. Bakk, J. S. Hoye, A. Hansen, *Biophys. J.* **82**, 713 (2002).
  - [24] P. Bruscolini and Lapo Casetti *Phys. Rev. E* **61**, R2208 (2000); *Phys. Rev. E* **64**, 051805 (2001); *J. Biol. Phys.* **27**, 243 (2001).
  - [25] P. Bruscolini, C. Buzano, A. Pelizzola and M. Pretti, *Phys. Rev. E* **64**, 050801 (2001).
  - [26] O. Collet, *Europhys. Lett.* **53**, 93 (2001).
  - [27] G. Franzese, M. I. Marqués and H. E. Stanley, *cond/mat0112341*; PRE in press.
  - [28] V. F. Petrenko and R. W. Whitworth, *Physics of Ice* (Oxford University Press, Oxford, 1999).
  - [29] G. Wilse Robinson and C. H. Cho, *Biophys. J.* **77**, 3311 (1999).
  - [30] F. Wang and D. P. Landau, *Phys. Rev. Lett* **86**, 2050 (2001).
  - [31] J. Zhang, X. Peng, A. Jonas and J. Jonas, *Biochemistry* **34**, 8631 (1995).