Generation of dipole squeezing in a two-mode system with entangled coherent states of a quantized electromagnetic field

Qin Rao

Department of Engineering Physics, Queen's University, Kingston, ON K7L 3N6, Canada

Rui-Hua Xie

Department of Chemistry, Queen's University, Kingston, ON K7L 3N6, Canada (November 10, 2018)

Abstract

Two-mode quantized electromagnetic fields can be entangled and admit a large number of coherent states. In this paper, we consider a two-mode system that consists of a two-level atom interacting with a two-mode quantized electromagnetic field, which is initially prepared in an entangled two-mode coherent state, via a nondegenerate two-photon process in a lossless cavity. We study the quantum fluctuations in the two-mode system and investigate in detail the effects of detuning, Stark shift and atomic coherence on atomic dipole squeezing (ADS). We show that ADS strongly depends on the atomic coherence. It is found that the stronger the correlations between the two modes are involved, the more the ADS could be generated. The detuning or Stark shift has a destructive effect on ADS, but the combined effect of the detuning and Stark shift may lead to a regular, periodical and strong ADS pattern.

Keywords: quantum fluctuations, squeezed state, detuning, Stark shift. PACS (numbers): 42.50.Dv, 42.50.Lc, 03.67.-a

I. INTRODUCTION

During the last few years, it has been shown that quantum entanglement of states [1–7] is considered to be a basic prerequisite for teleporation of quantum state, quantum communication, quantum computing, quantum information processing and experimentally testing quantum mechanics. Theoretical and experimental studies on quantum communication through the use of quantum entangled states have shown great promise for establishing new kinds of information processing with no classical counterparts. One example is the concept of quantum teleporation [4], where an entangled source was built from two single-mode phase squeezed vacuum states combined at a beamsplitter. Other examples of the application of entanglement states to quantum communication include dense coding [7] and cryptography [3]. It is believed that the use of entangled states can lead to a substantial decrease of computational resources [5]. Also, entangled states have been widely used to test fundamental quantum features such as nonlocality [6]. These extensive studies have paved a way toward a variety of useful quantum informational devices.

Two-mode quantized electromagnetic fields can be entangled and admit a large number of coherent states [8]. Most of these coherent states may be associated with low-order Lie algebra su(1,1) [8,9]. For example, two interesting types of su(1,1) coherent states are the Perelomov coherent states (PCS) [8,10,11] and Barut-Girardello coherent states [8,12]. The former involves strong correlations between the modes and is generated from the unitary evolution of two-mode number states driven by a nondegenerated parametric device [8] (for example, the type that gives rise to down-conversion or harmonic generation). The latter type also involves tight correlations with respect to the photon number states between the modes and may be generated in a process involving the competition between nondegenerate parametric amplification and nondegenerate two-photon absorption [8]. The other class of two-mode su(1,1) coherent states is the intelligent states [13], which equalize various possible uncertainty product that can be constructed as consequence of the su(1,1) commutation relation. Also, the two-mode squeezed pair-coherent state [14] is an intelligent state but

with unequal uncertainty. All of these two-mode su(1,1) coherent states may be written as superposition of the form $\sum_{n=0}^{\infty} C_n^q \mid n+q \rangle_1 \mid n \rangle_2$ [8], where the modes are labeled by 1 and 2, respectively, and the integer number q is the (fixed) difference in the photon numbers of the two-modes and the coefficient C_n^q depends on the specific su(1,1) coherent state. These states are of interest because they often display nonclassical characteristics such as squeezed light and photon antibunching as well as violations of the Cauchy-Schwarz inequality and Bell's inequality [11,15]. In this paper, we study further the properties of these correlated states of the field since the importance lies in their close connection to the two-photon nonlinear optical processes. On the other hand, because the study of the squeezed states can provide a fundamental understanding of quantum fluctuations and open the way for new schemes of quantum communication or imaging beating the standard quantum noise limit, we pay attention to investigating the generation of atomic dipole squeezing (ADS) [16] in a two-mode system that consists of a two-level atom interacting with a quantized electromagnetic field, which is initially prepared in an entangled two-mode coherent state, via a nondegenerate two-photon process in a lossless cavity. We find that entangled twomode coherent states of the field can also display ADS and that the stronger the correlations between the modes are involved, the more the ADS could be generated.

The organization of this paper is as follows. In section II, we describe the theoretical model of the two-mode system and give its general solutions. In section III, we briefly introduce the definition of atomic dipole squeezing. In section IV, assuming that the twolevel atom is initially prepared in a coherent superposition of its ground and excited states and that the quantized electromagnetic field is initially in a two-mode PCS, we investigate the generation of ADS in the two-mode system and examine in detail the effects of detuning, Stark shift and atomic coherence on ADS. Finally, our conclusion and discussions are presented in section V.

II. THEORETICAL MODEL

We consider a two-level atom interacting with two modes of a quantized electromagnetic field via a nondegenerate two-photon process in a lossless cavity. For convenience, we denote ω as the natural transition frequency between the excited state $|+\rangle$ and the ground state $|-\rangle$ of the two-level atom and Ω_j as the frequency of mode j of the field. It is known that some intermediate virtual states | **virtual** \rangle , which are assumed to be coupled to the excited and ground states of the two-level atom by a dipole-allowed transition, are involved in the two-photon process. We assume that Ω_1 is about several times Ω_2 and the two modes are tuned as near one-photon resonance as possible but still off the one-photon linewidth, i.e., the detuning δ_1 and δ_2 are larger than but as close as the linewidth of | **virtual** \rangle . For the case of $\delta_1 = -\delta_2 = \delta$ and by the method of adiabatic elimination of the virtual state | **virtual** \rangle [17], Gou [18] obtained an effective Hamiltonian for this system within a rotating-wave approximation

$$H = \Omega_1 a_1^{\dagger} a_1 + \Omega_2 a_2^{\dagger} a_2 + \omega S_z + \beta_2 S_+ S_- a_2^+ a_2 + \beta_1 S_- S_+ a_1^{\dagger} a_1 + \epsilon (a_1^{\dagger} a_2^{\dagger} S_- + a_1 a_2 S_+), \quad (1)$$

where $\epsilon = \sqrt{\beta_1 \beta_2}$. Here S_z and S_{\pm} are operators of the atomic pseudospin inversion and transition, respectively, and satisfy the commutation

$$[S_+, S_-] = 2S_z, (2)$$

$$[S_z, S_{\pm}] = \pm S_{\pm}.\tag{3}$$

 a_i^{\dagger} and a_i are the creation and annihilation operators for the photons of mode *i*, respectively, which obey the boson operators' commutation relation

$$[a_i, a_j^{\dagger}] = \delta_{ij}. \tag{4}$$

 ϵ is the coupling constant between the atom and the field. A Stark shift is caused by the intermediate virtual state | **virtual** \rangle , and the corresponding parameter β_1 and β_2 are given by $\beta_1 = \epsilon_1^2/\delta$, $\beta_2 = \epsilon_2^2/\delta$ and $\epsilon = \epsilon_1\epsilon_2/\delta$, where $\delta = \omega - (\Omega_1 + \Omega_2)$ and ϵ_1 and ϵ_2 are the

coupling strengths of the intermediate virtual state with the ground and excited states of the two-level atom, respectively. Throughout we employ the units of $\hbar = c = 1$.

We denote $|n\rangle$ as the Fock state of the field. Using the standard bare-state procedure introduced by Yoo and Eberly [19], we are able to calculate the expectation value of any operator in the two-mode system described above. In an ideal cavity, the two-level atom absorbs and emits one pair of photons, and the two bases in the bare-state representation are given by $|+, n_1, n_2\rangle$ and $|-, n_1 + 1, n_2 + 1\rangle$, respectively. Thus, the total Hamiltonian given by Eq.(1) can be written as:

$$H = \begin{bmatrix} \Omega_1 n_1 + \Omega_2 n_2 + \omega/2 + \beta_2 n_2 & \epsilon \sqrt{(n_1 + 1)(n_2 + 1)} \\ \epsilon \sqrt{(n_1 + 1)(n_2 + 1)} & \Omega_1 (n_1 + 1) + \Omega_2 (n_2 + 1) - \omega/2 + \beta_1 (n_1 + 1) \end{bmatrix}.$$
 (5)

The eigenvalues of the above Hamiltonian are

$$\sigma_{+} = \Omega_{1}n_{1} + \Omega_{2}n_{2} + \frac{1}{2}[\omega + \beta_{2}n_{2} + \beta_{1}(n_{1} + 1)] + \epsilon\Gamma(n_{1}, n_{2}), \tag{6}$$

$$\sigma_{-} = \Omega_1 n_1 + \Omega_2 n_2 + \frac{1}{2} [\omega + \beta_2 n_2 + \beta_1 (n_1 + 1)] - \epsilon \Gamma(n_1, n_2)$$
(7)

with

$$\Gamma(n_1, n_2) = \sqrt{\chi^2(\mu, n_1, n_2)/4 + (n_1 + 1)(n_2 + 1)},$$
(8)

$$\chi(\mu, n_1, n_2) = \begin{cases} \Delta + [\mu(n_1 + 1) - n_2]/\sqrt{\mu} & \mu \neq 0, \\ \Delta & \mu = 0, \end{cases}$$
(9)

where $\mu = \beta_1/\beta_2$ and $\Delta = \delta/\epsilon$ is the scaled detuning and the two-photon Rabi frequency is written as

$$\Omega(n_1, n_2) = \epsilon \Gamma(n_1, n_2). \tag{10}$$

It should be mentioned that $\mu = 0$ is the case in the absence of Stark shift. Then, the time evolution operator $U(n_1, n_2, t)$ of the two-mode system is given by

$$U(n_1, n_2, t) = \frac{1}{2} \begin{bmatrix} E_+(t) & \zeta(t) \\ \zeta(t) & E_-(t) \end{bmatrix},$$
(11)

where

$$E_{+}(t) = \left[1 + \frac{\chi(\mu, n_{1}, n_{2})}{2\Gamma(n_{1}, n_{2})}\right] \exp(-i\sigma_{+}t) + \left[1 - \frac{\chi(\mu, n_{1}, n_{2})}{2\Gamma(n_{1}, n_{2})}\right] \exp(-i\sigma_{-}t),$$

$$E_{-}(t) = \left[1 - \frac{\chi(\mu, n_{1}, n_{2})}{2\Gamma(n_{1}, n_{2})}\right] \exp(-i\sigma_{+}t) + \left[1 + \frac{\chi(\mu, n_{1}, n_{2})}{2\Gamma(n_{1}, n_{2})}\right] \exp(-i\sigma_{-}t),$$

$$\zeta(t) = \sqrt{(n_{1} + 1)(n_{2} + 1)} \left[\exp(-i\sigma_{+}t) - \exp(-i\sigma_{-}t)\right] / \Gamma(n_{1}, n_{2}).$$
(12)

In the resonant case (i.e, $\Delta = 0$), our solutions are in agreement with those obtained by Gou [18,20]. Once the matrix representation of $U(n_1, n_2, t)$ is obtained, the density operator of the two-mode system at time t can be calculated with an arbitrary initial condition $\rho(t = 0)$ by

$$\rho(t) = U(n_1, n_2, t)\rho(t=0)U^{\dagger}(n_1, n_2, t).$$
(13)

Thus, the expectation value of any physical operator O and its dependence on time can be obtained through the formula

$$\langle O(t) \rangle = Tr[\rho(t)O(0)]. \tag{14}$$

III. DEFINITION OF ATOMIC DIPOLE SQUEEZING

Squeezed states [1,21,22] are distinguished by the property that the quantum fluctuations in a dynamical observable may be reduced below the standard quantum noise limit at the expense of increased fluctuations in its canonical conjugated variable without violating the Heisenberg uncertain relation. Since the first observation of squeezed states of light, the generation of squeezed states has been the subject of an intense theoretical and experimental activity [1,23–25]. The study of these states provides a fundamental understanding of quantum fluctuations and opens a way for new schemes [1,23–25] of quantum communication or imaging beating the standard quantum noise limit. It has been predicted that a number of physical systems [1,23–27] (e.g., four-wave mixing, two-photon laser, parametric amplifiers and Raman scattering) can generate squeezed states of the radiation field. Also, it is known that atomic dipole squeezing (ADS) [16] could be generated through many kinds of nonlinear optical processes [16,28–31], such as resonance fluorescence, cooperative Dicke system, Rydberg atom maser, and so on. For the special case of a vacuum-field mode, Wodkiewicz et al. [32] examined the relationship between ADS and field squeezing. Very recently, a general relationship between field squeezing and ADS has been established under different initial conditions for both the field and atom in a high-Q micromaser cavity [33–38], and it has been shown that squeezed atom can generate squeezed light. In this paper, assuming that the field is initially prepared in an entangled two-mode coherent states, we study the generation of ADS in the two-mode system introduced above.

Here we briefly introduce the definition of atomic dipole squeezing [16]. Following the standard procedure, we define two slowly varying Hermitian quadrature operators [31]

$$S_x = \frac{1}{2} \left[S_+ \exp(-i\omega t) + S_- \exp(i\omega t) \right],$$
(15)

$$S_y = \frac{1}{2i} \left[S_+ \exp(-i\omega t) - S_- \exp(i\omega t) \right], \tag{16}$$

where S_x and S_y , actually, correspond to the dispersive and absorptive components of the slowly varying atomic dipole [39], respectively. They obey a commutation relation given by $[S_x, S_y] = iS_z$. Correspondingly, the Heisenberg uncertainty relation is [31]

$$(\Delta S_x)^2 (\Delta S_y)^2 \ge \frac{1}{4} \left| \langle S_z \rangle \right|^2, \tag{17}$$

where $(\Delta S_j)^2 = \langle S_j^2 \rangle - \langle S_j \rangle^2$ is the variance in the quadrature component S_j (j=x,y) of the atomic dipole. For convenience, we define the following functions [31]

$$F_1 = (\Delta S_x)^2 - \frac{1}{2} \left| \langle S_z \rangle \right|, \qquad (18)$$

$$F_{2} = (\Delta S_{y})^{2} - \frac{1}{2} |\langle S_{z} \rangle|.$$
(19)

Then, the atomic state is said to be squeezed if $F_1 < 0$ (or $F_2 < 0$). This is the general definition of ADS [16,28–31].

IV. RESULTS

From the two-photon Rabi frequency $\Omega(n_1, n_2)$ given by Eq.(10), we find that one of its terms, $\chi(\mu, n_1, n_2)$ given by Eq.(9), would not vanish for general two-mode field states due to the arbitrariness of photon number n_1 and n_2 which are presented in the term $\chi(\mu, n_1, n_2)$. Hence, in this sense, we cannot ignore the effect of the Stark shift in the two-mode system. Of course, it is possible to eliminate or weaken this effect. For example, if we take $\mu = 1$, then we have

$$\chi(\mu = 1, n_1, n_2) = \Delta + n_1 - n_2 + 1.$$
(20)

It is obvious that if there exists some kinds of correlation between the two-mode fields, for example, the difference between their photon numbers is fixed at a constant, i.e., $n_1 - n_2 = q$ (q, an invariant integer), then the term χ would be fixed at $\chi = \Delta + q + 1$ which is intensity independent. Moreover, if we choose the case of $q = -1 - \Delta$ (assuming that the scaled detuning Δ takes an integer value), we see that the term χ would vanish. This is only true for physical quantities which are not sensitive to the phase [18]. For phase-sensitive quantities, the Stark shift still plays an important role in determining their dynamical behaviours. We shall clarify this point in our following results.

In the two-mode system, we assume that the two-level atom is initially prepared in a coherent superposition of its ground state $|-\rangle$ and excited state $|+\rangle$ [31]

$$|atom\rangle = c_{-} |-\rangle + c_{+} |+\rangle \tag{21}$$

and that the two-mode field is initially in a two-mode PCS given by [8]

$$|field\rangle \equiv \sum_{n=0}^{\infty} (1 - |\xi|^2)^{(1+q)/2} \left[\frac{(n+q)!}{n!q!} \right]^{1/2} \xi^n |n+q\rangle_1 |n\rangle_2,$$
(22)

where $\xi = -\tanh(\theta/2) \exp(-i\phi) \equiv |\xi| \exp(i\Phi_2)$ with $0 < |\xi| < 1$ and the phase Φ_2 , and the number q (a positive integer or zero) is the eigenvalue of $|a_1^{\dagger}a_1 - a_2^{\dagger}a_2|$. Using Eqs.(14), we arrive at the expectation values of atomic inversion S_z , the quadrature components S_x and S_y of atomic dipole

$$\langle S_{z}(t) \rangle = \frac{1}{2} \sum_{n=0}^{\infty} P(n,q) \left\{ |c_{+}|^{2} \left[\cos^{2}[\nu(n,q)\tau] + \left(\frac{V^{2}(n,q)}{4} - U^{2}(n,q) \right) \sin^{2}[\nu(n,q)\tau] \right] - |c_{-}|^{2} \left[\cos^{2}[\nu(n-1,q)\tau] + \left(\frac{V^{2}(n-1,q)}{4} - U^{2}(n-1,q) \right) \sin^{2}[\nu(n-1,q)\tau] \right] - 4 |c_{+}|| c_{-} |\Pi(n,q)U(n,q) \times \left(\cos[\nu(n,q)\tau] \sin(\Phi_{1} - \Phi_{2}) - \frac{V(n,q)}{2} \sin[\nu(n,q)\tau] \cos(\Phi_{1} - \Phi_{2}) \right) \sin[\nu(n,q)\tau] \right\},$$
(23)

$$\langle S_{x}(\tau) \rangle = \sum_{n=0}^{\infty} P(n,q) \left\{ | c_{+} |^{2} \Pi(n,q)U(n,q) \sin[\nu(n,q)\tau] \times \left(\frac{V(n+1,q)}{2} \sin[\nu(n+1,q)\tau] \cos(\Phi_{\tau}-\Phi_{2}) + \cos[\nu(n+1,q)\tau] \sin(\Phi_{\tau}-\Phi_{2}) \right) \right. \\ \left. - | c_{-} |^{2} \Pi(n,q)U(n,q) \sin[\nu(n,q)\tau] \times \left(\frac{V(n-1,q)}{2} \sin[\nu(n-1,q)\tau] \cos(\Phi_{\tau}-\Phi_{2}) + \cos[\nu(n-1,q)\tau] \sin(\Phi_{\tau}-\Phi_{2}) \right) \right. \\ \left. + | c_{+} | | c_{-} | [U(n,q)U(n+1,q)\Pi(n,q)\Pi(n+1,q) \times \\ \left. \sin[\nu(n,q)\tau] \sin[\nu(n+1,q)\tau] \cos(\Phi_{\tau}+\Phi_{1}-2\Phi_{2}) + \cos(\Phi_{\tau}-\Phi_{1}) \times \left(\cos[\nu(n-1,q)\tau] \sin[\nu(n,q)\tau] - \frac{V(n,q)V(n-1,q)}{4} \sin[\nu(n-1,q)\tau] \sin[\nu(n,q)\tau] \right) \right. \\ \left. - \sin(\Phi_{\tau}-\Phi_{1}) \left(\frac{V(n,q)}{2} \cos[\nu(n-1,q)\tau] \sin[\nu(n,q)\tau] \right. \\ \left. + \frac{V(n-1,q)}{2} \cos[\nu(n,q)\tau] \sin[\nu(n-1,q)\tau] \right) \right] \right\},$$

$$\begin{split} \langle S_{y}(\tau) \rangle &= \sum_{n=0}^{\infty} P(n,q) \left\{ \mid c_{+} \mid^{2} \Pi(n,q) U(n,q) \sin[\nu(n,q)\tau] \times \\ & \left(\frac{V(n+1,q)}{2} \sin[\nu(n+1,q)\tau] \sin(\Phi_{\tau} - \Phi_{2}) - \cos[\nu(n+1,q)\tau] \cos(\Phi_{\tau} - \Phi_{2}) \right) \\ & - \mid c_{-} \mid^{2} \Pi(n,q) U(n,q) \sin[\nu(n,q)\tau] \times \\ & \left(\frac{V(n-1,q)}{2} \sin[\nu(n-1,q)\tau] \sin(\Phi_{\tau} - \Phi_{2}) - \cos[\nu(n-1,q)\tau] \cos(\Phi_{\tau} - \Phi_{2}) \right) \\ & + \mid c_{+} \mid\mid c_{-} \mid [U(n,q) U(n+1,q) \Pi(n,q) \Pi(n+1,q) \times \\ & \sin[\nu(n,q)\tau] \sin[\nu(n+1,q)\tau] \sin(\Phi_{\tau} + \Phi_{1} - 2\Phi_{2}) + \sin(\Phi_{\tau} - \Phi_{1}) \times \\ & \left(\cos[\nu(n-1,q)\tau] \cos[\nu(n,q)\tau] - \frac{V(n,q)V(n-1,q)}{4} \sin[\nu(n-1,q)\tau] \sin[\nu(n,q)\tau] \right) \\ & + \cos(\Phi_{\tau} - \Phi_{1}) \left(\frac{V(n,q)}{2} \cos[\nu(n-1,q)\tau] \sin[\nu(n,q)\tau] \end{split}$$

$$+ \frac{V(n-1,q)}{2} \cos[\nu(n,q)\tau] \sin[\nu(n-1,q)\tau] \bigg) \bigg] \bigg\}$$

$$(25)$$

with

$$U(n,q) = \sqrt{(n+1)(n+q+1)}/\nu(n,q),$$
(26)

$$V(n,q) = \chi(\mu, n_1 = n + q, n_2 = n) / \nu(n,q),$$
(27)

$$\nu(n,q) = \Gamma(n_1 = n + q, n_2 = n),$$
(28)

$$\Pi(n,q) = |\xi| \sqrt{(n+q+1)/(n+1)}$$
(29)

$$\Phi_{\tau} = \begin{cases}
(\mu + 1)\tau/(2\sqrt{\mu}) & \mu \neq 0, \\
0 & \mu = 0,
\end{cases}$$
(30)

$$P(n,q) = [1 - |\xi|^2]^{q+1} |\xi|^{2n} (n+q)! / [n!q!]$$
(31)

where $\tau = \epsilon t$ is the scaled interaction time, Φ_1 is the initial relative phase between the ground and excited states of the two-level atom, and Φ_{τ} is the induced phase due to the effect of Stark shift in the two-mode system.

Obviously, if we choose $\mu = 1$ and $q = -1 - \Delta$, then the term χ in the atomic inversion S_z given by Eq.(23) vanishs. Thus the effect of Stark shift on S_z can be totally eliminated. However, for phase-sensitive quantities, for example, the two components S_x and S_y of the atomic dipole given by Eq.(24) and (25), respectively, we see that the Stark-shift-induced phase Φ_τ could not vanish with the time evolution of the two-mode system, i.e., the effect of Stark shift on S_x or S_y cannot be eliminated. As an example, Fig.1(a-c) show the time evolution of S_z and S_y for the case of q = 1 when the two-level atom is initially prepared in its excited state. Comparing Fig.1(b) with Fig.1(a), one can see the distinguished effect of the Stark shift on the dynamical behaviours of S_z and S_y . Moreover, comparing Fig.1(c) with Fig.1(a), one can find that for the case of $\Delta = -2$ and $\mu = 1$ the effect of Stark shift on S_z is totally eliminated, but that on S_y still exists. Hence, we may conclude that the presence of Stark shift could bring phase information into the quadrature components S_x and S_y , but not to the atomic inversion S_z . Thus, we would expect that there is a significant effect of Stark shift on the squeezing properties of quantum fluctuations in one of the quadrature components of the atomic dipole.

In the following, taking the quantum fluctuations in the quadrature component S_x of the atomic dipole as an example, we investigate the generation of ADS in the two-mode system. Throughout, we set the phase of the parameter ξ at zero, i.e., $\Phi_2 = 0$. First, we examine the resonant case and consider that the Stark shift is absent. The numerical results are shown in Fig.2-4.

Assuming that the atom is initially prepared in a coherent superposition of its ground and excited states, as shown in Fig.2(a-c), we see that regular and periodical ADS patterns are observed for the cases of $|\xi| = 0.1$ and q = 1, 4, 8. As the value of the parameter $|\xi|$ is increased to 0.5, we find that the regular and periodical ADS pattern is destroyed greatly. Moreover, comparing Fig.2(b)(c) with Fig.2(a), we find that as the number q varies from 1 to 8, the squeezing duration is decreased gradually and the squeezing period is also shortened. In Fig.2(d), we display the relation between the function F_1 and the number q at a fixed interaction time $\tau = 0.8\pi$ for the case of $|\xi| = 0.1$. It is found that the squeeze depth (strength) is decreased as the number q varies from 1 to 200 and ADS can be shown only for q < 30. This implies that the stronger the correlations between the modes are involved, the more the atomic dipole squeezing could be generated.

Next, we examine the effect of atomic coherence on ADS exhibited in the two-mode system. Holding the phase Φ_1 at zero and the interaction time τ at 0.8π , we show in Fig.3 the relation between F_1 and the coefficient $|c_-|^2$ for $\xi = 0.1$ as the number q varies from 1 to 8. For q = 1, it is seen that ADS can be generated in a wide squeezing range of the parameter $|c_-|^2$ (i.e., $0.5 < |c_-|^2 < 1$). However, increasing the number q would narrow down the squeezing range, and ADS depth is decreased greatly.

Then, we turn to examine the phase dependence of ADS generated in the two-mode system. For $\xi = 0.1$, $\tau = 0.8\pi$ and $|c_{-}|^2 = 0.8$, Fig.4 presents the relation between F_1 and the relative phase Φ_1 as the number q varies from 1 to 8. We find that in the case of q = 1 ADS can be exhibited in a wide squeezing range of Φ_1 around $0, \pi, 2\pi$. However, the squeezing range is narrowed down as the number q increases.

Finally, we consider the nonresonant case and examine in detail the effects of detuning and/or Stark shift on ADS. The numerical results are shown in Fig.5(a-d) for q=1, $|\xi| = 0.1$ and different choices of parameter pair Δ and μ . Comparing Fig.5(b)(d) with Fig. 5(a), we find that the detuning or Stark shift has a destructive effect on the ADS pattern exhibited in Fig.5(a). The squeezing duration is decreased and the ADS pattern is irregular and nonperiodic. For the case of $\Delta = -2$ and $\mu = 1$, as discussed above, the effects of Stark shift on S_z can be eliminated totally but those on S_x and S_y cannot. Hence, we would expect a significant effect of the Stark shift on ADS. As shown in Fig.5(c), we really observe a regular, periodic and strong ADS pattern which is generated due to the combined effect of detuning and Stark shift. Compared to Fig.5(a), the squeezing duration in Fig.5(c) is decreased due to the combined effect of detuning and Stark shift.

V. CONCLUSION AND DISCUSSION

In this paper, we have considered a two-mode system that consists of a two-level atom interacting with a quantized electromagnetic field, which is initially prepared in an entangled two-mode PCS, via a nondegenerate two-photon process in a lossless cavity. We have investigated the generation of ADS in this system and examine in detail the effects of detuning, Stark shift and atomic coherence on ADS. We show that ADS strongly depends on the atomic coherence. It is found that the stronger the correlations between the modes are involved, the more the ADS can be generated. The detuning or Stark shift has a destructive effect on ADS. However, the combined effect of the detuning and Stark shift may lead to periodical, regular and strong ADS pattern but with decreased squeeze duration.

Gou [20] have shown that the correlation is responsible for generating the squeezing of the cavity fields. Our prediction of ADS in the two-mode system agrees well with this argument. In addition, it is obvious that two-mode squeezed vacuum [8] (i.e., the special case of the two-mode PCS for q = 0) is a strongly correlated quantum state since in the number-state expansion there are always an equal number of photons in the a_1 and a_2 modes. Hence,

a strong ADS pattern in the two-mode system would be expected if the field is initially prepared in a two-mode squeezed vacuum. This point can be clearly seen from Fig.6, where we compare the ADS pattern of q = 0 with that of q = 1.

Actually, in this paper, our discussion is restricted to an ideal case, i.e., without considering the cavity damping in the two-mode system. In the realistic situation, a high but finite cavity Q indeed causes some inevitable effects of photon dissipation in the cavity. Hence, it would be significant and interesting to include the cavity damping in the two-mode system and examine the effect of cavity damping on ADS predicted in this paper.

Finally, it should be mentioned that there are several significant experimental schemes which involve the nonresonant interaction of the atom with two nondegenerate cavity modes, for example, those illustrated in the works of Gou [20] and Abdalla *et al.* [40], and the recent one realized by Haroche's group [41] with a single circular Rydberg atom in a cavity. Certainly, these schemes could be helpful for realizing the two-mode system introduced in section II and for further understanding the ADS predicted in this paper.

ACKNOWLEDGEMENT

One of us (Q.R) thanks the Queen's University for a Reinhart Fellowship.

REFERENCES

- Y. S. Kim, M. A. Man'ko and A. Sergienko (eds.), J. Opt. B: Quantum Semiclass. Opt. 4, S91 (2002), one special issue on Uncertainty relations, quantum phase space, quantum optics, quantum information, imaging and computing.
- [2] P. Zanardi, Phys. Rev. A **65**, 042101 (2002).
- [3] T. Jennewein, C. Simon, G. Weihs, H. Weinfurter, and A. Zeilinger, Phys. Rev. Lett. 84, 4729 (2000).
- [4] A. Furusawa, J. Sorensen, S.L. Braunstein, C.A. Fuchs, H.J. Kimble, and E.S. Polzik, Science 282, 706 (1998).
- [5] L. K. Grover, Phys. Rev. Lett. **79**, 325 (1997).
- [6] X.H. Wu, R.H. Xie, X.D. Huang, and Y.F. Hsia, Phys. Rev. A 53, R1927 (1996).
- [7] C.H. Bennett, Phys. Rev. Lett. **69**, 2881 (1992).
- [8] C.C. Gerry and A. Benmoussa, Phys. Rev. A 62, 033812 (2000).
- [9] R.R. Puri, Mathematical Methods of Quantum Optics (Springer-Verlag, Berlin, 2001).
- [10] A.M. Perelomov, Commun. Math. Phys. 26, 222 (1972).
- [11] C.C. Gerry, J. Opt. Soc. Am. B 8, 685 (1991).
- [12] A.O. Barut and L. Girardello, Commun. Math. Phys. 21, 41 (1971).
- [13] C.C. Gerry and R. Grobe, Phys. Rev. A 51, 4123 (1995).
- [14] C.C. Gerry, J. Mod. Opt. **42**, 585 (1995).
- [15] M.D. Reid and D.F. Walls, Phys. Rev. A. **34**, 1260 (1986).
- [16] D.F. Walls and P. Zoller, Phys. Rev. Lett. 47, 709 (1981).
- [17] P. Alsing and M.S. Zubairy, J. Opt. Soc. Am. B 4, 177 (1987).

- [18] S.C. Gou, Phys. Lett. A **147**, 218 (1990).
- [19] H.I. Yoo and J.H. Eberly, Phys. Rep. **118**, 239 (1985).
- [20] S. C. Gou, Phys. Rev. A 40, 5116 (1989).
- [21] R.H. Xie, Phys. Rev. A **65**, 055801 (2002).
- [22] R.H. Xie and Q. Rao, Physica A **312**, 421 (2002).
- [23] R.H. Xie and Q. Rao, Phys. Lett. A **302**, 28 (2002).
- [24] J. Kim, S. Somani, and Y. Yamamoto, Nonclassical Light from Semiconductor Lasers and LEDs (Springer-Verlag, Berlin, 2001).
- [25] Q. Rao and R.H. Xie, Physica A **280**, 362 (2000).
- [26] R.H. Xie, Phys. Rev. A 65, 055801 (2002).
- [27] R.H. Xie and S. Yu, J. Opt. B: Quantum Semiclass. Opt. 4, 172 (2002).
- [28] R.H. Xie and Q. Rao, Physica A **315**, 386 (2002).
- [29] R.H. Xie and Q. Rao, Physica A **315**, 411 (2002).
- [30] R.H. Xie and Q. Rao, Physica A **315**, 427 (2002).
- [31] R.H. Xie, G.O. Xu, and D.H. Liu, Phys. Lett. A **202**, 28 (1995).
- [32] K. Wodkiewicz, P. L. Knight, S. J. Buckle, and S. M. Barnett, Phys. Rev. A 35, 2567 (1987).
- [33] P.K. Aravind and G. Hu, Physica C **150**, 427 (1988).
- [34] G. Hu and P.K. Aravind, J. Opt. Soc. Am. B 6, 1757 (1989).
- [35] R.H. Xie, Phys. Rev. A 53, 2897 (1996).
- [36] R.H. Xie and G.O. Xu, Phys. Rev. E 54, 1402 (1996).

- [37] A.B. Bhattacherjee, Phys. Rev. A 56, 796 (1997).
- [38] R.H. Xie, Phys. Rev. A **62**, 017801 (2001).
- [39] N. B. Narozhny, J. J. Sanchez-Mondragon, and J. H. Eberly, Phys. Rev. A. 23, 236 (1981).
- [40] M.S. Abdalla, M.M.A. Ahmed, and A.S.F. Obada, Physica A 170, 393 (1991).
- [41] P. Bertet, S. Osnaghi, P. Milman, A. Auffeves, P. Maioli, M. Brune, J.M. Raimond, and S. Haroche, Phys. Rev. Lett. 88, 143601 (2002).

CAPTION OF FIGURES

FIG.1: Time evolution of S_z and S_y for q = 1, $|\xi| = 0.9$, $\Phi_2 = 0$, $|c_-|^2 = 0$ and (a) $\Delta = 0, \mu = 0$; (b) $\Delta = 0, \mu = 1$; (c) $\Delta = -2, \mu = 1$.

FIG.2: Time evolution of F_1 for $\Delta = 0$, $\mu = 0$, $|c_-|^2 = 0.8$ and $\Phi_1 = 0$ when the two-mode field is in a two-mode PCS with (a) q=1, (b) q=4 and (c) q=8, where the solid and dot-dashed lines are for $|\xi| = 0.1, 0.5$, respectively. Case (d) is the relation between F_1 and q for $|\xi| = 0.1$ at a fixed time $\tau = 0.8\pi$.

FIG.3: Relation between F_1 and coefficient $|c_-|^2$ for $\Delta = 0$, $\mu = 0$, $|\xi| = 0.1$ and $\Phi_1 = 0$ at a fixed time $\tau = 0.8\pi$, where the solid, dashed and dot-dashed lines are for q = 1, 4, 8, respectively.

FIG.4: Relation between F_1 and phase Φ_1 for $\Delta = 0$ and $\mu = 0$, $|\xi| = 0.1$ and $|c_-|^2 = 0.8$ at a fixed time $\tau = 0.8\pi$, where the solid, dashed and dot-dashed lines are for q = 1, 4, 8, respectively.

FIG.5: Time evolution of F_1 for $|\xi| = 0.1$, $|c_-|^2 = 0.8$ and $\Phi_1 = 0$ when the two-mode field is initially prepared in a two-mode PCS with q=1: (a) $\Delta = 0$, $\mu = 0$; (b) $\Delta = 0$, $\mu = 1$; (c) $\Delta = -2$, $\mu = 1$; (d) $\Delta = -2$, $\mu = 0$.

FIG.6: Time evolution of F_1 for $\Delta = 0$, $\mu = 0$, $|c_-|^2 = 0.8$, $\Phi_1 = 0$ and $|\xi| = 0.1$ when the two-mode field is in a two-mode PCS, where the dot-dashed and solid lines are for q = 0, 1, respectively.

Rao & Xie: FIG.1/PHYSA



Rao & Xie: FIG.2/PHYSA



Rao & Xie: FIG.3/PHYSA



Rao & Xie: FIG.4/PHYSA



Rao & Xie: FIG.5/PHYSA



Rao & Xie: FIG.6/PHYSA

