

The Two-Component Model and Metallization of Van der Waals Crystals

V. N. Bogomolov

*A. F. Ioffe Physical & Technical Institute,
Russian Academy of Science,
194021 St. Petersburg, Russia**

(Dated: November 21, 2018)

The paper discusses a model of Van der Waals crystals in which band-gap structures do not form. An effect of strong and chaotic electron-electron repulsion, which was excluded from consideration in the traditional approach, is taken into account. A condensate exists as a result of a dynamic equilibrium among atoms acted upon by constant Van der Waals forces and periodically forming and disappearing covalent bonding. One part of atoms is, on the average, in the ground, and the other, in excited state, to form diatomic virtual molecules. Treated in terms of this pseudoclassical model, the interatomic distances, binding energies, volumes, and pressures at which metallization, for instance, of inert gases and hydrogen, sets in is described by simple relations involving only two spectroscopic parameters of atoms (molecules). Applying pressure to a VdW crystals transfers it from the insulator first to a Bose superconductor, and after that, to a Fermi metal. An empirical relation $T_c \sim N^{2/3}$ between the superconductivity transition temperature T_c and the particles concentration N in chalcogens under pressure is considered as an example of such situation.

PACS numbers:

The problem of metallization of Van der Waals (VdW) condensates is one of the oldest in condensed matter physics. It originated in XVIII century from the assumption that hydrogen can undergo metallization in transformation to a solid [1]. One of the latest attempts at obtaining metallic hydrogen experimentally and estimating the necessary pressures was undertaken in [2].

The properties of VdW condensates are dealt with in a large number of theoretical studies based on traditional band-gap concepts. Some of the complicated problems generated in this approach still remain unsolved [3]. One of the reasons that could account for these difficulties lies in that VdW condensates are possibly not band-gap materials at all. One may, however, conceive of a model of the condensate in which atomic interactions, rather than being constant, appear and disappear periodically between pairs of atoms over the whole lattice, i.e., they are localized, both in time and in space. The validity of this model is argued for by the experimental support obtained by the Herzfeld criterion, which defines the VdW condensate volume at metallization in a straightforward way through atomic properties as the atomic or molar refractivity (the molar volume should become equal to the gas-phase molar refractivity) [4]. This supportive evidence was obtained from xenon metallization ($V_m = 10.2\text{cm}^3/\text{mol}$, $P_m = 160\text{ GPa}$) [5].

This VdW condensate model with a flickering, pulsating coupling between neighboring atoms, which originates statistically throughout the lattice, was used to interpret the results of the metallization of Xe as evidencing its transition to a superconducting state with a very high $T_c > 300\text{ K}$ [6, 7]. In [5], Xe metallization (from optical characteristics) was treated as resulting from band clo-

sure and a transition to a conventional metal. The possibility that the "metallization" of the VdW condensates could be accompanied by the onset of superconductivity at a superhigh-temperature did not become a subject of serious consideration because of the traditional belief that VdW condensates are typical band-gap substances.

Due to its inherent simplicity, the VdW condensate model proposed in [6, 7] not only is in accord with Herzfeld's approach, but yields, even in a very rough approximation, as simple expressions for the condensate binding energies q and the critical volumes V_m and pressures P_m at metallization (through the equation of state [8]). The only things one needs here are two spectroscopic parameters of atoms or molecules, namely, the ionization potential E_0 and the excitation energy ΔE :

$$q = C(E_0 - \Delta E) \exp\{-1.73/[(E_0/\Delta E) - 1]\};$$

$$P_m = KE_0^4; \quad V_m = F/\Delta E^3.$$

The parameters $C = 1.44$, $K = 0.0074$, $F = 5880$ for the inert gases were derived from data available for Xe. For helium He and hydrogen H_2 , which have only one electronic shell, the parameter $C = 1.44/8$. Using the known values of E_0 and ΔE (in eV) for these gases

He (24.8 ;19.8); Ne (21.56; 16.6); Ar (15.76; 11.6);
Kr (14.02; 9.92); Xe (12.13; 8.32); H_2 (15.43; 11.2),

one obtains the following ratios of the calculated to experimental values q/q_e :

(a) He (0.00098/0.00098); Ne (0.021/0.018);
Ar (0.048/0.063); Kr (0.089/0.090);
Xe (0.126/0.126); H_2 (0.0080/0.0093),

and the volumes at metallization (cm^3/mol), V_m , relative to the values by Herzfeld, V_{mH} :

*Electronic address: V.Bogomolov@mail.ioffe.rssi.ru

(b) He (0.76/0.50); Ne (1.28/1.00); Ar (3.77/4.16);
Kr (6.02/6.22); Xe (10.2/10.2); H₂(4.2/2.0),

and the metallization pressures P_m (GPa):

(c) He (2800); Ne (1600); Ar (457);
Kr (286); Xe (160); H₂(420).

The metallization pressure of hydrogen is close to 450 GPa, the estimate obtained in [2]. In the model under discussion, the VdW condensate lattice exists due to a dynamic equilibrium between two virtual "phases". One of them, the "ground-state phase", comprises atoms in the ground state, which are "slowly" brought together by the constant forces of Van der Waals attraction. The band-gap properties are not efficient (in accordance with the Mott criterion) on account of the large interatomic distances, because the atomic electron shells are filled. The principal quantum numbers of the first free and last filled electronic levels in an atom differ by one, and the diameters of the orbitals differ by 2–3 Å. The other "phase" is actually a gas of diatomic virtual covalent-bonded molecules, which form and break up in a random manner within the ground-state phase of virtually excited atoms (excimers). Because of disorder and short lifetimes of the virtual states, band-gap structures do not form in this phase either. The lattice is maintained in equilibrium due to particles transferring continually from one phase to another. The fluctuations in electronic system $\sim (E_0 - \Delta E)$ eV are followed by the lattice vibrations (mean energy about $(E_0 - \Delta E)m_e/M_a \sim 0.7$ meV for He). A new type of localized excitations (~ 1 meV) were observed in solid helium [9].

This pseudoclassical pattern reminds one of a garland of balls suspended in air, which exists due to periodic pulses received by the balls from the juggler's hands. Ideas that could be treated as supporting such a VdW condensate scheme appeared after Ref.[4] as well. These are, for instance, the chemical bond resonance theory of Pauling and detection of "virtual Xe₂ molecules" [10] in solid Xe. The enigmatical high mobility of hydrogen atoms in palladium may be a result of resemblance the electronic structures of Pd($4d^{10}5s^0$) and atoms of inert gases and existence of the ground state atom sublattice in ordinary space.

The "juggler hand" mechanism producing virtual molecules is also very simple. The wave functions of atoms in the first phase are not coherent (there are no band gap structures). Therefore, the strong and random electron-electron interaction of average energy $\sim e^2/D$ between atoms is similar to the thermal vibration field in conventional condensates, where the excitation probability of a single impurity $p \sim \exp(-\Delta E/kT)$. Equating this energy to the average thermal vibration energy $3kT/2 = e^2/D$ yields for the probability of atom excitation in the first phase

$$X \sim \exp[-(1.5\Delta ED/e^2)], \quad (\Delta ED) = f(P).$$

This is the probability for an atom of the first phase to

transfer to the second phase, or the relative particle concentration in the second phase. The interatomic distance D can be expressed through the hydrogen-like diameter of an excited atom, with due account of its increase (1.15) in the condensate:

$$D = 1.15 e^2/(E_0 - \Delta E).$$

The diameter of atom in ground state is $d_0 = e^2/E_0$ and the Mott relation $D > 2d_0$ ($0.5 > 1 - (\Delta E/E_0)$) is valid for inert gases and H₂.

Assuming such excitations to form in pairs to produce virtual molecules, we obtain for the number of covalent molecules $\sim X \ll 1$, and for the number of atoms in the ground state and with a weak Van der Waals attraction, $\sim (1 - X)$. Therefore the average binding energy of a VdW condensate $q \sim X e^2/D$.

It is these simple expressions that underlie the above relations (at metallization X_m and $\Delta E_m D_m$ are const.), which describe in a consistent way the VdW condensate properties. One can derive in a similar manner a relation for the compressibility and the equation of state [8], which is in agreement with the results reported in [5]:

$$\begin{aligned} -[dV/VdP] &\sim V/q; \\ P &\sim E_0^4 \int y^{-7/3} \{\exp[-1.5(y^{1/3} - 1)]\} dy; \\ y = V/v; \quad V^{1/3} &\sim D; \quad v^{1/3} \sim e^2/E_0. \end{aligned}$$

Within this VdW condensate model, the only cooperative effect, as in the band model, is the establishment of an average interatomic distance through the inertia of atoms, that vibrate as a result of periodically forming and breaking-up covalent molecules at fixed separations and of the Van der Waals attraction. And it is only at "metallization" ($X_m \sim 1$) that cooperative effects of the electronic type become possible. It was suggested that the VdW "metallization" is actually superconductivity with a very high T_c [6, 7]. An electron pair in a virtual molecule, for instance, in Xe₂, may be considered as a tight-bound boson (~ 0.5 eV), a Cooper pair two lattice constants in size. Its bonding is realized through a "two-atom lattice". At the VdW condensate metallization, the concentration of such "molecular" small radius bosons in a gas reaches the Bose condensation point. There is no problem of boson formation or decay at real temperatures, as is the case with metals. We are aware of only one attempt at detecting superconductivity of Xe from conductivity characteristics [11]. In this experiment, however, one could not raise the pressure above 155 GPa. The change in the color of H₂ under compression [2], as well as in the compression of Xe at $P < P_m$ [5], could be due not to band closure, but rather to the absorption of light by particles of the forming metallic or Bose phase [6, 7, 12]. The diamagnetic "regions" (undetectable in conductivity [11]) were observed in HTSC materials at $T > T_c$ [13]. It is well known that the color of systems with conducting particles depends on both the particle size and their concentration (recall, for instance, the coloring of metal sols).

It may be conceived that an electron pair in a divalent atom ("atomic" boson) embedded in a metal, rather than breaking down completely because of the delocalization of its electrons, as is the case with electrons in univalent atoms, remains a "pseudomolecule", albeit very weakly bound ($E \ll E_0$). At $T = 0$, these "atomic" bosons, which are not completely "dissolved" in the metal (the Cooper pairs), form a Bose condensate to give rise to superconductivity, which usually does not set in in the case of univalent atoms. The BCS theory provides an explanation for what prevents the electrons of such "relic" bosons in metals from complete delocalization. For tight-bound "molecular", noninteracting bosons in a VdW condensate or for "atomic" ("relic") bosons (or "relic" atomic wave functions) in a dense gas of divalent atoms (Mg, for example), such danger apparently does not exist (before metallization).

Divalent atoms or diatomic molecules of univalent atoms dissolved in dielectric media, are diamagnetic. For instance, it is a Na_2 molecules in NH_3 (but not an "individual electrons self-trapped in physical cavities in solvent medium") [14]. A volume per atom in Na_2 molecule is larger of that in metallic Na.

Despite the simplicity of the structure of the VdW condensates, their most intriguing property, namely, the possibility of providing superhigh-temperature superconductivity (with the atoms being nonmagnetic) remains a mystery. There is, an experimental problem of increasing the boson concentration either through an increase in pressure or by physicochemical means [14]. Metallic $\text{Pd}(4d^{10};9s^{0;1})$ may represent roughly a final state of the VdW condensate under pressure. It is a paramagnetic ($+540 \times 10^{-6} \text{ cm}^3/\text{mol}$). Nevertheless, a compounds $\text{PdH}_{0.6}$, PdH and $(\text{Pd}_{0.7} \text{Ag}_{0.3})\text{H}_{0.8}$ are diamagnetic and superconductors (9 K and 16 K) respectively [15]. {tolerance=400 Applying pressure to a VdW condensate (a gas of noninteracting bosons) transfers it

from the insulator first to a Bose superconductor, and after that, to a Fermi metal [8, 14?] An example of such situation is, perhaps, behaviour of T_c in chalcogens under pressure [17]. The phase transition ($S_8 \rightarrow S_6$) in sulphur (close to insulator) is followed by increase in concentration N of molecules

$$(N_6 \sim N_8 \times (8/6)^2) \text{ and by increase in } T_c$$

$$T_{c6}/T_{c8} = 17.3/12.5 = 1.38 \sim (N_6/N_8)^{2/3} = 1.46).$$

Under pressure the molecular structures of Se and Te (close to a Fermi metal) disappears and the phase transition $(\text{Se}; \text{Te})_6 \rightarrow (\text{Se}; \text{Te})_1$ is followed by increase in concentration of particles ($N_1 \sim 6N_6$) and by increase in T_c :

$$\text{Se} : T_{c1}/T_{c6} = 10.3/3.4 = 3.0;$$

$$\text{Te} : T_{c1}/T_{c6} = 7.5/2.5 = 3.0. \quad \text{It is close to relation}$$

$$T_{c1}/T_{c6} \sim (N_1/N_6)^{2/3} \sim 6^{2/3} = 3.3 \quad \text{and do not}$$

contradict to the Bose condensation mechanism of superconductivity. Relatively low T_c may be a result of the virtual molecules paramagnetism.

Besides the hydrogen and inert gases, there is a large number of VdW condensates with a relatively low P_m and, perhaps, with very high T_c superconductivity [6, 7].

The VdW condensate is an example of an intermediate state of matter between a gas and a solid, between micro- and macroobjects with combined electron systems. The interactions in system are below a threshold value and band-gap structures (coherent state) do not form. It is the origin of difference between the chemical substances and the sorption compounds [18]. Such materials can be constructed of atoms with either filled or unfilled electron shells. Properties (a), (b), and (c) can be readily explained in terms of both the model under discussion and the band approximations. Only the concepts of VdW condensate "metallization" are quite different.

-
- [1] K. Mendelsson, in *The Quest for Absolute Zero*, (World University Library, Weidenfeld and Nicolson, 1968), 110.
- [2] P. Loubeyre, F.Occelli, R.Letoullec, *Nature* **416**, 613 (2002).
- [3] K.A. Johnson, N.W. Ashcroft, *Nature* **403**, 632 (2000).
- [4] K.F. Herzfeld, *Phys Rev.* **29**, 701 (1927).
- [5] E.A Goettel, J.H Eggert, I.F Silvera, W.C. Moss, *Phys. Rev. Lett.* **62**, 665 (1989).
- [6] V.N. Bogomolov, *Metallic Xenon. Conductivity or Superconductivity ?* Preprint 1734 Russian Academy of Sciences. A.F.Ioffe Physical Technical Institute (1999); e-print cond-mat/9902353.
- [7] V.N.Bogomolov, *Techn. Phys. Lett.***28**, 211 (2002).
- [8] V.N. Bogomolov, e-print cond-mat/9912034.
- [9] T.Markovich, E.Polturak, J.Bossy, E.Farhi, e-print cond-mat/0203086.
- [10] R.A.Tilton, C.P.Flynn, *Phys. Rev. Lett.* **34**, 20 (1975).
- [11] M.I.Eremets, E.A.Gregoryanz, V.V.Struzhkin, H.Mao, R.J.Hemley, N.Mulders, N.M.Zimmerman, *Phys. Rev. Lett.* **85**, 2797 (2000).
- [12] R.Braut, *Phase Transitions*, Univ. of Brusseles, N.Y.-Amsterdam, 251 (1965).
- [13] I. Iguchi, T.Yamaguchi, A.Sugimoto, *Nature* **412**, 420 (2001).
- [14] R.A. Ogg, Jr., *Phys. Rev.* **69**, 668 (1946).
- [15] G.Heim, B.Stritzker, *Appl. Phys.* **7**, 239 (1975).
- [16] M.Capone, M.Fabrizio, E.Tosatti, e-print cond-mat/0101402.
- [17] E.A.Gregoryanz, V.V.Struzhkin, R.J.Hemley, M.I.Eremets, H.Mao, Y.A.Timofeev, e-print cond-mat/0108267.
- [18] V.N. Bogomolov, *Phys. Rev.* **51**, 17040 (1995).