

Gauge Fields, Geometric Phases, and Quantum Adiabatic Pumps

Huan-Qiang Zhou^{1*}, Sam Young Cho², and Ross H. McKenzie^{1,2}

¹*Centre for Mathematical Physics, The University of Queensland, 4072, Australia and*

²*Department of Physics, The University of Queensland, 4072, Australia*

(Dated: April 26, 2019)

Quantum adiabatic pumping of charge and spin between two reservoirs (leads) has recently been demonstrated in nanoscale electronic devices. Pumping occurs when two or more system parameters are varied in a cyclic manner and sufficiently slowly that the quantum system always remains in its ground state. We show that quantum pumping has a natural geometric representation in terms of gauge fields (both Abelian and non-Abelian) defined on the space of system parameters. We make explicit the similarities and differences with Berry's geometric phase. Tunneling from a scanning tunneling microscope tip through a magnetic atom could be used to demonstrate the non-Abelian character of the gauge field.

PACS numbers: 03.65.Vf, 73.23.-b, 73.63.-b

Normally transport of electrical charge is dissipative (i.e., it produces heat). However, quantum adiabatic pumping [1] provides a means in nanoscale electronic devices to use novel quantum effects to transport single electrons with minimal dissipation [2]. Furthermore, it is also possible to pump electron spin without pumping charge [3, 4]. Both charge[5] and spin [6] pumping have been recently achieved experimentally, by cyclic variation of the gate voltages that control the shape of an open quantum dot. This motivated extensive theoretical research in this topic, especially on quantum charge pumping [7, 8, 9, 10, 11]. Quantum spin pumping opens the way for applications in spintronics. It is sometimes suggested, but not explicitly shown, that quantum pumping is related to Berry's phase. As first emphasized by Berry [12], discrete quantum systems have the counter-intuitive property that when some of the parameters controlling the system are slowly varied and brought back to their initial value the quantum state of the system is different to the initial state. That is, a quantum state may acquire a geometric phase $\exp(i\gamma_C)$ in addition to the normal dynamic phase $\exp(-(i/\hbar) \int E(t)dt)$. This remarkable discovery may be recast in the language of holonomy theory [13]. Subsequent work showed that non-Abelian gauge potentials can arise as a result of degeneracies of energy levels of the system [14, 15, 16]. However, it is unlikely that adiabatic pumping, characteristic of quantum *open* systems, results from Berry's phase for *isolated discrete* systems.

We present a systematic treatment of quantum adiabatic pumping in open systems in terms of parallel transport and gauge fields (both Abelian and non-Abelian) defined on the system parameter space, which reveals a unifying concept of geometric phase underlying scattering states. We make explicit the similarities and differences with Berry's phase associated with cyclic variations of isolated quantum systems (both degenerate and non-degenerate) (see Table 1). In the scattering approach developed by Brouwer [17], based on an earlier work of Büttiker, Thomas, and Prêtre [18], a compact formula was presented for the pumped charge (current) in terms

of the parametric derivatives of the time-dependent scattering matrix subjected to the modulating potential. We show that the pumped charge, given by Brouwer's formula [17], is essentially the geometric phase associated with the $U(1)$ subgroup of the gauge group $U(M)$ (M is the number of channels in a certain lead), whereas the non-Abelian sector $SU(M)$ describes the adiabatic pumping associated with the internal degrees of freedom such as spin. Expressions are given for the gauge potentials associated with tunneling from an STM (scanning tunneling microscope) through a magnetic atom. We suggest an experiment which can be used to illustrate the non-Abelian character of the gauge field.

The quantum system. Consider a mesoscopic system with N leads, and for the n -th lead there are M_n channels. Our aim is to study quantum pumping by periodically varying a set of the independent external parameters $X \equiv (X^1, \dots, X^\nu, \dots, X^p)$ slowly as a function of time t . In the scattering approach, the S matrix is an $\mathcal{N} \times \mathcal{N}$ matrix with $\mathcal{N}$ the total number of channels, $\mathcal{N} = \sum_{n=1}^N M_n$. We define vectors $\mathbf{n}_\alpha \equiv (S_{\alpha 1}, \dots, S_{\alpha \mathcal{N}})$ in terms of the rows of the scattering matrix $S[X(t)]$ associated with the n -th lead. The unitarity of the scattering matrix implies that these vectors are orthonormal

$$\mathbf{n}_\alpha^* \cdot \mathbf{n}_\beta = \delta_{\alpha\beta}, \quad \alpha, \beta = 1, \dots, M_n.$$

That is, this provides us with a smooth set of (local frame) bases $\mathbf{n}_\alpha(t)$.

The gauge potential. Assume that the parallel transport law

$$\Psi_\alpha^* \cdot d\Psi_\beta = 0 \quad (1)$$

holds, where $d\Psi$ is the variation in Ψ resulting from a variation dX in the external parameters. If $\Psi_\alpha(0) = \mathbf{n}_\alpha(0)$, i.e., the initial vector describing the scattering process in which the incident particle comes from the α -th channel in the n -th lead, then the indistinguishability of particles implies that the transported vector $\Psi_\alpha(t)$ must be a linear combination of all $\mathbf{n}_\alpha(t)$, $\Psi_\alpha(t) = \sum_\beta U_{\alpha\beta}(t) \mathbf{n}_\beta(t)$. Expressed another way, the

transported vector describes a combined scattering process in which particles come from all channels in the n -th lead. Obviously, $U(t)$ is unitary, i.e., $U(t) \in U(M_n)$. Physically, this means certain information about where the incident particles come from is lost during parallel transport, and is encoded in the unitary matrix $U(t)$. Inserting into the parallel transport law in Eq. (1), we have

$$(U^{-1}dU)_{\alpha\beta} = -\mathbf{n}_\beta^* \cdot d\mathbf{n}_\alpha. \quad (2)$$

Since $\mathbf{n}_\alpha$ varies as the parameters X^ν vary with time, we can thus define the gauge potential

$$A_{\alpha\beta\nu} \equiv \mathbf{n}_\beta^* \cdot \partial_\nu \mathbf{n}_\alpha, \quad (3)$$

where $\partial_\nu \equiv \partial/\partial X^\nu$ so that

$$(U^{-1}dU)_{\alpha\beta} = -\sum_\nu A_{\alpha\beta\nu} dX^\nu.$$

This can be integrated in terms of exponential integrals,

$$(U(t))_{\alpha\beta} = (\text{P exp}(-\oint A_\nu dX^\nu))_{\alpha\beta}, \quad (4)$$

where P denotes path ordering. Defining $A \equiv A_\nu dX^\nu$, one can see it is Lie algebra $u(M_n)$ valued and thus anti-Hermitian. A plays the role of a gauge potential, as in the case of Berry's phase [14] for closed (discrete) quantum systems.

Gauge transformation. The gauge group $U(M_n)$ originates from the unitary freedom in choosing local bases $\mathbf{n}_\alpha (\alpha = 1, \dots, M_n)$,

$$\mathbf{n}'_\alpha(t) = \sum_\beta \omega_{\alpha\beta}(t) \mathbf{n}_\beta(t).$$

This amounts to different choices of the scattering matrix: $S'(t) = \Omega(t)S(t)$ with $\Omega(t)$ a diagonal block matrix, the n -th block of which is an $M_n \times M_n$ unitary matrix $\omega(t)$. Physically speaking, left multiplication of the scattering matrix $S(t)$ by $\Omega(t)$ just redistributes the scattering particles among different incoming channels associated with a certain lead, which does not affect correlations at the scatterer and so the physics remains the same. The gauge potential $A(t)$ transforms as

$$A'(t) = d\omega\omega^{-1} + \omega A\omega^{-1}. \quad (5)$$

The gauge field strength defined by $F \equiv dA - A \wedge A$, transforms covariantly $F' = \omega F \omega^{-1}$. Therefore, a $U(M_n) \equiv U(1) \times SU(M_n)$ gauge field is defined on a p -dimensional parameter space which drives the quantum pumping. The trace of $U(t)$ given by Eq. (4) is gauge-invariant.

Justification of parallel transport. Now we need to justify the assumption of the parallel transport law in Eq. (1). Physically, by ‘‘adiabatic’’ we mean that the dwell time τ_p during which particles scatter off the scatterer is much shorter than the time period $\tau = 2\pi/\omega_a$

during which the system completes the adiabatic cycle. τ_p is related to (but not determined alone by) Wigner time delay matrix $\tau_w(s, E) \equiv -iS^\dagger(s, E)\partial S(s, E)/\partial E$ with E being the energy of scattering particles and s being the so called *epoch* defined as $s = \omega_a t$ [19]. Here ω_a is a slow frequency characterizing the adiabaticity. Then the response to the variation of the particle distribution in a certain channel is *only limited to* channels associated with the same lead. That means we ignore any responses which involve channels in different leads. Such a response can be treated as dissipation, a correction to the adiabatic limit. Then, the parallel transport law in Eq. (1) follows from the parallel transport law for the wave function. The latter is a solution of the time-dependent Schrödinger equation $i\hbar\partial_t|\Psi(t)\rangle = H(t)|\Psi(t)\rangle$. As is well known, the Schrödinger equation induces a parallel transport law $\text{Im}\langle\phi(t)|\partial_t\phi(t)\rangle = 0$ [20], with $|\phi(t)\rangle \equiv \exp(i\int h(t)dt)|\Psi(t)\rangle$, where $h(t) = \langle\Psi|H|\Psi\rangle/\langle\Psi|\Psi\rangle$. We can write the wave function $|\phi(t)\rangle$ as a linear combination of *all* scattering states associated with a certain lead in the adiabatic case. Formally, $|\phi(t)\rangle = \sum_\alpha c_\alpha|\psi_\alpha(t)\rangle$, with $|\psi_\alpha(t)\rangle$ denoting the scattering states in which the scattered particles come from channels associated with the n -th lead, and c_α being arbitrary constants. Then we have $\langle\psi_\alpha(t)|\partial_t\psi_\beta(t)\rangle = 0$. The adiabatic assumption implies that $|\psi_\alpha(t)\rangle$ may be expanded in terms of instantaneous asymptotic scattering states, $|\psi_\alpha(t)\rangle = \sum_\beta U_{\alpha\beta}(t)|\psi_\beta^S(t)\rangle$, with $|\psi_\alpha^S(t)\rangle = |\alpha\rangle_{in} + \sum_{\beta=1}^N S_{\alpha\beta}(t)|\beta\rangle_{out}$, $\alpha = 1, \dots, M_n$. Here, $|\alpha\rangle_{in}$ and $|\beta\rangle_{out}$ denote, respectively, the incoming and outgoing scattering states, which are normalized such that they carry a unit flux. Substituting into the parallel transport law for $|\psi_\alpha\rangle$, one gets Eq. (2) which is equivalent to the parallel transport law for row vectors of the scattering matrix.

Quantum adiabatic pumping. In order to establish the connection between the geometric phase above and the quantum pumping charge, we need to consider the time-reversed scattering states $|\hat{\psi}_\alpha^S(t)\rangle = |\hat{\alpha}\rangle_{in} + \sum_{\beta=1}^N S_{\beta\alpha}(t)|\hat{\beta}\rangle_{out}$ with $\hat{\cdot}$ denoting the counterparts under time reversal operation [21], which constitute a solution of the Schrödinger equation for the time-reversed Hamiltonian $\hat{H}$ at any given (frozen) time *at the epoch scale* [19]. This gives rise to another gauge potential $\hat{A}_{\alpha\beta\nu} \equiv \hat{\mathbf{n}}_\beta^* \cdot \partial_\nu \hat{\mathbf{n}}_\alpha$ with $\hat{\mathbf{n}}_\alpha \equiv (S_{1\alpha}, \dots, S_{N\alpha})$, i.e., the column vectors of the scattering matrix $S(t)$. In this case, the gauge group arises from redistribution of scattering particles among different outgoing channels. If $\hat{\mathbf{n}}'_\alpha(t) = \sum_\beta \hat{\omega}_{\alpha\beta} \hat{\mathbf{n}}_\beta(t)$, then the gauge transformation takes $\hat{A}'(t) = d\hat{\omega}\hat{\omega}^{-1} + \hat{\omega}\hat{A}\hat{\omega}^{-1}$. The gauge fields A and $\hat{A}$ are connected via time reversal operation. If we identify the emissivity into the α -th channel in the n -th lead as $\text{Im}[\hat{A}_{\alpha\alpha}/2\pi]$ [18], then we immediately reproduce Brouwer's formula [17] describing charge pumping, which turns out to be associated with the Abelian subgroup

$U(1)$,

$$Q = \frac{e}{2\pi} \text{Im} \oint \text{Tr} \hat{A}, \quad (6)$$

with Q being the charge transferred into the n -th lead during one cycle. That is, the charge transferred during adiabatic pumping is essentially the geometric phase associated with the charge sector $U(1)$. This also explains why Planck's constant $\hbar$ does not occur in the adiabatic quantum pumped charge (current), a peculiar feature different from the Landauer-Büttiker conductance. However, as is well known, the geometric phase is determined only up to a multiple of 2π . This concerns global geometric properties, i.e., the winding number of the overall phase of the gauge transformation in Eq. (5), $N \equiv 1/(2\pi i) \oint \text{Tr}(d\hat{\omega}\hat{\omega}^{-1})$. The requirement that all physical observables be invariant under the gauge transformation leads us to the conclusion that $O(Q) = O(Q - N)$, with O denoting any observable. This result has been noticed by Makhin and Mirlin [11], without proper justification, for the counting statistics in quantum charge pumps (see also, Ref. [9]). To see the effects caused by non-Abelian gauge potentials, we need to consider the gauge invariant quantity $\text{Tr}U(t)$.

In contrast to the Abelian $U(1)$ sector, there is no gauge-independent definition of pumping currents associated with the non-Abelian sector. However, if in a certain gauge, the gauge potential $\hat{A}(t)$ turns out to be diagonal, i.e., the so-called ‘‘Abelianized’’ non-Abelian gauge potential, it is legitimate to speak of pumping currents in the non-Abelian $SU(M_n)$ sector associated with internal degrees of freedom, such as spin and modes [22]. However, we stress that this is *only* true in a *fixed* gauge due to the gauge-dependence of the ‘‘spin’’ current so defined.

Tunneling through a single magnetic spin. Consider the Hamiltonian which describes a system consisting of two leads coupled to a single site, the spin of which has an exchange interaction J with a magnetic spin [23],

$$H = \sum_{k \in L, R, \sigma} \epsilon_{k\sigma} c_{k\sigma}^\dagger c_{k\sigma} + J \sum_{\sigma, \sigma'} d_\sigma^\dagger \Omega_{\sigma\sigma'} d_{\sigma'} + \sum_{k \in L, R, \sigma; \sigma'} (V_{k\sigma, \sigma'} c_{k\sigma}^\dagger d_{\sigma'} + \text{H.c.}). \quad (7)$$

Here $c_{k\sigma}^\dagger$ and $c_{k\sigma}$ are, respectively, the creation and destruction operators of an electron with momentum k and spin σ in either left (L) or the right (R) lead, and $d_\sigma^\dagger$ and d_σ are the counterpart of the single electron with spin σ at the spin site. The quantity $\epsilon_{k\sigma}$ are the single particle energies of conduction electrons in the two leads, which we will assume $\epsilon_{k\sigma} = v_F(|k| - k_F)$ with the convention that $v_F = 1$, and the momentum is measured from the Fermi surface for electrons in leads. The electrons on the spin site are connected to those in the two leads with the tunneling matrix elements $V_{k\sigma, \sigma'}$. For simplicity, we assume symmetric tunneling barriers between the local spin and the leads, and only keep the spin-conserved

coupling; viz. $V_{L++} = V_{L--} = V_{R++} = V_{R--} = V$ and $V_{L+-} = V_{L-+} = V_{R+-} = V_{R-+} = 0$. The entries of the coupling matrix Ω take the form $\Omega_{++} = -\Omega_{--} = \cos\theta$ and $\Omega_{+-} = \Omega_{-+}^* = \sin\theta \exp(-i\phi)$, where $\vec{n} = (\sin\theta \cos\phi, \sin\theta \sin\phi, \cos\theta)$ is the direction of the magnetic spin. The model is exactly soluble as far as the scattering matrix is concerned.

Once the scattering matrix is determined, our general formalism in Eq. (3) leads us to the non-Abelian gauge potential,

$$\hat{A} = \hat{A}_\theta d\theta + \hat{A}_\phi d\phi, \quad (8)$$

where $\hat{A}_\theta \equiv \hat{A}_\theta^1 \sigma^1 + \hat{A}_\theta^2 \sigma^2 + \hat{A}_\theta^3 \sigma^3$ with $\hat{A}_\theta^1 = i(\sin(\delta_1 - \delta_2) \cos\theta \cos\phi + (1 - \cos(\delta_1 - \delta_2)) \sin\phi)/4$, $\hat{A}_\theta^2 = i(\sin(\delta_1 - \delta_2) \cos\theta \sin\phi - (1 - \cos(\delta_1 - \delta_2)) \cos\phi)/4$, and $\hat{A}_\theta^3 = -i(\sin(\delta_1 - \delta_2) \sin\theta)/4$, and $\hat{A}_\phi \equiv \hat{A}_\phi^1 \sigma^1 + \hat{A}_\phi^2 \sigma^2 + \hat{A}_\phi^3 \sigma^3$ with $\hat{A}_\phi^1 = -i(\sin(\delta_1 - \delta_2) \sin\theta \sin\phi - (1 - \cos(\delta_1 - \delta_2)) \sin\theta \cos\theta \cos\phi)/4$, $\hat{A}_\phi^2 = i(\sin(\delta_1 - \delta_2) \sin\theta \cos\phi + (1 - \cos(\delta_1 - \delta_2)) \sin\theta \cos\theta \sin\phi)/4$, and $\hat{A}_\phi^3 = -i(1 - \cos(\delta_1 - \delta_2)) \sin^2\theta/4$. Here δ_i ($i = 1, 2$) are the phase shifts defined by $\delta_1 = -2 \tan^{-1}(\Gamma/(k - J))$ and $\delta_2 = -2 \tan^{-1}(\Gamma/(k + J))$ with the tunneling rate $\Gamma \equiv V^2$.

The gauge field strength F then takes the form $\hat{F} = -i(1 - \cos(\delta_1 - \delta_2)) \vec{n} \cdot \vec{\sigma} d\Omega/4$. Here the invariant area element $d\Omega = \sin\theta d\theta \wedge d\phi$. Obviously, this is just a simple rotation of the standard form $\hat{F} = -i(\alpha^2 - 1)\sigma^3 d\Omega/2$. Up to a gauge transformation, this is the same non-Abelian gauge potential, found by Moody *et al.* [15] for a diatomic molecule. This is consistent with a theorem, proved in [24], stating that the rotationally invariant connection on the sphere is essentially unique. To establish the relation between α and $\cos(\delta_1 - \delta_2)$, we need to calculate the gauge invariant quantity $\text{Tr} \hat{F} \wedge * \hat{F}$, with $*\hat{F}$ being the dual of $\hat{F}$. Then we have $\alpha^2 = (3 - \cos(\delta_1 - \delta_2))/2$. When $J = 0$, i.e., in the absence of the direct exchange interaction between electrons and the local spin, the gauge field is a pure gauge because $\hat{F} = 0$. Since the Pauli matrices are traceless, we have $\text{Tr} \hat{A} = 0$, meaning that charge pumping is absent in the model under consideration.

To observe the effects of the non-Abelian gauge field, it is necessary to choose a cycle which varies both θ and ϕ . The analysis of Zee [16] for the gauge field in Eq. (8) showed that one can compute a phase factor U_{SR} which is obtained from Eq. (4) for the ‘‘spherical rectangle’’ path starting from the point (θ_1, ϕ_1) , then going to (θ_2, ϕ_1) along a fixed- ϕ path, then to (θ_2, ϕ_2) along a fixed- θ path, then to (θ_1, ϕ_2) along a fixed- ϕ path, and finally back to (θ_1, ϕ_1) along a fixed- θ path (see Fig. 1). The quantum spin pumping phase $1/2 \text{Tr} U_{SR}$ is shown in Fig. 1 as a function of system parameters for two different paths, C_1 and C_2 . If the phase takes a value close to unity, spin pumping disappears. When the phase goes away from unity, the scattering matrix acquires a non-Abelian matrix phase. The spin pumping phase takes its maximum value around the resonant scattering lines $k = \pm J$.

Possible experiments. A scanning tunneling microscope (STM) has been used to detect a quantum mirage around a single magnetic cobalt atom placed on a non-magnetic metallic copper surface [25]. Electron spin resonance (ESR)-STM experiments [26] have advanced to the point that they have spatial resolution at the level of a few spins [27]. The STM setup as shown in Fig. 1 A should make it possible to observe gauge invariant spin pumping via a single magnetic atom on the surface of the

substrate. One way to do this is to observe the interference resulting from two different tunneling paths: in one path, electrons tunnel directly from the tip to the substrate, and in the other path, electrons tunnel through a single magnetic spin and couple to the adiabatic spin dynamics of the magnetic atom.

Acknowledgments. We thank Urban Lundin and Gerard Milburn for valuable discussions. This work was supported by the Australian Research Council.

-
- [*] E-mail: hqz@maths.uq.edu.au
- [1] D. J. Thouless, Phys. Rev. B **27**, 6083 (1983).
- [2] B. L. Altshuler and L. I. Glazman, Science **283**, 1864 (1999).
- [3] P. Sharma and C. Chamon, Phys. Rev. Lett. **87**, 096401 (2001); E. R. Mucciolo, C. Chamon, and C. M. Marcus, Phys. Rev. Lett. **89**, 146802 (2002).
- [4] T. Aono, cond-mat/0205395; M. Governale, F. Taddei, and R. Fazio, cond-mat/0211211.
- [5] M. Switkes, C. M. Marcus, K. Campman, and A. C. Gosard, Science **283**, 1905 (1999); For an alternative interpretation, see P. W. Brouwer, Phys. Rev. B **63**, 121303 (2001).
- [6] S. K. Watson, R. M. Potok, C. M. Marcus, and V. Umansky, cond-mat/0302492.
- [7] F. Zhou, B. Spivak, and B. Altshuler, Phys. Rev. Lett. **82**, 608 (1999).
- [8] T. A. Shutenko, I. L. Aleiner, and B. L. Altshuler, Phys. Rev. B **61**, 10366 (2000).
- [9] A. Andreev and A. Kamenev, Phys. Rev. Lett. **85**, 1294 (2000).
- [10] J. E. Avron *et al.*, Phys. Rev. B **62**, R10618 (2000); J. E. Avron *et al.*, Phys. Rev. Lett. **87**, 236601 (2001).
- [11] There is a difference of factor 2 between our result and equation (5) in Y. Makhin and A. D. Mirlin, Phys. Rev. Lett. **87**, 276803 (2001).
- [12] M. V. Berry, Proc. Roy. Soc. London. A **392**, 45 (1984).
- [13] B. Simon, Phys. Rev. Lett. **51**, 2167 (1983).
- [14] F. Wilczek and A. Zee, Phys. Rev. Lett. **52**, 2111 (1984).
- [15] J. Moody, A. Shapere, and F. Wilczek, Phys. Rev. Lett. **56**, 893 (1986).
- [16] A. Zee, Phys. Rev. A **38**, 1 (1988).
- [17] P. W. Brouwer, Phys. Rev. B **58**, R10135 (1998).
- [18] M. Büttiker, H. Thomas, and A. Prêtre, Phys. B **94**, 133 (1994).
- [19] J. E. Avron *et al.*, J. Math. Phys. **43**, 3415 (2002).
- [20] Y. Aharonov and J. Anandan, Phys. Rev. Lett. **58**, 1593 (1987); J. Samuel and R. Bhandari, Phys. Rev. Lett. **60**, 2339 (1998).
- [21] We emphasize that the time reversal operation here is carried out at the epoch scale, which may be or may not be broken, depending on the Hamiltonian concerned. However, there is another time reversal operation at large time scale, which is always broken during pumping.
- [22] H.-Q. Zhou, S. Y. Cho, and R. H. McKenzie, *unpublished*.
- [23] J. X. Zhu and A. V. Balatsky, Phys. Rev. Lett. **89**, 286802 (2002).
- [24] M. Forgacs and N. Manton, Commun. Math. Phys. **72**, 15 (1980); E. Witten, Phys. Rev. Lett. **38**, 121 (1977); C. H. Gu, Phys. Rep. **80**, 251 (1981).
- [25] H. C. Manoharan, C. P. Lutz, and D. M. Eigler, Nature **403**, 512 (2000).
- [26] M. Farle, Rep. Prog. Phys. **61**, 755 (1998); M. Bode, M. Getzlaff, and R. Wiesendanger, Phys. Rev. Lett. **81**, 4256 (1998); S. Heinze *et al.*, Science **288**, 1805 (2000).
- [27] D. Ralph, Science **291**, 999 (2001); W. Weber, S. Riesen, and H. C. Siegmann, Science **291**, 1015 (2001).
-

Berry's Phase	Quantum Pumping Phase
Closed systems	Open systems
Wave functions $ \psi_\alpha\rangle$	Row (column) vectors $\mathbf{n}_\alpha$ ($\hat{\mathbf{n}}_\alpha$) of the S matrix
Energy levels E_n	Leads n
M degeneracies	M channels
Discrete spectrum (bound states)	Continuous spectrum (scattering states)
Eigenstates in the n -th level	Channels in the n -th lead
Parallel transport due to adiabatic theorem	Parallel transport due to adiabatic charge and spin pumps
Gauge potential $A_{\alpha\beta\nu} = \langle \psi_\beta^* \partial_\nu \psi_\alpha \rangle$	Gauge potentials $A_{\alpha\beta\nu} = \mathbf{n}_\beta^* \cdot \partial_\nu \mathbf{n}_\alpha$ and $\hat{A}_{\alpha\beta\nu} = \hat{\mathbf{n}}_\beta^* \cdot \partial_\nu \hat{\mathbf{n}}_\alpha$
$U(M)$ bundle	$U(M)$ bundle
Gauge group $U(M)$ arising from different choices of bases	Gauge group $U(M)$ arising from redistribution of the scattering particles among different channels
External parameters $X = (X^1, \dots, X^p)$	External parameters $X = (X^1, \dots, X^p)$

TABLE I: Comparison of Berry's phase and the quantum pumping phase.




FIG. 1: The dependence of the quantum spin pumping phase on system parameters for two leads connected to a single magnetic spin whose direction is slowly varied around the path shown on the right. **A**. Left: Schematic of the magnetic spin coupled to left (L) and right (R) leads. The magnetic spin $\mathbf{S}$ precesses around the direction of the magnetic field $\mathbf{B}$. Right: An equivalent scanning tunneling microscope experimental setup. The pumping cycles on the parameter (θ, ϕ) -sphere are, respectively, taken to be $C_1(\theta_1, \phi_1, \theta_2, \phi_2) = (\pi/8, 0, \pi/2, \pi)$ for B and $C_2(\theta_1, \phi_1, \theta_2, \phi_2) = (\pi/8, 0, 7\pi/8, \pi)$ for C.