Kondo effect in the presence of itinerant-electron ferromagnetism studied with the numerical renormalization group method

J. Martinek, ^{1,4} M. Sindel, ² L. Borda, ^{2,3} J. Barnaś, ^{4,5} J. König, ^{1,6} G. Schön, ¹ and J. von Delft²

¹Institut für Theoretische Festkörperphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany

²Sektion Physik and Center for Nanoscience, LMU München, Theresienstrasse 37, 80333 München, Germany

³ Hungarian Academy of Sciences, Institute of Physics, TU Budapest, H-1521, Hungary

⁴Institute of Molecular Physics, Polish Academy of Sciences, 60-179 Poznań, Poland

⁵Department of Physics, Adam Mickiewicz University, 61-614 Poznań, Poland

⁶Institut für Theoretische Physik III, Ruhr-Universität Bochum, 44780 Bochum, Germany

(Dated: Submitted: April 16, 2003)

The Kondo effect in quantum dots (QDs) - artificial magnetic impurities - attached to ferromagnetic leads is studied with the numerical renormalization group method. It is shown that the QD level is spin-split due to presence of ferromagnetic electrodes, leading to a suppression of the Kondo effect. We find that the Kondo effect can be restored by compensating this splitting with a magnetic field. Although the resulting Kondo resonance then has an unusual spin asymmetry with a reduced Kondo temperature, the ground state is still a locally-screened state, describable by Fermi liquid theory and a generalized Friedel sum rule, and transport at zero temperature is spin-independent.

PACS numbers: 75.20.Hr, 72.15.Qm, 72.25.-b, 73.23.Hk

Introduction. – The prediction [1] and experimental observation of the Kondo effect in artificial magnetic impurities – semiconducting quantum dots (QDs) [2, 3] – renewed interest in the Kondo effect and opened new opportunities of research. The successful observation of the Kondo effect in molecular QDs like carbon nanotubes [4, 5] and single molecules [6] attached to metallic electrodes opened the possibility to study the influence of many-body correlations in the leads (superconductivity [7] or ferromagnetism) on the Kondo effect. Recently the question arose whether the Kondo effect in a QD attached to ferromagnetic leads can occur or not. Several authors have predicted [8, 9, 10, 11] that the Kondo effect should occur. However, it was shown recently [12] that the QD level will be spin-split due to the presence of ferromagnetic electrodes leading to a suppression of the Kondo effect, and that the Kondo resonance can be restored only by applying an external magnetic field. The analyses of Refs. [8, 9, 10, 11, 12] were all based on approximate methods.

In this Letter we resolve the controversy by adapting the numerical renormalization-group (NRG) technique [13, 14], one of the most accurate methods available to study strongly-correlated systems in the Kondo regime, to the case of a QD coupled to ferromagnetic leads with parallel magnetization directions. We find that in general the Kondo resonance is split, similar to the usual magnetic-field-induced splitting [15, 16]. However, we find that by appropriately tuning an external magnetic field, this splitting can be fully compensatedand and the Kondo effect can be restored [17] (confirming Ref. [12]). We point out that precisely at this field the occupancy of the local level is the same for spin up and down, $n_{\uparrow} = n_{\downarrow}$, a fact that follows from the Friedel sum rule [29]. Moreover, we show that the Kondo effect then has unusual

properties such as a strong spin polarization of the Kondo resonance and, just as for ferromagnetic materials, for the density of states (DOS). Nevertheless, despite of the spin assymetries in the DOS of the QD and the leads, the symmetry in the occupancy $n_{\uparrow} = n_{\downarrow}$ implies that the system's ground state can be tuned to have a fully compensated local spin, in which case the QD conductance is found to be the *same* for each spin channel, $G_{\uparrow} = G_{\downarrow}$.

The Model. – For ferromagnetic leads electron-electron interactions in the leads give rise to magnetic order and spin-dependent DOS $\rho_{r\uparrow}(\omega) \neq \rho_{r\downarrow}(\omega), r = L, R$. Magnetic order of typical band ferromagnets like Fe, Co, and Ni is mainly related to electron correlation effects in the relatively narrow 3d sub-bands, which only weakly hybridize with 4s and 4p bands [19]. We can assume that due to a strong spatial confinement of d electron orbitals, the contribution of electrons from d sub-bands to transport across the tunnel barrier can be neglected [20]. In such a situation the system can be modeled by noninteracting [21] s electrons, which are spin polarized due to the exchange interaction with uncompensated magnetic moments of the completely localized d electrons. In mean-field approximation one can model this exchange interaction as an effective molecular field, which removes spin degeneracy in the system of noninteracting conducting electrons, leading to a spin-dependent DOS. The Anderson model (AM) for a QD with a single energy level $\epsilon_{\rm d}$, which is coupled to ferromagnetic leads, is given by

$$\tilde{H} = \sum_{rk\sigma} \epsilon_{rk\sigma} c_{rk\sigma}^{\dagger} c_{rk\sigma} + \epsilon_{d} \sum_{\sigma} \hat{n}_{\sigma} + U \hat{n}_{\uparrow} \hat{n}_{\downarrow}
+ \sum_{rk\sigma} (V_{r,k} d_{\sigma}^{\dagger} c_{r,k\sigma} + V_{r,k}^{*} c_{r,k\sigma}^{\dagger} d_{\sigma}) - g \mu_{B} B S_{z} .$$
(1)

Here $c_{rk\sigma}$ and d_{σ} $(\hat{n}_{\sigma} = d_{\sigma}^{\dagger}d_{\sigma})$ are the Fermi operators for electrons with momentum k and spin σ in the

leads (r=L/R), and in the QD, V_{rk} is the tunneling amplitude, $S_z=(\hat{n}_{\uparrow}-\hat{n}_{\downarrow})/2$, and the last term is the Zeeman energy of the dot. In general, all information about spin asymmetry in the leads can be modeled by the spin-dependent hybridization function $\Gamma_{r\sigma}(\omega)=\pi\sum_k\delta(\omega-\epsilon_{k\sigma})V_{r,k}^2=\pi\rho_{r\sigma}(\omega)V_r^2$, where $V_{r,k}=\mathrm{const}\equiv V_r$ and $\rho_{r\sigma}(\omega)$ is the spin-dependent DOS.

In order to understand the Kondo physics of a QD attached to two identical ferromagnetic electrodes with parallel configurations, it suffices to study, instead of the above general model [Eq. (1)], a simpler one, which captures the same essential physics, namely the fact that $\Gamma_{r\uparrow}(\omega) \neq \Gamma_{r\downarrow}(\omega)$ will generate an effective local magnetic field, which lifts the degeneracy of the local level (even for B=0). A simple (but not unique) way of modeling this effect is to take the DOS in the leads to be constant and spin-independent, $\rho_{r\sigma}(\omega) \equiv \rho$, the bandwidths to be equal $D_{\uparrow} = D_{\downarrow}$, and lump all spin-dependence into the spin-dependent hybridization function, $\Gamma_{r\sigma}(\omega)$, which we take to be ω -independent, $\Gamma_{r\sigma}(\omega) \equiv \Gamma_{r\sigma}$. By means of a unitary transformation [1] the AM [Eq. (1)] can be mapped onto a model in which the correlated QD level couples only to one electron reservoir described by Fermi operators $\alpha_{k\sigma}$ with strength $\Gamma_{\sigma} = \sum_{r} \Gamma_{r\sigma}$,

$$H = \sum_{k\sigma} \epsilon_k \alpha_{k\sigma}^{\dagger} \alpha_{k\sigma} + \epsilon_{\rm d} \sum_{\sigma} \hat{n}_{\sigma} + U \hat{n}_{\uparrow} \hat{n}_{\downarrow}$$

$$+ \sum_{k\sigma} \sqrt{\frac{\Gamma_{\sigma}}{\pi \rho}} (d_{\sigma}^{\dagger} \alpha_{k\sigma} + \alpha_{k\sigma}^{\dagger} d_{\sigma}) - g \mu_B B S_z .$$
(2)

Finally, we parametrize the spin-dependence of Γ_{σ} in terms of a spin-polarization parameter $P \equiv (\Gamma_{\uparrow} - \Gamma_{\downarrow})/\Gamma$, by writing $\Gamma_{\uparrow(\downarrow)} \equiv \frac{1}{2}\Gamma(1 \pm P)$, where $\Gamma \equiv \Gamma_{\uparrow} + \Gamma_{\downarrow}$.

In the model of Eq. (2), we allowed for $\Gamma_{\uparrow} \neq \Gamma_{\downarrow}$ but not for $D_{\uparrow} \neq D_{\downarrow}$, as would be appropriate for real ferromagnets, whose spin-up and down bands always have a Stoner splitting $\Delta D \equiv D_{\uparrow} - D_{\downarrow}$, with typical values $\Delta D/D_{\uparrow} \lesssim 20\%$ (for Ni, Co, Fe). However, no essential physics is thereby lost, since the consequence of taking $D_{\uparrow} \neq D_{\downarrow}$ is the same as that of taking $\Gamma_{\uparrow} \neq \Gamma_{\downarrow}$, namely to generate an effective local magnetic field [24].

The occurrence of the Kondo effect requires spin fluctuations in the dot as well as zero-energy spin-flip excitations in the leads. Indeed, a Stoner ferromagnet without full spin polarization -1 < P < 1 provides zero-energy Stoner excitations [25], even in the presence of an external magnetic field.

Method. – The model [Eq. (2)] can be treated by Wilson's NRG method. This method, with recent improvements related to high-energy features and finite magnetic field [15, 16], is a well-established method to study the Kondo impurity (QD) physics. It allows one to calculate the level occupation $n_{\sigma} \equiv \langle \hat{n}_{\sigma} \rangle$ (a static property), the QD spin spectral function, Im $\chi_s^z(\omega) = \mathcal{F}\{i\Theta(t)\langle[S_z(t),S_z(0)]\rangle\}$, where \mathcal{F} denotes the Fourier transform, and the spin-resolved single-particle spec-




FIG. 1: Spin-dependent occupation of the dot level at (a) B=0 and (b) $B=-0.1\Gamma$, as a function of spin polarization P. (a) For B=0, the condition $n_{\uparrow}=n_{\downarrow}$ only holds at P=0. (b) For finite P it can be satisfied if a finite, finetuned magnetic field, $B_{\text{comp}}(P)$, is applied, whose dependence on P is shown in (c). As expected, it is approximately linear [12, 28]. Here U=0.12D, $\epsilon_{\rm d}=-U/3$, $\Gamma=U/6$, and T=0.




FIG. 2: a) QD spectral function $A(\omega) = \sum_{\sigma} A_{\sigma}(\omega)$ for several values of spin polarization P; inset: expanded scale of the spectral function around $\epsilon_{\rm F}$. b) The spin-resolved spectral function for fixed P. For $P \to -P$, we have $A_{\sigma} \to A_{-\sigma}$. c) Compensation of the spin splitting by fine-tuning an external magnetic field. Parameters U, $\epsilon_{\rm d}$, Γ , and T as in Fig. 1.

tral density $A_{\sigma}(\omega,T,B,P) = -\frac{1}{\pi} \mathrm{Im} \mathcal{G}^R_{d,\sigma}(\omega)$ for arbitrary temperature T, magnetic field B and polarization P [where $G^R_{\sigma}(\omega)$ denotes a retarded Green function]. From this we can find the spin-resolved conductance $G_{\sigma} = \frac{e^2}{\hbar} \frac{2\lambda}{(\lambda+1)^2} \Gamma_{\sigma} \int_{-\infty}^{\infty} d\omega A_{\sigma}(\omega) (-\frac{\partial f(\omega)}{\partial \omega})$ with $f(\omega)$ denoting the Fermi function and $\lambda = \Gamma_{L\sigma}/\Gamma_{R\sigma}$ describing the asymmetry in the couplings. We choose $\lambda = 1$ below.

Generation and restoration of spin splitting. – In this Letter, we focus exclusively on the properties of the system at T=0 in the local moment - Kondo regime, where the total occupancy of the local level, $n=\sum_{\sigma}n_{\sigma}\approx 1$. The occurence of charge fluctuations of course broadens and shifts the position of the QD levels (for both spin up and down), and hence changes their occupation. For

 $P \neq 0$, the charge fluctuations and hence level shifts and level occupations become spin-dependent, causing the QD level to split [12] and the dot magnetization $n_{\uparrow} - n_{\downarrow}$ to be finite (Fig. 1). As a result, the Kondo resonance, too, is spin-split [26, 27] and weakened (Fig. 2), similarly to the effect of an applied magnetic field [15, 16]. This means that Kondo correlations are reduced or even suppressed in the presence of ferromagnetic leads. However, for any fixed P, it is possible to compensate the splitting of the Kondo resonance [Fig. 2(c) and 3(c)] by fine-tuning the magnetic field to an appropriate value, $B_{\text{comp}}(P)$, defined as the field which maximizes the height of the Kondo resonance. This field is found to depend linearly [28] on P [Fig. 1(c)] (as predicted in [12] for $U \to \infty$). Remarkably, we also find (throughout the local moment regime) that at B_{comp} the local occupancies satisfy $n_{\uparrow} = n_{\downarrow}$ [Fig. 1(b)]. The fact that this occurs simultaneously with the disappearance of the Kondo resonance splitting suggests that the local spin is fully screened at B_{comp} .

Spectral functions. – We henceforth fix the magnetic field at $B = B_{\text{comp}}(P)$. To learn more about the properties of the corresponding ground state, we computed the spin spectral function $\operatorname{Im}\{\chi_s^z(\omega)\}\$ for several values of P at T=0 (Fig. 3). Its behavior is characteristic for the formation of a local Kondo singlet: as a function of decreasing frequency, the spin spectral function shows a maximum at a frequency ω_{max} which we associate with the Kondo temperature [i.e. $k_B T_{\rm K} \equiv \hbar \omega_{\rm max}$ at $B = B_{\text{comp}}(P)$, and then decreases linearly with ω , indicating the formation of the Fermi liquid state [14]. By determining $T_{\rm K}(P)$ (from $\omega_{\rm max}$) for different P-values, we find that $T_{\rm K}$ decreases with increasing P [Fig. 3(b)]. For metals like Ni, Co, and Fe, where P = 0.24, 0.35, and 0.40 respectively, the decrease of $T_{\rm K}$ is rather weak, so the Kondo effect should still be experimentally accessible. Remarkably, both $\operatorname{Im}\{\chi_s^z(\omega)\}\$ and the spectral function $A_{\sigma}(\omega)$ collapse rather well onto a universal curve if plotted in appropriate units [Figs. 3(a) and 3(c)]. This indicates that an applied magnetic field B_{comp} restores the universal behavior characteristic for the isotropic Kondo effect, in spite of the presence of spin-dependent coupling to the leads. Fig. 4(a) shows that the amplitude of the Kondo resonance is strongly spin dependent for ferromagnetic leads, which is unusual and unique. The nature of this asymmetry is related to the asymmetry of the DOS in the leads, and its value is exactly proportional to $\sim 1/\Gamma_{\sigma}$. As a result the total conductance G_{σ} is not spin dependent [Fig. 4(b)]. This indicates the robustness of the Kondo effect in this system: if the external magnetic field has been tuned appropriately, it is able to compensate the presence of a spin asymmetry in the leads by creating a proper spin asymmetry in the dot spectral density, thereby conserving a fully compensated local spin and achieving perfect transmission.

Friedel sum rule. – Further insights can be gained from the Friedel sum rule, an exact $T\,=\,0$ relation

[29] that holds for arbitrary values of P and B [30]. The interacting Green's function can be expressed as [14] $\mathcal{G}_{d,\sigma}^R(\omega) = [\omega - \epsilon_{\mathrm{d}\sigma} + i\Gamma_{\sigma} - \Sigma_{\sigma}(\omega)]^{-1}$, with spin-dependent $\epsilon_{\mathrm{d}\sigma}$ and Γ_{σ} ; the former due to Zeeman splitting ($\epsilon_{\mathrm{d}\sigma} = \epsilon_{\mathrm{d}} - 1/2 \, \sigma g \mu_B B$) and latter due to the ferromagnetic leads. Here $\Sigma_{\sigma}(\omega)$ denotes the spin-dependent self energy. Now, the Friedel sum rule [29] implies that at T=0, the occupancy n_{σ} and spectral functions can be written as

$$n_{\sigma} = \phi_{\sigma}/\pi = \frac{1}{2} - \frac{1}{\pi} \tan^{-1} \left(\frac{\epsilon_{d\sigma} - \epsilon_{F} + \Sigma_{\sigma}^{R}(\epsilon_{F})}{\Gamma_{\sigma}} \right), (3)$$

$$A_{\sigma}(\epsilon_{\rm F}) = \frac{\sin^2(\pi n_{\sigma})}{\pi \Gamma_{\sigma}} \,, \tag{4}$$

where $\Sigma_{\sigma}^{R}(\omega) \equiv \text{Re } \Sigma_{\sigma}(\omega)$, and $\phi_{\sigma}(\omega)$ is the phase of $\mathcal{G}_{d,\sigma}^{R}(\omega)$. Since $\Sigma_{\uparrow}^{R}(\epsilon_{\mathrm{F}}) \neq \Sigma_{\downarrow}^{R}(\epsilon_{\mathrm{F}})$, an equal spin occupation, $n_{\uparrow} = n_{\downarrow}$, is possible only for $(\epsilon_{\mathrm{d}\uparrow} - \epsilon_{\mathrm{F}} + \Sigma_{\uparrow}^{R}(\epsilon_{\mathrm{F}}))/\Gamma_{\uparrow} = (\epsilon_{\mathrm{d}\downarrow} - \epsilon_{\mathrm{F}} + \Sigma_{\downarrow}^{R}(\epsilon_{\mathrm{F}}))/\Gamma_{\downarrow}$, which can be obtained only for an appropriate external magnetic field $B = B_{\mathrm{comp}}$. For the latter, in the local moment regime $(n \approx 1)$ we have $n_{\uparrow} = n_{\downarrow} \approx 0.5$, so that $\phi_{\uparrow} = \phi_{\downarrow} \approx \pi/2$, which implies that the peaks of A_{\uparrow} and A_{\downarrow} are aligned. Thus, the Friedel sum rule clarifies why the magnetic field B_{comp} at which the splitting of the Kondo resonance disappears, coincides with that for which $n_{\uparrow} = n_{\downarrow}$. For $B = B_{\mathrm{comp}}$, the spindependent amplitude $A_{\sigma}(\epsilon_{\mathrm{F}})$ of Eq. (4) and the conductance $G_{\sigma} \sim \Gamma_{\sigma} A_{\sigma}(\epsilon_{\mathrm{F}})$ agree well with the abovementioned NRG results [Fig. 4(a), 4(b)].




FIG. 3: a) The spin spectral function $\operatorname{Im}\{\chi_s^z(\omega)\}$ as a function of energy. b) Dependence of $T_{\rm K}$ on spin polarization P. The solid line shows the prediction from Ref. [12] for the functional form $T_K(P)/T_K(0) = \exp[C \operatorname{arctanh}(P)/P]$, where the best fit is obtained for C = -5.98. Equation (6) from Ref. [12] with $(\rho_{\uparrow} + \rho_{\downarrow})J_0 = (\Gamma/\pi)U/[|\epsilon_{\rm d}|(U + \epsilon_{\rm d})]$ would lead to C = -4.19. c) QD spectral function for several values of P. Parameters $U, \epsilon_{\rm d}, \Gamma$, and T are as in Fig. 1, $B = B_{\rm comp}(P)$.




FIG. 4: (a) Spin resolved QD spectral function amplitude $A_{\sigma}(\omega=0)$ at the Fermi level, and (b) the QD conductance G_{σ} , as functions of the spin polarization P, for $B=B_{\rm comp}(P)$ and symmetric couplings $(\Gamma_{L\sigma}=\Gamma_{R\sigma})$, with U, $\epsilon_{\rm d}$, Γ , and T as in Fig. 1, implying $n_{\sigma}\approx 0.5$. The dashed line in (a) is $1/(1\pm P)$ (Eq. (4) with $n_{\sigma}=0.5$). As expected, we find $G_{\sigma}=e^2/h$, with a numerical error less than 1%.

In conclusion, applying the NRG technique, we showed that the Kondo effect in a QD attached to ferromagnetic leads is in general suppressed, because the latter induce a spin splitting of the QD level, which leads to an asymmetry in the occupancy $n_{\uparrow} \neq n_{\downarrow}$. Remarkably, the Kondo effect may nevertheless be restored by applying an external magnetic field $B_{\rm comp}$, tuned such that the splitting of the Kondo resonance is compensated and the condition $n_{\uparrow} = n_{\downarrow}$ is fulfilled. Although the Kondo resonance is strongly spin polarized, it then features a locally-screened state, a spin-independent conductance, and a Kondo temperature which decreases with increasing spin asymmetry.

Acknowledgements. – We thank R. Bulla, T. Costi, L. Glazman, W. Hofstetter, H. Imamura, B. Jones, S. Maekawa, A. Rosch, M. Vojta, and Y. Utsumi for discussions. This work was supported by the DFG under the CFN, 'Spintronics' RT Network of the EC RTN2-2001-00440, Project PBZ/KBN/044/P03/2001, OTKA T034243 and the Emmy-Noether program.

Note added. – After submission of our paper, a preprint [M. S. Choi et al. cond-mat/0305107] studying a similar problem using the NRG technique appeared, with conclusions consistent with our's.

- [4] J. Nygård et al., Nature 408, 342 (2000).
- [5] M. R. Buitelaar *et al.*, Phys. Rev. Lett. **88**, 156801 (2002).
- [6] J. Park et al., Nature 417, 722 (2002); W. Liang et al., Nature 417, 725 (2002).
- [7] M. R. Buitelaar et al., Phys. Rev. Lett. 89, 256801 (2002)
- [8] N. Sergueev et al., Phys. Rev. B 65, 165303 (2002).
- [9] P. Zhang et al., Phys. Rev. Lett. 89, 286803 (2002).
- 10] B. R. Bułka et al., Phys. Rev. B 67, 024404 (2003).
- [11] R. Lopez *et al.*, Phys. Rev. Lett. **90**, 116602 (2003)
- 12 J. Martinek et al., Phys. Rev. Lett. 91, 127203 (2003).
- [13] K.G. Wilson, Rev. Mod. Phys. 47 773 (1975); T. A. Costi et al., J. Phys.: Cond. Mat. 6, 2519 (1994).
- [14] A. C. Hewson, The Kondo Problem to Heavy Fermions, Cambridge Univ. Press (1993).
- [15] W. Hofstetter, Phys. Rev. Lett. 85, 1508 (2000).
- [16] T. A. Costi, Phys. Rev. Lett. 85, 1504 (2000); Phys. Rev. B 64, 241310 (2001).
- [17] The remarkable full recovery of the Kondo effect should be contrasted to the competition between Kondo physics and magnetic RKKY interactions in heavy-fermion systems [18]. There, depending on the relation between the relevant energy scales $T_{\rm RKKY}$ and $T_{\rm K}$, either the local spin is quenched and no magnetic order occurs (for $T_{\rm K} > T_{\rm RKKY}$), or the local molecular field removes spin degeneracy and suppresses the Kondo effect (for $T_{\rm K} < T_{\rm RKKY}$).
- [18] A. H. Castro Neto and B. A. Jones, Phys. Rev. B 62, 14975 (2000) and references there in.
- [19] W. Nolting et al., Z. Phys. B **96**, 357 (1995).
- [20] E. Yu. Tsymbal and D. G. Pettifor, J. Phys.:Condens. Matter 9, L411 (1997).
- [21] We expect that a finite s-electron interaction, $U_s > 0$, will lead only to a slight renormalization of $T_{\rm K}$. Recently, an Anderson impurity embedded in a correlated host (without magnetic order) was studied [22, 23], and an enhancement of the Kondo scale $T_{\rm K}$ was found for weak s-electron interaction, $U_s/D < 1$, whereas for intermediate and strong interactions, $T_{\rm K}$ was found to be reduced.
- [22] T. Schork, P. Fulde, Phys. Rev. B 50, 1345 (1994); G.
 Khaliullin and P. Fulde, Phys. Rev. B 52, 9514 (1995).
- [23] W. Hofstetter et al., Phys. Rev. Lett. 84, 4417 (2000).
- [24] A lowest-order perturbation calculation shows that for $|U + \epsilon_{\rm d}|$, $|\epsilon_{\rm d}| \ll D_{\sigma}$ the spin splitting of the bottom of the band ΔD (for s-bands we consider no spin splitting for the band top) generates an effective local magnetic field $B_{\rm Stoner} \approx \pi \Gamma/2 \ln(1 \Delta D/D_{\uparrow})$. To account for this in our model of Eq. (2), one could renormalize P (or else use a shifted compensation field $B_{\rm comp}$ below).
- [25] K. Yosida, Theory of Magnetism, Springer (1996).
- [26] A similar splitting was predicted [D. Boese, et al., Phys. Rev. B **64**, 125309 (2001); W. Hofstetter, cond-mat/0104297] for a two-level QD system which, in a special situation, can be mapped on our model.
- [27] For certain special (non-generic) cases, the spin up and down level positions and hence also the Kondo resonance does not split up in two, even for $P \neq 0$: this occurs, for example, at the symmetry point $\epsilon_d = -U/2$ of the Anderson model, provided the bands are also symmetric around ε_F (but this is unrealistic for ferromagnets).
- [28] A poor man's scaling analysis [12] yields a renormalization $\delta \epsilon_{\mathrm{d}\sigma}$ of the level $\epsilon_{\mathrm{d}\sigma}$ proportional to $\Gamma_{\overline{\sigma}}$, i.e., $B_{\mathrm{comp}} \propto (\delta \epsilon_{\mathrm{d}\uparrow} \delta \epsilon_{\mathrm{d}\downarrow}) \propto (\Gamma_{\downarrow} \Gamma_{\uparrow}) = -P\Gamma$.
- [29] D. C. Langreth, Phys. Rev. 150, 516 (1966).
- [30] Our NRG confirms Eq. (4) for arbitrary P and B.

L. I. Glazman and M. E. Raikh, JETP Lett. 47, 452 (1988); T. K. Ng and P. A. Lee, Phys. Rev. Lett. 61, 1768 (1988).

^[2] D. Goldhaber-Gordon et al., Nature **391**, 156 (1998).

 ^[3] S. M. Cronenwett et al., Science 281, 540 (1998); F. Simmel et al., Phys. Rev. Lett. 83, 804 (1999); J. Schmid et al., Phys. Rev. Lett. 84, 5824 (2000); W. G. van der Wiel et al., Science 289, 2105 (2000).