Pretransitional phenomena in dilute crystals with first-order phase transition

P. N. Timonin*

Physics Research Institute at Rostov State University 344090, Rostov - on - Don, Russia

(Dated: February 7, 2020)

Pretransitional phenomena at first-order phase transition in crystals diluted by 'neutral' impurities (analogue of nonmagnetic atoms in dilute magnets) are considered. It is shown that field dependence of order parameter becomes nonanalytical in the stability region of the ordered phase, while smeared jumps of thermodynamic parameters and anomalous (non-exponential) relaxation appear near transition temperature of pure crystal.

PACS numbers: 64.60.-i, 05.70.Jk

In last decades the intensive theoretical and experimental studies of phase transitions in disordered crystals reveal the drastic influence which frozen impurities and defects may have on the transition. They can also give rise to a variety of pretransitional phenomena. Such phenomena were first considered by Griffiths¹ in the framework of dilute Ising model, which has empty sites (without spins) randomly scattered over lattice. He had shown that in such dilute magnet the transition is preceded by specific phase, where magnetization is nonanalytical function of magnetic field.

While the bulk transition in this model takes place in the (infinite) percolation cluster of spins, the nature of pretransitional effects lies in the properties of finite magnetic clusters. Due to the short-range (nearest neighbor) interaction between spins all magnetic clusters are completely independent and give additive contributions to the thermodynamic parameters of the system. Immediately below the transition point of pure magnet large clusters begin to develop high thermodynamic barriers between the states with all spins up or down. With due reservations these processes can be called 'local phase transitions'. As there is a finite probability to find arbitrarily large magnetic clusters with arbitrarily high barriers this results in nonanalytical field dependence of magnetization 2,3,4 . Also same barriers give rise to anomalously slow nonexponential relaxation of the order parameter in this Griffiths' phase⁵.

Apparently, the existence of Griffiths' phase is not limited to the case of dilute Ising magnet. It should be also present in all crystalline solid solutions near generic second-order phase transition with short-range order parameter interaction, if it takes place in only one component of the solution. The other 'neutral' component(s) of the solution (impurities) is supposed to have no influence on the transition in the 'pure' copmponent being an analogue of nonmagnetic impurities or empty sites in the dilute Ising model. Hereafter we refer to this generalization of the notion of dilute magnet as 'dilute crystal with a (second-order) transition' or just 'dilute crystal'.

While the Griffiths' anomalies near second-order phase transitions are thoroughly investigated, now nothing is known about the pretransitional phenomena near firstorder transitions in such dilute crystals, in spite of the attention paid to a smearing of these transitions by frozen disorder^{6,7,8}. But it is evident that some similar anomalies could exist in this case, as high barriers between ordered and disordered states of finite clusters of pure component also appear above the transition point of pure percolation cluster (if it exists).

Let us consider the first-order transition with onecomponent order parameter φ , described in a pure crystal by the inequilibrium thermodynamic potential, which is even function of φ . We assume that the dimensionless (divided by the temperature T) density of such potential $f(\varphi)$ has a minimum at $\varphi = 0$ for T > 0 and two another minima at $\varphi \pm \varphi_s$ when $T < T_+, T_- < T_+$. Below some $T_0, T_- < T_0 < T_+, f_s \equiv f(\pm \varphi_s)$ becomes less than $f_0 \equiv f(0)$, so the first-order transition takes place at T_0 in pure crystal.

It can be shown that in dilute crystal the dependence of the order parameter on the conjugate field becomes nonanalitycal below T_+ . To do this we consider the contributions to the average order parameter from large compact clusters of pure component with a number of sites much greater than that of neighboring impurities and diameter much larger than order parameter correlation length. Such clusters can be described by the density of (inequilibrium) thermodynamic potential of infinite pure crystal. Their order parameter distribution in the external field h has the form

$$\rho_n\left(\varphi,h\right) = Z_n^{-1} \exp\left[-nf\left(\varphi\right) - nh\varphi\right] \tag{1}$$

$$Z_{n} = \int d\varphi \exp\left[-nf\left(\varphi - nh\varphi\right)\right]$$

where n is number of sites in a cluster. It is easy to obtain the average order parameter for a cluster

$$\langle \varphi \rangle_{T,n} \equiv \int d\varphi \varphi \rho_n \left(\varphi, h\right)$$

in a small field

$$|h| \le \varphi_s / \max(\chi_0, \chi_s),$$

$$\chi_0 = 1/f''(0), \qquad \chi_s = 1/f''(\varphi_s).$$

For n >> 1 at $T_- < T < T_+$ we have

$$\left\langle \varphi\left(h\right)\right\rangle_{T,n} = \frac{\chi_{0}h + 2\varphi_{s}\left(h\right)\left(\chi_{s}/\chi_{0}\right)^{1/2}e^{-n\delta f}\sinh\left(nh\varphi_{s}\right)}{1 + 2\left(\chi_{s}/\chi_{0}\right)^{1/2}e^{-n\delta f}\cosh\left(nh\varphi_{s}\right)}$$
(2)

while at $T < T_{-}$

$$\langle \varphi(h) \rangle_{T,n} = \varphi_s(h).$$
 (3)

Here $\delta f = f_s - f_0$ and

$$\varphi_s(h) = \varphi_s \tanh(nh\varphi_s) + \chi_s h.$$

Using the numbers of compact clusters per site, N(n), one can find the average order parameter

$$\left\langle \varphi \right\rangle = \sum_{n > n_0} N\left(n\right) \left\langle \varphi \right\rangle_{T,n}$$

Here sum is taken over sufficiently large $n > n_0$ for which Eqs. (2) and (3) is valid,

$$n_0 = c\xi_{max}^d \gg 1,$$

 ξ_{max} being the largest of correlation lengths of two phases (in lattice units) when $T_{-} < T < T_{+}$, $c \gg 1$ and d is space dimension.

When $\delta f < 0$ the expression in Eq. (2) has poles in the complex h plane which are close to that of $\tanh(nh\varphi_s)$

$$h_{n,m} \approx \pm (4m+1) \frac{i\pi}{2n\varphi_s}, \ m = 0, 1, \dots$$

They are arbitrarily close to h = 0 for large n. As $N(n) \neq 0$ at all n the expansion of average order parameter in powers of h would not converge at $T < T_0$.

Above T_0 the expression in Eq. (2)has poles at

$$h_{n,m} \approx \pm \delta f / \varphi \pm i (2m+1) \frac{2\pi}{n\varphi_s}, \ m = 0, 1, \dots$$

which are close to $\pm h_c = \pm \delta f/\varphi_s$ for large n. So at $T > T_0$ average order parameter $\langle \varphi(h) \rangle$ is nonanalytical at the fields $h = h_c$ and $h = -h_c$ in which transition in pure crystal takes place. We can also expect that the contributions to $\langle \varphi(h) \rangle$ from sparse clusters would have the similar property at larger h as they have, presumably, smaller φ_s and larger $|\delta f|$ at the same T. Then the field dependence of the average order parameter will be nonanalytical at all $|h| > h_c$.

The 'local transitions' in finite clusters also manifest themselves in the temperature dependence of thermodynamic parameters at h = 0. Consider the compact cluster contribution to the zero field susceptibility

$$\begin{split} \langle \chi \rangle &= \sum_{n > n_0} N\left(n\right) \langle \chi \rangle_{T,n} \\ \langle \chi \rangle_{T,n} &\equiv n \left[\left\langle \varphi^2 \right\rangle_{T,n} - \left\langle \varphi \right\rangle_{T,n}^2 \right] \end{split}$$

At $T_{-} < T < T_{+}$ we have

$$\langle \chi \rangle_{T,n} \approx \frac{\chi_0^{3/2} e^{n\delta f} + 2\chi_s^{3/2} + 2n^2 \varphi_s^2 \chi_s^{1/2}}{\chi_0^{1/2} e^{n\delta f} + 2\chi_s^{1/2}}, \qquad (4)$$

N(n) is given by the general formulae⁹,

$$N(n) = \sum_{s} g(n,s)p^{n} \left(1-p\right)^{s}$$

where 1-p is the concentration of impurities, and g(n, s) is the number of pure clusters with n sites and s neighboring impurities. The compactness of clusters considered can be accounted for by restriction of the sum to $s \ll n$. In the following we will use the lower estimate for N(n),

$$N(n) \approx p^n \left(1 - p\right)^{s_{\min}(n,d)} \tag{5}$$

$$s_{\min}(n,d) = e_d n^{(d-1)/d}, \ e_d = \left[2\pi^{d/2} d^{d-1} / \Gamma(d/2)\right]^{1/d}$$

with $s_{\min}(n, d)$ being the area of d - dimensional sphere, which bounds the volume n. According to Ref. (10) Eq. (5) describe correctly the main features of large nbehavior of N(n) either for $p \to 1$ or $p \to 0$ except for a change of p to some p' > p when $p \to 0$ (see Ref. (4)).

Near T_0 , when the condition

$$n_0^{-1} \ll |\delta f| \ll \min\left\{\ln\left(\frac{1}{p}\right), e_d^{d/(d-1)} \ln^{d/(d-1)}\left(\frac{1}{1-p}\right)\right\}$$

can be fulfilled, we get from Eqs. (4), (5)

$$\left\langle \chi \right\rangle = \begin{cases} \nu_0 \chi_0, \ T > T_0 + \delta T \\ \nu_0 \chi_s + \nu_2 \varphi_s^2, \ T < T_0 - \delta T \\ \nu_k \equiv \sum_{n > n_0} n^k N\left(n\right) \end{cases}$$

Here δT is defined by the condition $n_0 |\delta f| \gg 1$, so

$$\delta T \sim T_0 / n_0 \delta S,$$

where δS is entropy jump at the transition in pure crystal. Thus near transition point of pure crystal there is slightly smeared jump in the zero field susceptibility.

Also near T_0 the average entropy of large compact clusters,

$$\langle S \rangle = \frac{\partial}{\partial T} \sum_{n > n_0} N(n) T \ln Z_n,$$

mimics the behavior of pure crystal entropy with prefactor ν_0 .

Let us also consider the average modulus of the order parameter, $\langle |\varphi| \rangle$. This quantity can be interpreted as measure of the local order and could be observed in local experiments such as NMR. In pure (infinite) crystal it becomes nonzero below T_0 simultaneously with $\langle \varphi \rangle$. In dilute crystal $\langle |\varphi| \rangle \neq 0$ at all T. Indeed, for the compact clusters we have

$$\langle |\varphi| \rangle_{T,n} \approx (8\chi_0/\pi n)^{1/2}, \ T > T_+ \langle |\varphi| \rangle_{T,n} \approx \varphi_s, \ T < T_-$$

while at $T_{-} < T < T_{+}$

$$\langle |\varphi| \rangle_{T,n} = \frac{(8\chi_0/\pi n)^{1/2} + 2\varphi_s \left(\chi_s/\chi_0\right)^{1/2} e^{-n\delta f}}{1 + 2\left(\chi_s/\chi_0\right)^{1/2} e^{-n\delta f}} \quad (6)$$

Thus in dilute crystal $\langle |\varphi| \rangle = \sum_{n > n_0} N(n) \langle |\varphi| \rangle_{T,n}$ has small nonzero value at all temperatures above the transition point.

We cannot describe in the same manner the (additive) contributions of the sparse clusters with $s \geq n$ to the thermodynamic parameters. Yet we may suppose that they undergo similar 'local transitions', but at temperatures lower than T_0 due to the lower average coordination number. Then continuous sequence of these transitions below T_0 will result in steep increase of $\langle |\varphi| \rangle$ and decrease of $\langle S \rangle$ at $T < T_0$ for all concentrations p, while the bulk transition takes place at lower temperature (less than pT_0^{11}) or vanishes altogether below percolation threshold.

We can also show that in the region where phases coexist large compact clusters have anomalously slow dynamics with broad spectrum of relaxation times. Here the most slow relaxation process is the decay of metastable state. In case of one-component nonconserved order parameter it normally proceeds via nucleation and growth of droplets of stable phase^{12,13}. As the nucleation barrier must be overcome the time it takes is

$$\tau_c = \tau_1 \exp \Delta,$$
$$\Delta = \max \left(dl^{d-1}\sigma - l^d \left| \delta f \right| \right) = \sigma l_c^{d-1}$$

Here σ is dimensionless interphase surface tension (divided by T and multiplied by the area of unit cell face), l_c is the critical droplet size,

$$l_c = (d-1)\,\sigma/\left|\delta f\right|$$

and τ_1 is the time inverse proportional to the droplet growth rate¹².

The decay time τ_c does not depend on system size, which is implicitly assumed to be much greater than l_c . Yet for finite clusters with size less than that of critical droplet, $n < n_c \equiv l_c^d$, the above relaxation process transforms into the sweeping of domain wall through a cluster. Then size-dependent barrier,

$$\Delta_n = n^{(d-1)/d} \sigma - n \left| \delta f \right| / 2, \tag{7}$$

should be overcome, which takes the time

$$\tau_n = \tau_1 \exp \Delta_n. \tag{8}$$

Far from T_0 most clusters exhibit the exponential longtime relaxation with single relaxation time τ_c . When $T \to T_0$, $l_c \to \infty$ and the large part of clusters have sizedependent relaxation times. Thus in dilute crystal near T_0 the broad spectrum of relaxation times emerges. To describe this quantitatively let us consider a contribution of compact clusters to a long-time asymptotics of average dynamic correlator

$$\delta \langle C(t) \rangle = \sum_{n_0 < n} N(n) C_n(t),$$

$$C_{n}(t) = \langle \varphi(t) \varphi(0) \rangle_{T,n} - \langle \varphi \rangle_{T,n}^{2}.$$

Here $C_n(t)$ describes relaxation in *n*-site compact cluster and according to above consideration we have for $t \to \infty$

$$C_{n}(t) = \begin{cases} A_{n} \exp\left(-t/\tau_{c}\right), n > n_{c} \\ A_{n} \exp\left(-t/\tau_{n}\right), n < n_{c} \end{cases}$$

Thus near T_0 when $n_c > n_0$

$$\delta \langle C(t) \rangle = \langle A \rangle \exp(-t/\tau_c) + B(t)$$

$$\langle A \rangle = \sum_{n_c < n} N(n) A_n$$

$$B(t) = \sum_{n_0 < n < n_c} N(n) A_n \exp\left(-t/\tau_n\right)$$
(9)

Assuming A_n to be a slow varying function of n (slower than exp(an)), we can use the steepest descend method to find time dependence of B(t) for large t. From Eqs. (7)-(9) it follows that at $t >> \tau_c$ clusters with $n_* = (2/d)^d n_c$, having the largest barrier for a sweeping domain wall, give main contribution to B(t),

$$B(t) \sim \exp\left(-t/\tau_c'\right), \ t \gg \tau_c$$

$$\tau_c' = \tau_1 \exp\left[\left(\sigma/d\right) \left(2l_c/d\right)^{d-1}\right].$$

When

$$\tau_0 < t \ll \tau_c, \tag{10}$$

$$\tau_0 \equiv \tau_1 \sigma^{-1} \exp\left(\sigma n_0^{(d-1)/d}\right)$$

the main contribution to B(t) give clusters of size $n_* = \left[\sigma^{-1}\ln\left(\sigma t/\tau_1\right)\right]^{d/(d-1)}$ and we get

$$B(t) \sim \left(\frac{\tau_1}{t}\right)^{\lambda(p)} \exp\left\{\ln\left(p\right) \left[\sigma^{-1}\ln\left(\sigma t/\tau_1\right)\right]^{d/(d-1)}\right\}\right)$$
(11)
$$\lambda(p) = -e_d \sigma^{-1} \ln\left(1-p\right).$$

Exponential term in (11) falls slower than pure exponent but faster than all powers of 1/t. Yet for p close to 1 its variation can be small as compared to that of prefactor in all region (10) and this term should be dropped leaving only power-law relaxation. This could happen when

$$l_p \equiv \frac{\lambda(p)\sigma}{\ln(1/p)} = e_d \frac{\ln(1-p)}{\ln p} >> 1$$

Then, not very close to T_0 , for $l_c \ll l_p$ we have power-law relaxation in the interval (10) described solely by prefactor in (9), while in the immediate vicinity of T_0 for $l_c \gg l_p$ this prefactor can be dropped in (11).

Thus near T_0 there appears time interval (10) with non-exponential relaxation which expands to infinity right at T_0 . Owing to the common relaxation mechanism, same Eq. (11) holds for long time asymptotic in original Griffiths' phase near second-order transition with onecomponent nonconserved order parameter. It was first derived in Ref. 5 without power-law prefactor since author used the estimate $N(n) \sim p^n$, which does not hold for p close to 1.

Sparse pure clusters with $s \ge n$ would also exhibit nonexponential relaxation of a similar origin. We cannot give an exact description of their contributions to the dynamic correlator. Yet at every temperature below T_0 a 'local transition' takes place in some specific sort of sparse pure clusters and their relaxation would dominate the infinite time asymptotic. So the above results for compact clusters describe the order parameter relaxation

- * Electronic address: timonin@aaanet.ru
- ¹ R. B. Griffiths, Phys. Rev. Lett. **23**, 17 (1969).
- ² A. B. Harris, Phys. Rev. B **12**, 203 (1975).
- ³ M. M. Bakri and D. Stauffer, Phys. Rev. B **14**, 4215 (1976).
- ⁴ M. Schwartz, Phys. Rev. B **18**, 2364 (1978).
- ⁵ A. J. Bray, Phys. Rev. Lett **60**, 720 (1988).
- ⁶ Y. Imry and M. Wortis, Phys. Rev. B **19**, 3580 (1979).
- ⁷ J. Cardy and J. L. Jacobsen, Phys. Rev. Lett **79**, 4063 (1997).
- ⁸ C. Chatelain, B. Berche, W. Janke, and P.-E. Berche, cond-mat/0103377.

at $t \to \infty$ in close vicinity of T_0 only, while far below it relaxation of sparse clusters must be considered.

Here we may also note that quite similar dynamics will appear above the isostructural first-order transitions (of the liquid-vapor type), which is described by the potential with two minima, one at $\varphi = \varphi_0$ and another at $\varphi = \varphi_s$, corresponding to high- and low-temperature phases. Also the smeared jumps of the susceptibility and the entropy appear at T_0 in this case, while the field dependence of the average order parameter becomes nonanalytical near T_0 at $h_c = \delta f / (\varphi_s - \varphi_0)$.

To conclude, the present qualitative considerations show that pretransitional phenomena in dilute crystal with first-order phase transition has features rather similar to that of the original Griffiths' phase. They can be generalized to the first-order transitions with multicomponent order parameters and other types of random media, which also have cluster size distributions spreading up to $n = \infty$. So one may expect that analogous phenomena could be observed, for example, near melting points or the liquid crystal transitions in the substances confined in porous media.

Acknowledgments

This work was made under support from INTAS, grant 2001-0826. I gratefully acknowledge helpful and illuminating discussions with V. P. Sakhnenko.

- ⁹ D. Stauffer and A. Aharony, *Introduction into percolation theory* (Taylor and Francis, 1992).
- ¹⁰ H. Kesten, *Percolation theory for mathematicians* (Birkhauser, 1982).
- ¹¹ R. B. Stinchcomb, in Phase Transition and Critical Phenomena, eds. C. Domb, J. L. Lebowitz (Academic Press, 1983).
- ¹² M. Grant and J. D. Gunton, Phys. Rev. B **32**, 7299 (1985).
- ¹³ A. J. Bray, Adv. in Physics **43**, 357 (1994).