

# Larmor clock and the local density of states in a quantum wire

P. Singha Deo

*S. N. Bose National Centre for Basic Sciences, JD Block, Sector III, Salt Lake City, Kolkata 98, India.*

(Dated: October 28, 2018)

Buttiker-Thomas-Pretre (BTP) [Z. Phys B **94**, 133 (1994)] proposed that the concepts behind the Larmor clock tell us that it is possible to define exactly the local density of states (LDOS) in terms of the scattering matrix. However, we take into account evanescent modes and show that for an impurity in a quantum wire, this is in principle not exactly true. We also prove that the Wigner delay time gives correct superluminal times at the Fano resonances, in spite of the fact that the stationary phase approximation is not valid there.

PACS numbers:

Electronic states in a one-dimensional (1D) ring with an Aharonov-Bohm flux piercing the ring exhibit the parity effect according to which if the magnetization of the ring with  $N$  electrons is diamagnetic (or paramagnetic) then the magnetization with  $N + 1$  electrons is paramagnetic (or diamagnetic). Leggett conjectured that this is true even in presence of interaction between electrons as well as defects or disorder in the ring [1]. Now suppose another quantum system  $S$  is coupled to the 1D ring  $R$  such that the states of  $S$  can leak into  $R$  and become flux dependent. Ref. [2] shows that the states of the combined system ( $S+R$ ) does not always have the property of reversing the magnetization with the addition of a single electron. It was found that the electrons in the ring undergo usual phase changes associated with their quantum mechanical motion. These phase changes are i) Aharonov-Bohm phase, ii) statistical phase, which means electrons being fermion, acquire a phase change of  $\pi$  when they cross each other and iii) relative phase change due to the wave like property of the electrons, that depend on their wave vector or their kinetic energy. It was also shown [2] that apart from these phase changes, there are discontinuous phase changes by  $\pi$ , at the zeroes of the Fano resonances (say at energy  $E_0$ ) that will be there when the ring (with the system  $S$  attached to it) is severed at a point and two leads are attached to the two broken ends [2]. If Fermi energy of  $N$  body state is below  $E_0$ , and that of the  $N + 1$  body state is above  $E_0$ , then the magnetization of the  $N$  body state and the  $N + 1$  body state is the same, and otherwise opposite. Thus Leggett's arguments can be generalized to systems that are of the form  $S + R$ , with the conclusion that this new discontinuous phase change is a new phase different from the ones mentioned in i), ii) or iii) [2]. While phase of scattering matrix elements for Breit-Wigner resonance is often discussed in text books, that for Fano resonances was not known until very recently.

In an experiment [3], a quantum dot was coupled to a ring ( $S+R$  is now a dot+ring) and the conductance oscillations of the system with an Aharonov-Bohm flux was measured. Yeyati and Buttiker [4] tries to interpret the conductance oscillations in terms of the flux dependence (or magnetization) of the electronic levels of the combined system (dot + ring) using the Friedel sum

rule (FSR). The phase change at the resonances of the dot could be roughly understood but the phase change between the resonances of the dot could not be understood. Other works tried to assign this in between resonance phase change to processes like charge decapture (the system throws away unit charge) [5]. Ref. [6] and [7] predicted that the resonances are actually Fano resonances and the phase change between the resonances can be explained by the new phase at the transmission zeroes and it will be an abrupt drop. It was confirmed in a later experiment that this phase drop occur over an energy scale, that is much smaller than any energy scale present in the system [8]. Finite width of the leads and evanescent modes has to be considered to explain [7] the abrupt phase changes occurring between each consecutive resonances. Ref. [7] considers there is a pole between the zeroes and that is where a charge is captured and phase changes smoothly by  $\pi$  in agreement with FSR. But the discontinuous phase drops being a new kind of phase, does not have anything to do with charge capture or decapture. Later on this was proved to be true whenever we have time reversal symmetry [9, 10]. Also the fact that simple 1D calculations will not explain the phase drop between each two consecutive resonances was shown [11].

In fact unitarity is also required to produce such discontinuous phase changes [12]. It was subsequently seen that when an unitary channel continuously evolves into a non-unitarity channel, then along with it, the transmission zeroes evolve into minima (difference with Breit-Wigner resonance is that the minima has a complex zero, that is for a complex value of incident energy, transmission amplitude is zero), and discontinuous phase drops evolve into continuous phase drops [13]. As an example, one can consider a two channel quantum wire with a delta potential (see Fig. 1) at the middle of the quantum wire. Since the two channels of the quantum wire are opposite parity states like the ground state and the first excited state of a quantum well, the two channels are decoupled. If an electron is incident from the left in the first channel then amplitude of transition to the second channel i.e.,  $t_{12}$  (state 1 to state 4) or  $r_{12}$  (state 1 to state 2) is zero, making the first channel preserve unitarity. Whenever  $|t_{11}|^2$  has a zero, the phase of  $t_{11}$  drops

(or decreases) discontinuously by  $\pi$ . Now if the delta potential is shifted slightly from the center of the quantum wire, then parity is no longer a good quantum state and  $t_{12}$  etc are no longer zero. So by continuously displacing the delta potential from the center, one can destroy the individual unitarity of each channel and thus make the discontinuous phase drops continuously evolve into fast continuous phase drops. These phase drops can be seen in Fig. 2 at an energy  $Ea \approx 85$ . Even dephasing can result in the loss of unitarity [12] and can explain the small width of the phase drops observed in the experiment. For the two propagating channel case,

$$\begin{aligned} \frac{\partial \theta_f}{\partial E} = & \frac{1}{2\pi} [|r_{11}|^2 \frac{\partial \arg(r_{11})}{\partial E} + |r_{22}|^2 \frac{\partial \arg(r_{22})}{\partial E} \\ & + 4|r_{12}|^2 \frac{\partial \arg(r_{12})}{\partial E} + |t_{11}|^2 \frac{\partial \arg(t_{11})}{\partial E} + |t_{22}|^2 \frac{\partial \arg(t_{22})}{\partial E}] \end{aligned} \quad (1)$$

For the delta potential at the center of the wire, the discontinuous drops in  $\arg(t_{11})$  or  $\arg(t_{22})$  occur when  $|t_{11}|^2=0$  and  $|t_{22}|^2=0$  and so there are no drops in  $\theta_f$ . In that case, obviously the drops in  $\arg(t_{11})$  or  $\arg(t_{22})$  do not have anything to do with  $\theta_f$  and hence charge decapture. But for the  $\delta$  potential off center, since  $|t_{11}|^2$  or  $|t_{22}|^2$  are not zero when there are drops in their arguments, it is not obvious that these drops do not have to do anything with charge decapture. However only after all the simplification, one finds that [13]

$$\frac{\partial \theta_f}{\partial E} = \frac{\partial \arg(r_{\alpha\beta})}{\partial E} \quad (2)$$

That means  $\theta_f$  only depends on the phase of reflection amplitudes and hence is completely independent of the phase drops in the transmission amplitudes. So once again the phase drops do not imply a drop in  $\theta_f$  or charge decapture [13].

One also knows from earlier known results that FSR is valid only in semi-classical regimes. And so Fano resonance being a purely quantum interference phenomenon, one would expect larger violations of FSR at the Fano resonances. Instead what Refs. [14] shows is that FSR is exact at the energy corresponding to the quasi bound state, but there are large violations away from this energy. Refs. [14] also explained that the correctness of FSR at the Fano resonance is due to the fact that there is a quasi bound state here, and hence the self energy due to the leads become minimum and hence its energy derivative becomes 0. This is obviously true for any potential that supports Fano resonances. The exactness of the FSR at the Fano resonance ( $Ea \approx 85$ ) and the deviation away from it is also shown in Fig. 2.

Apart from the DOS we know that time scales associated with a particle crossing a quantum mechanical potential can also be determined from the scattering phase shifts. For example, in the stationary phase approximation (i.e., phase shifts do not strongly depend on energy)  $\frac{d}{dE} \arg(t_{\alpha\beta})$  gives the Wigner delay time (WDT)

for the particle to be transmitted from state  $\alpha$  to state  $\beta$ . A negative slope in  $\arg(t_{\alpha\beta})$ , like that in the solid and dash-dotted curve at  $Ea \approx 85$  in Fig. 2, means superluminality, i.e., the particle can travel faster than light across the potential, according to the WDT. However, once again, since Fano resonance is a quantum interference effect, dispersion will be very strong and stationary phase approximation will not hold good at the Fano resonances, and one cannot be sure if these negative slopes actually correspond to super-luminality. To be sure of superluminality, established theories say that one should see if we get negative delay times from the Buttiker-Thomas-Pretre (BTP) formalism. They proposed that the Larmor precession time can be determined exactly from the scattering matrix and give the correct local delay times in all regimes [15] and delay time can be determined by integrating the local delay times. It gives

$$\nu(\alpha, E, r, \beta) = -\frac{1}{4\pi i} \text{Tr} [S_{\alpha\beta}^\dagger \frac{\partial S_{\alpha\beta}^\dagger}{\partial V(r)} - HC] \quad (3)$$

where  $\nu(\alpha, E, r, \beta)$  is proportional to the time spent by the particle going from state  $\alpha$  to state  $\beta$ , while encountering the potential at  $r$ . If we sum it up for all  $\alpha$  and  $\beta$  then it should give the LDOS at  $r$ , exactly [15, 16]. Originally, it was derived by considering the effect of a small magnetic field on the outgoing spin wave function. But more generally, for any potential, to obtain the LDOS at  $r$ , we have to create a  $\delta$  function potential like local perturbation at  $r$ , and see the change in the scattering matrix  $S$  of the entire system. Also, the scattering matrix  $S$  of an extended system can be written as [17]  $S = S_1 \otimes S_2 \otimes S_3$ , where  $S_2$  is the scattering matrix of an infinitesimal region at  $r$ ,  $S_1$  and  $S_3$  are the scattering matrices of the regions to the left and right of  $r$ , respectively. Hence  $S_2$  is the scattering matrix of a  $\delta$  function potential that we have to further perturb infinitesimally to see the changes in  $S_2$ . All the change in  $S$  will be due to this change in  $S_2$ . Thus complications of the BTP formula will depend on its complications for a  $\delta$  function potential.

The derivation of Eq. 3 assumes that a small perturbation to the actual system, allows us to expand a scattering matrix element as

$$\begin{aligned} S_{\alpha\beta}^\pm(E, V(x_i, y_i) \mp \delta V(x_i, y_i)) &= S_{\alpha\beta}(E, V(x_i, y_i)) \\ &\mp \int dx'_i dy'_i [\delta S_{\alpha\beta}(E, V(x'_i, y'_i)) / \delta V(x'_i, y'_i)] \delta V(x'_i, y'_i) + \dots \end{aligned} \quad (4)$$

If we make a substitution of the type

$$- \int dx'_i dy'_i \frac{\delta}{\delta V(x'_i, y'_i)} \rightarrow \frac{d}{dE} \quad (5)$$

then we derive FSR. But this substitution being an approximation, the FSR is also approximate.

In this work, we will show the expansion in 4 is not defined at all energies and thus the concept of Larmor clock fails. That means LDOS cannot be defined in terms of S matrix. We shall also show that the expansion in 4 is valid at the Fano resonance and also explicit calculations of LDOS prove the correctness of BTP formula only at the Fano resonance. Then we will show that WDT give the correct delay time at the Fano resonances in spite of all non-stationary behavior ( a very counter intuitive result) and in agreement with the delay time calculated from Larmor precession. If it predicts superluminality then such superluminality can be observed quantitatively.

The first result of this work (that BTP formula is correct only at the Fano resonance), imply the LHS of 5, operating on  $S_{\alpha\beta}$ , gives the correct delay time at the Fano resonance. The substitution in 5, is exact at the Fano resonance follows from earlier results [14] that the FSR is exact at Fano resonance. So the RHS in 5, operating on  $S_{\alpha\beta}$  will give the same quantity as the delay time calculated from BTP formalism, and that means WDT gives the correct delay time at the Fano resonance (our second result).

For the system in Fig. 1 [18],

$$t_{\alpha\beta} = r_{\alpha\beta} = -\frac{i\Gamma_{\alpha\beta}}{2d\sqrt{k_\alpha k_\beta}} \quad (6)$$

where for the transmissions,  $\alpha \neq \beta$  and

$$t_{\alpha\alpha} = r_{\alpha\alpha} + 1 \quad (7)$$

$$d = 1 + \sum_e \frac{\Gamma_{ee}}{2\kappa_e} + i \sum_\alpha \frac{\Gamma_{\alpha\alpha}}{2k_\alpha} \quad (8)$$

For a quantum wire with hard wall confinement,  $\Gamma_{nn} = (2m_e\gamma/\hbar^2)\sin^2[\frac{n\pi}{w}(y_i + w/2)]$ ,  $m_e$  is electron mass,  $k_n = \sqrt{(2m_e/\hbar^2)(E - E_n)}$ ,  $\kappa_n = \sqrt{(2m_e/\hbar^2)(E_n - E)}$ ,  $E_n = (\hbar^2/2m_e)(n^2\pi^2/w^2)$ ,  $E$  is incident energy,  $\sum_e$  is sum over all the evanescent modes.

*Explicit calculations of LDOS:* We derive LDOS explicitly from the internal wavefunctions for unit incident flux by using [10]

$$\rho(E, x = 0, y = y_i) = \sum_\alpha \frac{2}{\hbar v_\alpha} |\psi_{nk}(x = 0, y = y_i)|^2 \quad (9)$$

$\psi_{nk}(x, y)$  can be taken from Ref. [18] and can be further simplified to give

$$\rho(E, 0, y_i) = \sum_\alpha \frac{2}{\hbar v_\alpha} \left| \sum_m t_{\alpha m} \sin\left(\frac{m\pi}{w}(y_i + w/2)\right) \right|^2 \quad (10)$$

where  $v_\alpha = \hbar k_\alpha / m_e$  and [14],

$$t_{\alpha e} = \frac{-\frac{\Gamma_{\alpha e}}{2\kappa_e}}{1 + \sum_{e>1} \frac{\Gamma_{ee}}{2\kappa_e} + i \sum_{\alpha\alpha} \frac{\Gamma_{\alpha\alpha}}{2k_\alpha}} \quad (11)$$

In Eq. 10, sum over  $\alpha$  runs over the propagating modes only while that for  $m$  is for all modes.

*LDOS from BTP formula:* We consider two propagating channels to present our results but we have checked that the results are the same for any number of channels including just one channel. For this case, Eq. 3 gives

$$\nu(E, 0, y_i) = 2[r_{11}^* r'_{11} + r_{22}^* r'_{22} + 4r_{12}^* r'_{12} + t_{11}^* t'_{11} + t_{22}^* t'_{22} - HC] \quad (12)$$

where primes mean derivatives wrt  $\gamma$ .

First of all, note that the role of evanescent modes on the scattering matrix elements only is that it renormalizes the strength of the potential i.e.,  $\gamma$  according to the following relation.

$$\gamma \rightarrow \gamma' = \gamma \left(1 + \sum_e \frac{\Gamma_{ee}}{2\kappa_e}\right) \quad (13)$$

Note that the sum is not a converging series. It diverges as  $\log[N]$  as  $N \rightarrow \infty$  [19]. Here  $N$  is the total number of evanescent modes. And so derivative w.r.t  $\gamma$  will not exist for any arbitrary number of modes and the BTP formula will be undefined. But if we truncate the series then what happens to the BTP formula? For example if we truncate the series at the 3rd term, that is we are considering 2 evanescent modes only then we show in Fig. 3 that the BTP formula holds good. To calculate the derivative we have taken  $\delta\gamma=0.001$ . However this kind of agreement does not occur for example when we consider 5000 evanescent modes with  $\delta\gamma=0.001$  (see Fig. 4). This can be cross checked by using Eq. 13. Note that now although the two curves do not coincide with each other exactly, they do coincide exactly at the energy corresponding to the bound state (i.e., where both curves peak). This is because it can be shown that at the bound state  $(1 + \sum_e \frac{\Gamma_{ee}}{2\kappa_e}) = 0$ , irrespective of the number of evanescent modes. Of course one can take a smaller value of  $\gamma$  and get a better agreement between the two curves, but then again the same disagreement will be there if more evanescent modes are considered. Since the sum is not a converging sum, this will be a never ending story. One can see that  $\frac{\partial S_{\alpha\beta}}{\partial\gamma} = \frac{\partial S_{\alpha\beta}}{\partial\gamma'} \frac{\partial\gamma'}{\partial\gamma}$  do not exist for an infinite number of evanescent modes as  $\frac{\partial\gamma'}{\partial\gamma}$  diverge as  $\log[N]$ . If we want to integrate LDOS over an extended region then one has to take  $\delta$  function potential like perturbations at many many places of the region and sum the changes they produce on the S matrix. The error will be cumulative.

However, since the BTP formula is exact at the energy corresponding to the Fano resonance, the WDT give the correct superluminal time at the phase drops associated with the Fano resonance. So far, experiments to observe superluminality, only considered systems with Breit Wigner type resonances [20]. Features of superluminality in case of these Fano type resonances should be explored in the future. Some of its features that we have checked are that to enhance superluminality, we have to tune just one parameter, that is the coupling between the bound

state and the continuum. Also, this enhancement is not at the cost of decreasing the transmission probability, as is the case with Breit-Wigner resonance.

Fano resonance is a purely quantum effect that has no classical or semi-classical analogue. As can be seen from Fig. 2, at the energy corresponding to the Fano resonance, the scattering phase shifts are extremely energy dependent. So are the scattering probabilities. At the Fano resonance there is no classical path for reflection but when we solve the Schrodinger equation we find that reflection dominates. All these make it clear that it is a purely quantum effect. Yet, we have shown that semi-classical laws like Friedel sum rule and Wigner delay time are applicable here exactly. This motivated us to study how a wave packet is dispersed by typical potentials like the stub and the delta function potential in a quantum wire, that supports Fano resonances. We took two wave packets (Gaussian or square), incident on such a potential from the two sides, one being left moving and the other being right moving. Each of them will get strongly scattered, but after scattering we will get two wave packets, one of them left moving and the other being right moving. Exactly at the Fano resonance, these two wave packets after scattering, are similar to the incident wave packets. This normally happens in the case of solitonic solutions of non-linear differential equations. Hence such

behavior for solutions of Schrodinger equation is quite novel. Away from the Fano resonances, the wave packets after scattering is different from the incident wave packets.

The BTP formalism is very crucial to understand mesoscopic transport beyond the Landauer conductance formula, that is beyond the linear and DC response. So far it has been verified for simple 1D systems [16]. We have shown that BTP formalism is not exact in Quasi-one-dimension (Q1D) due to the presence of evanescent modes. These evanescent modes make the scattering matrix singular in nature and the series expansions in Eq. 5. required to show the equality between  $\rho$  and  $\nu$  breaks down. If there are only a few evanescent modes in the system, then the formula may be acceptable for practical purposes. For extended potentials and large number of evanescent modes, it will not be practical. It definitely cannot be used as a definition for DOS. DOS has to be defined in terms of the internal wave function and Hamiltonian. Scattering matrix will not contain all the information. We have also proved that in spite of all non-stationary phase effects, WDT ( $\frac{d}{dE}arg(t_{\alpha\alpha})$ ) correctly give the superluminal times and there is superluminality at the energy where the phase drops of transmission amplitudes occur (for example at  $Ea=85$  in Fig. 2) that can be detected experimentally.

- 
- [1] A.J. Leggett, in Granular Nano-Electronics, Vol.251 of NATO Advanced Studies Institute, Series B: Physics, edited by D.K. Ferry, J.R. Barker and C. Jacoboni (Plenum, New York, 1991), p.297
- [2] P. S. Deo, Phys. Rev. B **53**, 15447(1996)
- [3] A.Yacoby et al., Phys. Rev. Lett. **74**, 4047 (1995)
- [4] L.Yeyati and M.Büttiker, Phys. Rev.B **52**, R14360 (1995)
- [5] G. Hackenbroich, W.D. Heiss and H.A. Weidenmuller, Phys. Rev. Lett. **79** 127 (1997) and references therein; P.G. Silvestrov and Y. Imry, Phys. Rev. Lett. **85**, 2565 (2000).
- [6] P. Singha Deo and A.M.Jayannavar Mod. Phys. Lett. B **10**, 787 (1996)
- [7] P. Singha Deo, Physica E **1**, 301 (1997); P. Singha Deo, Solid State Commun. **107**, 69 (1998)
- [8] R. Schuster et al., Nature(London) **385**, 417 (1997)
- [9] H.W.Lee, Phys. Rev. Lett. **82**, 2358 (1999)
- [10] T.Taniguchi and M.Buttiker, Phys. Rev.B **60**, 13814 (1999)
- [11] A.L. Yeyati and M. Büttiker, PRB, **62**, 7307 (2000)
- [12] P. Singha Deo, cond-mat/0005123.
- [13] P. S. Deo, S. Bandopadhyay and S. Das, IJMP B, **16**, 2247 (2002)
- [14] S. Bandopadhyay and P. S. Deo, Phys. Rev. B (in press); PhD thesis of S. Bandopadhyay (under preparation).
- [15] M. Büttiker, H. Thomas, and A Pretre, Z. Phys B **94**, 133 (1994); M. Büttiker, Pramana Journal of Physics,**58**, 241 (2002).
- [16] T. Gramespacher and M. Buttiker, Phys. Rev. B **56**, 13026 (1997); V. Gasparian, T. Christen and M. Buttiker, Phys. Rev. A **54** 4022 (1996).
- [17] S. Datta, Electronic transport in mesoscopic systems, Cambridge university press, 1995, page - 125.
- [18] P.F.Bagwell, Phys. Rev.B **41**, 10354(1990)
- [19] D. Boese, M. Lischka, and L.E. Reichl, Phys. Rev. B **61**, 5632 (2000).
- [20] L. J. Wang et al, Nature **406**, 277 (2000); A. M. Steinberg, P. G. Kwiat and R. Y. Chiao, Phys. Rev. Lett. **71**, 708 (1993); S. Chu and S. Wong, Phys. Rev. Lett. **48**, 738 (1982).

### Figure Captions

Fig. 1. Here we show a quantum wire of width  $w$  with a delta function potential  $V(x, y) = \gamma\delta(x)\delta(y - y_i)$  situated at  $\times$ . We consider scattering effects when the incident electron is from the left. The sub-bands on the left of the impurity is denoted as 1 for the first mode and 2 for the second mode. Similarly the sub-bands on the right of the impurity is denoted as 3 for the first mode and 4 for the second mode.

Fig. 2. The dotted curve gives  $G=\pi$  times the change in DOS due to the defect considered in Fig. 1, in units of  $g = (m_e w^2 / \hbar^2)$ . The dashed curve gives  $Q = d\theta_f / dE$  in units of  $q = g$ . The solid curve gives  $arg(t_{11})$  in radians, displaced on the y-axis by 0.1. The dash-dotted curve gives  $arg(t_{22})$  in radians, displaced on the y-axis by 0.8. We have taken  $\gamma = -15$ ,  $y_i=0.45$  and 2 propagating modes along with 2 evanescent modes. The x-axis is energy in units of  $a = w^2$ .

Fig. 3. The system under consideration is shown in

Fig. 1. with two propagating modes. Solid line gives  $H = \rho(E, 0, y_i)$ , that LDOS calculated from internal wavefunction, in units of  $h = (2mw/\hbar^2)$ . Dashed line gives  $J = \nu(E, 0, y_i)$ , i.e., LDOS calculated using BTP formula, in units of  $j = h$ . All parameters are the same as that considered in Fig. 2. The x-axis is energy in units

of  $a = w^2$ .

Fig. 4. Here we consider 5000 evanescent modes and  $\gamma = -1.5$ . Everything else is the same as in Fig. 3.







