Spin-resolved second-order correlation energy of the two-dimensional uniform electron gas

Michael Seidl, Institute of Theoretical Physics, University of Regensburg, D-93040 Regensburg, Germany

Abstract. For the two-dimensional electron gas, the exact high-density limit of the correlation energy is evaluated here numerically for all values of the spin polarization. The result is spin-resolved into $\uparrow\uparrow$, $\uparrow\downarrow$, and $\downarrow\downarrow$ contributions and parametrized analytically. Interaction-strength interpolation yields a simple model (LSD) for the correlation energy at finite densities.

In recent years, two-dimensional (2D) electron systems have become the subject of extensive research [1]. The 2D version of density functional theory (DFT) has proven particularly successful in studying quantum dots [2, 3, 4]. The local spin-density approximation (LSD) of DFT requires the correlation energy of the spin-polarized uniform electron gas. This quantity in 2D is known accurately for a wide range of densities and spin polarizations from fixed-node diffusion Monte Carlo simulations [5]. Its high-density limit is known exactly in terms of six-dimensional momentum-space integrals [6]. Resolved into contributions due to $\uparrow\uparrow$, $\uparrow\downarrow$, and $\downarrow\downarrow$ excitation electron pairs, these integrals are evaluated here numerically. The analytical parametrization of the results, Eqs. (16) and (17) below, is a crucial ingredient for the construction of the spin-resolved correlation energy at finite densities, performed recently for the 3D electron gas [7]. It is also required for studying the magnetic response of the spin-polarized 2D electron gas [8, 9]. Generally, it provides a fundamental test for numerical parametrizations of the correlation energy [5].

In the 2D uniform electron gas, the electrons are moving on a plane at uniform density $\rho = [\pi(r_s a_B)^2]^{-1}$, where $a_B = 0.529$ Å is the Bohr radius and r_s is the dimensionless density parameter (Seitz radius). We consider lowest-energy states with a given spin polarization

$$\zeta \equiv \frac{\rho_{\uparrow} - \rho_{\downarrow}}{\rho} \tag{1}$$

where ρ_{\uparrow} and $\rho_{\downarrow} \equiv \rho - \rho_{\uparrow}$, respectively, are the (uniform) densities of spin-up and spin-down electrons. Including a neutralizing positive background, the total energy per electron is a unique function of the dimensionless parameters r_s and ζ ,

$$e_{tot}(r_s,\zeta) = t_s(r_s,\zeta) + e_x(r_s,\zeta) + e_c(r_s,\zeta). \tag{2}$$

The non-interacting kinetic and exchange energies,

$$t_s(r_s,\zeta) = \frac{1+\zeta^2}{2} \frac{1}{r_s^2}, \qquad e_x(r_s,\zeta) = -\frac{4\sqrt{2}}{3\pi} \frac{(1+\zeta)^{3/2} + (1-\zeta)^{3/2}}{2} \frac{1}{r_s}$$
 (3)

(all energies are given in units of 1 Ha $\equiv e^2/a_B = 27.21$ eV in the following), may be understood as the 0th- and the 1st-order terms of a perturbation expansion for the electron-electron interaction (where r_s turns out to be the expansion parameter).

The remaining *correlation energy* in Eq. (2) appears to have the perturbation (high-density) expansion [10, 11]

$$e_c(r_s,\zeta) = \sum_{n=0}^{\infty} \left[a_n(\zeta) \ln(r_s) + b_n(\zeta) \right] r_s^n \qquad (r_s \ll 1).$$
 (4)

For the 2D electron gas (but not for the 3D one), the first coefficient vanishes, $a_0(\zeta) \equiv 0$. Consequently, the second-order (n=0) term is $e_c^{(2)}(\zeta) \equiv b_0(\zeta)$, representing the high-density $(r_s \to 0)$ limit of $e_c(r_s, \zeta)$. It can be split into an exchange ("2b") and a ring-diagram ("2r") term [6],

$$e_c^{(2)}(\zeta) = e_c^{(2b)} + e_c^{(2r)}(\zeta).$$
 (5)

The exchange term has only equal-spins contributions, $e_c^{(2b)} = e_{c\uparrow\uparrow}^{(2b)}(\zeta) + e_{c\downarrow\downarrow}^{(2b)}(\zeta)$, given by the $\delta_{\sigma_1\sigma_2}$ term of Eq. (14) in Ref. [6] (we choose the k_x axis in the direction of \mathbf{q}),

$$e_{c,\sigma\sigma}^{(2b)}(\zeta) = \frac{1}{8\pi^2} \int_0^\infty \frac{dq}{q} \int_{A[\kappa_{\sigma}(\zeta),q]} d^2k_1 \int_{A[\kappa_{\sigma}(\zeta),q]} d^2k_2 \frac{1}{|q \mathbf{e}_x + \mathbf{k}_1 + \mathbf{k}_2|} \frac{1}{q + k_{1x} + k_{2x}}.$$
 (6)

Here, q, \mathbf{k}_1 , and \mathbf{k}_2 are dimensionless, $\sigma \in \{\uparrow, \downarrow\}$, and the domain of the 2D integrals is

$$A[\kappa, q] \equiv \left\{ \mathbf{k} \in \mathbb{R}^2 \mid |\mathbf{k}| < \kappa, \ |\mathbf{k} + q \, \mathbf{e}_x| > \kappa \right\}, \qquad \kappa_{\sigma}(\zeta) \equiv \left[1 + \operatorname{sgn}(\sigma) \zeta \right]^{1/2}. \tag{7}$$

 $[\kappa_{\sigma}(\zeta)]$ is the Fermi wave vector for spin- σ electrons in units of its value at $\zeta = 0$.] Scaling the integration variables by some constant κ , $q = \kappa Q$ and $\mathbf{k} = \kappa \mathbf{K}$, we have generally

$$\int_0^\infty \frac{dq}{q} \int_{A[\kappa,q]} d^2k \, f(q,\mathbf{k}) = \kappa^2 \int_0^\infty \frac{dQ}{Q} \int_{A[1,Q]} d^2K \, f(\kappa q, \kappa \mathbf{K}). \tag{8}$$

Applying this rule to the integrals in Eq. (6), we find [6]

$$e_{c,\sigma\sigma}^{(2b)}(\zeta) = \left[1 + \operatorname{sgn}(\sigma)\zeta\right]J^{(2b)}.$$
(9)

Consequently [6], the full second-order exchange term $e_c^{(2b)} = e_{c\uparrow\uparrow}^{(2b)}(\zeta) + e_{c\downarrow\downarrow}^{(2b)}(\zeta) \equiv 2J^{(2b)}$ is ζ -independent. A Monte Carlo integration yields

$$J^{(2b)} \equiv e_{c\uparrow\uparrow}^{(2b)}(0) = (57.15 \pm 0.05) \,\text{mHa} \qquad (1 \text{mHa} = 10^{-3} \text{Ha}).$$
 (10)

The ring-diagram term $e_c^{(2r)}(\zeta)$ is the remaining part of expression (14) in Ref. [6], with the contributions

$$e_{c,\sigma_{1}\sigma_{2}}^{(2r)}(\zeta) = -\frac{1}{8\pi^{2}} \int_{0}^{\infty} \frac{dq}{q^{2}} \int_{A[\kappa_{\sigma_{1}}(\zeta),q]} d^{2}k_{1} \int_{A[\kappa_{\sigma_{2}}(\zeta),q]} d^{2}k_{2} \frac{1}{q + k_{1x} + k_{2x}}.$$
 (11)

The equal-spins terms $(\sigma_1 = \sigma_2)$ can be treated in the same way as the integral (6),

$$e_{c,\sigma\sigma}^{(2r)}(\zeta) = -\left[1 + \operatorname{sgn}(\sigma)\zeta\right]J^{(2r)}, \qquad J^{(2r)} = (76.69 \pm 0.03) \,\mathrm{mHa}.$$
 (12)

The only non-trivial ζ -dependence is in the opposite-spins term $e_{c\uparrow\downarrow}^{(2r)}(\zeta) \equiv e_{c\downarrow\uparrow}^{(2r)}(\zeta)$,

$$e_{c\uparrow\downarrow}^{(2r)}(\zeta) = e_{c\uparrow\downarrow}^{(2r)}(0) \left[1 - f(\zeta) \right]. \tag{13}$$

By definition, f(0) = 0, and, since $A[\kappa_{\downarrow}(1), q] = \emptyset$, f(1) = 1. Moreover, $e_{c\uparrow\downarrow}^{(2r)}(0) = -J^{(2r)}$. When the results of a Monte Carlo evaluation of $f(\zeta)$ at different values of ζ are compared with the functions $f_{\alpha}(\zeta) \equiv [(1+\zeta)^{\alpha} + (1-\zeta)^{\alpha} - 2]/(2^{\alpha} - 2)$, particularly good agreement (specially for $\zeta \to 0$ and $\zeta \to 1$) is found in the limit $\alpha \to 1$ (Fig. 1a),

$$f(\zeta) = f_1(\zeta) + \delta f(\zeta),$$
 $f_1(\zeta) \equiv \frac{(1+\zeta)\ln(1+\zeta) + (1-\zeta)\ln(1-\zeta)}{2\ln 2}.$ (14)

[Note that $f_{\alpha}(\zeta)$ also represents the ζ -dependence of t_s ($\alpha = 2$) and e_x ($\alpha = \frac{3}{2}$) in Eq. (3).] The small deviation $\delta f(\zeta)$ is accurately fitted by a polynomial (Fig. 1b)

$$\delta f(\zeta) \approx 0.0636 \,\zeta^2 - 0.1024 \,\zeta^4 + 0.0389 \,\zeta^6 \,.$$
 (15)

The small minimum of $\delta f(\zeta)$ indicated by the numerical data (dots in Fig. 1b) at $\zeta \approx 0.98$ is probably real, since a similar peculiarity is observed for the 3D electron gas (see the inset in Fig. 1 of Ref. [12]).

In summary, the second-order correlation energy $e_c^{(2)}(\zeta)=e_c^{(2b)}+e_c^{(2r)}(\zeta)$ is

$$e_c^{(2)}(\zeta) \equiv e_{c\uparrow\uparrow}^{(2)}(\zeta) + 2e_{c\uparrow\downarrow}^{(2)}(\zeta) + e_{c\downarrow\downarrow}^{(2)}(\zeta) = \left[153.38 f(\zeta) - 192.46\right] \text{ mHa},$$
 (16)

where $f(\zeta)$ is given by Eqs. (14) and (15). The spin resolution is fixed by

$$e_{c\uparrow\uparrow}^{(2)}(\zeta) \equiv e_{c\downarrow\uparrow}^{(2)}(-\zeta) = -(1+\zeta) \times 19.54 \,\text{mHa}.$$
 (17)

 $e_c^{(2)}(\zeta) \equiv e_c(0,\zeta)$ is the high-density limit of the general correlation energy $e_c(r_s,\zeta)$. To illustrate the relevance of this limit for finite densities $(r_s > 0)$, the present result can be used in the interaction-strength interpolation (ISI) of Ref. [13]. This approach does not require the higher-order $(n \geq 1)$ terms of the expansion (4) (which is expected

to have only a finite radius of convergence). Instead, information beyond the second order is taken from the low-density (strong-interaction or Wigner-crystal) limit of the exchange-correlation energy $e_{xc} \equiv e_x + e_c$ (per electron),

$$e_{xc}(r_s,\zeta) \rightarrow \frac{a_\infty}{r_s} + \frac{b_\infty}{r_s^{3/2}} \qquad (r_s \rightarrow \infty).$$
 (18)

The coefficients [14] $a_{\infty} \approx -1.1061$ and $b_{\infty} \approx \frac{1}{2}$ are independent of ζ , since any spatial overlap between two electrons is strongly suppressed in this limit, no matter whether their spins are parallel or not [15]. The resulting ISI expression for the exchange-correlation energy at finite densities reads [13]

$$e_{xc}^{ISI}(r_s,\zeta) = \frac{a_\infty}{r_s} + \frac{2X}{Y} \left[(1+Y)^{1/2} - 1 - Z \ln\left(\frac{(1+Y)^{1/2} + Z}{1+Z}\right) \right]. \tag{19}$$

Using $b_{\infty} = \frac{1}{2}$ and writing $e_x(r_s, \zeta) = c_x(\zeta)/r_s$, we have explicitly [13]

$$X(r_{s},\zeta) = \frac{-b_{0}(\zeta)}{[c_{x}(\zeta) - a_{\infty}]^{2}} \frac{1}{r_{s}},$$

$$Y(r_{s},\zeta) = \frac{4 b_{0}(\zeta)^{2}}{[c_{x}(\zeta) - a_{\infty}]^{4}} r_{s},$$

$$Z(\zeta) = \frac{-b_{0}(\zeta)}{[c_{x}(\zeta) - a_{\infty}]^{3}} - 1.$$
(20)

Eq. (19) provides a simple explicit LSD,

$$E_{xc}^{LSD}[\rho_{\uparrow}, \rho_{\downarrow}] = \int d^2r \, \rho(\mathbf{r}) \, e_{xc}^{ISI}(r_s(\mathbf{r}), \zeta(\mathbf{r})), \tag{21}$$

for treating arbitrary 2D electron systems (also finite ones such as quantum dots) by the Kohn-Sham Equations of DFT. In Eq. (21), $r_s(\mathbf{r}) = a_B^{-1}[\pi \rho(\mathbf{r})]^{-1/2}$ and $\zeta(\mathbf{r}) = [\rho_{\uparrow}(\mathbf{r}) - \rho_{\downarrow}(\mathbf{r})]/\rho(\mathbf{r})$.

In Fig. 2a, the ISI prediction $e_c^{ISI}(r_s,\zeta)=e_{xc}^{ISI}-e_x$ for the correlation energy of the unpolarized uniform electron gas $(\zeta=0)$ is compared with the accurate parametrization of the fixed-node diffusion Monte Carlo results in Ref. [5]. e_c^{ISI} differs slightly from the latter by up to 4%. This mild deviation might be cured by including in the ISI a simple model for the next-order coefficient of expansion (4) [16]. In the high-density limit $(r_s \to 0)$, however, where the present result is exact, the parametrization in Ref. [5] has for $0.7 < \zeta < 0.95$ a small positive deviation [5], shown in Fig. 2b.

Figure captions:

Fig. 1. (a) Numerical results (dots) for the function $f(\zeta)$ of Eq. (13) obtained by Monte Carlo integrations of expression (11) (with $\sigma_1\sigma_2 = \uparrow \downarrow$) at selected values of ζ . The analytical function $f_1(\zeta)$ of Eq. (14) is plotted as a dashed curve. The solid curve represents the accurate fit $f_1(\zeta) + \delta f(\zeta)$, using Eq. (15) for $\delta f(\zeta)$. (b) The fit (15) (solid curve) compared to the true deviation (dots) of the Monte-Carlo-integration results from $f_1(\zeta)$.

Fig. 2. The correlation energy of Ref. [5] (dotted curves) versus the present ISI results (solid curves).

References

- [1] E. Abrahams, S. V. Kravchenko, and M. P. Sarachik, Rev. Mod. Phys. 73, 251 (2001).
- [2] S.M. Reimann and M. Manninen, Rev. Mod. Phys. 74, 1283 (2002).
- [3] H. Jiang, H.U. Baranger, and W. Yang, Phys. Rev. B 68, 165337 (2003).
- [4] H. Saarikoski, E. Räsänen, S. Siljamäki, A. Harju, M.J. Puska, R.M. Nieminen, Phys. Rev. B 67, 205327 (2003).
- [5] C. Attaccalite, S. Moroni, P. Gori-Giorgi, and G. B. Bachelet, Phys. Rev. Lett. 88, 256 601 (2002).
- [6] A.K. Rajagopal and J.C. Kimball, Phys. Rev. B 15, 2819 (1977).
- [7] P. Gori-Giorgi and J.P. Perdew, Phys. Rev. B (to appear; cf. cond-mat/0305250).
- [8] J. Moreno and D.C. Marinescu, Phys. Rev. B 68, 195210 (2003)
- [9] M. Polini and M.P. Tosi, Phys. Rev. B **63**, 045118 (2001).
- [10] W. Macke, Z. Naturforsch. A 5A, 192 (1950).
- [11] W.J. Carr and A.A. Maradudin, Phys. Rev. 133, A371 (1964).
- [12] G.G. Hoffman, Phys. Rev. B 45, 8730 (1992).
- [13] M. Seidl, J.P. Perdew, and S. Kurth, Phys. Rev. Lett. 84, 5070 (2000).
- [14] L. Bonsall and A.A. Maradudin, Phys. Rev. B 15, 1959 (1977).
- [15] M. Seidl, J.P. Perdew, and S. Kurth, Phys. Rev. A 62, 012502 (2000).
- [16] M. Seidl and J.P. Perdew (unpublished).




Figure 1: (a) Numerical results (dots) for the function $f(\zeta)$ of Eq. (13) obtained by Monte Carlo integrations of expression (11) (with $\sigma_1\sigma_2=\uparrow\downarrow$) at selected values of ζ . The analytical function $f_1(\zeta)$ of Eq. (14) is plotted as a dashed curve. The solid curve represents the accurate fit $f_1(\zeta) + \delta f(\zeta)$, using Eq. (15) for $\delta f(\zeta)$. (b) The fit (15) (solid curve) compared to the deviation (dots) of the Monte-Carlo-integration results from $f_1(\zeta)$.




Figure 2: The correlation energy of Ref. [5] (dotted curves) versus the present ISI results (solid curves).