

Comment on “Current fluctuations in non-equilibrium diffusive systems: an additivity principle”

The main purpose of this comment is to point out that the results of Ref. [1] hold without the need of either an additivity principle, or scaling assumption which are foundational for Ref. [1]. We further comment on the range of validity of the theory and the advantages of our approach [2].

In a recent Letter [1], Bodineau and Derrida find that the “distribution of current fluctuations through a large one dimensional system in contact with two reservoirs at unequal densities” depends on only two parameters, the diffusion constant $D(\rho)$ and the noise density $\sigma(\rho)$, where ρ is the charge density. Their result is based on the assumptions of an “additivity principle” and a “scaling hypothesis”. The additivity principle postulates that a system may be decomposed into two independent subsystems, except that they are allowed to adjust the density ρ at their contact in order to maximize the probability to observe a given integrated current. The scaling hypothesis postulates that the current cumulant generating function scales as $1/N$, where N is the large system size.

While the assumptions are interesting, it is natural to ask about their microscopic origin, and in what situations the principles will successfully describe the physics. Therefore, it is important to have an alternative approach which is based on microscopic conditions for transport. Such a theoretical approach has been formulated in Ref. [2], where the results of the paper [1] have been derived independently with no assumption of the additivity principle or of the scaling hypothesis. Our method is related to an earlier technique [3], although they did not consider transport statistics and assume Gaussian fluctuations a priori.

The above additivity principle may be *derived* as the continuum limit of our theory of transport statistics in an arbitrary classical network, consisting of nodes storing charge and connectors carrying statistically distributed currents [4]. Our approach represents the current probability distribution function by a stochastic path integral, whose validity depends on (i) a separation of time scales requirement, prompting the Markov approximation, (ii) the assumption of an effectively continuous charge, and (iii) the saddle-point approximation. The saddle-point approximation is well controlled by a large parameter γ , the number of elementary charges participating in the transport. The dominant contribution to the generator of current cumulants, given by the saddle-point action S , is proportional to γ , with corrections of order 1. Thus the additivity principle may fail if $\gamma \sim 1$. For exclusion processes the assumption of effectively continuous charge is an approximation, which may fail close to critical points.

Considering a lattice of $N \gg 1$ of nodes and taking a

continuum limit of the stochastic path integral, i.e. expanding the action S to leading order in $1/N$, we have arrived [2] at the following 1D diffusive field theory:

$$S = - \int_0^t dt' \int_0^L dz [\lambda \dot{\rho} + D \rho' \lambda' - (F/2)(\lambda')^2], \quad (1)$$

where λ is the “counting field”, $D(\rho)$ is the diffusion constant, and $F(\rho)$ is the noise density in our notation. In the stationary limit, the solutions of the saddle-point equations of (1) with the boundary conditions $\lambda(0) = 0$, $\lambda(L) = \chi$ must be substituted back into the action to give the generating function $S(\chi)$ of cumulants of transmitted charge. This result is described in Eqs. (79-81) of [2] and is mathematically equivalent to the solution of Ref. [1] giving the generator (7,8) as coupled, implicit integral equations, after the said assumptions have been made. For example, the field equation (16) of Ref. [1] for the density ρ is the same as our equation (79) in Ref. [2].

We would like to stress that the additivity principle does not need to be postulated. From the field-theoretic point of view, it is the direct consequence of the space-time locality of the elementary transport transitions (sources of noise), and of the existence of a large parameter, the number of particles participating in transport. The fact that the transport statistics is characterized by only two parameters $D(\rho)$ and $F(\rho)$ originates from the central limit theorem, and is a consequence of the large N limit which removes high powers of λ from the action and leads to the Gaussian character of the local noise.

The advantage of the field-theoretic approach is not only that it shows the limits of the applicability of the theory, but also that it provides a framework for the perturbation expansion of current cumulants [2], investigation of time-dependent phenomena [5], and treatment of multiple dimensions [2].

E. V. Sukhorukov and A. N. Jordan,
Département de Physique Théorique
Université de Genève
CH-1211 Genève 4, Switzerland

Date: June 15, 2004

PACS numbers: 05.40.-a, 72.70.+m, 73.23.-b, 74.40.+k

-
- [1] T. Bodineau and B. Derrida Phys. Rev. Lett. **92**, 180601 (2004); cond-mat/0402305.
 - [2] A.N. Jordan, E.V. Sukhorukov, and S. Pilgram, cond-mat/0401650.
 - [3] L. Bertini *et al.*, J. Stat. Phys. **107**, 635 (2002); cond-mat/0108040.
 - [4] S. Pilgram *et al.*, Phys. Rev. Lett. **90**, 206801 (2003).
 - [5] S. Pilgram, K.E. Nagaev, M. Büttiker, cond-mat/0401632.