

# Ultracold two-component Fermi gases with a magnetic field gradient near a Feshbach resonance

Hongwei Xiong, Shujuan Liu, and Mingsheng Zhan

*State Key Laboratory of Magnetic Resonance and Atomic and Molecular Physics,  
Wuhan Institute of Physics and Mathematics, Chinese Academy of Sciences, Wuhan 430071, P. R. China*

Weiping Zhang

*Department of Physics, East China Normal University*

(Dated: May 23, 2019)

We study theoretically the ultracold two-component Fermi gases when a gradient magnetic field is used to tune the scattering length between atoms. For  ${}^6\text{Li}$  at the narrow resonance  $B_0 = 543.25$  G, it is shown that the gases would be in a coexistence of the regimes of BCS, BEC and unitarity limit with the present experimental technique. In the case of thermal and chemical equilibrium, we investigate the density distribution of the gases and show that a double peak of the density distribution can give us a clear evidence for the coexistence of BCS, BEC and unitarity limit.

PACS numbers: 03.75.Ss, 05.30.Fk, 05.30.Jp, 03.75.Hh

With the remarkable development of the cooling technique for two-component Fermi gases and Feshbach resonance, the recent experiments [1, 2, 3] have finally realized molecular Bose-Einstein condensation (BEC) and the condensation of fermionic atom pairs on the side of attractive interaction has also been investigated by several experiments [4, 5, 6, 7, 8, 9, 10, 11]. Together with intensive theoretical investigations [12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30] such as resonance superfluid [16, 17] and universal behavior for the gases with divergent scattering length [24, 25, 26, 27, 28, 29], these experimental advances provide us a quite unique system which will even contribute largely to our understanding on the mechanism of high-temperature superconductors.

For two-component Fermi gases, when the magnetic field is tuned so that the energy of a quasibound molecular state in a closed channel matches the total energy in an open channel, there is a magnetic-field Feshbach resonance [24, 31] which can change the scattering length between atoms. In the present experiments on the BCS-BEC crossover, a uniform magnetic field is used to tune the scattering length through the magnetic-field Feshbach resonance. Below the Feshbach resonant magnetic field (BEC side), the interaction between atoms is repulsive and there exists molecule which is a short-range fermionic atom pairs. Above the Feshbach resonant magnetic field (BCS side), the interaction between atoms is attractive and there would be a superfluid behavior due to the atomic Cooper pairs at sufficient low temperature. On resonance, the scattering length is divergent and one expects a universal behavior for the system in the unitarity limit. By tuning the uniform magnetic field near the resonant magnetic field, several experiments have investigated the BCS-BEC crossover such as condensate fraction [4, 5], collective excitation [8, 9], pairing gap [10], and heat capacity [11].

For ultracold two-component Fermi gases confined in an optical trap with axial symmetry along the  $z$  axis, in

the present work, we will investigate the unique property of the system when the magnetic field to tune the scattering length has a gradient of  $\alpha$ . The angular frequencies of the optical trap are  $\omega_r (= \omega_x = \omega_y)$  and  $\omega_z$  in the radial and axial directions, respectively. For Fermi gases with an equal incoherent mixture of the internal states  $|1\rangle$  and  $|2\rangle$ , near a Feshbach resonant magnetic field  $B_0$ , the scattering length is  $a(B) = Aa_0(1 - w/(B - B_0))$  with  $a_0$  being the Bohr radius. For the magnetic field in  $x$ -direction with gradient  $\alpha$  along  $z$ -direction, we have  $\vec{B} = (B_0 + \alpha z)\vec{e}_x$ . In this case, the scattering length becomes  $z$ -dependent and takes the following form:

$$a(z) = Aa_0 \left(1 - \frac{w}{\alpha z}\right). \quad (1)$$

If the magnetic field gradient  $\alpha$  is positive, when  $z < 0$  and  $a(z)n^{1/3} < 1$  ( $n$  is the total density distribution of the ultracold gases), there is repulsive interaction between fermionic atoms which corresponds the BEC side, while when  $z > 0$  and  $|a(z)|n^{1/3} < 1$  there is attractive interaction between fermionic atoms which corresponds to the BCS side. In the regime determined by  $|a(z)|n^{1/3} \gg 1$ , the ultracold gases are in the unitarity limit where the scattering length can be regarded as divergent and will not appear in the final result of a physical quantity such as chemical potential. Thus, by using the gradient magnetic field and appropriate parameters, there would be a coexistence of the gases in the form of BCS, BEC and unitarity limit.

When the magnetic field with gradient  $\alpha$  is applied to the ultracold gases, the overall external potential for the fermionic atom in the internal state  $|i\rangle$  is

$$V_{|i\rangle} = \frac{1}{2}m\omega_r^2(x^2 + y^2) + \frac{1}{2}m\omega_z^2(z - \Delta z_{|i\rangle})^2 - \mu_{|i\rangle}^{mag} B_0 + \Delta V_{|i\rangle}, \quad (2)$$

where  $i = 1, 2$ . Due to the presence of the magnetic field, for different internal state  $|i\rangle$ , there is a different shift to the optical trap which is determined by

$\Delta z_{|i\rangle} = \alpha \mu_{|i\rangle}^{mag} / m\omega_z^2$  with  $\mu_{|i\rangle}^{mag}$  being the magnetic moment of the internal state  $|i\rangle$ . In the above expression,  $\Delta V_{|i\rangle} = -\alpha^2 \left( \mu_{|i\rangle}^{mag} \right)^2 / 2m\omega_z^2$ . One can also get the overall external potential for a molecule

$$V_{mol} = m\omega_z^2 (x^2 + y^2) + m\omega_z^2 (z - \Delta z_{mol})^2 - \sum_i \mu_{|i\rangle}^{mag} B_0 + \Delta V_{mol}, \quad (3)$$

where  $\Delta z_{mol} = \alpha \left( \sum_i \mu_{|i\rangle}^{mag} \right) / 2m\omega_z^2$  and  $\Delta V_{mol} = -\alpha^2 \left( \sum_i \mu_{|i\rangle}^{mag} \right)^2 / 4m\omega_z^2$ .

The regime of the gases in the unitarity limit can be roughly determined by  $|a(z)|n^{1/3} \sim 1$ . Thus, the length scale for the regime in the unitarity limit can be estimated to be  $2l_z^{res}$  with  $l_z^{res} = Awn^{1/3}a_0/\alpha$ . Because there is a different shift for the external potential of the internal state  $|1\rangle$  and  $|2\rangle$ , the difference of the shift should be much smaller than  $2l_z^{res}$  so that one can omit safely the effect of the shift  $\Delta z_{|i\rangle}$ . After a simple calculation, this condition  $|\Delta z_{|1\rangle} - \Delta z_{|2\rangle}| \ll 2l_z^{res}$  means that

$$Aw \ll \frac{2(l_z^{res})^2 m\omega_z^2}{n^{1/3}a_0 \left| \mu_{|1\rangle}^{mag} - \mu_{|2\rangle}^{mag} \right|}. \quad (4)$$

For the typical parameters that  $\omega_z/2\pi \sim 100$  Hz,  $n \sim 10^{13}$  cm $^{-3}$  and  $l_z^{res} \sim 10$   $\mu$ m, the condition given by Eq. (4) requests that  $Aw$  is smaller than 10 G. This means that very narrow resonance for an element is an appropriate choice to investigate the coexistence of the regimes of BCS, BEC and unitarity limit. A careful investigation shows that the mixture of  $|1\rangle \equiv |F=1/2, m_F=1/2\rangle$  and  $|2\rangle \equiv |F=1/2, m_F=-1/2\rangle$  for  ${}^6\text{Li}$  at the narrow resonance located at  $B_0 = 543.25$  G [33] can satisfy our request. For this narrow resonance,  $Aw$  is estimated to be 6 G [33]. Thus, in the present work, we will consider two-component Fermi gases of  ${}^6\text{Li}$  at the narrow resonance  $B_0 = 543.25$  G.

For two-component Fermi gases of  ${}^6\text{Li}$  confined in the optical trap, we will investigate the density distribution of the gases by using the condition that the system is in thermal and chemical equilibrium. Our researches show that with appropriate choice of parameters and element for the system, the coexistence of the gases in the form of BCS, BEC and unitarity limit can be clearly shown through the density distribution. In the present experiments, the gases can be cooled far below the critical temperature of the molecular gas and Fermi temperature of the Fermi gases. Thus, we will consider the system at zero temperature to give a clear presentation.

To calculate the density distribution of the system, we firstly investigate the chemical potential of the system. In the regime of molecular BEC ( $a(z)n^{1/3} < 1$ ) with density distribution  $n_m^{BEC}$  and molecular scattering length

$a_m = 0.6a$  [34], the chemical potential of the molecular BEC is given by

$$\mu_m^{BEC} = \frac{2\pi\hbar^2 a_m n_m^{BEC}}{m} + V_{mol} + \varepsilon_m, \quad (5)$$

where  $\varepsilon_m = -E_m$  with  $E_m = \hbar^2/ma^2$  being the binding energy of the molecule. In this regime, the chemical potential of the Fermi gas in the internal state  $|i\rangle$  takes the form

$$\mu_{f|i\rangle}^{BEC} = \frac{\hbar^2 (6\pi^2)^{2/3}}{2m} \left( n_{f|i\rangle}^{BEC} \right)^{2/3} + \frac{2\pi\hbar^2 a}{m} n_{f|i\rangle}^{BEC} + V_{|i\rangle}, \quad (6)$$

where  $n_{f|i\rangle}^{BEC}$  is the density distribution of the Fermi gas in the internal state  $|i\rangle$ .

In the regime of the unitarity limit where the absolute value of the scattering length is much larger than the average distance between atoms, as pointed out in [29], we assume that the gases is in the mixture of Fermi gases and dimeric gas in chemical equilibrium. Based on the local density approximation at zero temperature, the chemical potential of the Fermi gas with density distribution  $n_{f|i\rangle}^{UL}$  takes the form

$$\mu_{f|i\rangle}^{UL} = (1 + \beta_1) \frac{\hbar^2 (6\pi^2)^{2/3}}{2m} \left( n_{f|i\rangle}^{UL} \right)^{2/3} + V_{|i\rangle}, \quad (7)$$

where  $\beta_1$  is firstly measured in [24] and  $\beta_1 = -0.56$  based on a quantum Monte Carlo calculation [26]. Omitting the binding energy of the dimer in the unitarity limit, based on the dimensionality analysis, the chemical potential of the dimeric gas with density distribution  $n_d^{UL}$  can be assumed as [29]

$$\mu_d^{UL} = (1 + \beta_2) \frac{\hbar^2 (6\pi^2)^{2/3}}{2 \times 2m} \left( n_d^{UL} \right)^{2/3} + V_d. \quad (8)$$

For the dimeric gas, we assume that the external potential  $V_d$  takes the same form as the external potential  $V_{mol}$  of the molecular gas. Based on the condition of the chemical equilibrium  $\sum_i \mu_{f|i\rangle}^{UL} = \mu_d^{UL}$ , one can get the ratio  $n_d^{UL}/n_{f|i\rangle}^{UL} = [4(1 + \beta_1)/(1 + \beta_2)]^{3/2}$  on resonance. From the experimental result  $n_d^{UL}/n_{f|i\rangle}^{UL} \approx 4$  in [5],  $\beta_2$  is estimated to be  $-0.3$ .

In the regime of BCS, the chemical potential can be approximated to be

$$\mu_{f|i\rangle}^{BCS} = \frac{\hbar^2 (6\pi^2)^{2/3}}{2m} \left( n_{f|i\rangle}^{BCS} \right)^{2/3} + V_{|i\rangle}, \quad (9)$$

where  $n_{f|i\rangle}^{BCS}$  is the density distribution of the fermionic atoms on the BCS side.

In the case of thermal and chemical equilibrium for the system, the chemical equilibrium gives us the following important relation:

$$\mu_m^{BEC} = \sum_i \mu_{f|i}^{BEC} = \sum_i \mu_{f|i}^{UL} = \mu_d^{UL} = \sum_i \mu_{f|i}^{BCS} \equiv \mu. \quad (10)$$

To illustrate clearly the density distribution and evidence for an experiment to show the coexistence of the regime of BCS, BEC and unitarity limit, we use the parameters  $\omega_z/2\pi = 120$  Hz and  $n_{f|1}^{UL} = 0.2 \times 10^{12}$  cm $^{-3}$  at the center  $z = 0$ . In addition, the gradient of the magnetic field is chosen to be  $\alpha = 0.21$  G/cm. Based on these parameters,  $|\Delta z_{|i}|/2l_z^{res} \ll 1$ ,  $|\Delta V_{|i}|/\mu \ll 1$ ,  $\Delta z_{mol} = 0$  and  $\Delta V_{mol}/\mu = 0$ . Thus, one can safely omit  $\Delta z_{|i}$ ,  $\Delta V_{|i}$ ,  $\Delta z_{mol}$  and  $\Delta V_{mol}$ .

From the equilibrium condition for the chemical potential given by Eq. (10), one gets the following expression for the density distribution:

$$n_m^{BEC} = \frac{(\mu_{eff} + \hbar^2/ma^2 - 2V_{ext})m}{2\pi\hbar^2 a_m}, \quad (11)$$

$$n_{f|1}^{BEC} = n_{f|2}^{BEC} = \left[ \frac{2m(\mu_{eff}/2 - V_{ext})}{\hbar^2 (6\pi^2)^{2/3}} \right]^{3/2}, \quad (12)$$

$$n_{f|1}^{UL} = n_{f|2}^{UL} = \left[ \frac{2m(\mu_{eff}/2 - V_{ext})}{(1 + \beta_1)\hbar^2 (6\pi^2)^{2/3}} \right]^{3/2}, \quad (13)$$

$$n_d^{UL} = \left[ \frac{4m(\mu_{eff} - 2V_{ext})}{(1 + \beta_1)\hbar^2 (6\pi^2)^{2/3}} \right]^{3/2}, \quad (14)$$

$$n_{f|1}^{BCS} = n_{f|2}^{BCS} = \left[ \frac{2m(\mu_{eff}/2 - V_{ext})}{\hbar^2 (6\pi^2)^{2/3}} \right]^{3/2}, \quad (15)$$

where  $V_{ext}$  takes the form

$$V_{ext} = \frac{1}{2}m\omega_r^2(x^2 + y^2) + \frac{1}{2}m\omega_z^2 z^2. \quad (16)$$

In getting Eqs. (11)-(15), the term  $\mu_{|i}^{mag} B_0$  in the final expression of the chemical potential given by Eqs. (5)-(9) has been cancelled out when the condition of the chemical equilibrium given by Eq. (10) is used. In this situation, the relation between the effective chemical potential  $\mu_{eff}$  and the density  $n_{f|1}^{UL}$  (or  $n_{f|2}^{UL}$ ) at  $z = 0$  is

$$\mu_{eff} = 2(1 + \beta_1) \frac{\hbar^2 (6\pi^2)^{2/3}}{2m} \left( n_{f|1}^{UL}(z=0) \right)^{2/3}. \quad (17)$$




FIG. 1: Shown is the density distribution of the gases when a gradient magnetic field is used to tune the atomic scattering length. After the molecules in the regime of BEC, the dimers in the regime of unitarity limit, and the atomic Cooper pairs in the regime of BCS are converted into deeply bound molecules by decreasing non-adiabatically the magnetic field, the thick dashed line illustrate the density distribution of the deeply bound molecules (See also the inset). We see that there is a double peak for the density distribution of the deeply bound molecules which can give us a clear evidence for an experiment to make the system in the coexistence of BEC, BCS and unitarity limit. In this figure, the density distribution is in unit of  $n_{f|1}^{UL}(z=0)$ , while the coordinate  $z$  is in unit of  $R_z = \sqrt{\mu_{eff}/m\omega_z^2}$ .

Using the parameters in this paper, shown in Fig.1 is the density distribution of the gases where the density is in unit of  $n_{f|1}^{UL}(z=0)$ , while the coordinate  $z$  is in unit of  $R_z = \sqrt{\mu_{eff}/m\omega_z^2}$ . The arrow shows the location  $\pm 0.7R_z$  which is determined by  $|a(z)|n^{1/3} = 1$ . The solid line in Fig.1 shows the density distribution of molecular BEC  $n_m^{BEC}(z < -0.7)$ , while the dot-dashed line in this figure shows the density distribution  $n_d^{UL}(-0.7 < z < 0.7)$  of the dimers in the unitarity limit. The dotted line shows the density distribution of  $n_{f|1}^{BCS}(z > 0.7)$ . Analogously to the experiments in [4, 5], we consider the process that the magnetic field is decreased non-adiabatically so that the fermionic atom pairs (i.e. molecules on the BEC side, dimers in the unitarity limit, and atomic Cooper pairs on the BCS side before the decreasing of the magnetic field) are converted into deeply bound molecules. After this non-adiabatical process, from the density distribution of molecular BEC (solid line) and dimers in the unitarity limit (dot-dashed line), we see that the coexistence of the gases in the form of BCS, BEC and unitarity limit can be clearly shown through the double peak in the density distribution of the

deeply bound molecules. The thick dashed line (see also the inset) shows a schematic interpolation of the density distribution of the deeply bound molecules. The evidence for the coexistence of the regimes of BCS, BEC, and unitarity limit can be also shown through the non-symmetric density distribution of the deeply bound molecules about  $z = 0$  after the non-adiabatic process.

In summary, by using a gradient magnetic field to change the scattering length, an experimental scheme is proposed to make the gases become the coexistence of the regimes of BCS, BEC and unitarity limit. A careful investigation shows that for  ${}^6\text{Li}$  at the narrow resonance  $B_0 = 543.25$  G, one can realize this suggestion with the present experimental technique. In the case of thermal and chemical equilibrium, we calculate the density distribution of the system and shows that the double peak in the density distribution of the deeply bound molecules af-

ter the non-adiabatic decreasing of the magnetic field can give us a clear evidence for the coexistence of the regimes of BCS, BEC and unitarity limit. It is quite promising that this novel system can give us many opportunities to investigate the dynamics of the ultracold gases such as a new type of Josephson effect between gases in the form of BCS and BEC.

### Acknowledgments

We acknowledge the useful discussions with Dr. Cheng Chin. This work was supported by the National Natural Science Foundation of China under Grant Nos.10474117 and 10205011.

- 
- [1] S. Jochim, M. Bartenstein, A. Altmeyer, G. Hendl, S. Riedl, C. Chin, J. H. Denschlag, and R. Grimm, *Science* **302**, 2101 (2003).
  - [2] M. Greiner, C. A. Regal, and D. S. Jin, *Nature* **426**, 537 (2003).
  - [3] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, S. Gupta, Z. Hadzibabic, and W. Ketterle, *Phys. Rev. Lett.* **91**, 250401 (2003).
  - [4] C. A. Regal, M. Greiner, and D. S. Jin, *Phys. Rev. Lett.* **92**, 040403 (2004).
  - [5] M. W. Zwierlein, C. A. Stan, C. H. Schunck, S. M. F. Raupach, A. J. Kerman, and W. Ketterle, *Phys. Rev. Lett.* **92**, 120403 (2004).
  - [6] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. H. Denschlag, and R. Grimm, *Phys. Rev. Lett.* **92**, 120401 (2004).
  - [7] T. Bourdel, L. Khaykovich, J. Cubizolles, J. Zhang, F. Chevy, M. Teichmann, L. Tarruell, S. J. J. M. F. Kokkelmans, and C. Salomon, *Phys. Rev. Lett.* **93**, 050401 (2004).
  - [8] J. Kinast, S. L. Hemmer, M. E. Gehm, A. Turlapov, and J. E. Thomas, *Phys. Rev. Lett.* **92**, 150402 (2004).
  - [9] M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, C. Chin, J. H. Denschlag, R. Grimm, *Phys. Rev. Lett.* **92**, 203201 (2004).
  - [10] C. Chin, M. Bartenstein, A. Altmeyer, S. Riedl, S. Jochim, J. H. Denschlag, R. Grimm, *Science* **305**, 1128 (2004).
  - [11] J. Kinast, A. Turlapov, J. E. Thomas, preprint cond-mat/0409283 (2004).
  - [12] A. J. Leggett, *J. Phys. C. (Paris)* **41**, 7 (1980).
  - [13] P. Nozières and S. Schmitt-Rink, *J. Low Temp. Phys.* **59**, 195 (1985).
  - [14] H. T. C. Stoof, M. Houbiers, C. A. Sackett, and R. G. Hulet, *Phys. Rev. Lett.* **76**, 10 (1996).
  - [15] Y. Ohashi and A. Griffin, *Phys. Rev. A* **67**, 033603 (2003); *Phys. Rev. A* **67**, 063612 (2003); *Phys. Rev. Lett.* **89**, 130402 (2002).
  - [16] E. Timmermans, K. Furuya, P. W. Milonni, and A. K. Kerman, *Phys. Lett. A* **285**, 228 (2001).
  - [17] M. Holland, S. J. J. M. F. Kokkelmans, M. L. Chiofalo, and R. Walser, *Phys. Rev. Lett.* **87**, 120406 (2001); J. N. Milstein, S. J. J. M. F. Kokkelmans and M. J. Holland, *Phys. Rev. A* **66**, 043604 (2002).
  - [18] J. Stajic, J. N. Milstein, Q. Chen, M. L. Chiofalo, M. J. Holland, and K. Levin, preprint cond-mat/0309329 (2003).
  - [19] X. Yang, and Y. Wu, *Commun. Theor. Phys.* **39**, 543 (2003).
  - [20] L. D. Carr, G. V. Shlyapnikov, and Y. Castin, *Phys. Rev. Lett.* **92**, 150404 (2004).
  - [21] G. M. Bruun, preprint cond-mat/0401497 (2004).
  - [22] G. M. Falco and H. T. C. Stoof, *Phys. Rev. Lett.* **92**, 130401 (2004).
  - [23] H. Xiong, S. Liu, M. Liu, K. Gao, and M. Zhan, preprint cond-mat/0404565 (2004).
  - [24] K. M. O'Hara, S. L. Hemmer, M. E. Gehm, S. R. Granade, and J. E. Thomas, *Science* **298**, 2179 (2002).
  - [25] H. Heiselberg, *Phys. Rev. A* **63**, 043606 (2001).
  - [26] J. Carlson, S. Y. Chang, V. R. Pandharipande, and K. E. Schmidt, *Phys. Rev. Lett.* **91**, 050401 (2003).
  - [27] T. L. Ho and E. J. Mueller, *Phys. Rev. Lett.* **92**, 160404 (2004).
  - [28] T. L. Ho, *Phys. Rev. Lett.* **92**, 090402 (2004).
  - [29] H. Xiong and S. Liu, preprint cond-mat/0403652 (2004).
  - [30] J. Kinnunen, M. Rodriguez, P. Törmä, *Science* **305**, 1131 (2004).
  - [31] W. C. Stwalley, *Phys. Rev. Lett.* **37**, 1628 (1976); E. Tiesinga, B. J. Verhaar, and H. T. C. Stoof, *Phys. Rev. A* **47**, 4114 (1993).
  - [32] S. Inouye, M. R. Andrews, J. Stenger, H.-J. Miesner, D. M. Stamper-Kurn, and W. Ketterle, *Nature* **392**, 151 (1998); P. Courteille, R. S. Freeland, D. J. Heinzen, F. A. van Abeelen, and B. J. Verhaar, *Phys. Rev. Lett.* **81**, 69 (1998).
  - [33] K. E. Strecker, G. B. Partridge, and R. G. Hulet, *Phys. Rev. Lett.* **91**, 080406 (2003).
  - [34] D. S. Petrov, C. Salomon, and G. V. Shlyapnikov, *Phys. Rev. Lett.* **93**, 090404 (2004).