Geometrical and Algebraic structures in Quantum Hall Systems

J.C. Wallet*

20 novembre 2018

Groupe de Physique Théorique, Institut de Physique Nucléaire, F-91406, ORSAY CEDEX (France)

 $Abstract^{1}$: We review the main features of a mathematical framework encompassing some of the salient quantum mechanical and geometrical aspects of Hall systems with finite size and general boundary conditions. Geometrical as well as algebraic structures controlling possibly the integral or fractional quantization of the Hall conductivity are discussed.

PACS : 71.10.-w; 02.40.-k

^{*}e-mail address : wallet@ipno.in2p3.fr

¹Talk given at the "Conference on Higher Dimensional Quantum Hall Effect, Chern-Simons Theory and Non-commutative Geometry in Condensed Matter Physics and Field Theory", Trieste, 1-4 March 2005.

1 Introduction

The integral quantization of the Hall conductivity σ_H can be understood as stemming from topological features underlying the Quantum Hall systems [1]. The pioneering argument in that direction proposed by Laughlin [2] for a system with a cylindrical geometry has been refined in subsequent works dealing with non interacting electrons in a periodic potential [3] and further extended to take into account disorder and/or electron mutual interaction [4], [5], [6], [7]. Soon after [2], it has been observed [8] that the Kubo formula from which σ_H is obtained can be related to the integral of the first Chern class of some Line Vector Bundle. This latter structure shows up because two parameters have to be introduced within the quantum mechanical description of the considered systems. Attempts to explain the observed fractional quantization of σ_H have also appeared, combining the above scheme with additional physical plausible assumptions (e.g. the possible degeneracy of the ground state, the existence of a gap above it and/or the validity of averaging σ_H over the above two parameters) [4-7].

The above "Topological Approach to the Quantum Hall Effect" has been recognized as providing a possible explanation 2 for the observed robustness of the quantization of σ_H but it is not quite fully satisfactory. It appears that this approach does not permit one to really predict the value for σ_H starting from *basic* ingredients pertaining to the quantum mechanical description of the systems. This may be due to the fact that the above mentionned Line Vector Bundle does not capture (in most cases) a sufficient amount of the physical properties shared by the real experiments (e.g. the ("engineering") geometry of the experimental devices). Notice that many quantum mechanical models used so far rely on assumptions made on the geometry of the system (e.g. starting from a torus or annular geometry). While these assumptions are convenient from a mathematical viewpoint (since they simplify the analyzis and/or permits one to deal rather easily with a suitable choice for the boundary conditions), they are not very natural from an experimental viewpoint. Ultimately, it would be useful to incorporate into a single framework (hopefully) all the geometrical and physical constraints characterizing the real experiments and control the actual validity of the assumptions. This, together with a proper inclusion of disorder, would permit one to describe in

 $^{^2\}mathrm{even}$ when disorder is included which however requires the actual validity of some additional hypothesis

a single formalism integral and fractional quantization (and by the way to fully reconcile localization with the topological machinery).

This talk summarizes some of the results derived in [9] and [10] in collaboration with Y. Georgelin and T. Masson. The physical system we consider is a rectangle coupled to a four-point probe and is similar to the one considered in [6], [11]. It is depicted on figure 1. This is a natural choice regarding the actual experimental devices as discussed in particular in [11], [9]. In most of the following discussion, the disorder is left aside. In section 2, we outline the main features of the formalism introduced in [9] which permits us to deal with general boundary conditions for the physical system. Consistency of the necessary self-adjointness for the Hamiltonian description with general boundary conditions can only be achieved through the introduction of two real parameters³ (that will generate some "extended" Brillouin zone). In section 3, we point out the occurrence of a symmetry for the system which can be identified with the irrational rotation algebra of the Non-Commutative Torus [12]. This symmetry basically connects the physical space to the "extended" Brillouin zone, that we call reciprocical space in the following. This symmetry is also briefly discussed in the light of the Non-Commutative Torus and Area-Preserving Diffeomorphism algebra. In section 4, we review the case where $\frac{NB}{2\pi}$ is integer. Here N (resp. B) is the number of charge carriers (resp. the external magnetic field) and $\frac{NB}{2\pi}$ is the relevant control parameter for the system. The formalism of section 2 combined with the relevant version of the irrational rotation algebra for $\frac{NB}{2\pi}$ integer permits one to compute entirely σ_H from the Kubo formula which is found to take integer or fractional values. In section 5, we discuss the case where $\frac{NB}{2\pi}$ is not an integer for which the situation becomes far more complicated. When it takes rational values, the Hall conductivity can again be computed leading again to integer or fractional values. The extension of the analyzis to the case where $\frac{NB}{2\pi}$ is irrational does not lead to a reliable characterization of the reciprocical space, therefore signaling the limitation of the present approach. A possible way to circumvent this limitation (which could also possibly lead to a proper incorporation of the disorder) is discussed in section 6 where promising indications favoring the choice of a more algebraic framework are breifly described.

³These parameters generalize the so called "boundary condition angles" which appear e.g. when dealing with "Periodic or Toroidal boundary conditions".



FIG. 1 - The geometry of the system

2 Hall system on a finite size sample

We consider the system depicted on figure 1 where N interacting spinless particles with mass m are confined on a rectangular sample (that we assume in the following to be a square of unit length) submitted to an external magnetic field B and whose opposite edges are connected by wires. Units are $e = \hbar = c = 1$. (x_i, y_i) (resp. (p_{x_i}, p_{y_i})) denote the coordinates (resp. momenta) for particle *i*.

The electrons located in the planar sample are ruled by Quantum Mechanics while the currents circulating in the wires (which are connected to macroscopic devices in the experiments) should be regarded as classical quantities. General boundary conditions can be obtained by constraining these classical quantities. Namely, we assume that the currents circulating in the wires are conserved and that the charge located on each edge of the rectangle is equal to the one located on the corresponding opposite edge. These conditions will be translated into constraints on the densities and currents expressed in the Schrödinger representation. In this way, the presence of the wires reflects itself as boundary conditions constraining the Quantum Mechanics describing the dynamics of the electron in the sample.

As recalled in [9], the Hamiltonian for the system, in the absence of random potential, can be conveniently written in term of collective (center of mass) variables as

$$H = \frac{1}{2Nm} \left(\left(P_x + \frac{1}{2} NBy \right)^2 + \left(P_y - \frac{1}{2} NBx \right)^2 \right) + H_I \equiv H_0 + H_I \qquad (1)$$

where $x = \frac{1}{N} \sum_{i=1}^{N} x_i$, $P_x = \sum_{i=1}^{N} p_{x_i}$, $y = \frac{1}{N} \sum_{i=1}^{N} y_i$, $P_y = \sum_{i=1}^{N} p_{y_i}$ and H_I depends only on the "internal" variables ($\tilde{x}_i = x_i - x$, $\tilde{p}_{x_i} = p_{x_i} - \frac{1}{N} P_x$ and similar expressions for the *y*-counterpart). In (1), the symmetric gauge for the external gauge potential has been used. The corresponding Hilbert space is then a tensor product $\mathcal{H}_0 \otimes \mathcal{H}_I$ where the collective (resp. internal) operators act only on \mathcal{H}_0 (resp. \mathcal{H}_I). In the following, we focus the discussion

on the H_0 part of the Hamiltonian together with the \mathcal{H}_0 part of the Hilbert space since the observables we are interested in depends only on the center of mass variables and the relevant operators act only on \mathcal{H}_0 .

To proceed further, one needs to built a representation of the abstract operator algebra built from H_0 , P_x , P_y . As we know from general operator theory, care must be taken here because we work on a sample with finite size so that, in order to obtain a self-adjoint representation of the operators, one has to specify carefully the space of functions on which they act. Therefore, requiring self-adjointness and using boundary conditions constraints expressed as constraints on the edge densities and currents in the Schödinger representation, we find the following result [9] :

$$P_x = -i\partial_x + \frac{NB}{2}y \; ; \; P_y = -i\partial_y - \frac{NB}{2}x \tag{2}$$

must act on the linear space of functions verifying

$$\Phi(1,y) = e^{i(\gamma + \frac{NB}{2}y)} \Phi(0,y) \; ; \; \Phi(x,1) = e^{i(\eta - \frac{NB}{2}x)} \Phi(x,0) \tag{3}$$

where γ and η are two real arbitrary parameters which *must* necessarely be introduced in the course of the construction and label inequivalent representations for P_x and P_y [9] modulo 2π . In mathematical words, self-adjointness combined with boundary conditions preserving current conservation in the wires and edge charge conservation force the representation of the relevant operator algebra to be reducible. The space generated by γ and η is called the *reciprocical space* and will be characterized more closely in a while. Note that, at this point, γ and η are not constrained. Constraints on these parameters will appear by further assuming specific values for $\frac{NB}{2\pi}$. According to the above discussion, the Hilbert space \mathcal{H}_0 is a direct sum (actually a direct Hilbertian integral) of each Hilbert space indexed by each pair (γ , η) and from (3) any general wave function $\psi(x, y; \gamma, \eta)$ must satisfy

$$\psi(1, y; \gamma, \eta) = e^{i(\gamma + \frac{NB}{2}y)}\psi(0, y; \gamma, \eta) \; ; \; \psi(x, 1; \gamma, \eta) = e^{i(\eta - \frac{NB}{2}x)}\psi(x, 0; \gamma, \eta)$$
(4).

The various differences between the present framework and other related works are discussed in the section 4 of [9] to which we refer for more details. Before going further, some comments are in order. Equation (4) defines ψ as being a section of a Line Vector Bundle over a Torus in the (x, y)-space only when $\frac{NB}{2\pi}$ is integer. In this case, the corresponding base space, a (x, y)-Torus with periodicity 1, has nothing to do with the physical sample (whose topology is that of a Torus with one puncture). When $\frac{NB}{2\pi}$ takes rational values, the base space (in the (x, y)-space) of the Line Vector Bundle becomes a Rieman surface of higher genus with punctures [10].

3 The irrational rotation algebra

It appears that the system built from H_0 and \mathcal{H}_0 has a symmetry [9], [10]. It is generated by a family of unitary operators $\mathcal{U}_{\sigma,\theta}$ indexed by two parameters σ and θ verifying

$$\mathcal{U}_{\sigma,\theta}\psi(x,y;\gamma,\eta) = e^{-\frac{i}{2}NB(\sigma x + \theta y)}\psi(x+\theta,y-\sigma;\gamma+NB\sigma,\eta+NB\theta)$$
(5)

$$\mathcal{U}_{\sigma_1,\theta_1}\mathcal{U}_{\sigma_2,\theta_2} = e^{-iNB(\sigma_2\theta_1 - \sigma_1\theta_2)}\mathcal{U}_{\sigma_2,\theta_2}\mathcal{U}_{\sigma_1,\theta_1} \tag{6}$$

where (5) holds provided the wave functions admit (monovalued) extensions on some suitable space related to R^2 , a condition which can be always achieved [10]. Here, we point out that this symmetry connects both the (x, y) space and the reciprocical space as it can be realized from (5). It appears that this observation proves usefull in the computation of the Hall conductivity when $\frac{NB}{2\pi}$ takes integer values. Equation (6) is reminiscent of the magnetic algebra obeyed by the magnetic translation operators in Landau type problems. In fact, when $\frac{NB}{2\pi}$ is irrational, (6) is known in the mathematical litterature as the "irrational rotation algebra" of the Non-Commutative Torus [12]. It appears that it can be related to the abstract algebra defining the Non-Commutative Torus. This abstract algebra is defined as an associative (non commutative) unital *-algebra with involution † generated by two elements Z_1 and Z_2 satisfying

$$Z_1 Z_2 = e^{i2\pi\Theta} Z_2 Z_1 , \ Z_i^{\dagger} = Z_i^{-1}, \ i = 1, 2 , \ \Theta \in [0, 1]$$
(7)

(Θ is usually called the non commutativity parameter) with smooth completion obtained from all elements f of the form⁴

$$f = \sum_{m_1, m_2 \in Z^2} f_{(m_1, m_2)} e^{i\pi\Theta m_1 m_2} Z_1^{m_1} Z_2^{m_2}$$
(8)

where the (Schwartz sequences) $f_{(m_i,m_j)}$'s decrease sufficiently fast for $|m_i| \to \infty$. One observes that (6) reduces to (7) upon setting $Z_1 \equiv \mathcal{U}_{\sigma,0}, Z_2 \equiv \mathcal{U}_{0,\theta}$

⁴which in some sense generalizes the usual Fourier series expansion

and $NB\sigma\theta = 2\pi\Theta$. Notice that, by defining $T_{(n_1,n_2)} \equiv \frac{1}{\Theta}e^{in_1n_2\Theta/2}Z_1^{n_1}Z_2^{n_2}$, one obtains from (8)

$$[T_{(n_1,n_2)}, T_{(m_1,m_2)}] = \frac{i2}{\Theta} \sin(\Theta(n_1m_2 - n_2m_1))T_{(n_1+m_1,n_2+m_2)}$$
(9).

This infinite dimensional Lie algebra is the "trigonometric algebra" initially advocated in [13] which, in the limit $\Theta \to 0$ reduces to $[T_{(n_1,n_2)}, T_{(m_1,m_2)}] = i2(n_1m_2 - n_2m_1)T_{(n_1+m_1,n_2+m_2)}$, that is the W_{∞} algebra of the Area-Preserving Diffeomorphisms of the Torus.

4 Computing the Hall conductivity

When $\frac{NB}{2\pi} = l$, l integer, that we call the integer case, we find that (4) can be extended as

$$\psi(x+1,y;\gamma,\eta) = e^{i(\gamma + \frac{NB}{2}y)}\psi(x,y;\gamma,\eta) \tag{10a}$$

$$\psi(x, y+1; \gamma, \eta) = e^{i(\eta - \frac{NB}{2}x)}\psi(x, y; \gamma, \eta)$$
(10b)

which, for fixed (γ, η) , permits us to interpret ψ as a section of a Line Vector Bundle (LVB) over a torus in the (x, y)-space with periodicity 1. Notice that the quantization condition $\frac{NB}{2\pi} = l$ simply expresses the fact that the first Chern class of this LVB, $\frac{NB}{2\pi}$, is an integer l. Recall that the above torus has nothing to do with the physical sample! It is convenient at this point to perform the following gauge transformation $\tilde{\psi}(x, y; \gamma, \eta) = e^{-i(\gamma x + \eta y)}\psi(x, y; \gamma, \eta)$. The relevant symmetry stemming from (5) and (6) is generated by

$$\mathcal{U}_{1/l,0} \equiv \tilde{U} , \ \mathcal{U}_{0,1/l} \equiv \tilde{V} ; \ \tilde{V}\tilde{U} = e^{i\frac{2\pi}{l}}\tilde{U}\tilde{V}$$
(11)

and the (gauge transformed) wave functions $\tilde{\psi}$ can be splitted into l components $\tilde{\psi}_I$, I = 1, ..., l which, upon using (10), (11) and (5) in which the \mathcal{U} 's are replaced by their \tilde{U}, \tilde{V} counterparts, must satisfy $(q = e^{i\frac{2\pi}{l}})$

$$\tilde{\psi}_I(x,y;\gamma,\eta) = q^{1-I} e^{i(\pi y + \frac{\gamma}{l})} \tilde{\psi}_I(x + \frac{1}{l},y;\gamma,\eta + 2\pi)$$
(12a)

$$\tilde{\psi}_{I+1}(x,y;\gamma,\eta) = e^{i(\pi x + \frac{\eta}{l})} \tilde{\psi}_I(x,y - \frac{1}{l};\gamma + 2\pi,\eta)$$
(12b)

and

$$\tilde{\psi}_I(x,y;\gamma+2\pi l,\eta) = e^{-i2\pi lx} e^{i\eta} \tilde{\psi}_I(x,y;\gamma,\eta)$$
(13a),

$$\tilde{\psi}_I(x,y;\gamma,\eta+2\pi l) = e^{-i2\pi l y} e^{-i\gamma} \tilde{\psi}_I(x,y;\gamma,\eta)$$
(13b).

The relation (12) expresses the fact that the function $\tilde{\psi}_I$ (resp. $\tilde{\psi}_{I+1}$) restricted to the domain $\eta \in [2\pi n, 2\pi (n+1)], 1 \leq n \leq l-1$ (resp. $\gamma \in [2\pi m, 2\pi (m+1)], 1 \leq m \leq l-1$) is completely determined by the restriction of $\tilde{\psi}_I$ on $\eta \in [0, 2\pi]$ (resp. $\gamma \in [2\pi (m-1), 2\pi m]$. More importantly, the relation (13), obtained from (10) with the help of (11), permits us to interpret each $\tilde{\psi}_I$ as a section of a LVB over a torus in the (γ, η) reciprocical space with periodicity $2\pi l$.

Now we can use the Kubo formula to compute the Hall conductivity σ_H . By using standard manipulations, σ_H can be expressed as [9]

$$\sigma_H = \frac{N^2}{i(2\pi)^2} \frac{1}{l} \sum_{I=1}^l \int_{[0,2\pi]^2} d\gamma d\eta \int_{[0,1]^2} dx dy \left(\frac{\partial \tilde{\psi}_I^*}{\partial \gamma} \frac{\partial \tilde{\psi}_I}{\partial \eta} - \frac{\partial \tilde{\psi}_I^*}{\partial \eta} \frac{\partial \tilde{\psi}_I^*}{\partial \gamma}\right)$$
(14).

Set now

$$\Omega_I \equiv \int_{[0,1]^2} dx dy \left(\frac{\partial \tilde{\psi}_I^*}{\partial \gamma} \frac{\partial \tilde{\psi}_I}{\partial \eta} - \frac{\partial \tilde{\psi}_I^*}{\partial \eta} \frac{\partial \tilde{\psi}_I^*}{\partial \gamma}\right), \ I = 1, ..., l$$
(15).

By further making use of (12), one can show [9] that

$$\frac{1}{l} \sum_{I=1}^{l} \int_{[0,2\pi]^2} d\gamma d\eta \Omega_I = \frac{1}{l^2} \int_{[0,2\pi l]^2} d\gamma d\eta \Omega_1$$
(16)

where in the RHS of (16), one can easily recognizes [9] the first Chern number for the relevant LVB over the Torus in the reciprocical space with periodicity $2\pi l$, namely $C_1^R = i2\pi \int_{[0,2\pi l]^2} d\gamma d\eta \Omega_1$. It appears that this Chern number can be computed entirely. The derivation can be found in the appendix B of [9] and the result is $C_1^R = i4\pi l$. This, combined with (16) yields

$$\sigma_H = \frac{1}{2\pi} \left(\frac{2N^2}{l}\right) \tag{17}.$$

Therefore, when $\frac{NB}{2\pi} = l$, l integer, we find that the Hall conductivity can take integer or fractional values.

5 The non-integer case

The situation becomes far more complicated when $\frac{NB}{2\pi}$ is not an integer value. When $\frac{NB}{2\pi} = l/k$, the Hall conductivity σ_H can again be explicitly

computed [10], provided some additional assumptions on the initial interaction potential is made. We find that σ_H takes again integer or fractional values, namely

$$\sigma_H = \frac{1}{2\pi} \frac{2(kN)^2}{kl}$$
(18).

If the above assumption is relaxed, one has first to extend (4) to some suitable domain of the (x, y)-space built from R^2 . We find that this can be done on a Riemann surface S_k with genus $g = \frac{1}{2}(2 + k^2(k - 1))$ with k^2 punctures. The proof (given in appendix A of [10]) is rather involved and amounts to combine surface surgery to the van Kampen theorem. Roughly speaking, S_k is obtained by gluing k^3 copies of the physical sample and receives a magnetic flux equal to $2\pi lk^2$, each puncture carrying a magnetic flux equal to $2\pi l$. The relation (4) (with the gauge transformed wave functions $\tilde{\psi}$ as defined in section 4) becomes

$$\tilde{\psi}(x+k,y;\gamma,\eta) = e^{ik\gamma + \frac{i}{2}2\pi ly}\tilde{\psi}(x,y;\gamma,\eta)$$
(19a)

$$\tilde{\psi}(x, y+k; \gamma, \eta) = e^{ik\eta - \frac{i}{2}2\pi lx} \tilde{\psi}(x, y; \gamma, \eta)$$
(19b),

so that it can be now interpreted a section of a LVB over S_k in the (x, y)space. The rest of the computation of σ_H , although very involved, could be done in principle by relating the above LVB to its counterpart in the reciprocical space through (5) and (6) and using the representations of the relevant algebra stemming from (6). When $\frac{NB}{2\pi}$ takes irrational values, the relevant space on which (4) can be extended is very complicated : it is the universal covering space of R^2/Z^2 [10] and no conclusion can be obtained from the present formalism.

6 Discussion and outlook

Provided some reliable identification of the reciprocical space (i.e the Brillouin zone) can be done, integer as well as fractional quantization for σ_H can be shown within the present approach. This latter however loses its efficiency when no Brillouin zone can be characterized which is the case when $\frac{NB}{2\pi}$ is irrational or when the disorder is taken into account. We note that plausible phenomenological conclusions can be obtained in the presence of (at least weak) disorder by supplementing the present analyzis with reasonable physical assumptions somehow similar to those initialy proposed in [5]. We will not discuss this point here, referring to [10] for more details. The conclusions we obtain are consistent either with the physics stemming from the global phase diagram proposed in [14] or with the phase diagram that we proposed in [15].

The Chern numbers are not the only topological invariants playing a salient role in the Topological approach for the Quantum Hall Effect. When no Brillouin zone can be identified, it appears that σ_H can also be related to another topological invariant [16], namely the Fredholm index which is stable under (small) deformations and can be viewed as some non-commutative version of the Chern number. The introduction of Fredholm index permits one to deal with disorder but only integer quantization for σ_H is recovered. So far, a self-contained mathematical framework dealing with interacting electrons in the presence of a random potential (in a realistic geometry similar to the one depicted on fig.1) and giving rise to integer as well as fractional quantization for σ_H is still missing.

The analyzis presented in [9], [10] involves encouraging indications favoring a more algebraic approach that could reach this goal and appears to be related to the Kasparov K-K Theory [17]. To get a flavor of this, one first notices that deformations can be better treated within an algebraic framework. One further notices that LVB can be indeed completely characterized by their sections. This stemms basically from the Serre-Swan Theorem arising in Bundle Theory which asserts that there is a one-to-one correspondence between LVB over some base space and projective modules over the algebra of functions describing this base space. Projective modules are classified by K-Theory. Now, sections in the (x, y)-space must be connected to their counterpart in the reciprocical space through transformations associated with the irrational rotation algebra. This amounts to connect K-theory in the (x, y)space to its K-Theory counterpart in the reciprocical space through the above transformations which is reminiscent of (Kasparov) K-K Theory. The actual implementation of these ideas in Quantum Hall systems as well as in other physical situations are presently under study [18].

Acknowledgements : The content of this talk has been greatly influenced by many stimulating discussions with my collaborators Y. Georgelin and T. Masson. I am grateful to the organizers of the "Conference on Higher Dimensional Quantum Hall Effect, Chern-Simons Theory and Non-commutative Geometry in Condensed Matter Physics and Field Theory" for invitation. Hospitality of ICTP (Trieste) is also acknowledged.

Références

- For a review, see "The Quantum Hall Effect", R.E. Prange, S.M. Girvin Eds., Springer Verlag, Berlin (1990); see also "Quantum Hall Effect", M. Stone Ed., World Scientific, 1992.
- [2] : R.B. Laughlin, Phys. Rev. B23 (1981) 5632.
- [3] : D.J. Thouless, M. Kohmoto, M.P. Nightingale, M. de Nijs, Phys. Rev. Lett.49 (1982) 405; see also M. Kohmoto, Ann. Phys. 160 (1985) 343.
- [4] : Q. Niu, D.J. Thouless, J. Phys. A17 (1984) 2453.
- [5] : Q. Niu, D.J. Thouless, Y.-S. Wu, Phys. Rev. B31 (1985) 3372.
- [6] : J.E. Avron, R. Seiler, Phys. Rev. Lett.54 (1985) 259.
- [7] : D.J. Thouless, Phys. Rev. Lett.57 (1983) 57.
- [8] : J. Avron, R. Seiler, B. Simon, Phys. Rev. Lett.51 (1983) 51; see also 2nd of ref. [3].
- [9] : F. Chandelier, Y. Georgelin, T. Masson, J.C. Wallet, Ann. Phys. 305 (2003) 60.
- [10] : F. Chandelier, Y. Georgelin, T. Masson, J.C. Wallet, Ann. Phys. 314 (2004) 476.
- [11] : Q. Niu, D.J. Thouless, Phys. Rev. B35 (1987) 2188.
- [12] : M.A. Rieffel, Pacific J. Math. 93 (1981) 415.
- [13] : D.B. Fairlie, P. Fletcher, C.K. Zachos, Phys. Lett. B218 (1989) 203.
- [14] : S. Kivelson, D.H. Lee, S.C. Zhang, Phys. Rev. B46 (1992) 2223.
- [15] : F. Chandelier, Y. Georgelin, T. Masson, J.C. Wallet, Phys. Lett. A301 (2002) 451.
- [16]: J. Bellissard, A. van Elst, H. Schultz-Baldes, J. Math. Phys. 35 (1994) 5373; see also J. Avron, R. Seiler, B. Simon, Comm. Math. Phys. 159 (1994) 399.
- [17]: for a review see e.g G. Skandalis, Exp. Math. 9 (1991) 193.
- [18] : T. Masson, J.C. Wallet, in preparation.