## How Multivalency controls Ionic Criticality

Michael E. Fisher, Jean-Noël Aqua,<sup>\*</sup> and Shubho Banerjee<sup>†</sup>

Institute for Physical Science and Technology, University of Maryland, College Park, Maryland 20742, USA

(Dated: August 13, 2018)

To understand how multivalency influences the reduced critical temperatures,  $T_c^*(z)$ , and densities,  $\rho_c^*(z)$ , of z:1 ionic fluids, we study equisized hard-sphere models with z=1-3. Following Debye, Hückel and Bjerrum, association into ion clusters is treated with, also, ionic solvation and excluded volume. In good accord with simulations but contradicting integral-equation and field theories,  $T_c^*$  falls when z increases while  $\rho_c^*$  rises steeply: that 80 - 90% of the ions are bound in clusters near  $T_c$  serves to explain these trends. For  $z \neq 1$  interphase Galvani potentials arise and are evaluated.

PACS numbers: 05.70.Fh,61.20.Qg,64.60.Fr,64.70.Fx

Multivalent ions play a significant role in condensedmatter, physicochemical, biophysical and, via the plasma transition, astrophysical contexts [1]. The effects of multivalency are, however, often hard to comprehend. One central issue—relevant to electrolyte solutions, molten salts, liquid metals, and dense plasmas [1]—arises in Coulomb-driven phase separation. The most basic model for such ionic fluids consists of  $N = \rho V$  hard-core spherical ions of various species  $\sigma$  in a volume V of uniform dielectric constant D, with  $N_{\sigma} = \rho_{\sigma} V$  ions of diameter  $a_{\sigma}$  carrying charges  $q_{\sigma} = z_{\sigma}q_0$ , where  $q_0$  is an elementary charge. In the simple equisized z:1 charge asymmetric primitive models ( $C_z$ APMs), on which we focus here, one has  $\sigma = +, -, a_+ = a_-$ , and  $q_+ = zq_0, q_- = -q_0$ . The basic energy scale and associated reduced temperature and density are then  $\varepsilon = zq_0^2/Da$ ,  $T^* = k_B T/\varepsilon$ ,  $\rho^* = \rho a^3$ .

Monte Carlo simulations [2] show that (at least for  $z \leq 5$ ) the C<sub>z</sub>APMs exhibit "gas-liquid" phase separation; furthermore, the critical parameters,  $T_c^*(z)$  and  $\rho_c^*(z)$ , are found to reasonable precision : see Table I and the open circles in Figs. 1 and 2. One observes that  $T_c^*(z)$ falls with increasing z, while  $\rho_c^*(z)$  rises sharply. But we ask : How can these trends be understood? Or accounted for semiquantitatively? To address this issue we review briefly previous work, including a pioneering fieldtheoretic attack [3], and then report on a recent study [4] which we believe provides significant insight. This extends an earlier analysis I [5] for the symmetric z=1 restricted primitive model (RPM) that was founded on the original Debye-Hückel (DH) approach but incorporated (i) Bjerrum ion pairs and (ii) their *solvation* in the residual ionic fluid. For z=2 and 3 larger ion clusters, trimers and tetramers, must be included [4, 6]; but then explicit results are also obtained for the interphase Galvani potential [7] that appears in any two-phase nonsymmetric ionic system [4, 7].

The field-theoretic analysis of Netz and Orland (NO) [3] was designed to address z:1 ionic fluids and colloids  $(z \gg 1)$  and to include correlations in a systematic manner. The Coulomb interaction,  $q_{\sigma}q_{\tau}/r$ , was transformed to yield a functional integral over an auxiliary potential  $\phi(\mathbf{r})$ . At the  $\langle \phi^2 \rangle$  level the DH effective interaction,



FIG. 1: Reduced critical temperatures for z:1 charge asymmetric equisized hard-core primitive model electrolytes ( $C_z$  APMs) according to Monte Carlo (MC) simulations [2]; Debye-Hückel (DH) theory; field-theoretic approaches : NO [3] (with a factor  $\frac{1}{10}$ ) and NMF [8]; approximate integral equations : MSA [9], SPB, and MPB [10]; and the present DHBjCI and DHBjCIHC solvated ion-cluster theory [4]; See text.

TABLE I: Monte Carlo (MC) estimates [2] for the reduced critical parameters for z:1 equisized hard-sphere electrolytes, values calculated from DHBjCIHC theory (CI) [4], and approximate estimates based on ion cluster statistics : see text.

	critical temp.		$10^2 T_c^*(z)$		critical density		$10^2  ho_c^*(z)$	
z	MC	CI	$E_{\rm DH}$	$E_{\rm MC}$	MC	CI	$\mathbf{E}_{\rho}$	$\mathbf{E}_{\kappa}$
1	$4.93_{3}$	$5.56_{9}$	5.45	$4.93_{5}$	7.50	$2.61_{4}$	2.72	2.37
<b>2</b>	4.70	$4.90_{7}$	5.11	4.65	9.3	$6.26_{1}$	4.27	3.49
3	4.10	$4.33_{4}$	4.85	4.44	12.5	11.90	6.96	5.40

 $v_{\rm DH} \propto e^{-\kappa r}/r$  is captured with

$$\kappa^2(T; \{\rho_\sigma\}) = 4\pi (q_0^2/Dk_B T) \sum_{\sigma} z_{\sigma}^2 \rho_{\sigma} \,. \tag{1}$$

The reduced free energy density,  $\bar{f}(T;\rho) \equiv -F/Vk_BT$ , was computed to eighth order in  $\phi$  but a momentum cutoff is essential: NO adopted  $|\mathbf{k}_{\Lambda}| = 2\pi/a$  thereby incorporating the ionic diameter and, for the z:1 case, leading to



FIG. 2: Reduced critical densities  $\rho_c^*(z)$ , for the C<sub>z</sub>APM electrolyte as in Fig. 1 (except that the NO plot is not rescaled).

 $\kappa^2 a^2 = 4\pi \rho^*/T^*$ . Since this treatment of the hard cores is approximate, accurate predictions for  $T_c^*(z)$  and  $\rho_c^*(z)$ are not expected. Nevertheless, one might anticipate reliable *trends* when z varies in contrast to DH theory which yields *no dependence* on z with (after I)

DH: 
$$\kappa_c a = 1$$
,  $T_c^* = \frac{1}{16}$ ,  $\rho_c^* = 1/64\pi \simeq 0.005$ . (2)

In fact, as NO report, "the [predicted] deviations from DH theory are pronounced" for z > 1: see the bold dashed plots in Figs. 1 and 2.

But evidently the NO results are not merely quantitatively wrong; the *trends* are quite incorrect since  $T_c^*$  is asserted to rise rapidly (instead of falling) while  $\rho_c^*$  falls sharply for small z-1 (instead of rising) and *then* increases but much too slowly. While one may blame the approximate treatment of the hard cores, we believe this is *not* the primary culprit. Indeed, a recent field-theoretic analysis paid closer attention to the ion-ion repulsions [8]; but the subsequent "new mean-field" (NMF) results still exhibit strong increases in  $T_c^*$  and an overly weak variation of  $\rho_c^*$ : see the NMF plots in Figs. 1 and 2 [8].

Integral equation theories are hardly better : see Figs. 1 and 2. The mean spherical approximation (MSA), like DH theory, predicts no variation of  $T_c^*$  and  $\rho_c^*$  with z [9]. A symmetric Poisson-Boltzmann (SPB) theory [10] does predict the correct falling and rising trends for  $T_c^*$ and  $\rho_c^*$ , but the degree of variation is woefully inadequate. Moreover, the modified Poisson-Boltzmann (MPB) approximation, that the same authors [10] argue should be more reliable, yields the wrong trend for  $T_c^*$ .

In order to better understand the effects of multivalency we turn to recent calculations [4, 6] based on the solvated ion-cluster view [5] of the C<sub>z</sub>APM near criticality that is supported 'pictorially' by simulations [2]. In brief, the aim is to construct the free energy density,  $\bar{f}(T; \{\rho_{\sigma}\})$ , for ionic species  $\sigma$  consisting (i) of + and monomers, i.e., isolated,  $n_{+}=n_{-}=1$  single, unassociated



FIG. 3: Coexistence curves predicted for z:1 equisized primitive models by the DHBjCI and DHBjCIHC theories (solid and dashed lines, respectively) together with Monte Carlo estimates (MC) based on [11].

ions of valency  $z_+ = z$  and  $z_- = -1$ ; (ii) a set of associated *primary* clusters,  $\sigma = 2, 3, \ldots$ , dimers, trimers,  $\ldots$ , each consisting of one "central" + ion and  $m_{\sigma} = \sigma - 1$  "satellite" counterions for a total of  $n_{\sigma} = m_{\sigma} + 1$  ions in a cluster of valency  $z_{\sigma} = z - m_{\sigma}$ ; up to (iii), the largest primary cluster, the *neutral* or 'molecular' (z+1)-mer of one  $z_+$  ion and z negative ions [4, 6].

For each species,  $\overline{f}$  contains an ideal-gas term  $\bar{f}^{\text{Id}}(T,\rho_{\sigma})$ , and an electrostatic term  $\bar{f}^{\text{El}}_{\sigma}(T,\{\rho_{\tau}\})$ , that, following DH, incorporates Cluster solvation in the partially associated Ionic fluid: this description is thus dubbed "DHBjCI" [4]. By adding a Hard Core freevolume term,  $\bar{f}^{\text{HC}}(\{\rho_{\sigma}\})$ , as in **I**, one may also account for those excluded volume effects not already encompassed in the basic solvation and association calculations [4, 5], so generating a "DHBjCIHC" theory [4]. (The effective HC virial coefficient  $B_{\sigma}^{\rm bcc} = 4a_{\sigma}^{3/3^{1/2}}$  has been adopted [4, 5].) Examination of Figs. 1 and 2 reveals that these solvated ion-cluster theories are surprisingly successful! Not only are both the downward trend in  $T_c^*(z)$  and the rapid rise of  $\rho_c^*(z)$  well captured, but the quantitative agreement with each of the MC estimates is significantly better than achieved by other approaches.

One must recognize that (all) these theories are of mean-field character: thus 5 to 15% over-estimates of  $T_c^*(z)$  are to be expected. Indeed, neglected fluctuations typically depress  $T_c$  by such amounts and also flatten the *coexistence curves* as seen in Fig. 3. Second, note that the hard-core terms have a small effect on  $\rho_c^*(z)$  while reducing  $T_c^*(z)$  values by only 5–10%. Nevertheless, Fig. 3 reveals that the liquid phases, especially for  $\rho^* \gtrsim 0.15$ , are sensitive to  $\bar{f}^{\rm HC}$ : but, recall the discussion in I. In fact, the crucial feature of DHBj-type theories—not represented in field-theoretic or standard integral-equation treatments—is the *chemical equilibrium* maintained be-

tween the cluster species via the Law of Mass Action:

$$\rho_{\sigma} = K_{m,z}(T) \,\rho_{+} \,\rho_{-}^{m} \,\exp\left[\mu_{+}^{\text{Ex}} + m\mu_{-}^{\text{Ex}} - \mu_{\sigma}^{\text{Ex}}\right] \,, \qquad (3)$$

for  $\sigma = m + 1 \ge 2$ , with the excess chemical potentials  $\mu_{\sigma}^{\text{Ex}} = -(\partial/\partial \rho_{\sigma})[\bar{f}^{HC} + \sum_{\sigma} \bar{f}_{\sigma}^{El}]$ , while the association constants are taken as [4, 5]

$$K_{m,z}(T;R) = \frac{1}{m!} \prod_{i=1}^{m} \int_{a}^{R} d\mathbf{r}_{i} \exp\left(-\frac{\mathcal{E}_{m,z}(\{\mathbf{r}_{i}\})}{k_{B}T}\right), \quad (4)$$

in which  $\mathcal{E}_{m,z}(\{\mathbf{r}_i\})$  is the electrostatic energy of an isolated (m + 1)-mer with satellite coordinates  $\{\mathbf{r}_i\}$ . The lower limits *a* and the condition  $\mathcal{E}_{m,z} = +\infty$  for  $|\mathbf{r}_i - \mathbf{r}_j| < a$ , represent hard cores. Following Bjerrum [5], the necessary cut-off radius *R* is chosen so that  $(\partial K_{m,z}/\partial R)$  is minimal. The resulting 3-fold  $K_{2,z}$  integral is managable but the 6-fold integral for  $K_{3,3}$  requires a Padé approximant study of the low-*T* expansion cross-checked to a part in 10<sup>3</sup> by MC evaluations [4]. It transpires, however, that  $T_c^*$  and  $\rho_c^*$  are *insensitive* to the  $K_{m,z}$  values [4].

Lastly, one needs to account for the solvation of *all* the ion species,  $\sigma$ , by the free ions and charged clusters via the electrostatic terms [4, 5, 6]

$$\bar{f}_{\sigma}^{\scriptscriptstyle El}(T;\{\rho_{\tau}\}) = \frac{4\pi\rho_{\sigma}}{Dk_{\scriptscriptstyle B}T} \sum_{l=0}^{\infty} \frac{u_{2l}(\kappa a_{\sigma})}{a_{\sigma}^{2l+1}} \sum_{m=-l}^{l} \langle |Q_{lm}^{\sigma}|^2 \rangle , \quad (5)$$

where the  $u_{2l}(x)$  are related to the spherical Bessel functions  $k_l(x)$  [4]; the second sum requires the cluster electric multipole moments,  $Q_{l,m}^{\sigma}$ , thermally averaged [4] over the ionic configurations that already enter in the  $K_{m,z}(T)$ .

Finally,  $a_{\sigma}$  is an effective cluster diameter, i.e. the radius of the approximating sphere (centered to minimize  $\bar{f}_{\sigma}^{El}$ ) that substitutes for the true, thermally fluctuating, hard-core exclusion domain: see I and [4]. One concludes, as in I, that a most reasonable choice for  $a_{\sigma}$  is the average over solid angle of the radial distance to the true exclusion surface of the ground-state cluster : this yields  $a_2 = (\frac{3}{4} + \frac{3}{8} \ln 3 \simeq 1.162)a, a_3 = 1.250a$  and  $a_4 = 1.375a$ . For z = 1 the values of  $T_c^*$  and  $\rho_c^*$  vary by less than  $\pm 2\%$ over plausible alternatives for  $a_2$  [4]; but the sensitivity to  $a_3$  and  $a_4$  for z = 2 and 3 is greater. As a result, this hard-to-avoid approximation contributes significantly to the overall quantitative uncertainties.

From the total free energy  $f(T, \{\rho_{\sigma}\})$ , all thermodynamic properties follow [4, 5]. One may then conclude from Figs. 1 and 2 that the principal defect of the fieldtheoretic and integral-equation approaches is a failure to account effectively for strong ionic association near criticality. But can the actual *trends* of  $T_c^*$  and  $\rho_c^*$  with z be demonstrated in a direct, transparent way? To answer, consider the *fractions*,  $y_{\sigma} = n_{\sigma} N_{\sigma}/N$ , of ions bound in clusters of  $n_{\sigma}$  ions with  $\rho_{\sigma} = (y_{\sigma}/n_{\sigma})\rho$ . The critical point

TABLE II: Inverse screening length  $\kappa$  and fractions,  $y_{\sigma} = n_{\sigma}N_{\sigma}/N$ , of ions in clusters of  $n_{\sigma}$  ions at criticality, as percentages, according to DHBjCIHC theory [4].

z	$\kappa_c a$	$y^c_+$	$y^c$	$y_2^c$	$y_3^c$	$y_4^c$
1	1.04	9.14	9.14	81.72	_	_
2	1.37	1.31	10.33	15.43	72.93	_
3	1.57	0.34	8.04	3.32	11.13	77.17

values that result from DHBjCIHC theory [4, 12] are displayed in Table II. A significant fact is the rapid decrease in  $y_{+}^{c}$ , the fraction of unassociated  $z_{+}$  ions, from 9.1 to 1.3 to 0.3%. But more can be learned!

To understand the variation of  $T_c^*(z)$  let us regard the electrolyte in the critical region as a mixture of clusters with fixed mole fractions  $x_{\sigma} = (y_{\sigma}/n_{\sigma})/\sum_{\tau}(y_{\tau}/n_{\tau})$ . A pair  $(\sigma, \tau)$  will either mutually repel or attract with pairwise binding energies, say,  $\varepsilon_{\sigma\tau}$ . Thus unlike monomers attract with  $\varepsilon_{\pm} = \varepsilon$ . However, a dimer attracts only negative monomers with  $\varepsilon_{2-} = (z - \frac{1}{2})\varepsilon/z$ ; but repels all  $z_+ \ge +2$  ions. Two dimers repel when  $z \ge 3$ ; but one has  $\varepsilon_{2,2}/\varepsilon \simeq 0.586$  and 0.345 for z = 1 and 2. And so on.

To estimate  $T_c^*$  for this mixture we adopt a van-der-Waals approach as in [2(d)]. Thus, for the overall cluster density  $\hat{\rho} (= \rho \sum_{\sigma} y_{\sigma}/n_{\sigma})$ , we take  $p/\hat{\rho}k_BT \simeq Z(B_0\hat{\rho}) + B_1(T^*)\hat{\rho}$  with  $Z(u) = 1+u+\ldots$  in which the second virial coefficient has been decomposed as  $B(T^*) = B_0 + B_1(T^*)$ where  $B_0$  (=  $b_0 a^3$ , say) represents the hard-core repulsions while  $B_1(T^*)$  embodies the attractions. Solving  $\partial_{\rho}p = \partial_{\rho}^2 p = 0$ , as usual, yields  $\rho_c^*$  and  $B_c^* \equiv B_1(T_c^*)/b_0 a^3$ . At low T, which is relevant here, one has

$$B_1(T^*) \approx -\sum_{\sigma,\tau} b_{\sigma\tau} a^3 x_{\sigma} x_{\tau} \exp(\varepsilon_{\sigma\tau}^*/T^*), \quad (6)$$

where  $\varepsilon_{\sigma\tau}^* \equiv \varepsilon_{\sigma\tau}/\varepsilon$ , while  $b_{\sigma\tau}a^3$  specifies the volume of mutual attractions: this *vanishes* if  $\sigma$  and  $\tau$  repel.

Now, the  $x_+x_-$  term dominates in  $B_1(T^*)$  at low Twith corrections of relative order  $(x_2^2/x_+x_-)e^{-0.414/T^*}$ for z = 1 and  $2(x_2/x_+)e^{-1/2zT^*}$  for  $z \ge 2$ . We may then calibrate  $B_1(T_c^*)/a^3$  by using pure DH theory (2) for which, since association is not considered,  $x_+ = x_- = \frac{1}{2}$ . Thereby we obtain the  $E_{\rm DH}$  estimates

$$T_c^*(z) \simeq 1/(16 + |\ln 4x_+^c(z)x_-^c(z)|), \qquad (7)$$

in which  $x_{+}^{c} \propto y_{+}^{c}$  and  $x_{-}^{c} \propto y_{-}^{c}$  follow from Table II.

The resulting predictions are listed in Table I under  $E_{DH}$ . In light of the heuristic nature of the arguments, they reflect the trend of the MC and CI values surprizingly well. Certainly the contention that association is a prime factor is well confirmed. By replacing 16 by 20.27 (or 17.96) in (7), and the factor 4 by  $1/x_{+}^{c}(1)x_{-}^{c}(1)$ , one calibrates  $B_{1}(T_{c}^{*})$  on the MC (or CI) values for the RPM. Column  $E_{MC}$  in Table I lists the MC-calibrated values : for z = 2 and 3 these match the Monte Carlo estimates to within 1% and 8%, respectively [13].



FIG. 4: Reduced Galvani potentials,  $\Delta \bar{\phi} = q_0 \Delta \phi / k_B T$ , vs.  $T/T_c$  for z:1 electrolytes according to pure DH theory (dotted) and DHBjCI(HC) theories : solid (dashed) plots.

Now, for the critical density, the significance of ion pairing is already clear in *pure* DHBj theory for the RPM [5]. The heavy depletion of the free ions (which, in DHBj theory, drive the transition *alone*) means that to reach criticality the overall density  $\rho (= \rho_+ + \rho_- + 2\rho_2)$  must be increased until the DH criterion  $\rho_+^* + \rho_-^* = \rho_{\text{DH}}^{*c} = 1/64\pi$ is met : see (2). Does the same depletion-by-association mechanism account for the z-dependence of  $\rho_c^*(z)$ ?

To progress, rewrite (1) generally as  $\kappa^2 a^2 = 4\pi \rho^{\dagger}/T^*$ , with the effective, depleted ionic density

$$\rho^{\dagger} \equiv \rho^* \sum_{\sigma} z_{\sigma}^2 y_{\sigma}(T; \{\rho_{\sigma}\}) / z n_{\sigma} \,. \tag{8}$$

If one accepts the DH criterion and uses Table II, the estimates  $E_{\rho}$ , in Table I, result. Although these fall short of the Monte Carlo values by 74, 54 and 44% for z = 1-3, they reproduce the accelerating increase with z (by factors 1.57, 1.63 vs. 1.24, 1.34).

An alternative approach adopts the DH value  $\kappa_c a = 1$ : see (2) but note from Table II that DHBjCIHC theory implies that  $\kappa_c a$  rises from 1.04 for the RPM to 1.57 for z = 3. Then using the  $E_{DH}$  values for  $T_c^*$ , in Table I, leads to the  $E_{\kappa}$  predictions for  $\rho_c^*(z)$ : these are all rather low but the increases with z, by factors 1.47, 1.55, again reflect the correct behavior.

Finally, we note that the Galvani potential,  $\Delta \phi$ , that arises between coexisting phases in charge asymmetric fluids is readily calculated [4, 7]. The predictions from pure DH theory are shown dotted in Fig. 4 : one finds  $\Delta \phi_{\rm DH} \propto (1-z^{-1})$ . The other plots result from the DH-BjCI and DHBjCIHC theories [4]. Surprisingly, the calculations suggest no clear trend with z. It is natural to conjecture that  $\Delta \phi$  vanishes as  $G_0(T_c - T)^{\beta}$ ; moreover to the extent that the expected mean-field value  $\beta = \frac{1}{2}$  is realized, the present results support this.

In conclusion, we have elucidated the mechanisms underlying how multivalency influences critical behavior. Specifically, we have summarized briefly analytical calculations for 3:1, 2:1 and 1:1 equisized charged hardsphere fluids [4] that, for the first time, reasonably reflect the true variation of critical temperatures and densities,  $T_c^*(z)$  and  $\rho_c^*(z)$  (as revealed by simulations [2]). On that basis, supported by analysis that correlates  $T_c^*(z)$ and  $\rho_c^*(z)$  with the increasingly depleted populations of free ions and charged clusters as z increases, it is clear that recognizing ionic association is inescapable for a successful theory. Previous treatments [3, 8, 9, 10], lacking allowance for ion clusters fail seriously. The ion-cluster solvation theories also yield quantitative results for the interphase Galvani potentials.

National Science Foundation support via Grants CHE 99-81772 and 03-01101 is gratefully acknowledged.

- \* Present address: École normale supérieure de Lyon, 69364, Lyon, France
- <sup>†</sup> Present address: Department of Physics, Rhodes College, Memphis TN 38112
- See, (a) articles in Elementary Processes in Dense Plamas, edited by S. Ichimaru and S. Ogata (Addison-Wesley, New-York, 1995); (b) A. A. Likal'ter, Phys. Uspekhi 43, 777 (2000); (c) A. Yu. Grosberg et al., Rev. Mod. Phys. 74, 329 (2000); (d) H. Weingärtner and W. Schröer, Adv. Chem. Phys. 116, 1 (2001); (e) Y. Levin, Rep. Prog. Phys. 65, 1377 (2002).
- [2] (a) P. J. Camp and G. N. Patey, J. Chem. Phys. 111, 9000 (1999); (b) Q. Yan and J. J. de Pablo, Phys. Rev. Lett. 86, 2054 (2001); (c) A. Z. Panagiotopoulos and M. E. Fisher, Phys. Rev. Lett. 88, 045701 (2002); (d) Y. C. Kim and M. E. Fisher, Phys. Rev. Lett. 92, 185703 (2004).
- [3] R. R. Netz and H. Orland, Europhys. Lett. 45, 726 (1999).
- [4] J.-N. Aqua, S. Banerjee, and M. E. Fisher, condmat/0410692 [submitted for publication].
- [5] (a) M. E. Fisher and Y. Levin, Phys. Rev. Lett. 71, 3826 (1993); (b) Y. Levin and M. E. Fisher, Physica A 225, 164 (1996); to be denoted I.
- [6] M. N. Artyomov et al., J. Chem. Phys. 118, 6394 (2003).
- [7] M. J. Sparnaay, *The Electrical Double Layer* (Pergamon Press, Oxford, 1972), pp. 4, 338; P. B. Warren, J. Chem. Phys. **112**, 4683 (2000); I. Iosilevski and A. Chigvintsev, J. Phys. IV France **10**, Pr5-451 (2000); A. R. Muratov, Zh. Éksp Teor. Fiz. **120**, 104 (2001) [Sov. Phys. JETP **93**, 89 (2001)].
- [8] J.-M. Caillol, Mol. Phys. 103, 1271 (2005); we quote the (most favorable) "MSA regularization" results.
- [9] E. Waisman and J. L. Lebowitz, J. Chem. Phys. 56, 3086 (1972); E. Gonzalez-Tovar, Mol. Phys. 97, 1203 (1999).
- [10] A. K. Sabir *et al.*, Mol. Phys. **93**, 405 (1998); C. W. Outhwaite, Cond. Matt. Phys. **7**, 719 (2004).
- [11] Y. C. Kim, M. E. Fisher, and E. Luijten, Phys. Rev. Lett. 91, 065701 (2003).
- [12] While the  $y_{\sigma}^{c}$  are not very sensitive to the  $a_{\sigma}$ , they cannot be determined unequivocally via simulations.
- [13] The arguments based on (6) extend naturally to charged

hard dumbbell models and their variants : see J. M. Romero-Enrique et~al., Phys. Rev. E 66,~041204~(2002).