

Spectral microscopic mechanisms and quantum phase transitions in a 1D correlated problem

J. M. P. Carmelo

Department of Physics, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139-4307, USA

K. Penc

Research Institute for Solid State Physics and Optics,
H-1525 Budapest, P.O.B. 49, Hungary

Abstract. In this paper we study the dominant microscopic processes that generate more than 99% of the one-electron removal and addition spectral-weight distributions of the one-dimensional Hubbard model for all values of the on-site repulsion U . We find that for the doped Mott-Hubbard insulator there is a competition between the microscopic processes that generate the one-electron upper-Hubbard band spectral-weight distributions of the Mott-Hubbard insulating phase and finite-doping-concentration metallic phase, respectively. The spectral-weight distributions generated by the non-perturbative processes studied here are shown elsewhere to agree quantitatively for the whole momentum and energy bandwidth with the peak dispersions observed by angle-resolved photoelectron spectroscopy in quasi-one-dimensional compounds.

PACS numbers: 71.20.-b, 71.10.Pm, 72.15.Nj, 71.27.+a

1. INTRODUCTION

The description of the microscopic mechanisms behind the unusual finite-energy spectral properties observed in low-dimensional materials remains until now an interesting open problem. For energies larger than the transfers integrals for electronic hopping between the chains, the one-dimensional (1D) Hubbard Hamiltonian is the simplest model for the description of electronic correlation effects on the spectral properties of quasi-1D compounds [1, 2, 3, 4]. It reads,

$$\hat{H} = -t \sum_{j,\sigma} [c_{j,\sigma}^\dagger c_{j+1,\sigma} + h.c.] + U \sum_j \hat{n}_{j,\uparrow} \hat{n}_{j,\downarrow}, \quad (1)$$

where $c_{j,\sigma}^\dagger$ and $c_{j,\sigma}$ are spin-projection $\sigma = \uparrow, \downarrow$ electron operators at site $j = 1, 2, \dots, N_a$, $\hat{n}_{j,\sigma} = c_{j,\sigma}^\dagger c_{j,\sigma}$, and t is the transfer integral. In contrast to other interacting models [5] and in spite of the model exact solution [6, 7], for finite values of the on-site repulsion U little is known about the non-perturbative microscopic processes that control its finite-energy spectral properties.

Recently, the problem was studied in Refs. [8, 9] by the use of a pseudofermion description. The preliminary predictions of the method introduced in these references agree quantitatively for the whole momentum and energy bandwidth with the peak dispersions observed by angle-resolved photoelectron spectroscopy in the quasi-1D conductor TTF-TCNQ [1, 2] and are consistent with the phase diagram observed for the (TMTTF)₂X and (TMTSF)₂X series of compounds [4]. More recently, results for the TTF-TCNQ spectrum consistent with those of Refs. [1, 2] were obtained by the dynamical density matrix renormalization group method [3]. Within the method of Refs. [8, 9], the finite-energy spectral properties are controlled by the overall pseudofermion phase shifts, through the pseudofermion anticommutators [10]. Such a method is a generalization for all values of the on-site repulsion U of the technique introduced in Refs. [11, 12] for $U \rightarrow \infty$.

The studies of this paper are complementary to those of Refs. [8, 9] and focus on the specific case of the one-electron spectral weight. Our motivation is to contribute to the further understanding of the non-perturbative microscopic processes which generate the finite-energy one-electron experimental spectral features. The pseudofermion description used in the derivation of the general expressions of Ref. [8] for the one- and two-electron spectral weight distributions of the model metallic phase is directly related to the holon, spinon, and c pseudoparticle representation introduced in Ref. [13]. Such objects emerge as a result of the electron - rotated-electron unitary transformation. On the other hand, the c pseudofermions and composite pseudofermions are generated from the c pseudoparticles and composite pseudoparticles of Ref. [13] by a second unitary transformation [8]. Such a transformation introduces shifts of order $1/L$ in the pseudoparticle discrete momentum values and leaves all other pseudoparticle properties invariant, where L is the electronic lattice length. As a result of such momentum shifts and in contrast to the c pseudoparticles and composite pseudoparticles, the corresponding pseudofermions have no residual-interaction energy terms. This

greatly simplifies the evaluation of matrix elements between energy eigenstates [8, 9]. However, it is required that the involved electronic operators are expressed in terms of pseudofermion operators.

In this paper we study the generators of the one-electron dominant microscopic processes defined in Refs. [8, 9] and express them in terms of pseudofermion operators. Our results confirm that also for finite values of the lattice length L more than 99% of the one-electron removal and addition spectral weight distributed over the whole (k, ω) plane is generated by such dominant processes. The quantum phase transitions of the model are controlled by the interaction dependence of the local entanglement [14]. Our results provide further information about the interplay between the microscopic processes that generate the one-electron spectral-weight distributions and the metal - Mott-Hubbard insulator quantum phase transition. Since the mechanisms found here are expected to occur in other correlated models, our results are of interest for the further general understanding of the microscopic mechanisms associated with the quantum phase transitions.

The model (1) describes N_\uparrow spin-up electrons and N_\downarrow spin-down electrons in a chain of N_a sites. We denote the electronic number by $N = N_\uparrow + N_\downarrow$. The number of lattice sites N_a is even and very large. For simplicity, we use units such that both the lattice spacing a and the Planck constant are one. In these units the chain length L is such that $L = N_a a = N_a$. Our results refer to periodic boundary conditions. We consider an electronic density $n = n_\uparrow + n_\downarrow$ in the range $0 < n \leq 1$ and a spin density $m = n_\uparrow - n_\downarrow = 0$, where $n_\sigma = N_\sigma/L$ and $\sigma = \uparrow, \downarrow$. We introduce the Fermi momenta which, in the thermodynamic limit $L \rightarrow \infty$, are given by $\pm k_{F\sigma} = \pm \pi n_\sigma$ and $\pm k_F = \pm [k_{F\uparrow} + k_{F\downarrow}]/2 = \pm \pi n/2$

Our results refer to the one-electron spectral function $B^l(k, \omega)$ such that $l = -1$ (and $l = +1$) for electron removal (and addition). It is given by,

$$\begin{aligned} B^{-1}(k, \omega) &= \sum_{\sigma} \sum_f |\langle f | c_{k,\sigma} | GS \rangle|^2 \delta(\omega + E_{f,N-1} - E_{GS}), \quad \omega < 0; \\ B^{+1}(k, \omega) &= \sum_{\sigma} \sum_{f'} |\langle f' | c_{k,\sigma}^\dagger | GS \rangle|^2 \delta(\omega - E_{f',N+1} + E_{GS}), \quad \omega > 0. \end{aligned} \quad (2)$$

Here $c_{k,\sigma}$ (and $c_{k,\sigma}^\dagger$) are electron annihilation (and creation) operators of momentum k and $|GS\rangle$ denotes the initial N -electron ground state. The f and f' summations run over the $N - 1$ and $N + 1$ -electron excited states, respectively, and $[E_{f,N-1} - E_{GS}]$ and $[E_{f',N+1} - E_{GS}]$ are the corresponding excitation energies.

The Hamiltonian (1) commutes with the generators of the η -spin and spin $SU(2)$ algebras [13, 15, 16]. Here we call the η -spin and spin of the energy-eigenstates η and S , respectively, and the corresponding projections η_z and S_z . All such states can be described in terms of occupancy configurations of η -spin 1/2 holons, spin 1/2, spinons, and η -spin-less and spin-less $c0$ pseudoparticles [13]. (The latter objects are called c pseudoparticles in Refs. [13, 17].) We use the notation $\pm 1/2$ holons and $\pm 1/2$

spinons according to the values of the η -spin and spin projections, respectively. For large values of U/t the $+1/2$ holons and $-1/2$ holons become the holons and doublons, respectively, used in the studies of Ref. [18]. The electron - rotated-electron unitary transformation [13] maps the electrons onto rotated electrons such that rotated-electron double occupation, unoccupancy, and spin-up and spin-down single occupation are good quantum numbers for all values of U . The $\pm 1/2$ holons of charge $\pm 2e$ and zero spin and the charge-less $\pm 1/2$ spinons are generated from the electrons by that unitary transformation. The corresponding holon and spinon number operators $\hat{M}_{c,\pm 1/2}$ and $\hat{M}_{s,\pm 1/2}$, respectively, are of the form given in Eq. (24) of Ref. [13] and involve the electron - rotated-electron unitary operator.

While the $-1/2$ and $+1/2$ holons refer to the rotated-electron doubly occupied and unoccupied sites, respectively, the $-1/2$ and $+1/2$ spinons correspond to the spin degrees of freedom of the spin-down and spin-up rotated-electron singly occupied sites, respectively. The charge degrees of freedom of the latter sites are described by the spinless and η -spin-less $c0$ pseudoparticles, which are composite objects of a chargeon and a antichargeon, and thus carry charge $-e$ or $+e$ for the description of the transport of charge in terms of electrons and electronic holes, respectively [13]. Here $-e$ denotes the electronic charge. The $c\nu$ pseudoparticles (and $s\nu$ pseudoparticles) such that $\nu = 1, 2, \dots$ are η -spin singlet (and spin singlet) 2ν -holon (and 2ν -spinon) composite objects. Thus, $M_{\alpha,\pm 1/2} = L_{\alpha,\pm 1/2} + \sum_{\nu=1}^{\infty} \nu N_{\alpha\nu}$ where $\alpha = c, s$, $N_{\alpha\nu}$ denotes the number of $\alpha\nu$ pseudoparticles, and $L_{c,\pm 1/2} = \eta \mp \eta_z$ and $L_{s,\pm 1/2} = S \mp S_z$ gives the number of $\pm 1/2$ Yang holons and $\pm 1/2$ HL spinons, respectively. Those are the holons and spinons that are not part of composite pseudoparticles. All energy eigenstates can be described by occupancy configurations of $\alpha\nu$ pseudoparticles, $-1/2$ Yang holons, and $-1/2$ HL spinons [13]. For the ground state, $N_{c0} = N$, $N_{s1} = N_{\downarrow}$, and $N_{c\nu} = N_{s\nu'} = L_{\alpha,-1/2} = 0$ for $\alpha = c, s$, $\nu > 0$, and $\nu' > 0$. The deviations $\Delta N_{\alpha\nu}$, $\Delta N_{\alpha\nu}^h$, $\Delta L_{c,-1/2}$, $\Delta L_{s,-1/2}$, $\Delta M_{c,-1/2}$, $\Delta M_{s,-1/2}$, ΔM_c , ΔM_s of the above numbers which result from the ground-state - excited-energy-eigenstate transitions play a major role in our study. It follows from the values of the ground-state numbers that $\Delta N_{c\nu} = N_{c\nu}$, $\Delta N_{s\nu'} = N_{s\nu'}$, $\Delta L_{c,-1/2} = L_{c,-1/2}$, $\Delta L_{s,-1/2} = L_{s,-1/2}$, $\Delta M_{c,-1/2} = M_{c,-1/2}$, and $\Delta M_{s,-1/2} = M_{s,-1/2}$ for $\nu > 0$ and $\nu' > 1$. Thus, often we replace the latter deviations by the corresponding numbers. A concept widely used in our studies is that of a CPHS ensemble subspace [8, 9]. (Here CPHS stands for $c0$ pseudofermion, holon, and spinon.) Such a subspace is spanned by all energy eigenstates with fixed values for the $-1/2$ Yang holon number $L_{c,-1/2}$, $-1/2$ HL spinon number $L_{s,-1/2}$, and for the sets of $\alpha\nu$ pseudofermion numbers $\{N_{c\nu}\}$ and $\{N_{s\nu'}\}$ corresponding to the $\nu = 0, 1, 2, \dots$ and $\nu' = 1, 2, \dots$ branches.

It is useful for our study to consider the pseudofermion subspace (PS). In our case it is spanned by an initial ground state $|GS\rangle$ and all excited energy eigenstates contained in the one-electron excitations $c_{j,\sigma}^{\dagger}|GS\rangle$ and $c_{j,\sigma}|GS\rangle$ [8]. The above unitary transformation which maps the $\alpha\nu$ pseudoparticle onto the $\alpha\nu$ pseudofermion is defined in the PS. The $\alpha\nu$ pseudoparticle has discrete bare-momentum values $q_j = [2\pi/L]I_j^{\alpha\nu}$ such that $I_j^{\alpha\nu}$ are consecutive integers or half-odd integers [13]. These values are good quantum numbers

whose allowed occupancies are one and zero only [13]. The $\alpha\nu$ pseudofermion has discrete canonical-momentum values given by [8, 9, 10],

$$\bar{q}_j = \bar{q}(q_j) = q_j + Q_{\alpha\nu}^{\Phi}(q_j)/L, \quad (3)$$

where $j = 1, 2, \dots, N_{\alpha\nu}^*$, $N_{\alpha\nu}^* = N_{\alpha\nu} + N_{\alpha\nu}^h$, and $N_{\alpha\nu}^h$ denotes the number of $\alpha\nu$ pseudofermion (and $\alpha\nu$ pseudoparticle) holes whose value is defined in Ref. [13]. Such a canonical-momentum pseudofermion is related to the *local $\alpha\nu$ pseudofermion* by a Fourier transform. The latter object occupies the sites of the effective $\alpha\nu$ lattice [17]. Except for the discrete momentum values, the above pseudoparticle and pseudofermion have the same properties. Thus, all the energy eigenstates that span the PS can be described by occupancy configurations of $\alpha\nu$ pseudofermions, $-1/2$ Yang holons, and $-1/2$ HL spinons. The functional,

$$Q_{\alpha\nu}^{\Phi}(q_j)/2 = \pi \sum_{\alpha'\nu', j} \Phi_{\alpha\nu, \alpha'\nu'}(q_j, q_{j'}) \Delta N_{\alpha'\nu'}(q_{j'}), \quad (4)$$

of equation (3) is a pseudofermion or pseudofermion hole scattering phase shift [10]. Here $\Delta N_{\alpha\nu}(q_j) \equiv N_{\alpha\nu}(q_j) - N_{\alpha\nu}^0(q_j)$ is the $\alpha\nu$ branch bare-momentum distribution-function deviation relative to the ground state value and $\pi \Phi_{\alpha\nu, \alpha'\nu'}(q, q')$ is an elementary *two-pseudofermion phase shift* [10]. Note that $Q_{\alpha\nu}^{\Phi}(q_j) = 0$ for the initial ground state and thus $\bar{q}_j = q_j$ for that state.

For the $\mathcal{N} = 1$ electron problem, the operator $\hat{O}_{\mathcal{N}, j}^l = \hat{O}_{1, j}^l$ of the spectral-function expressions of Eq. (7) of Ref. [9] is the operator $c_{j, \sigma}$ for $l = -1$ and $c_{j, \sigma}^{\dagger}$ for $l = +1$. These expressions can be re-expressed in terms of the operator $\hat{\Theta}_{\mathcal{N}, j}^l$, which plays an important role in our study and is defined in terms of the original \mathcal{N} -electron operator $\hat{O}_{\mathcal{N}, j}^l$ in Eqs. (27) and (28) of Ref. [9]. Its expression in terms of local creation and annihilation rotated-electron operators is of the general form,

$$\hat{\Theta}_{\mathcal{N}, j}^l = \tilde{\Theta}_{\mathcal{N}_0, j}^l + \sum_{i=1}^{\infty} \sqrt{c_i^l} \tilde{\Theta}_{\mathcal{N}_i, j}^l; \quad j = 1, 2, \dots, N_a; \quad l = \pm 1, \quad (5)$$

where for $i > 0$ the index $i = 1, 2, \dots$ gives the number of extra pairs of creation and annihilation rotated-electron operators in the expressions of the operators $\tilde{\Theta}_{\mathcal{N}_i, j}^l$ relative to that of $\tilde{\Theta}_{\mathcal{N}_0, j}^l$ and the value of the constant c_i^l reads $c_0^l = 1$ and for $i > 0$ is a function of n , m , and U/t such that $c_i^l \rightarrow 0$ as $U/t \rightarrow \infty$ [9]. Moreover, the operators $\hat{\Theta}_{\mathcal{N}, j}^l$ and $\tilde{\Theta}_{\mathcal{N}_0, j}^l$ of Eq. (5) have the same expression in terms of local creation and annihilation electron and rotated-electron operators, respectively, and thus $\mathcal{N}_0 = \mathcal{N}$ and $\mathcal{N}_i = [\mathcal{N} + 2i]$. For the $\mathcal{N} = 1$ electron problem considered here, let us call $\tilde{\theta}_{i, j, \sigma}$ and $\tilde{\theta}_{i, j, \sigma}^{\dagger}$ the operators $\tilde{\Theta}_{\mathcal{N}_i, j}^{-1}$ and $\tilde{\Theta}_{\mathcal{N}_i, j}^{+1}$, respectively, on the right-hand side of Eq. (5), whose number of elementary rotated-electron operators is $\mathcal{N}_i = [1 + 2i]$. According to the general studies of Refs. [8, 9], the $i = 0$ operator $\tilde{\Theta}_{\mathcal{N}_0, j}^l$ generates nearly the whole spectral weight of the corresponding \mathcal{N} -electron problem for all values of U/t and $L \rightarrow \infty$.

In the ensuing section we confirm that for finite values of the chain length L the processes generated by application onto the ground state of the $i = 0$ operators $\tilde{\theta}_{0, j, \sigma}$

and $\tilde{\theta}_{0,j,\sigma}^\dagger$ lead to more than 99% of the one-electron spectral weight for all values of U/t . Our results refer both to the Mott-Hubbard insulating and metallic phases. Similar results are reached below for these phases for $L \rightarrow \infty$ and large values of U/t , where the problem can be handled analytically. Since such operators generate the dominant microscopic processes that control the one-electron weight distribution, in Sec. III we calculate their expressions. In Sec. IV we study the main microscopic effects of the metal - Mott-Hubbard insulator quantum phase transition onto the one-electron spectral properties. Finally, the concluding remarks are presented in Sec. V.

2. ONE-ELECTRON SPECTRAL-WEIGHT DOMINANT PROCESSES

According to the results of Ref. [9] for general \mathcal{N} -electron spectral functions, application onto the ground state of the operator $\tilde{\Theta}_{\mathcal{N}_0,j}^l$ generates transitions to excited energy eigenstates whose $M_{c,-1/2}$, $L_{c,-1/2}$, $N_{c\nu}$ values obey for $\nu > 0$ the following exact charge selection rule,

$$M_{c,-1/2} = L_{c,-1/2} + \sum_{\nu=1}^{\infty} \nu N_{c\nu} = 0, 1, 2, \dots, \sum_{l_s=\pm 1} \bar{\mathcal{N}}_{-1,l_s}^l. \quad (6)$$

For $l = -1$ (and $l = +1$) the numbers $\bar{\mathcal{N}}_{-1,l_s}^l$ are those specific to the \mathcal{N} -electron operator $\hat{\Theta}_{\mathcal{N},j}^l$. $\bar{\mathcal{N}}_{-1,l_s}^l$ is for $l = -1$ (and $l = +1$) the number of local electron creation and annihilation operators for $l_c = -1$ and $l_c = +1$, respectively, and with spin down and spin up for $l_s = -1$ and $l_s = +1$, respectively, of the expression of the operator $\hat{\Theta}_{\mathcal{N},j}^l$.

Moreover, the values of the numbers of $-1/2$ Yang holons and $-1/2$ HL spinons generated by application onto the ground state of the corresponding \mathcal{N} -electron operator $\hat{O}_{\mathcal{N},j}^l$ are restricted to the following ranges,

$$\begin{aligned} L_{c,-1/2} &= 0, 1, 2, \dots, \sum_{l_s=\pm 1} \mathcal{N}_{-1,l_s}^l; \\ L_{s,-1/2} &= 0, 1, 2, \dots, \sum_{l_c,l_s=\pm 1} \delta_{l_c,l_s} \mathcal{N}_{l_c,l_s}^l, \end{aligned} \quad (7)$$

respectively. Here the numbers \mathcal{N}_{l_c,l_s}^l have the same definition as the corresponding numbers $\bar{\mathcal{N}}_{l_c,l_s}^l$, but refer to the original \mathcal{N} -electron operator $\hat{O}_{\mathcal{N},j}^l$.

While the selection rules (6) and (7) are exact for the \mathcal{N} -rotated-electron and \mathcal{N} -electron problems, respectively, direct evaluation of the weights by the method introduced of Refs. [8, 9] reveals that 94% to 98% of the spectral weight generated by the operator $\tilde{\Theta}_{\mathcal{N}_0,j}^l$ corresponds to transitions to excited energy eigenstates whose $L_{s,-1/2}$ and $N_{s\nu}$ values for $\nu > 1$ are in the following range,

$$L_{s,-1/2} + \sum_{\nu=2}^{\infty} (\nu - 1) N_{s\nu} = 0, 1, 2, \dots, \sum_{l_c,l_s=\pm 1} \delta_{l_c,l_s} \bar{\mathcal{N}}_{l_c,l_s}^l, \quad (8)$$

where the numbers $\bar{\mathcal{N}}_{l_c,l_s}^l$ are the same as above.

One-electron processes associated with excited energy eigenstates whose deviations do not obey the rule (6) are generated by the operators $\tilde{\theta}_{i,j,\sigma}$ or $\tilde{\theta}_{i,j,\sigma}^\dagger$ such that $i > 0$. We confirm below that application onto the ground state of all such $i > 0$ operators amounts for less than 1% of the (k, ω) -plane one-electron removal or addition spectral weight.

We start by confirming that the one-electron processes which generate excited energy eigenstates obeying the ground-state charge selection rule (6) are dominant for finite values of L and correspond to 99.75% - 100.00% of the whole electronic spectral weight for all values of U/t . For finite values of L the thermodynamic Bethe-ansatz equations introduced by Takahashi [7] do not apply, whereas the Bethe-ansatz solution and the concept of a rotated electron remain valid. Therefore, for finite values of L we can characterize the excited energy eigenstates in terms of rotated-electron occupancy configurations and thus of the related $\pm 1/2$ holon, $\pm 1/2$ spinon, and $c0$ pseudofermion number deviations [17]. Moreover, for zero-magnetization initial ground states the number of excited-energy-eigenstate $s1$ pseudofermion holes equals the number of holes in the corresponding Bethe-ansatz spin excitation spectrum. Thus, one can classify the finite- L processes by the deviation values ΔN_{c0} , $M_{c,-1/2}$, $\Delta M_{s,-1/2}$, ΔM_c , ΔM_s , and ΔN_{s1}^h .

The correlation energy 2μ , defined in Eq. (21) of Ref. [8], plays an important role in the model finite-energy spectrum. The limiting values of such a correlation energy are given in Eq. (13) of Ref. [9]. At half filling it equals the Mott-Hubbard gap, $2\mu = E_{MH}$. Our considerations refer mostly to one-electron addition. For simplicity, let us assume that the initial ground state has zero spin density. In this case the spectral-weight distribution associated with creation of a spin-up electron has the same form as that associated with creation of a spin-down electron. Here we consider the former case. The spin-up electron creation operator is related by Eq. (5) with $l = +1$ and $\mathcal{N} = \mathcal{N}_0 = 1$ to the spin-up rotated-electron creation operator. For creation of a spin-up rotated electron, the selection rule (6) only allows the two values $\Delta M_{c,-1/2} = M_{c,-1/2} = 0, 1$. From the relation of the electron and rotated-electron numbers to the holon, spinon, and $c0$ pseudofermion numbers [13, 17], we find that for creation of a spin-up rotated electron the following transitions obey the selection rule (6):

(i) Lower-Hubbard band (LHB) transitions such that $\Delta N_{c0} = 1$, $\Delta M_{c,-1/2} = 0$, $\Delta M_{s,-1/2} = 0$, $\Delta M_c = -1$, and $\Delta M_s = 1$. The minimal excitation energy for such transitions is zero. The sub-class of these transitions that also obey the restrictions of Eq. (8) are such that $\Delta N_{s1}^h = 1$.

(ii) Upper-Hubbard band (UHB) transitions such that $\Delta N_{c0} = -1$, $\Delta M_{c,-1/2} = 1$, $\Delta M_{s,-1/2} = -1$, $\Delta M_c = 1$, and $\Delta M_s = -1$. The minimal excitation energy for such transitions is 2μ . The sub-class of these transitions that also obey the restrictions of Eq. (8) correspond to a $s1$ pseudofermion-hole deviation value $\Delta N_{s1}^h = 1$.

The simplest transitions that do not obey the selection rule (6) involve creation of three $c0$ pseudofermion holes and three holons:

(iii) Second-UHB transitions such that $\Delta N_{c0} = -3$, $\Delta M_{c,-1/2} = 2$, $\Delta M_{s,-1/2} = -2$, $\Delta M_c = 3$, and $\Delta M_s = -3$. The minimal excitation energy for such transitions is 4μ . The sub-class of these transitions that also obey the restrictions of Eq. (8) are such that $\Delta N_{s1}^h = 1$.

The simplest transitions of types (i) and (ii) that do not obey the restrictions (8) involve creation of three $s1$ pseudofermion holes:

(i') and (ii') transitions with the same values for the deviations ΔN_{c0} , $\Delta M_{c,-1/2}$, $\Delta M_{s,-1/2}$, ΔM_c , ΔM_s as for the above general (i) and (ii) transitions, respectively, and $\Delta N_{s1}^h = 3$. The minimal excitation energy for such transitions is zero and 2μ , respectively.

For the Mott-Hubbard insulator quantum phase such that $n = 1$ one often shifts the ground-state zero-energy level to the middle of the Mott-Hubbard gap. Then the excitation energies 0 , E_{MH} , and $2E_{MH}$ become 0 , $E_{MH}/2$, and $3E_{MH}/2$, respectively. Here we call *three-holon states* the excited energy eigenstates of the transitions (iii) because these involve the creation of three holons. Such excited energy eigenstates also involve the annihilation of three $c0$ pseudofermions. Moreover, we call *three-s1-hole states* the excited energy eigenstates of transitions (i') and (ii'). These involve creation of three $s1$ pseudofermion holes. Finally, LHB and UHB excited energy eigenstates belonging to the sub-class of the general states generated by both the transitions (i) and (ii) such that $\Delta N_{s1}^h = 1$ are called one-holon and one-s1-hole states.

We start by evaluating the relative weight of the excited states of type (iii) (or states of types (iii), (i'), and (ii')) relative to the weight of the states of types (i) and (ii) (or states of types (i) and (ii) such that $\Delta N_{s1}^h = 1$) when all these states are generated by application onto the ground state of the spin-up electron creation operator. (The types of excited energy eigenstates refer to the above transitions which generate these states from the ground state.) To assess the importance for finite values of L of the three-holon final states of type (iii) and three-s1-hole states of types (i') and (ii'), we perform an exact diagonalization of small chains. There is no one-to-one correspondence between the small-chain weight associated with each specific excited energy eigenstate and the excited-energy-eigenstate weight for $L \rightarrow \infty$. However, there is such a correspondence for the spectral-weight sum rules of the states of types (i) and (ii) generated by dominant processes, relative to the sum rule of the states of type (iii). Moreover, within the excited energy eigenstates generated by dominant processes, we consider the relative weight of the states of types (i') and (ii'). For these spectral-weight sum rules the small-chain results provide values for the relative weights which agree up to 99% with the corresponding thermodynamic-limit values.

The full electron addition and removal spectrum for six sites with six electrons (half filling) is shown in Fig. 1 for $U = 12t$. The first Hubbard bands at $\pm E_{MH}/2$ are generated by dominant processes, whereas the second Hubbard bands at $+3E_{MH}/2$ and $-3E_{MH}/2$ result from processes generated by the above $i > 0$ operators $\tilde{\theta}_{i,j,\sigma}^\dagger$ and $\tilde{\theta}_{i,j,\sigma}$, respectively. Note that the first and second Hubbard bands are well separated. The weights of the latter bands are orders of magnitude smaller than the contribution




Figure 1. Electron addition ($\omega > 0$) and removal ($\omega < 0$) spectrum for the half-filled six-site ring, with $U = 12t$. Note the logarithmic scale for the weights. The five-holon states are not shown. Indeed, the contributions of these states are extremely small and their energies are out of the shown energy window.

from the first Hubbard bands. The states centered around $3E_{MH}/2$ are three-holon states of type (iii). As a result of the half-filling particle-hole symmetry, there is a corresponding structure for electron removal centered around $-3E_{MH}/2$. The latter structure is associated with creation of two rotated-electron unoccupied sites. (We recall that the $-1/2$ holon number deviation selection rule (6) refers to excited-energy-eigenstate electronic densities such that $n < 1$. There is a corresponding $+1/2$ holon number deviation selection rule for $n > 1$.)

In figure 2 we plot the contribution of different excited energy eigenstates to the sum rule for half filling. For that, we have followed adiabatically the weights of different states as the value of U/t is reduced, and summed the weight over the particular family of states. Analysis of the figure reveals that the contribution of the three-holon excited energy eigenstates of type (iii) to the total sum rule is largest at intermediate values $U \approx 4t$. It does not exceed 0.25% in the total sum rule. For large values of U/t it decreases as $(t/U)^4$, while for small values as $(U/t)^4$. The three- s 1-hole contribution of the above states of type (ii') is also less than 0.25 %, and for small values of U/t it decreases as $(U/t)^2$. (These states belong to a sub-class of the excited energy eigenstates of type (ii).) While figure 2 corresponds to $n = 1$, the relative spectral weight of the excited energy eigenstates of type (iii) decreases for decreasing values of the electronic density n . For instance, at quarter filling such a weight is 2% of that of half filling and vanishes as $n \rightarrow 0$. Thus, at quarter filling and $U \approx 4t$ the excited energy eigenstates of types (i) and (ii) (including the states of types (i') and (ii')) correspond to $\approx 99.99\%$ of the total spectral weight and the states (iii) to $\approx 0.005\%$ of such a weight.

Evaluation for the model metallic phase of the available spectral-weight contributions by the method of Refs. [8, 9], confirms that the above results remain valid for $L \rightarrow \infty$. The exception is the relative weight of the excited energy eigenstates




Figure 2. The contribution of different states to the electron-addition sum rule for half filling. Over 99% of the sum rule is exhausted by one-holon and one- $s1$ pseudofermion hole excited energy eigenstates of type (ii). For larger systems this remains true if we also consider the states of type (ii'), which as the states of type (ii), are also associated with dominant processes generated by the $i = 0$ operator $\tilde{\theta}_{0,j,\sigma}^\dagger$.

of types (i') and (ii'), which increases for increasing values of L . This is confirmed by the values given in the Table, which displays the relative weights generated by the transitions to the one- $s1$ -pseudofermion-hole excited energy eigenstates that obey the relation (8) both for the one-electron removal and addition spectral functions. These weights were obtained for the one-electron spectral functions as $U/t \rightarrow \infty$ [12]. Thus, they provide the one-rotated-electron relative weights for all values of U/t and the number of rotated electrons N given in the Table [17]. For $L \rightarrow \infty$ the preliminary use of the methods of Refs. [8, 9] leads to values of the relative weights for one-rotated-electron removal and addition of approximately 98% and 94%, respectively. The numbers provided in the Table confirm that also for smaller values of L the deviation value restrictions of Eq. (8) refer to a sub-class of excited energy eigenstates associated a substantial part of the full one-rotated-electron spectral weight. On the other hand, the set of excited energy eigenstates which obey the exact selection rule (6) is larger: it corresponds to the whole one-rotated-electron spectral weight. The excited energy eigenstates of types (i') and (ii') are associated with dominant processes generated by the $i = 0$ operator $\tilde{\theta}_{0,j,\sigma}^\dagger$. In contrast, the states of type (iii) are not generated by dominant processes and remain having very little spectral weight as $L \rightarrow \infty$.

Let us next consider the limit of large U/t , where the $L \rightarrow \infty$ problem can be handled analytically by the method of Ref. [12]. In that case the general $c0$ pseudofermions of Refs. [8, 9] become the spin-less fermions used in the studies of Refs. [11, 12]. Our first goal is the confirmation that the contribution from the three-holon states of type (iii) is very small and leads nearly to the same relative weight both

for $L \rightarrow \infty$ and finite values of L . For large values of U/t one can derive a systematic t/U expansion for the electron - rotated-electron unitary operator defined by Eqs. (21)-(23) of Ref. [13]. (See reference [19], which studies that transformation for large values of U/t .) By use of the inverse of the relation between the one-electron and one-rotated-electron operators given in Eq. (19) of Ref. [13] in such a t/U expansion, we find after some algebra the following partial sum rule,

$$\int \mathcal{A}^{2\text{UHB}}(\omega) d\omega = \frac{t^4}{U^4} \left\langle \frac{3}{2} - 2\mathbf{S}_0\mathbf{S}_1 - 2\mathbf{S}_1\mathbf{S}_2 - 2\mathbf{S}_0\mathbf{S}_2 \right\rangle_{\text{spin}} \times \langle \hat{n}_{x_0, c0} \hat{n}_{x_1, c0} \hat{n}_{x_2, c0} \rangle, \quad (9)$$

where $\hat{n}_{x_j, c0} = f_{x_j, c0}^\dagger f_{x_j, c0}$ and $f_{x_j, c0}^\dagger$ and $f_{x_j, c0}$ are the local $c0$ pseudofermion creation and annihilation operators, respectively [8, 9]. The expectation value $\left\langle \frac{3}{2} - 2\mathbf{S}_0\mathbf{S}_1 - 2\mathbf{S}_1\mathbf{S}_2 - 2\mathbf{S}_0\mathbf{S}_2 \right\rangle_{\text{spin}}$ refers to the spin degrees of freedom. The partial sum rule (9) corresponds to the second UHB. This band is generated by transitions to the above three-holon states of type (iii) and involves annihilation of three $c0$ pseudofermions. The expectation value to find three neighboring local $c0$ pseudofermions is,

$$\begin{aligned} \langle \hat{n}_{x_0, c0} \hat{n}_{x_1, c0} \hat{n}_{x_2, c0} \rangle &= \left| \begin{array}{ccc} \langle f_{x_0, c0}^\dagger f_{x_0, c0} \rangle & \langle f_{x_0, c0}^\dagger f_{x_1, c0} \rangle & \langle f_{x_0, c0}^\dagger f_{x_2, c0} \rangle \\ \langle f_{x_1, c0}^\dagger f_{x_0, c0} \rangle & \langle f_{x_1, c0}^\dagger f_{x_1, c0} \rangle & \langle f_{x_1, c0}^\dagger f_{x_2, c0} \rangle \\ \langle f_{x_2, c0}^\dagger f_{x_0, c0} \rangle & \langle f_{x_2, c0}^\dagger f_{x_1, c0} \rangle & \langle f_{x_2, c0}^\dagger f_{x_2, c0} \rangle \end{array} \right| \\ &= n^3 - \frac{2n \sin^2(\pi n)}{\pi^2} + \frac{\sin^2(\pi n) \sin(2\pi n)}{\pi^3} \\ &\quad - \frac{n \sin^2(2\pi n)}{4\pi^2}. \end{aligned} \quad (10)$$

This expectation value has the following limiting behavior,

$$\langle \hat{n}_{x_0, c0} \hat{n}_{x_1, c0} \hat{n}_{x_2, c0} \rangle = \begin{cases} \frac{4\pi^6}{135} n^9 & \text{if } n \ll 1; \\ 1 - 3(1 - n) & \text{if } 1 - n \ll 1. \end{cases} \quad (11)$$

Note that the spectral weight generated by the transitions to the three-holon states of type (iii) decreases rapidly away from half filling. At quarter filling ($n = 1/2$) it is about 2% of that at half filling.

The expectation values associated with the spin degrees of freedom can in the thermodynamic limit and for large values of U/t be evaluated by use of the relation between the 1D Hubbard model and the spin 1/2 isotropic Heisenberg chain [11, 12]. This leads to the following values [20],

$$\langle \mathbf{S}_0\mathbf{S}_1 \rangle_{\text{spin}} = \frac{1}{4} - \ln 2 \approx -0.443147 \quad (12)$$

$$\langle \mathbf{S}_0\mathbf{S}_2 \rangle_{\text{spin}} = \frac{1}{4} - 4 \ln 2 + \frac{9}{4} \zeta(3) \approx 0.182039, \quad (13)$$

N	rotated-electron removal	rotated-electron addition
6	0.998792	0.977515
8	0.997486	0.972141
10	0.996277	0.968088
12	0.995178	0.964847
14	0.994176	0.962156
16	0.993258	0.959862
18	0.992409	0.957867
20	0.991622	0.956105
22	0.990886	0.954531
24	0.990196	0.953109

Table 1. The relative weight of the one $s1$ pseudfermion hole contributions that obey relation (8) in the $U/t \rightarrow \infty$ limit for finite-size systems of N electrons. This also gives the same relative weight in the case of excitations generated by one-rotated-electron operators for any value of U/t .

so that,

$$\left\langle \frac{3}{2} - 2\mathbf{S}_0\mathbf{S}_1 - 2\mathbf{S}_1\mathbf{S}_2 - 2\mathbf{S}_0\mathbf{S}_2 \right\rangle_{\text{spin}} = 12 \ln 2 - \frac{9}{2} \zeta(3) \approx 2.91. \quad (14)$$

The sum rule of the second UHB is then given by,

$$\int \mathcal{A}^{2\text{UHB}}(\omega) d\omega \approx 2.91 \frac{t^4}{U^4} \langle \hat{n}_{x_0, c0} \hat{n}_{x_1, c0} \hat{n}_{x_2, c0} \rangle. \quad (15)$$

For the six-site finite-size cluster, the expectation value (14) is,

$$\left\langle \frac{3}{2} - 2\mathbf{S}_0\mathbf{S}_1 - 2\mathbf{S}_1\mathbf{S}_2 - 2\mathbf{S}_0\mathbf{S}_2 \right\rangle_{\text{spin}} = (169 + 17\sqrt{13})/78 \approx 2.95, \quad (16)$$

which is about 1% off from the thermodynamic-limit value given in Eq. (14). The asymptotic behavior $2.95 t^4/U^4$ is shown in Fig. 2 as a dashed line.

Our above numerical results for all values of U/t and a small system lead to the same general picture as the results for $L \rightarrow \infty$. For electron addition the relative spectral weight of the excited energy eigenstates of types (i) and (ii) generated by dominant processes is minimum for $U \approx 4t$. This *minimum* value decreases with decreasing density. For half filling it is given by $\approx 99.75\%$, whereas for quarter filling it reads $\approx 99.99\%$ and in the limit of vanishing density it becomes $\approx 100.00\%$. The extremely small amount of missing one-electron addition spectral weight corresponds mainly to the three-holon states of type (iii) generated by the $i > 0$ operators $\tilde{\theta}_{i,j,\sigma}^\dagger$. Transitions from the ground state to higher-order five-holon/five- c -hole states generated by the latter operators lead to nearly vanishing spectral weight.

The three- $s1$ -hole states of types (i') and (ii') are generated by a sub-class of dominant processes. The relative weight of these excited energy eigenstates increases for increasing values of the system length L . For one-electron addition its maximum

value occurs at half filling and is of about 6% as $L \rightarrow \infty$. (At $n = 1$ there are no LHB one-electron addition excited energy eigenstates of types (i) and (i').) For half filling, all values of U/t , and $L \rightarrow \infty$ over 99% of the one-electron addition spectral weight corresponds to generation of one holon and one and three $s1$ pseudofermion holes. This is similar to the relative weights of Fig. 2 for L finite. For $L \rightarrow \infty$ the amount of one-electron spectral weight generated by dominant processes increases for decreasing values of the electronic density for all values of U/t .

While for electron addition the states of types (i) and (ii) such that $\Delta N_{s1}^h = 1$ correspond to at least 94% of the spectral weight, for electron removal the $\Delta N_{s1}^h = 1$ excited energy eigenstates amount for at least 98% of the total weight. However, we note that concerning the small amount of spectral weight generated by the excited energy eigenstates such that $\Delta N_{s1}^h > 1$, the case of the 1D Hubbard model is different from that of the $t - J_{XY}$ model considered in Ref. [21]. The significant difference is the $SU(2)$ symmetry in the spin sector of the 1D Hubbard model, which by standard selection rules prohibits matrix elements that are present in the $t - J_{XY}$ case. For instance, for low energy such a symmetry *protects* the lower-edge of the removal and addition spectrum of being dressed by $s1$ pseudofermion-like particle-hole excited energy eigenstates in the main conformal tower (see Fig. 5 in Ref. [12]). The low-energy $\Delta N_{s1}^h = 3$ excited energy eigenstates are for the 1D Hubbard model mostly associated with the next conformal tower centered at $3k_F$ for the spin excitations. In the $t - J_{XY}$ case, there is no similar protecting mechanism and, therefore, the weight is more easily redistributed among the $s1$ pseudofermion-like particle-hole excited energy eigenstates. Let us also note that the studies of Ref. [22] have addressed the same question for other models. The authors of that reference found that the weight coming from $\Delta N_{s1}^h = 1$ excited states is also dominant.

3. ROTATED-ELECTRON GENERATORS OF THE ONE-ELECTRON SPECTRAL WEIGHT

In this section we study the pseudofermion microscopic processes associated with the excitations $c_{j,\sigma}^\dagger |GS\rangle$ and $c_{j,\sigma} |GS\rangle$ which can be written as $\sum_{i=0}^{\infty} \sqrt{c_i^{+1}} \tilde{\theta}_{i,j,\sigma}^\dagger |GS\rangle$ and $\sum_{i=0}^{\infty} \sqrt{c_i^{-1}} \tilde{\theta}_{i,j,\sigma} |GS\rangle$, respectively, where the constants c_i^l are the same as in Eq. (5). Fortunately, for all values of U/t more than 99% of the spectral weight associated with such excitations corresponds to the $i = 0$ contributions $\tilde{\theta}_{0,j,\sigma}^\dagger |GS\rangle$ and $\tilde{\theta}_{0,j,\sigma} |GS\rangle$, respectively. Therefore, here we limit our study to the $i = 0$ operators $\tilde{\theta}_{0,j,\sigma}^\dagger$ and $\tilde{\theta}_{0,j,\sigma}$, which generate nearly the whole one-electron spectral weight measured in photoemission experiments [1, 2]. Indeed, for the electronic densities of the TTF and TCNQ stacks of molecules considered in the photoemission experiments of Ref. [2] the studies of the previous section reveal that the dominant processes generated by the operators $\tilde{\theta}_{0,j,\sigma}^\dagger$ and $\tilde{\theta}_{0,j,\sigma}$ account for more than $\approx 99.9\%$ of the total one-electron spectral weight. We note that in spite of the recent improvements in the resolution of photoemission experiments [2], it is difficult to measure the finest details of the electronic structure experimentally,

in part due to the extrinsic losses that occur on very anisotropic conducting solids such as the organic compound TTF-TCNQ [23]. Therefore, the less of 0.01% of missed *theoretical spectral weight* is irrelevant for the description of the spectral features measured by photoemission experiments.

Following the exact selection rules of Eq. (7), the CPHS ensemble subspaces of the excited energy eigenstates generated by application of the operators $c_{j,\sigma}^\dagger$ and $c_{j,\sigma}$ onto the ground state can have the following values for the numbers $\{L_{c,-1/2}, L_{s,-1/2}\}$,

$$\begin{aligned} c_{j,\downarrow}^\dagger \text{ operator: } & \{0, 0\}, \{1, 0\}, \{0, 1\}, \{1, 1\}; & c_{j,\downarrow} \text{ operator: } & \{0, 0\}; \\ c_{j,\uparrow}^\dagger \text{ operator: } & \{0, 0\}, \{1, 0\}; & c_{j,\uparrow} \text{ operator: } & \{0, 0\}, \{0, 1\}. \end{aligned}$$

We emphasize that the expressions of the operators $\tilde{\theta}_{0,j,\sigma}$ and $\tilde{\theta}_{0,j,\sigma}^\dagger$ only depend on the values $\{L_{c,-1/2}, L_{s,-1/2}\}$ of the CPHS ensemble subspace they refer to. Different CPHS ensemble subspaces with the same values for $\{L_{c,-1/2}, L_{s,-1/2}\}$ have the same expressions for the operators $\tilde{\theta}_{0,j,\sigma}$ and $\tilde{\theta}_{0,j,\sigma}^\dagger$. Evaluation of the commutators given in Eq. (27) of Ref. [9] for the one-electron case considered here together with the property that the $i = 0$ operators $\tilde{\theta}_{0,j,\sigma}$ and $\tilde{\theta}_{0,j,\sigma}^\dagger$ have the same expressions in terms of rotated-electron creation and annihilation operators as the corresponding operators on the left-hand side of the first equality of that equation in terms of creation and annihilation electronic operators, leads to the following expressions,

$$\begin{aligned} \tilde{\theta}_{0,j,\downarrow}^\dagger &= \tilde{c}_{j,\downarrow}^\dagger, & \{0, 0\}, \\ \tilde{\theta}_{0,j,\downarrow}^\dagger &= \frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} \tilde{c}_{j,\uparrow}, & \{1, 0\}, \\ \tilde{\theta}_{0,j,\downarrow}^\dagger &= \frac{1}{\sqrt{N_\uparrow^0 - N_\downarrow^0 + 1}} \tilde{c}_{j,\uparrow}^\dagger, & \{0, 1\}, \\ \tilde{\theta}_{0,j,\downarrow}^\dagger &= -\frac{(-1)^j}{\sqrt{(N_a - N^0 + 1)(N_\uparrow^0 - N_\downarrow^0 + 1)}} \tilde{c}_{j,\downarrow}, & \{1, 1\}, \end{aligned} \quad (17)$$

and

$$\begin{aligned} \tilde{\theta}_{0,j,\uparrow}^\dagger &= \tilde{c}_{j,\uparrow}^\dagger, & \{0, 0\}, \\ \tilde{\theta}_{0,j,\uparrow}^\dagger &= -\frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} \tilde{c}_{j,\downarrow}, & \{1, 0\}, \\ \tilde{\theta}_{0,j,\uparrow} &= \tilde{c}_{j,\uparrow}, & \{0, 0\}, \\ \tilde{\theta}_{0,j,\uparrow} &= -\frac{1}{\sqrt{N_\uparrow^0 - N_\downarrow^0 + 1}} \tilde{c}_{j,\downarrow}, & \{0, 1\}, \\ \tilde{\theta}_{0,j,\downarrow} &= \tilde{c}_{j,\downarrow}, & \{0, 0\}. \end{aligned} \quad (18)$$

Here the values of the numbers $\{L_{c,-1/2}, L_{s,-1/2}\}$ are those provided above and N^0 , N_\uparrow^0 , and N_\downarrow^0 are the electron numbers of the initial ground state. The expressions given in

Eqs. (17) and (18) are valid for ground-state electron numbers such that $0 \leq N^0 \leq N_a$ and $0 \leq N_{\downarrow}^0 \leq N_{\uparrow}^0$.

The CPHS ensemble subspaces associated with η -spin value deviations ΔS_c and spin value deviations ΔS_s such that $\Delta S_c = -1/2$ and $\Delta S_s = -1/2$ are not permitted for initial ground states such that $N^0 = N_a$ and $N_{\downarrow}^0 = N_{\uparrow}^0$, respectively. The reason is that for such ground states the η -spin and spin values are $S_c^0 = 0$ and $S_s^0 = 0$, respectively, and thus negative deviations ΔS_c and ΔS_s are not allowed. It follows that for $N_{\downarrow}^0 = N_{\uparrow}^0$ ground states, the transitions to CPHS ensemble subspaces such that $L_{s,-1/2} = 0$ cannot be generated by application of the operators $c_{j,\downarrow}^{\dagger}$ and $c_{j,\uparrow}$ onto such initial ground states. The case of the $N_{\downarrow}^0 = N_{\uparrow}^0$ ground states plays an important role in the applications of our results and those of Refs. [8, 9]. In the following, we provide the expression of the operators $\tilde{\theta}_{0,j,\sigma}$ and $\tilde{\theta}_{0,j,\sigma}^{\dagger}$ of Eqs. (17) and (18) in terms of pseudofermion creation and annihilation operators for the set of CPHS ensemble subspaces associated with the $N_{\downarrow}^0 = N_{\uparrow}^0$ ground states.

The one-electron processes generated by the rotated-electron operator $\tilde{c}_{j,\sigma}^{\dagger}$ give rise to the LHB when $M_{c,-1/2} = 0$ and UHB when $M_{c,-1/2} = 1$. According to the exact selection rules (6), those are the only permitted values for the excited-energy-eigenstate $-1/2$ holon numbers. Since the numbers of the initial ground-state CPHS subspace are known and well defined, here we characterize the CPHS ensemble subspaces of the excited states by the deviation numbers ΔN_{c0} and ΔN_{s1} and by the set $\{N_{\alpha\nu}\}$ such that $\alpha\nu \neq c0, s1$ and $\{L_{\alpha,-1/2}\}$ where $\alpha = c, s$ whose values are finite for the latter states. Thus, it is considered that the numbers of the set $\{N_{\alpha\nu}\}$ such that $\alpha\nu \neq c0, s1$ and $\{L_{\alpha,-1/2}\}$ where $\alpha = c, s$ which are not provided, vanish. As discussed above, our aim is the study of the dominant microscopic processes that generate the excitations $c_{j,\uparrow}^{\dagger}|GS\rangle$ and $c_{j,\downarrow}|GS\rangle$ where $|GS\rangle$ is a $N_{\downarrow}^0 = N_{\uparrow}^0$ ground state.

The whole $M_{c,-1/2} = 0$ LHB spectral weight of the excitation $\tilde{\theta}_{0,j,\uparrow}^{\dagger}|GS\rangle$ corresponds to $\{L_{c,-1/2}, L_{s,-1/2}\} = \{0, 0\}$ CPHS ensemble subspaces. According to Eq. (18), in such subspaces the above excitation reads $\tilde{\theta}_{0,j,\uparrow}^{\dagger}|GS\rangle = \tilde{c}_{j,\uparrow}^{\dagger}(1 - \tilde{n}_{j,\downarrow})|GS\rangle$ where $(1 - \tilde{n}_{j,\downarrow})$ is the LHB projector. We recall that the LHB processes do not exist for the $N^0 = N_a$ half-filling ground state. Otherwise, the spectral weight of this excitation is generated by transitions to the excited energy eigenstates which span the set of CPHS ensemble subspaces such that,

$$\Delta N_{c0} = 1, \quad \Delta N_{s1} = - \sum_{\nu'=2}^{\infty} \nu' N_{s\nu'}, \quad \{N_{s\nu}\}; \quad \nu > 1. \quad (19)$$

Thus, the deviation $\Delta N_{s1}^h = \Delta N_{c0} - 2\Delta N_{s1} - 2\sum_{\nu'=2}^{\infty} N_{s\nu'}$ in the number of $s1$ pseudofermion holes is $\Delta N_{s1}^h = -1 + 2\sum_{\nu'=2}^{\infty} (\nu' - 1)N_{s\nu'}$ for these subspaces. We note that for the present case of zero-magnetization the $s1$ pseudofermion band is full for the ground state. Therefore, the deviation ΔN_{s1}^h equals the number of $s1$ pseudofermion holes of the CPHS ensemble subspace excited energy eigenstates.

At least 94% of the spectral weight of $\tilde{\theta}_{0,j,\uparrow}^{\dagger}|GS\rangle = \tilde{c}_{j,\uparrow}^{\dagger}(1 - \tilde{n}_{j,\downarrow})|GS\rangle$ corresponds

to the CPHS ensemble subspace such that,

$$\Delta N_{c0} = 1, \quad \Delta N_{s1} = 0, \quad (20)$$

and $\Delta N_{s1}^h = -1$. These numbers obey the relation (8). In this subspace the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger (1 - \tilde{n}_{j,\downarrow})$ is given in Eq. (47) of Ref. [9]. Nearly the whole of the remaining spectral weight of the above LHB excitation is associated with the CPHS ensemble subspace whose numbers are,

$$\Delta N_{c0} = 1, \quad \Delta N_{s1} = -2, \quad N_{s2} = 1, \quad (21)$$

and $\Delta N_{s1}^h = 3$. Such values do not obey the relation (8). In this subspace, the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger (1 - \tilde{n}_{j,\downarrow})$ is given by,

$$\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger (1 - \tilde{n}_{j,\downarrow}) = f_{x_{j''',s2}}^\dagger f_{x_{j'',s1}} f_{x_{j',s1}} f_{x_j,c0}^\dagger. \quad (22)$$

Following the description of the energy eigenstates introduced in Ref. [17], the effective $s1$ lattice indices j' and j'' and the effective $s2$ lattice index j''' have a well-defined relation to the lattice index j , as for the pseudofermion operators appearing in Eqs. (46)-(53) of Ref. [9]. For simplicity, here we do not provide such index relations. We recall that the effective $c0$ lattice index j equals that of the rotated-electron lattice [17].

Next, we consider the excitation $c_{j,\downarrow}|GS\rangle$. Again, more than 99% of the spectral weight corresponds to the excitation $\tilde{\theta}_{0,j,\downarrow}|GS\rangle$. Such an excitation is associated with $\{L_{c,-1/2}, L_{s,-1/2}\} = \{0, 0\}$ CPHS ensemble subspaces where according to Eq. (18), $\tilde{\theta}_{0,j,\downarrow}|GS\rangle = \tilde{c}_{j,\downarrow} (1 - \tilde{n}_{j,\uparrow})|GS\rangle$. Here the projector $(1 - \tilde{n}_{j,\uparrow})$ is associated with the ground-state lack of rotated-electron double occupation [13]. The spectral weight of this excitation is generated by transitions to the states which span the set of CPHS ensemble subspaces such that,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1 - \sum_{\nu=2}^{\infty} \nu' N_{s\nu'}, \quad \{N_{s\nu}\}; \quad \nu > 1, \quad (23)$$

and $\Delta N_{s1}^h = 1 + 2 \sum_{\nu'=2}^{\infty} (\nu' - 1) N_{s\nu'}$. However, up to 98% of the spectral weight of $\tilde{\theta}_{0,j,\downarrow}|GS\rangle = \tilde{c}_{j,\downarrow} (1 - \tilde{n}_{j,\uparrow})|GS\rangle$ corresponds to the CPHS ensemble subspace whose deviations read,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1, \quad (24)$$

and $\Delta N_{s1}^h = 1$. The values of the above deviations obey the relation (8). The pseudofermion expression of the operator $\tilde{\theta}_{0,j,\downarrow} = \tilde{c}_{j,\downarrow} (1 - \tilde{n}_{j,\uparrow})$ in this subspace, is given in Eq. (46) of Ref. [9]. Nearly the whole of the remaining spectral weight of the above electron-removal excitation is associated with the CPHS ensemble subspace whose numbers are given by,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -3, \quad N_{s2} = 1, \quad (25)$$

and $\Delta N_{s1}^h = 3$. These do not obey the relation (8). In this subspace the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\downarrow} = \tilde{c}_{j,\downarrow} (1 - \tilde{n}_{j,\uparrow})$ is given by,

$$\tilde{\theta}_{0,j,\downarrow} = \tilde{c}_{j,\downarrow} (1 - \tilde{n}_{j,\uparrow}) = f_{x_{j''',s2}}^\dagger f_{x_{j'',s1}} f_{x_{j',s1}} f_{x_j,c0}. \quad (26)$$

For the $M_{c,-1/2} = 1$ UHB excitations, let us start by considering electronic densities such that $0 < \delta \leq 1$, where $\delta = [1 - n]$ is the doping concentration. We are assuming that δ can be small, but not vanishing. The case of vanishing doping concentrations corresponds to the Mott-Hubbard insulator phase where $(N_a - N^0) = 0$ and to the *doped Mott-Hubbard insulator* such that $(N_a - N^0)$ is finite and will be addressed in the ensuing section. For finite doping concentrations the whole UHB spectral weight of the excitation $\tilde{\theta}_{0,j,\uparrow}^\dagger |GS\rangle$ corresponds to $\{L_{c,-1/2}, L_{s,-1/2}\} = \{0, 0\}$ CPHS ensemble subspaces. According to Eq. (18), such an excitation can be written as $\tilde{\theta}_{0,j,\uparrow}^\dagger |GS\rangle = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow} |GS\rangle$ where $\tilde{n}_{j,\downarrow}$ is the UHB projector. For finite values of δ , the spectral weight of this excitation is generated by transitions to the excited energy eigenstates which span the set of CPHS ensemble subspaces such that,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1 - \sum_{\nu'=2}^{\infty} \nu' N_{s\nu'}, \quad N_{c1} = 1, \quad \{N_{s\nu'}\}; \quad \nu > 1. \quad (27)$$

The deviation in the number of $s1$ pseudofermion holes of these subspaces is $\Delta N_{s1}^h = 1 + 2 \sum_{\nu'=2}^{\infty} (\nu' - 1) N_{s\nu'}$. At least 94% of the spectral weight of the excitation $\tilde{\theta}_{0,j,\uparrow}^\dagger |GS\rangle = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow} |GS\rangle$ corresponds to the CPHS ensemble subspace such that,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1, \quad N_{c1} = 1, \quad (28)$$

and $\Delta N_{s1}^h = 1$. These obey the relation (8). In this subspace, the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow}$ is given by,

$$\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow} = f_{x_{j\mu},c1}^\dagger f_{x_j,c0} f_{x_{j'},s1}. \quad (29)$$

Nearly the whole of the remaining spectral weight of the above UHB excitation is associated with the CPHS ensemble subspace whose numbers are,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -3, \quad N_{c1} = 1, \quad N_{s2} = 1, \quad (30)$$

and $\Delta N_{s1}^h = 3$. These however do not obey the relation (8). In this subspace the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow}$ is given by,

$$\tilde{\theta}_{0,j,\uparrow}^\dagger = \tilde{c}_{j,\uparrow}^\dagger \tilde{n}_{j,\downarrow} = f_{x_{j\mu\mu},s2}^\dagger f_{x_{j\mu\mu},c1}^\dagger f_{x_{j\mu},s1} f_{x_{j'},s1} f_{x_{j'},s1} f_{x_j,c0}. \quad (31)$$

4. THE METAL - MOTT-HUBBARD INSULATOR QUANTUM PHASE TRANSITION

The further understanding of the microscopic processes behind the quantum phase transitions is a problem of great physical interest, as mentioned in Sec. I. For the 1D Hubbard model there is only one quantum phase transition as a function of the on-site repulsion U , which occurs at $U = 0$ for half filling. As a function of the interaction, such transitions are controlled by the dependence on that interaction of the local quantum entanglement [14]. On the other hand, for all finite values of U there is a metal - insulator quantum phase transition which occurs as a function of the electronic density at $n = 1$.

As a function of the doping that quantum phase transition occurs at $\delta = 0$. Here we study the microscopic effects of that quantum phase transition on the one-electron spectral properties.

Let us consider the case when δ is vanishing, *i.e.* there are no $c0$ pseudofermion holes (half-filling) or there is a finite number of such holes in the initial ground state (doped Mott-Hubbard insulator). As the number of holes decreases and one reaches the doped Mot-Hubbard insulator regimen, there arises a competition of the UHB processes considered in the previous section with other UHB processes. The latter processes correspond to the excitation $\tilde{\theta}_{0,j,\uparrow}^\dagger|GS\rangle$ for CPHS ensemble subspaces such that $\{L_{c,-1/2}, L_{s,-1/2}\} = \{1, 0\}$. Thus, according to Eq. (18), one has that $\tilde{\theta}_{0,j,\uparrow}^\dagger|GS\rangle = -[(-1)^j/\sqrt{N_a - N^0 + 1}] \tilde{c}_{j,\downarrow}^\dagger \tilde{n}_{j,\uparrow}|GS\rangle$, where $\tilde{n}_{j,\uparrow}$ is the UHB projector. Note that for finite values of the doping concentration δ the one-electron spectral weight associated with such an excitation vanishes in the themodynamic limit $L \rightarrow \infty$. It is generated by transitions to the excited energy eigenstates which span the set of CPHS ensemble subspaces such that,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1 - \sum_{\nu'=2}^{\infty} \nu' N_{s\nu'}, \quad L_{c,-1/2} = 1, \quad \{N_{s\nu'}\}; \quad \nu' > 1, \quad (32)$$

and $\Delta N_{s1}^h = 1 + 2 \sum_{\nu'=2}^{\infty} (\nu' - 1) N_{s\nu'}$. For vanishing values of the doping concentration at least 94% of the spectral weight of the excitation $\tilde{\theta}_{0,j,\uparrow}^\dagger|GS\rangle$ corresponds to the CPHS ensemble subspace with numbers,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -1, \quad L_{c,-1/2} = 1, \quad (33)$$

and $\Delta N_{s1}^h = 1$. Such numbers obey the relation (8). In this subspace the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger$ reads,

$$\tilde{\theta}_{0,j,\uparrow}^\dagger = -\frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} \tilde{c}_{j,\downarrow}^\dagger \tilde{n}_{j,\uparrow} = -\frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} f_{x_{j',s1}} f_{x_j,c0}. \quad (34)$$

Nearly the whole of the remaining spectral weight of such a UHB excitation is associated with the CPHS ensemble subspace whose numbers are,

$$\Delta N_{c0} = -1, \quad \Delta N_{s1} = -3, \quad L_{c,-1/2} = 1, \quad N_{s2} = 1, \quad (35)$$

and $\Delta N_{s1}^h = 3$. These do not obey the relation (8). In this subspace the pseudofermion expression of the operator $\tilde{\theta}_{0,j,\uparrow}^\dagger$ is given by,

$$\begin{aligned} \tilde{\theta}_{0,j,\uparrow}^\dagger &= -\frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} \tilde{c}_{j,\downarrow}^\dagger \tilde{n}_{j,\uparrow} \\ &= -\frac{(-1)^j}{\sqrt{N_a - N^0 + 1}} f_{x_{j'''',s2}}^\dagger f_{x_{j''',s1}} f_{x_{j'',s1}} f_{x_{j',s1}} f_{x_j,c0}. \end{aligned} \quad (36)$$

For finite values of δ , the UHB processes associated with CPHS ensemble subspaces such that $\{L_{c,-1/2}, L_{s,-1/2}\} = \{1, 0\}$ are allowed. However, for finite doping concentrations the relative UHB weight of the $\{L_{c,-1/2}, L_{s,-1/2}\} = \{0, 0\}$ and

$\{L_{c,-1/2}, L_{s,-1/2}\} = \{1, 0\}$ processes is approximately given by $1/\delta L$ and thus vanishes in the thermodynamic limit, as mentioned above. Therefore, the weight of the UHB $\{L_{c,-1/2}, L_{s,-1/2}\} = \{1, 0\}$ processes associated with the CPHS ensemble subspaces of numbers (32) vanishes for finite values of δ .

In contrast, for half filling only the UHB $\{1, 0\}$ processes contribute, whereas for finite yet small values of $(N_a - N^0)$ both the $\{0, 0\}$ and $\{1, 0\}$ processes contribute to the UHB spectral weight. If the value of $(N_a - N^0)$ further increases so that δ becomes finite, the weight of the UHB $\{1, 0\}$ processes vanishes. For $N^0 \neq N_a$ and $(N_a - N^0)$ finite but small, the relative weight of the $\{0, 0\}$ and $\{1, 0\}$ processes is approximately given by $1/[N_a - N^0 + 1]$.

We thus conclude that the collapse of the LHB processes and the interplay between the $\{0, 0\}$ and $\{1, 0\}$ UHB processes for decreasing values of $(N_a - N^0)$ are the main effects of the $(N_a - N^0) \rightarrow 0$ metal - Mott-Hubbard insulator quantum phase transition onto the one-electron spectral properties. The ground-state - excited-energy-eigenstate transitions associated with the $\{0, 0\}$ and $\{1, 0\}$ UHB processes change the value of η spin by $\Delta\eta = -1/2$ and $\Delta\eta = +1/2$, respectively. For the metallic phase at finite values of the doping concentration the whole UHB weight corresponds to excited states with deviation $\Delta\eta = -1/2$, whereas for the Mott-Hubbard insulator phase only the excited states with deviation $\Delta\eta = +1/2$ are allowed. However, the physics of the doped Mott-Hubbard insulator, which corresponds to a finite number of holes $(N_a - N^0)$, is different: It involves a competition between the two above classes of states. Such a competition is mainly controlled by the form of the operators given in Eqs. (29) and (34), which generate excited energy eigenstates with η -spin deviations $\Delta\eta = -1/2$ and $\Delta\eta = +1/2$, respectively.

5. CONCLUDING REMARKS

In this paper we used the holon and spinon description introduced in Ref. [13] and expressed in terms of composite pseudofermions in Ref. [17] in the study of the non-perturbative microscopic processes that control the unusual one-electron spectral-weight distributions of the 1D Hubbard model.

The results of this paper are complementary to those of Refs. [8, 9] and are useful for the further understanding and description of the microscopic mechanisms behind the unusual finite-energy spectral properties observed in quasi-1D compounds [1, 2]. Our results also provide new insights about the microscopic mechanisms of quantum phase transitions in electronic correlated problems. We found that for the doped Mott-Hubbard insulator there is a competition between the microscopic processes which determine the UHB one-electron spectral distributions of the Mott-Hubbard insulator phase and metallic phase for finite values of the doping concentration.

Acknowledgments

We thank Daniel Bozi, Ralph Claessen, Patrick A. Lee, Tiago C. Ribeiro, and Pedro D. Sacramento for stimulating discussions. J.M.P.C. thanks the hospitality and support of MIT and the financial support of the Gulbenkian Foundation, Fulbright Commission, and FCT under the grant POCTI/FIS/58133/2004 and K.P. thanks the support of the OTKA grants T038162 and T049607.

References

- [1] Carmelo JMPC, Penc K, Martelo LM, Sacramento PD, Lopes dos Santos JMB, Claessen R, Sing M and Schwingenschlögl U 2004 *Europhys. Lett.* **67**, 233; Carmelo JMP, Bozi D, Sacramento PD and Penc K 2005 submitted for publication.
- [2] Sing M, Schwingenschlögl U, Claessen R, Blaha P, Carmelo JMP, Martelo LM, Sacramento PD, Dressel M and Jacobsen CS 2003 *Phys. Rev. B* **68**, 125111.
- [3] Benthien H, Gebhard F and Jeckelmann E 2004 *Phys. Rev. Lett.* **92**, 256401.
- [4] Carmelo JMP, Guinea F, Penc K and Sacramento PD 2004 *Europhys. Lett.* **68**, 839.
- [5] Simons BD, Lee PA and Altshuler BL 1993 *Phys. Rev. Lett.* **70**, 4122; Penc K and Shastry BS, 2002 *Phys. Rev. B* **65**, 155110.
- [6] Lieb Elliott H and Wu FY 1968 *Phys. Rev. Lett.* **20**, 1445; Ramos PB and Martins MJ 1997 *J. Phys. A* **30**, L195.
- [7] Takahashi M 1972 *Prog. Theor. Phys.* **47**, 69.
- [8] Carmelo JMP, Penc K and Bozi D, in press in *Nucl. Phys. B* [cond-mat/0505601] and references therein.
- [9] Carmelo JMP and Penc K, cond-mat/0311075.
- [10] Carmelo JMP, in print in *J. Phys.: Cond. Matt.* [cond-mat/0504271].
- [11] Penc K, Hallberg K, Mila F and Shiba H 1996 *Phys. Rev. Lett.* **77**, 1390.
- [12] Penc K, Hallberg K, Mila F and Shiba H 1997 *Phys. Rev. B* **55**, 15 475.
- [13] Carmelo JMP, Román JM, and Penc K 2004 *Nucl. Phys. B* **683**, 387.
- [14] Gu SJ, Deng SS, Li YQ and Lin HQ 2004 *Phys. Rev. Lett.* **93**, 086402.
- [15] Heilmann OJ and Lieb EH 1971 *Ann. N. Y. Acad. Sci.* **172**, 583.
- [16] Yang CN 1989 *Phys. Rev. Lett.* **63**, 2144.
- [17] Carmelo JMP, cond-mat/0508141.
- [18] Mizuno Y, Tsutsui K, Tohyama T and Maekawa S 2000 *Phys. Rev. B* **62**, R4769.
- [19] MacDonald AH, Girvin SM and Yoshioka D 1988 *Phys. Rev. B* **37**, 9753.
- [20] Takahashi M 1977 *J. Phys. C* **10**, 1289.
- [21] Sorella S and Parola A 1998 *Phys. Rev. B*, **57** 6444.
- [22] Talstra JC, Strong SP and Anderson PW 1995 *Phys. Rev. Lett.* **74**, 5256; Talstra JC and Strong SP 1997 *Phys. Rev. B* **56**, 6094.
- [23] Joynt R 2000 *Science* **284**, 777.