# Anyonic Excitations in Fast Rotating Bose Gases<sup>\*</sup>

J.C. Wallet

Groupe de Physique Théorique, Institut de Physique Nucléaire, F-91406 Orsay CEDEX, France

The role of anyonic excitations in fast rotating harmonically trapped Bose gases in a fractional Quantum Hall state is examined. Standard Chern-Simons anyons as well as "non standard" anyons obtained from a statistical interaction having Maxwell-Chern-Simons dynamics and suitable non minimal coupling to matter are considered. Their respective ability to stabilize attractive Bose gases under fast rotation in the thermodynamical limit is studied. Stability can be obtained for standard anyons while for non standard anyons, stability requires that the range of the corresponding statistical interaction does not exceed the typical wavelength of the atoms.

PACS numbers: 03.75.Lm, 73.43.-f

#### INTRODUCTION.

The experimental realization of Bose-Einstein Condensation (BEC) of atomic gases [1] has given rise to a rich variety of phenomena and motivated numerous investigations focussed on ultracold atomic Bose gases in rotating harmonic traps. BEC confined to two dimensions have been created [2] and their response under rotation has been studied. Basically, the rotation of a BEC produces vortices in the condensate [3],[4]. When the rotation frequency increases, the BEC state is destroyed and for sufficiently high frequency, a state corresponding to Fractional Quantum Hall Effect (FQHE) [5] is expected to possibly occur [6]. In particular, when the rotation frequency is tuned to the characteristic frequency of the harmonic confining potential in the radial plane, FQHE states have been predicted to become possible ground states for the system [7]. This observation has been followed by studies focused on the FQHE for bosons with short range interactions [8], [9]. It is known that FQHE states involve anyonic excitations [10]. Starting from the standard Chern-Simons (CS) realization for anyons [10], it has been shown in [11] that in the thermodynamical limit, a 2D harmonically trapped rotating Bose gas with *attractive* interactions can be stabilized (in a FQHE state) thanks to its anyonic (quasiparticle) excitations [12].

The above mentionned description of anyons is not unique. An alternative realization has been investigated in [13] where the minimal coupling of the statistical gauge potential to matter, which is now constrained to have Maxwell-Chern-Simons (MCS) dynamics, is supplemented by a Pauli-type couling, as recalled below. While the effective Landau-Ginzburg (LG) theories stemming from each anyon realization both reproduce the (so far observed) low energy features of the Quantum Hall Fluids (QHF), slight differences do exist between CS and MCS anyons. These latter stemm from a statistical interaction having finite range (whereas CS interaction has zero range) and have an additional attractive mutual interaction (not present for CS anyons). I point out that the existence of this attractive mutual interaction may have an impact on the ability of (MCS) anyonic excitations to stabilize an attractive Bose gas in a FQHE state. In particular, stability requires that the range of the MCS statistical interaction does not exceed the typical wavelength for the atoms. The description of QHF within the MCSLG theory stemming from the MCS description of anyons is also discussed.

### CHERN-SIMONS AND MAXWELL-CHERN-SIMONS ANYONS.

Consider an interacting Bose gas at zero temperature in a rotating harmonic trap with strong confinement in the direction of the rotation axis so that the system is actually two dimensional. The Hamiltonian in the rotating frame [14] and the corresponding action quoted here for further convenience are (for conventions, see [15])

$$H = \sum_{A=1}^{N} \frac{1}{2m} (\mathbf{p}_A - \mathbf{A}(\mathbf{x}_A))^2 + \sum_{A=1}^{N} \frac{m}{2} (\omega_T^2 - \omega^2) |\mathbf{x}_A|^2 + \sum_{A < B} V(\mathbf{x}_{AB}) + \dots$$
(1),

<sup>\*</sup> Presented at the School on Quantum Phase Transitions and Non-Equilibrium Phenomena in Cold Atomic Gases, ICTP Trieste, 11-22 july 2005 – e-mail: wallet@ipno.in2p3.fr

$$S_0 = \int_t \left(\sum_{A=1}^N \frac{m}{2} \dot{\mathbf{x}}_A^2 - \sum_{A=1}^N \frac{m}{2} (\omega_T^2 - \omega^2) |\mathbf{x}_A|^2 - \sum_{A < B} V(\mathbf{x}_{AB})\right) + \int_x \mathbf{A}(\mathbf{x}) \cdot \mathbf{J}(\mathbf{x}, t) + \dots$$
(2),

where  $\mathbf{J}(\mathbf{x},t) = \sum_A \dot{\mathbf{x}}_A(t) \delta(\mathbf{x} - \mathbf{x}_A(t))$ , the external gauge potential  $A_i(\mathbf{x}_A) = m\omega\epsilon_{ij}x_A^j$  yields the Coriolis force,  $\frac{\omega}{2\pi}$  (resp.  $\frac{\omega_T}{2\pi}$ ) is the rotation (resp. trapping) frequency),  $V(\mathbf{x}_{AB})$  is the two-body potential felt by the bosons in the plane. It can be well approximated by  $V(\mathbf{x}_{AB}) = g_2 \delta(\mathbf{x}_{AB})$  since the scattering between ultracold bosons is dominated by the s-wave [14]. In (1)-(2), the ellipses denote possible multibody interaction terms. We now consider the "limit"  $\omega \simeq \omega_T$  for which the trapping and centrifugal potential (nearly) balance each other so that W can be neglected.

The standard CS description of anyons [10] obtained from (1)-(2) is achieved by minimally coupling to  $S_0$  a statistical CS gauge potential  $a_{\mu}$ :  $S_0 \rightarrow S_1 = S_0 + \int_x \frac{\eta}{4} \epsilon_{\mu\nu\rho} a^{\mu} f^{\nu\rho} + a_{\mu} J^{\mu}$  where the CS parameter  $\eta$  is dimensionless,  $J_{\mu} = (\rho; \mathbf{J})$  with  $\rho = \rho(\mathbf{x}, t) = \sum_A \delta(\mathbf{x} - \mathbf{x}_A(t))$  and  $\mathbf{J}$  as given above  $(f_{\mu} \equiv \frac{1}{2} \epsilon_{\mu\nu\rho} f^{\nu\rho}, f_{\mu\nu} = \partial_{\mu} a_{\nu} - \partial_{\nu} a_{\mu})$ ,  $f_0$  (resp.  $f_i$ ) is the statistical magnetic (resp. electric) field). The equations of motion for  $a_{\mu}$  take the field-current identity form  $f_{\mu} = -\frac{1}{\eta} J_{\mu}$  (i), insuring that particle-statistical magnetic flux composite anyonic objects are formed and permit one to determine  $a_{\mu}$  in term of the particle current [10]. Particle-flux coupling responsible for the formation of the above composite quasiparticles leads to the occurence of Aharonov-Bohm type interactions. The anyonic character of the wave functions for the quasiparticles is determined by the Aharonov-Bohm phase [10]  $\exp(i \int_C \mathbf{a}.d\mathbf{x}) = \exp \frac{i}{\eta} (C$  is some closed curve) that is induced when one quasiparticle moves adiabatically around another one, equivalent to a *double* interchange of identical quasiparticles in the wave function. Then, the statistics of the quasiparticles is controlled by the value of  $\eta$ : they have Fermi(resp. Bose) statistics whenever  $\eta = \frac{1}{(2k+1)2\pi}$  (resp.  $\frac{1}{(2k)2\pi}$ ),  $k \in \mathbb{Z}$ . The statistics is anyonic otherwise. Notice that, when  $\eta = \frac{1}{(2k+1)2\pi}$ , the initial Bose statistical transmutation [10]. This, plugged into a second quantized formalism, leads to the usual CSLG action [16] underlying the main part of the analyzis presented in [11]. Since the statistical interaction is mediated by a CS gauge potential, its range is zero.

An alternative realization for anyons has been proposed and discussed in [13]. It is obtained by coupling a MCS statistical gauge potential  $a_{\mu}$  to  $S_0$  through minimal and non minimal (Pauli-type) coupling term, the strength of this latter being fixed to a unique specific value [13], namely:  $S_0 \rightarrow S_2 = S_0 + \int_x -\frac{1}{4e^2} f_{\mu\nu} f^{\mu\nu} + \frac{\eta}{4} \epsilon_{\mu\nu\rho} a^{\mu} f^{\nu\rho} + a_{\mu} J^{\mu} - \frac{1}{\eta e^2} f_{\mu\nu} J^{\mu}$  ( $e^2$  has mass dimension 1), where the coupling constant of the Pauli-term (last term in  $S_2$ ) has been already fixed to the above mentionned specific value. Then, the equations of motion for  $a_{\mu}$  can be written as  $-\frac{1}{e^2} \epsilon_{\alpha\mu\rho} \partial^{\mu} (f^{\rho} + \frac{1}{\eta} J^{\rho}) + \eta (f_{\alpha} + \frac{1}{\eta} J_{\alpha}) = 0$  which is solved by (i). This latter observation has been used as a starting point to construct an effective theory reproducing the usual anyonic behaviour [13]. The Aharonov-Bohm phase defining the statistics of the resulting quasiparticles still verifies  $\exp(i \int_C \mathbf{a}.d\mathbf{x}) = \exp \frac{i}{\eta}$  so that  $\eta$  again controls the statistics. However, anyons obtained through this construction have an additional attractive contact mutual interaction (not present in the CS case) [13]. Combining  $S_2$  with the second quantization machinery, one obtains the corresponding MCSLG action

$$S = \int_{x} i\phi^{\dagger} \mathcal{D}_{0}\phi - \frac{1}{2m} |\mathcal{D}_{i}\phi|^{2} - g_{2}(\phi^{\dagger}\phi)^{2} - U(\phi) - \frac{1}{4e^{2}} f_{\mu\nu}f^{\mu\nu} + \frac{\eta}{4}\epsilon_{\mu\nu\rho}a^{\mu}f^{\nu\rho}$$
(3),

$$\mathcal{D}_0 = \partial_0 - ia_0 + \frac{if_0}{\eta e^2} ; \ \mathcal{D}_i = \partial_i - i(a_i + A_i) + \frac{if_i}{\eta e^2}$$
(4*a*; *b*).

In (3),  $\phi(=\phi(\mathbf{x},t))$  is the order parameter,  $\phi^{\dagger}\phi=\rho$ ,  $U(\phi)$  denotes the LG potential for multibody interactions,  $U(\phi)=g_3|\phi|^6+\dots$  ( $g_3>0$  as in [11]). The non minimal coupling terms have been included in the extended covariant derivative (4). The statistical MCS gauge potential has a finite mass [17]  $M = |\eta|e^2$ . Correspondingly, the statistical interaction has a finite range  $\Lambda_{st}=\frac{1}{M}$ . The CSLG action [16] is obtained from (3)-(4) by taking the limit  $e^2 \rightarrow \infty$ . While CSLG action is believed to encode the (so far observed) low energy physics of QHF, it appears that MCSLG action also reaches this goal and could therefore be used as an alternative description of QHF. This is discussed more closely at the end of this letter. For the moment, we assume that the system is in a FQHE state ( $\eta\neq 0$ ) described the MCSLG action (3)-(4).

The equations of motion derived from (3) are

$$i\mathcal{D}_0\phi + \frac{1}{2m}\mathcal{D}_i\mathcal{D}_i\phi - 2g_2(\phi^{\dagger}\phi)\phi = \frac{\delta U(\phi)}{\delta\phi^{\dagger}}; -\frac{1}{e^2}\epsilon_{\alpha\mu\rho}\partial^{\mu}(f^{\rho} + \frac{1}{\eta}\mathcal{J}^{\rho}) + (f_{\alpha} + \frac{1}{\eta}\mathcal{J}_{\alpha}) = 0$$
(5*a*; *b*)

where  $\mathcal{J}_0 = \rho$ ,  $\mathcal{J}_i = \frac{i}{2m} (\phi^{\dagger} \mathcal{D}_i \phi - (\mathcal{D}_i \phi)^{\dagger} \phi)$  and  $\mathcal{D}_{\mu}$  is defined in (4) while anyonic configurations, on which we now focus, are obtained from (5b) when  $f_{\mu} = -\frac{1}{\eta} \mathcal{J}_{\mu}$  [18]. From the field's conjugate momenta  $\Pi_{\phi^{\dagger}} = \Pi_{a_0} = 0$ ,  $\Pi_{\phi} = i\phi^{\dagger}$ ,  $\Pi_{a_i} = \frac{1}{e^2} f_{0i} - \frac{\eta}{2} \epsilon_{ij} a^j + \frac{1}{\eta e^2} \epsilon_{ij} \mathcal{J}^j$ , one obtains the Hamiltonian

$$H = \int_{\mathbf{x}} \frac{1}{2e^2} (f_0 + \frac{1}{\eta}\rho)^2 + \frac{1}{2e^2} \Theta f_i^2 + \frac{1}{2m} |D_i\phi|^2 + (g_2 - \frac{1}{2\eta^2 e^2})\rho^2 + U(\phi)$$
(6)

where  $D_i = \partial_i - i(a_i + A_i)$  and we defined  $\Theta = 1 - \frac{\rho}{m\eta^2 e^2}$ . The positivity of the model requires  $\rho \leq m\eta^2 e^2$  [19]. Restoring  $\hbar$  and c, this translates into  $\rho \leq \rho_{lim}$ ,  $\rho_{lim} = |\eta| (\frac{\Lambda}{\Lambda_{st}}) \frac{1}{\Lambda^2}$  where  $\Lambda = \frac{\hbar}{mc}$  is a typical (de Broglie) wavelength for the atoms and  $\Lambda_{st} = \frac{\hbar}{|\eta|\tilde{\mu}c}$  ( $e^2 = \tilde{\mu}c^2$ ). This condition should be fulfilled by current experimental values, provided  $\Lambda_{st} \lesssim \mathcal{O}(\Lambda)$ , a condition that we now assume [20]. Namely, for <sup>7</sup>Li, <sup>23</sup>Na, <sup>87</sup>Rb, one obtains respectively  $\rho_{lim} \simeq |\eta| (\frac{\Lambda}{\Lambda_{st}}) 10^{29}$ ,  $10^{30}$ ,  $10^{31} \text{ cm}^{-2}$ , which, for possibly reachable current Quantum Hall states, should be larger by several orders of magnitude than the experimental values reached by the matter density so that  $\frac{\rho}{m\eta^2 e^2} \ll 1$ .

Compared to the CSLG Hamiltonian [11], H involves additional contributions from the Maxwell part of (3) and the non minimal coupling terms in (4). The interaction energy  $g_2\rho^2$  receives contributions from  $\frac{1}{2e^2}f_0^2$  and the Pauli-type coupling in (4a): these latter combine to yield the first term in (6) together with the additional attractive (magnetic) contribution  $-\frac{\rho^2}{2\eta^2 e^2}$  to  $g_2\rho^2$ . This, depending on the relative magnitude of  $\Lambda$  and  $\Lambda_{st}$ , may somehow alter the conclusion obtained in [11] from a CSLG description of anyons about the ability of statistical interaction to stabilize an attractive Bose gas, as we now show.

#### STABILIZING AN ATTRACTIVE 2D BOSE GAS.

We set  $\phi = \gamma e^{i\varphi} \equiv \sqrt{\rho} e^{i\varphi}$ ,  $\hat{a}_i = \partial_i \varphi - (a_i + A_i)$ ,  $\eta > 0$  (here, one has  $\nu = 2\pi\eta$  where  $\nu > 0$  is the filling factor; see below). For N atoms in the trap, one has  $\int_{\mathbf{x}} \phi^{\dagger} \phi = N$ . Using  $f_{\mu} = -\frac{1}{\eta} \mathcal{J}_{\mu}$  and assuming that fields vanish at infinity so that boundary terms disappear, the static energy stemming from (6) can be conveniently written as

$$H = \omega N + \int_{\mathbf{x}} \frac{1}{2m} (\partial_i \gamma - \epsilon_{ij} \frac{\hat{a}^j \gamma}{\sqrt{\Theta}})^2 + (g_2 - g_0)\rho^2 + \int_{\mathbf{x}} m\eta^2 e^2 \sum_{k=3}^{\infty} (\eta e^2 C_{k-1} + 2\omega C_k) (\frac{\rho}{m\eta^2 e^2})^k + U(\gamma)$$
(7).

In (7),  $\omega > 0$ , the positive constants  $C_k$ 's are given by  $(1 - \frac{\rho}{m\eta^2 e^2})^{\frac{1}{2}} = 1 - \sum_{k=1}^{\infty} C_k (\frac{\rho}{m\eta^2 e^2})^k (C_k = \frac{\Gamma(k - \frac{1}{2})}{2k!\sqrt{\pi}}$  where  $\Gamma$  is the Euler function) and we have defined

$$g_0 = -\frac{\hbar^2}{2m\eta} \left(1 - \frac{\Lambda_{st}}{\Lambda} \left(1 - \frac{\hbar\omega}{2mc^2}\right)\right) \simeq -\frac{\hbar^2}{2m\eta} \left(1 - \frac{\Lambda_{st}}{\Lambda}\right) \tag{8}$$

where  $\hbar$  and c have been reinstalled and the rightmost relation stemms from  $\hbar\omega \ll mc^2$  which holds for current experimental values for  $\omega$  (taking  $\frac{\omega}{2\pi} \sim \mathcal{O}(10) - \mathcal{O}(10^3)$  Hz as a benchmark). In the same way, the term  $\sim \omega C_k$  in (7) can also been neglected (since  $\hbar\omega \ll \eta \tilde{\mu} c^2$  in view of  $\Lambda_{st} \lesssim \mathcal{O}(\Lambda)$ ).

The quartic interaction terms in (7) can be eliminated provided  $g_2$  is chosen to be

$$g_2 = g_0 \tag{9}.$$

Then, neglecting in (7) the small terms of order  $\sim \mathcal{O}(\gamma^6)$  (and higher) as in [11], the ground state of (7) is obtained for those configurations satisfying  $(\partial_i \gamma - \epsilon_{ij} \frac{\hat{a}^j \gamma}{\Theta^{\frac{1}{2}}}) = 0$ , i=1, 2. This, combined with (5a), (8), (9), further expanding the various contributions depending on  $\Theta$  in powers of  $\frac{\rho}{m\eta^2 e^2}$  and using the fact that  $\frac{\rho}{m\eta^2 e^2} \ll 1$  yields

$$\partial_i^2 \gamma - \frac{(\partial_i \gamma)^2}{\gamma} = -\gamma \left(\frac{\gamma^2}{\eta} \left(1 + \frac{\omega}{\eta e^2}\right) + 2m\omega\right) \tag{10a},$$

where, in the RHS,  $\frac{\omega}{ne^2} \ll 1$  still holds so that (10a) can be accurately approximated by

$$\partial_i^2 \gamma - \frac{(\partial_i \gamma)^2}{\gamma} = -\gamma (\frac{\gamma^2}{\eta} + 2m\omega) \tag{10b}.$$

When  $\omega$  goes to zero, the solutions of (10b) reduces smoothly [21] to the solutions of the Liouville equation

$$\Delta \ln \rho = -\frac{2}{\eta}\rho \tag{11}$$

These latter can be parametrized as  $\rho(z)=4\eta |h'(z)|^2(1+|h(z)|^2)^{-2}$   $(z=x_1+ix_2)$  for any holomorphic function h(z). The particular choice  $h(z)=(z_0/z)^n$   $(z=re^{i\theta})$ ,  $n \in \mathbb{N}$ , gives rise to radially symmetric vortex-type solutions  $\rho(r)=\frac{4\eta n^2}{r^2}((\frac{r_0}{r})^n+(\frac{r}{r_0})^n)^{-2}$  where  $r_0$  is some arbitrary length scale and  $\phi=\sqrt{\rho}e^{in\theta}$ .

Let us discuss the above analyzis. From (7)-(9), one realizes that the initial two-body coupling term  $\sim g_2$  can be compensated by the statistical interaction. In view of (7), for fixed  $\eta$ , the particular value  $g_0$  (8) represents the limiting value for the two-body coupling constant below which the Bose gas cannot be stabilized by the statistical interaction. When this latter is ruled by a CS dynamics for which  $\Lambda_{st}=0$ ,  $g_0$  is negative while the ground state of the system is exactly described (neglecting the small  $|\phi|^6$  terms in  $U(\phi)$ ) by (10b) giving rise to non singular finite energy matter distribution [21]. In that situation, one would conclude as in [11] that an attractive Bose gas may be stabilized against collapse in the thermodynamical limit by anyonic excitations stemming from a CS (zero range) statistical interaction. Notice that [11] can be viewed as a nice reinterpretation in the context of the trapped atomic Bose gases of the initial work [22]. This conclusion is somehow altered when the statistical interaction has a MCS dynamics as described by (3) corresponding now to a (non zero) finite range. Indeed, in view of (8), (up to small terms  $\sim \hbar \omega$ ), the sign of  $g_0$  depends now on the relative magnitude of  $\Lambda$  and  $\Lambda_{st}$ . Compared to the CS case, the limiting value of the two-body coupling constant  $g_0$  receives a positive contribution stemming from the (statistical) magnetic energy and the magnetic coupling in (4a) in addition to the (negative) contribution of the kinetic energy that can be converted into interaction energy. As a consequence, negative values for  $g_0$  can only be obtained when  $\Lambda_{st} < \Lambda$  while a statistical interaction whose range  $\Lambda_{st}$  exceeds slightly  $\Lambda$  gives rise only to positive values for  $g_0$  (non singular finite energy matter distribution for the ground state is now well approximated by the solutions of (10b), again up to small  $\gamma^6$  terms). This can be rephrased by stating that anyonic excitations stemming from a MCS interaction with finite range  $\Lambda_{st}$  cannot protect an attractive Bose gas from collapse (in the thermodynamical limit) unless  $\Lambda_{st} \lesssim \Lambda$ .

## MCS LANDAU-GINZBURG DESCRIPTION OF QUANTUM HALL FLUIDS.

QHF are usually described in the low energy regime by a CSLG theory [16]. It turns out that QHF could alternatively be described by a MCSLG theory such as the one defined in (3)-(4). To make more explicit the comparition to the CS case [16], it is convenient to add a chemical potential term to (3), namely  $S'=S+\int_x \mu \phi^{\dagger} \phi$  ( $\mu>0$ ) and set  $U(\phi)=0$  in all. (5a) becomes  $i\mathcal{D}_0\phi+\frac{1}{2m}\mathcal{D}_i\phi-2g_2(\phi^{\dagger}\phi)\phi=\mu\phi$  while the relevant anyonic configurations are still obtained when  $f_{\mu}=-\frac{1}{\eta}\mathcal{J}_{\mu}$  solving (5b). Then, the equations of motion admit the uniform (constant) density solution minimizing the energy  $H'=H-\int_x \mu \rho$ , given by

$$\phi = \sqrt{n_0}; \ \mathbf{a} + \mathbf{A} = 0; \ a_0 = 0; \ n_0 = \frac{\mu}{2\hat{g}_2}$$
 (12*a*; *b*; *c*; *d*),

where  $\hat{g}_2 = g_2 - \frac{1}{2\eta^2 e^2}$  which is similar to the usual uniform ground state solution supported by the CSLG action for QHF[16]. Recall that (12b) implies that the external magnetic field  $B = \epsilon_{ij}\partial^i A^j$  is screened by the statistical magnetic field (which signals (the analogue of) a Meissner type effect). This, further combined with  $f_0 = -\frac{1}{\eta}\rho$ , yields  $\eta = \frac{n_0}{B}$  so that  $\eta$  is related to the filling factor  $\nu$  by  $2\pi\eta = \nu$ . In particular, when statistical transmutation occurs as expected in Hall systems,  $\nu$  satisfies  $\nu = \frac{1}{2k+1}$ ,  $k \in \mathbb{Z}$ . Note that (12c) implies that the (analog of) the electric field seen by the particles is zero. Furthermore, introducing polar coordinates  $(r, \theta)$  and setting  $\phi = \sqrt{\rho}e^{in\theta}$ ,  $\mathbf{a} = \mathbf{e}_{\theta}\frac{\alpha(r)}{r}$  $(\mathbf{e}_{\theta} = (\sin \theta, -\cos \theta)), a_0 = a_0(r)$ , where  $n \in \mathbb{Z}$  is the winding number, one finds after some algebra that H' admits static finite energy vortex solutions satisfying the boundary conditions

$$\lim_{r \to \infty} \rho = n_0; \quad \lim_{r \to \infty} \frac{\alpha(r)}{r^2} = m\omega; \quad \lim_{r \to 0} \rho = 0; \quad \lim_{r \to 0} \alpha(r) = -n \tag{13a; b; c; d}$$

so that, in view of (13a), far away from the vortex center r=0 the order parameter  $\phi$  behaves as  $\phi \sim \sqrt{n_0} e^{in\theta}$ . The vortex charge is given by  $Q = \int_{\mathbf{x}} (\rho - n_0)$  where the second term represents the external (background) charge which must be substracted [21]. This, combined with  $n_0 = \eta B$ ,  $f_0 = -\frac{\rho}{\eta}$  and the above boundaries yields  $Q = 2\pi\eta n$ . Therefore,

when statistical transmutation occurs

$$Q = \frac{n}{2k+1} \tag{14}$$

so that the vortices can be interpreted as the fractionally charged Laughlin quasiparticles, as it is the case in the CS description of QHF [16]. Finally, expanding S' around the (mean field) uniform solution (12) up to the quadatric order in the fields fluctuations, fixing the gauge freedom and integrating out the fluctuations of the statistical field gives rise to the low energy effective action for the matter degree of freedom from which the dispersion relation for the system is easily extracted. It can be written in obvious notations as

$$\omega_0^2(\mathbf{p}) = \frac{\mathbf{p}^4}{4m^2} (1 - 2\Theta_0 [1 - 2(\frac{m-\mu}{\eta e^2})^2]) + \mathbf{p}^2 (\frac{\mu}{m} + \Theta_0 + 2\Theta_0^2 (\frac{m-\mu}{m})) + \frac{n_0^2}{m^2 \eta^2} (1 - 2\Theta_0^2)$$
(15)

with  $\Theta_0 = \frac{n_0}{m\eta^2 e^2}$  indicating that the system has a gap (last term in (15)), as it could have been expected, so that the fluid described by S' is incompressible. Notice that the terms involving  $\Theta_0$  can be safely neglected, since  $\Theta_0 \ll 1$  holds in the present situation, so that (15) reduces to  $\omega_0^2(\mathbf{p}) = \frac{\mathbf{p}^4}{4m^2} + \mathbf{p}^2(\frac{\mu}{m}) + + \frac{n_0^2}{m^2\eta^2}$ .

### CONCLUSION.

The role of anyonic excitations in fast rotating harmonically trapped Bose gases in a fractional Quantum Hall state has been examined. Standard CS anyons as well as "non standard" anyons obtained from a statistical interaction having MCS dynamics and suitable modified coupling to matter have been considered. Their respective ability to stabilize attractive Bose gases under fast rotation in the thermodynamical limit has been studied. Stability can be obtained for standard anyons [11] while for non standard anyons, stability requires that the range of the corresponding statistical interaction does not exceed the typical wavelength for the atoms.

A part of the material involved in this note has been obtained in collaboration with A. Lakhoua, M. Lassaut and T. Masson. I am grateful to the organizers of the "School on Quantum Phase Transitions and Non-Equilibrium Phenomena in Cold Atomic Gases" for invitation. Hospitality of ICTP Trieste is also acknowledged.

- M.H. Anderson et al., Science 269 (1995) 198; K.B. Davis et al., Phys. Rev. Lett. 75 (1995) 3969; W. Ketterle, Physics Today 52 (2000) 30.
- [2] : A. Görlitz et al., Phys. Rev. Lett. 87 (2001) 130402.
- [3] : M.R. Matthews et al., Phys. Rev. Lett. 83 (1999) 2498.
- [4] : K.W. Madison, F. Chevy, W. Wohlleben, J. Dalibard, Phys. Rev. Lett. 84 (2000) 806; F. Chevy, M. Madison, J. Dalibard, Phys. Rev. Lett. 85 (2000) 2223; J.R. Abo-Shaeer, C. Raman, J.M. Vogels, W. Ketterle, Science 292 (2001) 476.
- [5] : For a review on FQHE see e.g. "The Quantum Hall Effect", R.E. Prange, S.M. Girvin eds., Springer-Verlag, Berlin 1990 and references therein.
- [6] : When the trapping and centrifugal potential balance each other, the bosons feel only the Coriolis force. Then, the system is formally equivalent to that of 2D (bosonic) particles in a magnetic field, corresponding to the condition for occurence of Quantum Hall regime.
- [7]: N.K. Wilkin, J.M.F. Gunn, R.A. Smith, Phys. Rev. Lett. 80 (1998) 2265; N.R. Cooper, N.K. Wilkin, Phys. Rev. B60 (1999) R16279; N.K. Wilkin, J.M.F. Gunn, Phys. Rev. Lett. 84 (2000) 6; N.R. Cooper, N.K. Wilkin, J.M.F. Gunn, Phys. Rev. Lett. 87 (2001) 120405.
- [8]: A.D. Jakson et al., Phys. Rev. Lett. 86 (2001) 945; B. Paredes, P. Fedichev, J.I. Cirac, P. Zoller, Phys. Rev. Lett. 87 (2001) 010402; T.-L. Ho, Phys. Rev. Lett. 87 (2001) 060403; U.R. Fischer, G. Baym, Phys. Rev. Lett. 90 (2003) 140402.
- [9]: M. Manninen, S. Viefers, M. Koskinen, S.M. Reimann, Phys. Rev. B64 (2001) 245322; J. Sinova, C.B. Hanna, A.H. MacDonald, Phys. Rev. Lett. 89 (2002) 030403; V. Schweikhard et al., Phys. Rev. Lett. 92 (2004) 040404; N. Regnault, T. Jolicoeur, Phys. Rev. Lett. 91 (2003) 030402, Phys. Rev. B69 (2004) 235309.
- [10] : For reviews, see e.g. A. Lerda, "Anyons: Quantum Mechanics of Particles with Fractional Statistics", Lecture Notes in Physics, vol.14 (Springer Verlag, Berlin, 1992); A. Khare, "Fractional Statistics and Quantum Theory" (World Scientific, Singapore, 1997).
- [11] : U.R. Fischer, Phys. Rev. Lett. 93 (2004) 160403.
- [12] : Note that in the thermodynamical limit, a system of attractive untrapped bosons is unstable against collapse and is stable only for a finite number of trapped particles; see P. Nozières, D. Saint James, J. Phys. (Paris) 43 (1982) 1133.
- [13]: J. Stern, Phys. Lett. B265 (1991) 119; I.I. Kogan, Phys. Lett. B 262 (1991) 83; Y. Georgelin, J.C. Wallet, Mod. Phys. Lett. A7 (1992) 1149, Phys. Rev. D50 (1994) 6610.

- [14] : F. Dalfovo, S. Giorgini, L.P. Pitaevskii, S. Stringari, Rev. Mod. Phys. 71 (1999) 463; A. J. Leggett, Rev. Mod. Phys. 73 (2001) 307.
- [15] : Our conventions are  $\hbar = c = 1$  when not explicitly written; indices A, B = 1, ..., N label bosons; i, j = 1, 2 label coordinates in the plane,  $\mathbf{x} \equiv (x_1, x_2), \epsilon_{12} = 1, \mathbf{x}_{AB} = \mathbf{x}_A \mathbf{x}_B$ ; spacetime metric is  $g_{\mu\nu} = \text{diag}(+, -, -)$ , Greek indices  $\mu, \nu, ... = 0, 1, 2, \epsilon_{012} = 1,$  $\partial_i^{(A)} \equiv \frac{\partial}{\partial x_A^i}, \ \partial_0 \equiv \partial_t, \ \int_{\mathbf{x}} \equiv \int dt d\mathbf{x}, \ \int_{\mathbf{x}} \equiv \int dt d\mathbf{x}, \ \int_t \equiv \int dt.$  Summation over repeated indices is understood.
- [16]: S.-C. Zhang, T.H. Hanson, S. Kivelson, Phys. Rev. Lett. 62 (1989) 82.
- [17] : S. Deser, R. Jackiw, S. Templeton, Ann. Phys. 140 (1982) 372.
- [18] : It can be realized that the general (static) solutions of (5b) such that  $f_{\mu} \neq -\frac{1}{n} \mathcal{J}_{\mu}$  have infinite energy.
- [19] : S. Latinski, D. Sorokin, Mod. Phys. Lett. A6 (1991) 3525.
- $\begin{bmatrix} 20 \end{bmatrix} : \frac{\Lambda}{\Lambda_{st}} \ll 1 \text{ corresponds to a statistical interaction with long range which seems to be unlikely. A large <math>\Lambda_{st}$  would correspond either to the small  $e^2$  regime where MCS theory enters a strong coupling regime for which a description of anyons as given by (3) is questionable or to a fractional Quantum Hall state with  $\eta \ll 1$ , e.g. state with extremely large denominator.
- [21] : Z.F. Ezawa, H. Hotta, A. Iwazaki, Phys. Rev. D44 (1991) 452.
- [22] : R. Jackiw, S.Y. Pi, Phys. Rev. Lett. 64 (1990) 2969.