

Phase transition in liquids with directed intermolecular bonding

R.Ryltcev, L.Son

Ural State Pedagogical University, 620219 Ekaterinburg, Russia.

Abstract

Liquids with quasi - chemical bonding between molecules are described in terms of vertex model. It is shown that this bonding results in liquid - liquid phase transition, which takes place between phases with different mean density of intermolecular bonds. The transition may be suggested to be a universal phenomena for those liquids.

keywords: Liquid - liquid phase transition, intermolecular bonds, vertex model.

1 Introduction.

The idea that bonding in molecular liquids may result in existence of two structural states, was first suggested for water [1]-[6]. At low temperatures, supercooled water demonstrates two possible local arrangements of molecules at some intermediate length scale, high density (HD) and low density (LD) ones. This results in a competition between these two states, so the supercooled water may be treated as a binary mixture of HD and LD liquids [2]. The liquid - liquid phase transition [1, 2] in supercooled water is the ultimate manifestation of the HD and LD local states. Now, the water HD and LD local structures are well established both in experiment and in computer simulations (see, for example, [3]). Intermolecular coupling by hydrogen bonds occurs to be a mechanism for HD and LD local states formation [5], original idea was published in [6].

Recently, the unexpected physical behavior was found in Benzene, Quinoline, and other complex fluids [7]-[14]. Namely, the anomalies (discontinuities, kinks, peaks) on temperature dependencies of a variety of physical properties were observed. These anomalies may be treated as a manifestation of changes in the local structure and may be observed, under special experimental conditions, in many equilibrium and supercooled liquids. The molecules of those liquids demonstrate some common features. For considerations related to the structure of the liquid, these molecules may be treated as rigid bodies without internal degrees of freedom. The relatively rigid electronic configurations of the molecules are characterized by a significant separation of positive and negative electric charges at the molecular length scale. At very short distance between molecules, strong repulsion determines the interaction, and at the only slightly larger length-scale, attraction become important. At this distance, the electric charge distribution may be characterized in terms of rigid electrical multipoles. Interactions of these multipoles, together with the very short-range repulsion, determine the set of mutual positions and orientations of two

neighboring molecules corresponding to potential energy minima, which provide intermolecular bonds.

Thus, the intermolecular coupling seems to be a mechanism for two or more structures formation at some intermediate length scale. In present paper, we apply known method of description of the polymer solution to the statistics of couplings in molecular liquids. The structure of paper is as follows.

- In section 2, we give a brief introduction into formalism applied.
- Then, in section 3, we analyze the mean - field solution of the model, which demonstrates liquid - liquid phase transition.
- In section 4, exact solution of the model on the Bethe lattice is analyzed. It is shown that the phase transition takes place in this case too.
- In Conclusion, general evaluation of the model is given.

The aim of the paper is to demonstrate that local structure separation is rather universal phenomena for liquids with hydrogen - like bonding.

2 The model.

Statistical description of condensed substances with the possibility of association and aggregation of molecules has been intensively developed for a long time. The theory of associated solutions [15], the Flory theory of polymer solutions [16], are the milestones on this way. The idea of modelling of chemically aggregated systems in terms of scalar order parameter has a long history also. Probably I.Lifshitz was first who offered this approach [17]. Then, Flory mean - field theory [16] and spin - polymer analogy [18] were applied to describe polymer systems. In the present paper, we consider the model which is a step on this way.

Quasi - chemical bonding between molecules has been characterized by two main features. First, the bond is short - range and directed (i.e., may be drawn as the vector connecting two neighboring molecules). Second, the bonding is saturated, i.e. no more than some characteristic number of bonds N may be permitted per one molecule. The same features may be suggested for the bonds in polymers, so the statistical description of polymer solutions may be applied to the molecular liquids with bonding. The difference is that in the theory of polymers, one supposes the polymer chain very long, while in our case, the chain of bonded molecules may have an arbitrary length.

The idea of the model arises from the works of Baxter [25], Nikomarov and Obukhov [19], and may be formulated as follows.

Consider spatial net of sites which approximately corresponds to the spacing of molecules in the liquid. Neighboring sites are connected with each other by edges. The number of nearest neighbors we denote as γ . Let us numerate the sites by Greece indexes, and the edges by Latin ones. Each site α is characterized by variable n_α , which is the filling number

$$n_\alpha = \begin{cases} 1, & \text{if the site } \alpha \text{ is occupied by the molecule} \\ 0, & \text{if the site } \alpha \text{ is empty} \end{cases} \quad (1)$$

Also, real scalar variable ψ_i corresponds to the edge numbered by i . Let us consider the partition

$$Z = \sum_{\{n\}} \int D\psi \exp[-F\{n, \psi\}], \quad (2)$$

where the summation and integration goes over all possible configurations $\{n, \psi\}$ of variables n_α, ψ_i . The "effective Hamiltonian" $F\{n, \psi\}$ is defined by

$$F\{n, \psi\} = -\frac{1}{2} \sum_{\alpha, \beta} n_\alpha J_{\alpha\beta} n_\beta - \frac{\mu}{T} \sum_{\alpha} n_\alpha + \frac{1}{2} \sum_i K \psi_i^2 - \sum_{\alpha} \ln [1 + n_\alpha R_\alpha(\psi)] \quad (3)$$

$$R_\alpha(\psi) = a_1 \sum_{i_\alpha^1} \psi_{i_\alpha^1} + a_2 \sum_{i_\alpha^1 \neq i_\alpha^2} \psi_{i_\alpha^1} \psi_{i_\alpha^2} + \dots + a_N \sum_{i_\alpha^1 \neq i_\alpha^q} \psi_{i_\alpha^1} \psi_{i_\alpha^2} \dots \psi_{i_\alpha^N}. \quad (4)$$

Here, indexes $i_\alpha^1, i_\alpha^2, \dots, i_\alpha^p$, numerate the edges which are connected with the site α . Spatial matrix $J_{\alpha\beta}$ is

$$J_{\alpha\beta} = \begin{cases} J/T, & \text{if } \alpha, \beta \text{ are the nearest neighbors} \\ 0, & \text{otherwise} \end{cases}, \quad (5)$$

where J is the energy of attractive non - directed (Van der Waals) interaction between molecules, and μ is their chemical potential. Thus, the first and the second terms in (3) correspond to the lattice - gas model [25] and allow one to describe the density of the system. The value K is

$$K = \exp(U/T), \quad (6)$$

where U is the energy of quasi - chemical bond. In (5, 6), $T = k_B T_k$, where T_k, k_B are the temperature and the Boltzmann constant respectively. The last two terms in (3) allow one to describe bonding. To demonstrate this fact, let us rewrite (2)

$$\begin{aligned} Z &= \sum_{\{n\}} e^{-F\{n\}} \int D\psi \exp \left[-\frac{1}{2} \sum_i K \psi_i^2 \right] \prod_{\alpha} (1 + n_\alpha R_\alpha(\psi)), \\ F\{n\} &= -\frac{1}{2} \sum_{\alpha\beta} n_\alpha J_{\alpha\beta} n_\beta - \frac{\mu}{T} \sum_{\alpha} n_\alpha, \end{aligned} \quad (7)$$

and consider the functional integral over $\{\psi\}$ at some given configuration of filling numbers $\{n\}$. Releasing the brackets in the product, one gets the sum over all possible series of products:

$$\prod_{\alpha} (1 + n_\alpha R_\alpha(\psi)) = 1 + n_1 a_1 \psi_{1_1} + n_1 a_1 \psi_{2_1} + \dots n_1 n_2 a_1^2 \psi_{1_1} \psi_{2_1} + \dots \quad (8)$$

Then, the integration with the Gaussian weight $\exp \left[-\frac{1}{2} K \sum \psi_i^2 \right]$, produces all possible couplings between pairs of neighboring sites. The coupling takes place when the mathematical power of ψ on corresponding edge is two. Zero power corresponds to the edge without coupling. The ratio of the weight of coupled edge to the weight of uncoupled one is $K^{-1} = \exp(-U/T)$. Thus, the partition (8) generates the sum over all possible configurations of occupied sites, connected by all possible couplings between nearest neighbors, which correspond to intermolecular bonds. Each site with k bonds along its edges has an additional weight a_k . No more than one bond per edge, and no more than N bonds per site are permitted due to the structure of polynomial $R_\alpha(\psi)$, see (4).

Besides, each configuration of filling numbers $\{n\}$ has the standard weight of the lattice - gas theory, $\exp \left[\frac{1}{2} \sum_{\alpha\beta} n_\alpha J_{\alpha\beta} n_\beta - \frac{\mu}{T} \sum_{\alpha} n_\alpha \right]$. Thus, in the frames of formalism considered, one can easily model an arbitrary molecular liquid by an appropriate choice of model parameters. Those are:

U	The energy of quasi - chemical bond
N	The maximal number of bonds per molecule
J	The energy of non - directed (Van der Waals) interaction
μ	The chemical potential of molecules
a_m	The weight of m - bonded molecule

Instead of filling numbers, it is more useful to deal with scalar variables without any limitations imposed. To do so, Hubbard - Stratonovich transform to conjugated field may be used. Then, the summation over filling numbers leads to

$$Z = \int D\psi D\varphi \exp[-F\{\varphi, \psi\}]$$

$$F\{\varphi, \psi\} = \frac{1}{2}K \sum_i \psi_i^2 + \frac{1}{2} \sum_{\alpha\beta} \varphi_\alpha J_{\alpha\beta}^{-1} \varphi_\beta - \sum_\alpha \ln \left[1 + e^{\varphi_\alpha + \frac{\mu}{T}} (1 + R_\alpha(\psi)) \right]. \quad (9)$$

To analyze functional integral (9), one can apply traditional method: to find most probable configuration which minimizes the $F\{\varphi, \psi\}$, then to investigate fluctuations around it, etc. In the next section, we describe the first step, which is the mean - field approximation.

3 Mean - field approximation.

The mean - field configuration $\langle \varphi_\alpha \rangle, \langle \psi_i \rangle$ provides the minima of (9) and obeys following equations:

$$\frac{\partial F}{\partial \psi_i} |_{\langle \varphi_\alpha \rangle, \langle \psi_i \rangle} = 0, \quad \frac{\partial F}{\partial \varphi_\alpha} |_{\langle \varphi_\alpha \rangle, \langle \psi_i \rangle} = 0. \quad (10)$$

Instead of φ_α , it is better to use the variable

$$q_\alpha = \sum_\beta J_{\alpha\beta}^{-1} \varphi_\beta,$$

because its mean - field value,

$$w_\alpha = \langle q_\alpha \rangle, \quad (11)$$

coincides with the mean value of filling number $\langle n_\alpha \rangle$ and should be understood as the mean concentration of the molecules.

For homogeneous system, $\langle \psi_i \rangle = \Psi$, $w_\alpha = w$, equations (10) may be written in explicit form

$$w = \frac{\exp[\frac{J\gamma}{T}w + \frac{\mu}{T} + \ln(1 + R(\Psi))]}{1 + \exp[\frac{J\gamma}{T}w + \frac{\mu}{T} + \ln(1 + R(\Psi))]} \quad (12)$$

$$\Psi = \frac{n_s}{n_e} e^{-\frac{\mu}{T}} w \frac{R'(\Psi)}{1 + R(\Psi)} \quad (13)$$

$$R(\Psi) = a_1 \gamma \Psi + a_2 \frac{\gamma(\gamma-1)}{2} \Psi^2 + \dots + a_N C_\gamma^N \Psi^N \quad (14)$$

where n_s/n_e is the ratio of the number of sites to the number of edges in the lattice and γ is the number of nearest neighbors. Solutions of system (12-14) determine the equilibrium values

of Ψ, w . For nonequilibrium Ψ, w , expression (9) gives the density of thermodynamic potential in the Landau theory [20]:

$$f(\Psi, w) = \frac{n_e}{2n_s} e^{\frac{U}{T}} \Psi^2 + \frac{J\gamma}{2T} w^2 - \ln \left[1 + e^{\frac{J\gamma}{T} w + \frac{\mu}{T}} (1 + R(\Psi)) \right]. \quad (15)$$

Solutions of system (12-14) correspond to different phases of the system. The stable phases correspond to minimums of (15).

The first of equations (12) is exactly the equation which arises in the lattice - gas theory of critical point:

$$\ln \frac{w}{1-w} = \frac{J\gamma}{T} w + B, \quad B = \frac{\mu}{T} + \ln(1 + R(\Psi)). \quad (16)$$

Depending on parameters T, B , this equation may have from one to three solutions on the interval $(0, 1)$. The solution with $w \simeq 1$ corresponds to the condensed phase (liquid), while the solution $w \simeq 0$ represents the gas. We are interested in the condensed phase behavior, far from the liquid - gas critical point. Thus, we consider the case when the equation (16) has only one solution, $w \simeq 1$. This may always be provided by an appropriate choice of parameter μ in (16). If the equation (16) has only one solution, then one gets the Landau theory with single order parameter Ψ and with thermodynamic potential

$$f(\Psi) = \frac{1}{2b} \Psi^2 + \frac{a}{2} w^2(\Psi) - \ln \left[1 + e^{aw(\Psi) + \frac{\mu}{T}} (1 + R(\Psi)) \right], \quad (17)$$

where $w(\Psi)$ is given by (16), $b = \frac{n_s}{n_e} e^{-\frac{U}{T}}$, $a = \frac{J\gamma}{T}$. Function $f(\Psi)$ may have several minima, which corresponds to different local arrangements of bonds. These minima are determined by equation (13):

$$\Psi = bw(\Psi) \frac{R'(\Psi)}{1 + R(\Psi)}. \quad (18)$$

The formalism considered allows to describe a wide variety of intermolecular bonding in a liquid. The bonding has been modelled by the choice of the polynomial under the logarithm in (15):

$$R(\Psi) = a_1 \gamma \Psi + a_2 \frac{\gamma(\gamma-1)}{2} \Psi^2 + \dots + a_N C_\gamma^N \Psi^N \quad (19)$$

The number N is the maximal number of bonds per molecule, and the coefficient a_m is the weight of m -bonded molecule. Note, that $P(\Psi)$ is positive for most physical cases. Since the coupling arises due to the charge redistribution in the molecule, the maximal number of bonds is even. The odd number of bonds means that the charge distribution in the molecule is not symmetric, so

$$a_{2n} \geq a_{2n-1}.$$

It can be easily shown, that this provides positive values of $R(\Psi)$ at any Ψ .

At low temperatures, $w(\Psi) \simeq 1$, and (18) may be rewritten as

$$B\Psi = \frac{R'(\Psi)}{1 + R(\Psi)}, \quad B \sim \exp \left[\frac{U}{T} \right]. \quad (20)$$

Its graphic solution is presented on fig.1a. The right side of the equation,

$$\frac{R'(\Psi)}{1 + R(\Psi)} \xrightarrow{\Psi \rightarrow \pm\infty} 0,$$

with some intermediate behavior, as it presented on fig.1a. Since $U < 0$, the left side is a line which slope lowers with temperature lowering. Thus, if the temperature is low enough, there always exist nontrivial set of solutions, which starts from the minimum of $f(\Psi)$ at large negative Ψ , and completes with another minimum at large positive Ψ .

Here, some comments should be done. The mean - field approximation does not answer the question what is the physical difference between these two low temperature structures. To do so, one needs higher approximations. Similar situation takes place for the Ising model in two dimensions [25]. (In 2D, the Ising model may be formulated in terms of scalar field which corresponds to magnetization. At the same time, it may be understood as a statistics of self - closed contours which divide areas with inverted spins. Mean magnetization gives poor information about contours). Nevertheless, some assertions may be done. In our case, absolute value of Ψ is proportional to the mean length of quasi - polymers [19]. The structure $\Psi < 0$ is metastable, i.e. $f(\Psi < 0) > f(\Psi > 0)$ (fig.1b). It may be stabilized by external field s conjugated to ψ , i.e. by adding the term $s\psi$ to the Hamiltonian. Such a term produces spatial points, where the quasi - polymers have been terminated. Thus, the difference between $\Psi > 0$ and $\Psi < 0$ structures lies in the statistics of polymer tails. For the Bethe lattice, the metastable state does not exist (see the next section). Since for the Bethe lattice closed paths are prohibited, then it is natural to suggest that the metastable state may be characterized by a set of self - closed quasi - polymers.

When the maximal number of bonds per molecule $N \geq 4$, additional solutions of equation (18) arise. Mathematically, these solutions arise due to kinks in the right hand side of (18), provided by high powers in the polynomial $R(\Psi)$. As an example, the graphic solution of eq.(18) at some certain model parameters is presented at fig.1c, together with corresponding $F(\Psi)$ (fig.1d) dependence. As can be easily understood, the existence of such solutions may result in the liquid - liquid phase transition. The model predicts possible phase diagrams in the $\mu - T$ plane, which are presented at fig.2. Phase diagrams of that type were suggested earlier [22, 23]. Note, that the diagram at fig.2b demonstrates an interesting possibility for the system behavior: there is no first order transition in density parameter w , and Ψ occurs to be the relevant order parameter for the liquid - gas critical point.

Thus, the mean - field approximation predicts two non - trivial features. First, there exists metastable state with negative Ψ . This state appears at rather low temperatures. Second, high powers of $R(\Psi)$ which correspond to the case $N \geq 4$, may provide liquid - liquid phase transition at some sets of weights a_n . The transition is the first order phase transition, which line terminates in the critical point at the $\mu - T$ plane. At this line, order parameter jumps from Ψ_1 to Ψ_2 . In the critical point, the jump vanishes. Since the absolute value of Ψ is proportional to the mean length of quasi - polymers, then higher Ψ corresponds to the higher number of active bonds. The density difference is small, but higher Ψ corresponds to the higher density also.

To verify these results, in the next section we analyzed the exact solution of the model on the Bethe lattice.

4 The model on the Bethe lattice.

In this section, we put $\omega_\alpha = 1$ for simplicity. All the results obtained may be easily derived for arbitrary concentration, but we are interested mostly in the high density limit. Thus, we deal with the partition

$$Z = \int D\psi e^{-H_0\{\psi, s\}} \prod_{\alpha} [1 + R_{\alpha}(\psi)], \quad (21)$$




Figure 1: The graphic solution of equation (18), i.e. left and right sides of this equation, and corresponding plot of thermodynamic potential (17) at $\tilde{T} = T/U = 0.5$ (a - graphic solution, b - thermodynamic potential) and $\tilde{T} = T/U = 1.019$ (c - graphic solution, d - thermodynamic potential). The calculations are done in reduced variables: $n_e/n_s = \lambda = 3$, $\gamma = 6$, $a_1 = \dots = a_4 = a = 0.015$, $\tilde{J} = J/U = 0.2$, $\tilde{\mu} = \mu/U = 1$. Parameter $N = 4$ in both cases.

on the Bethe lattice. Here,

$$H_0 \{ \psi \} = \frac{1}{2} \sum_i K \psi_i^2 - \sum_i s_i \psi_i$$

In H_0 , we used the source term with $s_i = s$. Partition (21) may be treated as the generating function for mean values

$$\langle \psi_i \rangle = \frac{1}{Z} \frac{\partial Z}{\partial s_i} \Big|_{s=0}, \quad \langle \psi_i \psi_j \rangle = \frac{1}{Z} \frac{\partial^2 Z}{\partial s_i \partial s_j} \Big|_{s=0}, \quad \dots \quad (22)$$

For each edge i let us define function $Q(\psi_i)$, which is the partition (21) for sublattice, obtained from the initial Bethe lattice by cut of the edge i . In the thermodynamic limit, the partition for initial lattice depends no on the choice of the edge and is given by

$$Z = \int d\psi_i e^{-\frac{K}{2} \psi_i^2 + s_i \psi_i} Q^2(\psi_i). \quad (23)$$

Let us introduce the values

$$A = \int d\psi e^{-\frac{K}{2} \psi^2 + s\psi} Q(\psi) \psi, \quad (24)$$




Figure 2: The $\tilde{\mu} - \tilde{T}$ phase diagrams at $\tilde{J} = 0.7$ (a) and $\tilde{J} = 0.3$, (b). Other parameters are $n_e/n_s = \lambda = 3$, $\gamma = 6$, $a_1 = \dots = a_4 = a = 0.015$, $N = 4$. Liquid - liquid and liquid - gas critical points are denoted as C_1 and C_2 respectively

$$B = \int d\psi e^{-\frac{K}{2}\psi^2 + s\psi} Q(\psi). \quad (25)$$

At $s = 0$, these values are proportional to the probabilities of the edge to be with or without bond, respectively. Obviously,

$$Q(\psi) = F_1(A, B)\psi + F_2(A, B), \quad (26)$$

where explicit form of functions $F_1(A, B)$ and $F_2(A, B)$ depends on model details (the number of nearest neighbors and the set of coefficients a_n). Integrating (26), one gets system of equations on values A, B :

$$A = F_1(A, B)J_2(K, s) + F_2(A, B)J_1(K, s) \quad (27)$$

$$B = F_1(A, B)J_1(K, s) + F_2(A, B)J_0(K, s).$$

Here,

$$J_n(K, s) = \int_{-\infty}^{+\infty} d\psi e^{-\frac{K}{2}\psi^2 + s\psi} \psi^n = \sqrt{2\pi} K^{-\frac{3}{2}} \frac{d^{n-1}}{ds^{n-1}} \left(s e^{\frac{s^2}{2K}} \right)$$

From (22), (23), (26),(27) one obtains

$$\langle \psi \rangle = \frac{2F_1 F_2}{F_1^2 + F_2^2 K}, \quad \langle \psi^2 \rangle = \frac{3F_1^2 + F_2^2 K}{F_1^2 K + F_2^2 K^2}. \quad (28)$$

At $s = 0$ system (27) takes the form

$$A = F_1(A, B)J_2(K, 0) = F_1(A, B)\sqrt{2\pi}K^{-\frac{3}{2}} \quad (29)$$

$$B = F_2(A, B)J_0(K, 0) = F_2(A, B)\sqrt{2\pi}K^{-\frac{1}{2}},$$

and

$$\langle \psi \rangle = \frac{2AB}{KA^2 + B^2}, \quad \langle \psi^2 \rangle = \frac{3KA^2 + B^2}{K^2A^2 + KB^2}. \quad (30)$$

As can be seen from (29), (30), any mean value may be presented as a function of the parameter $x = A/B$:

$$\langle \psi \rangle = \frac{2x}{Kx^2 + 1}, \quad \langle \psi^2 \rangle = \frac{3Kx^2 + 1}{K^2x^2 + K}. \quad (31)$$

The equation on x is

$$Kx = y(x), \quad (32)$$

where $y(x) = F_1(x)/F_2(x)$.

Any thermodynamic function may be reconstructed using $\langle \psi^2 \rangle$. For example,

$$\frac{\partial \ln Z}{\partial T} = -\frac{U}{2T^2} K \langle \psi^2 \rangle.$$

Using known relation for internal energy,

$$E = T^2 \frac{\partial \ln Z}{\partial T} \quad (33)$$

one gets

$$E = -\frac{U}{2} K \langle \psi^2 \rangle. \quad (34)$$

Inserting $F = -T \ln Z$ into (33) and integrating, one gets the free energy thermodynamic potential

$$F(T) - F(T_0) = \frac{T}{2} \int_{T_0}^T d\tau \frac{K(\tau)}{\tau^2} \langle \psi^2(\tau) \rangle, \quad (35)$$

where $\langle \psi^2(\tau) \rangle$ is defined by (31) and (32).

Thus, in the thermodynamic limit, any thermodynamic characteristic may be calculated using $y(x)$. Now, let us determine this function for arbitrary model parameters. Those are the maximal number of bonds per molecule N , the number of nearest neighbors γ , and the set of coefficients a_n . The only limitation imposed is $\gamma \geq N$. Consider some site α and suggest that the integration in (8) is done over all ψ except those which correspond to the site α . If one of these edges is cut (for definition, the edge ψ_1), then initial lattice occurs to be divided in two sublattices (fig.3).

Then, $Q(\psi_1)$ may be written as

$$Q(\psi_1) = \int \left(\prod_{i=2}^{\gamma} d\psi_i e^{-\frac{K}{2}\psi_i^2} Q(\psi_i) \right) [1 + a_1(\psi_1 + \dots + \psi_{\gamma}) + a_2(\psi_1\psi_2 + \dots + \psi_{\gamma-1}\psi_{\gamma}) + \dots]. \quad (36)$$

In the thermodynamic limit, $Q(\psi_i)$ depends no on i . Integration together with (24), (25) and (26) gives

$$\begin{aligned} F_1(A, B) &= a_1 B^{\gamma-1} + a_2(\gamma-1)AB^{\gamma-2} + \dots + a_N C_{N-1}^{\gamma-1} A^{N-1} B^{\gamma-N} \\ F_2(A, B) &= B^{\gamma-1} + a_1(\gamma-1)AB^{\gamma-2} + \dots + a_N C_N^{\gamma-1} A^N B^{\gamma-N-1}, \end{aligned} \quad (37)$$




Figure 3: The cut of the initial Bethe lattice across the edge ψ_1 leads to the equation (36)

where $C_m^n = n!/m!(n-m)!$ - combinatoric coefficients. Since $x = A/B$, $y(x) = F_1(x)/F_2(x)$, then

$$y(x) = \frac{\sum_{m=0}^{N-1} a_{m+1} C_m^{\gamma-1} x^m}{\sum_{m=0}^N a_m C_m^{\gamma-1} x^m}. \quad (38)$$

Thus, equation (32) is algebraic with power $N-1$. Qualitatively, its structure coincides with mean - field equation (18). It should be mentioned, however, that (32) may be treated as a recurrence relation $Kx_n = y(x_{n-1})$, which connects x values for n and $n-1$ levels of the Cayley tree, taken in the thermodynamic limit. Since any initial $x = A/B$ is positive, then all stationary points of this recurrence are positive too. Thus, on the Bethe lattice, the low - temperature metastable state does not exist.

5 Conclusion

Here, we point out some general results and features which make the model rather attractive.

1. In mean field approximation, the metastable low - temperature state is predicted. This state is universal, i.e. appears for any set of model parameters. It becomes stable only at zero temperature, but may be stabilized by an appropriate external field. For example, phase separation in two phases with positive and negative Ψ may be achieved by impurity addition. Recently, similar separation was observed experimentally [24]. For the Bethe lattice, the metastable state does not exist. Since for the Bethe lattice closed paths are prohibited, then it is natural to suggest that the metastable state may be characterized by a set of self - closed quasi - polymers.
2. Liquid - liquid phase transition between phases with different density of bonds is predicted both in mean - field approximation and on the Bethe lattice. Possible phase diagrams correspond to those predicted by computer simulations [22, 23]. The phase transition is

the first order phase transition which takes place at some special sets of model parameters. Also, the set of model parameters may provide not phase transition but sharp crossover from one mean density of bonds to another. In that case, the order parameter temperature dependence demonstrates a kink. Thus, all Ψ - dependent properties should demonstrate a kink too. Such a kinks have often been observed in molecular liquids [7]-[14].

3. The model considered allows one to describe an arbitrary set of molecules and their bonding. For example, different conformations of bonds may be easily accounted by an appropriate choice of $R(\psi)$.
4. Actually, the model considered is the vertex model. Such models have often been used in statistical physics. For example, Ising model may be formulated as the vertex model [25]. Thus, the methods to investigate the model behavior are developed rather well.

The work is supported by RFBR (grants N 04-02-96095, 06-02-17269).

References

- [1] P. H. Poole, F. Sciortino, U. Essman, and H. E. Stanley, *Nature* (London) 360, 324 (1992).
- [2] E.G.Ponyatovskii, V.V.Sinitsyn, T.A.Pozdnyakova. *JETP Lett.*, **60**, 360 (1994).
- [3] G.G.Malenkov. *Computer simulation of supercooled water and amorphous ice*. Proceedings of the NATO Advances Research Workshop, Volga River, on *New Kinds of Phase Transitions: Transformations in Disordered Substances*, edited by V. V. Brazhkin, S. V. Buldyrev, V. N. Ryzhov, and H. E. Stanley. Kluwer Academic Publishers, Dordrecht, 2002.
- [4] C.M. Davis and J. Jarzynski, *Mixture Models of Water*, ch.10 in *Water and Aqueous Solutions. Structure, Thermodynamics, and Transport Processes*, R.A. Horne, ed., Wiley-Interscience, New York (1972).
- [5] G.Franzese, M.I.Marques, H.E.Stanley. Intramolecular coupling as a mechanism for a liquid - liquid phase transition. *Phys. Rev. E* 67 (2003) 011103, cond-mat/0112341.
- [6] G.Franzese and H.E.Stanley. Liquid - liquid critical point in a Hamiltonian model for water: analytic solution. *J.Phys. Cond. Matter* 14 (2002) 2201.
- [7] E. W. Fischer, *Physica A* 201 (1993) 183, and references therein.
- [8] S.A. Kivelson, X. Zao, D. Kivelson T. M. Fisher and C. Knobler. *J. Chem. Phys.* 101 (1994) 2391.
- [9] N.B.Rozhdestvenskaya and L.V. Smirnova, *JETP Lett.* 44 (1986) 1166.
- [10] F. J. Bermejo, M. Garcia-Hernandez, W. S. Howells, R. Burriel, F. J. Mompeán, D. Martin, *Phys. Rev. E* 48 (1993) 2766.
- [11] D. Jalabert, J. B. Robert, H. Roux-Buisson, J. P. Kintzinger, J. M. Lehn, R. Zinzius, D. Canet and P. Tekely, *Europhys. Lett.* 115 (1991) 435; J. B. Robert, J. C. Boubel and D. Canet, *Molecular Physics* 90 (1997) 399; L. Gauthier, J. B. Robert, and D. Canet, *Journal of Molecular Liquids* 85 (2000) 77.

- [12] N. B. Rozhdestvenskaya and L. V. Smirnova, J. Chem. Phys. 195 (1991) 1223 .
- [13] L. Letamendia, M. Belkadi, O. Eloutassi, G. Nouchi, C. Vaucamps, S. Iakovlev, N. Rozhdestvenskaya, L V. Smirnova, M. Runova, Phys. Rev. E48 (1993) 3572.
- [14] L.Letamendia, M.Belkadi, O. Eloutassi, E.Pru-Lestret, G. Nouchi, J. Rouch, D. Blaudez, F.Mallamace, N.Micali, C.Vasi, Phys. Rev. E54 (1996) 5327.
- [15] I.Prigogine. *The molecular theory of solutions*. North - Holland and Interscience, Amsterdam - New York, 1957.
- [16] P.Flory. *Statistical mechanics of chain molecules*.
- [17] I.M.Lifshitz. Selected papers, Part II. Nauka, Moscow, 1994.
- [18] P.G. de Gennes. *Scaling Concepts in Polymer Physics*. Cornell University Press, Ithaca and London, 1983.
- [19] E.Nikomarov, S.Obukhov. An extended description of linear polymer solution in terms of polymer-magnet analogy. JETP 80 (1980) 651-665 .
- [20] L.D.Landau, E.M.Lifshitz. *The Course of Theoretical Physics, vol.5 (Statistical physics, Part 1)*. Nauka, Moscow, 1988.
- [21] F.Firouzi, H.Modarres, G. Ali Mansoori. Predicting Liquid-Liquid Transitions in Polymer Solutions using the GCLF Equation of State. Eur. Polym. J. Vol.34, N.10 (1998) 1489-1498.
- [22] G.Franzese, G.Malescio, A.Skibinsky, S.V.Buldyrev and H.E.Stanley. Generic mechanism for generating a liquid - liquid phase transition. Nature 409 (2001) 692.
- [23] G.Malescio, G.Franzese, G.Pellicane, A.Skibinsky, S.V.Buldyrev and H.E.Stanley. Liquid - liquid phase ptransition in one-component fluids. J. Phys. Cond. Matter 14 (2002) 2193-2200.
- [24] Hono K. Phase separation and nanocrystallization of bulk-forming metallic glasses. Abstracts of RQ12, 21-26 Aygust, Jeju, Korea, 2005, P. 4.
- [25] R.Baxter. Exactly solved models in statistical mechanics. Acad.Press, London, 1982.