## Comment on "Entropy of Classical Systems with Long-Range Interactions"

In a recent letter [1], T.M. Rocha Filho and coworkers address the very interesting issue of the entropic form to be used for Hamiltonians with long-range interactions. In our opinion the letter misses several points which are of fundamental importance. Moreover it contains several statements which are not correct as explained below.

First of all, contrary to several statements in [1], the generalization of standard statistics [2] is neither "meaningless and may lead to wrong conclusions" nor "limited in scope". Actually, it has proved to be very useful in a variety of physical situations and many other applications beyond physics [3]. Standard Boltzmann-Gibbs (BG) statistical mechanics is for sure limited to equilibrium situations where all the possible microstates are equally probable. On the other hand, nonextensive statistics represents one possible generalization for those frequent situations where this is not true. Moreover, although the form and extensivity of the entropy are addressed in [1], the authors ignore results [4] centrally relevant to precisely these questions. Finally, the criticisms cited in Refs. [13,14] of [1] have since long been replied in [5]. Coming back now to the more specific argument of long-range Hamiltonian systems discussed in [1], the authors do not focus on the main situations where nonextensive statistics has been applied, i.e. on the nonhomogeneous quasi-stationary states found in these systems and more specifically on the Hamiltonian Mean Field (HMF) model [6]. In [1] in fact the homogenity of the quasi-stationary state is always a-priori assumed together with an equiprobability of microstates. At variance, the numerical evidence [6-8] indicates a strong hierarchical microscopic structure (very sensitive to the initial conditions) which can also be interpreted within a glassylike formalism [7]. In a very recent paper by Morita and Kaneko [9], a metastable collective oscillation beyond Vlasov analysis has also been observed. These facts, together with the crucial importance of the neglected finitesize effects, invalidates the general conclusions of [1]. In [6] it has been clearly shown that the generalized statistics allows to predict, in a quantitative way and within a coherent frame, the q-exponential decay of the velocity correlation functions and the anomalous diffusion observed. This is true also for a generalized version of the model [10], for several system sizes and many kind of initial conditions. Homogeneous quasi-stationary states, which have an almost exponential velocity correlation function decay, can be also explained within this same scenario as a particular case. In general the prediction  $\gamma = 2/(3-q)$  for the anomalous diffusion coefficient has been successfully verified [6]. Although further investigation is needed, it is by now certain that the dynamical anomalies observed in the HMF model and long-range Hamiltonian systems go beyond the possible explanation of standard statistical mechanics, and that a new kind of kinetic theory is needed, as claimed also in [11]. Generalized statistics offers a quite plausible perspective in this direction. Let us finally summarize some of the relevant

statements made in [1] that are incorrect for the HMF model: (i) In the limit  $N \to \infty$  the interparticle correlations are negligible is not always true for long-range interactions and depends on the order of the limits; (ii) The supposition that all microstates compatible with the given constraints are equally probable is invalid, even for the microcanonical ensemble, for the quasi-stationary state in the presence of long-range interactions for the physically important case where  $t \to \infty$  after  $N \to \infty$ ; strong indications of nonergodicity are available in the literature; (iii) The BG entropy is then the correct form to be used is trivially correct for  $N \to \infty$  after  $t \to \infty$ , and clearly wrong the other way around, since the distribution of velocities is not Maxwellian in the quasi-stationary state.

## A. Rapisarda \*, A. Pluchino \* and C. Tsallis \*\*

- \* Dipartimento di Fisica e Astronomia and Infn sezione di Catania Universitá di Catania, Via S. Sofia 64, I-95123 Catania, Italy.
- \*\* Santa Fe Institute, 1399 Hyde Park Road, Santa Fe, NM 87501, USA, and CBPF, Rio de Janeiro, Brazil.

Electronic addresses: andrea.rapisarda@ct.infn.it, alessandro.pluchino@ct.infn.it, tsallis@santafe.edu

- T.M. Rocha Filho et al., Phys. Rev. Lett. 95, 190601 (2005).
  - [2] C. Tsallis, J. Stat. Phys. **52**, 479(1988).
- [3] J.P. Boon and C. Tsallis eds, *Nonextensive Statistical Mechanics:New Trends, New Perspectives*, Europhysics News **36**, number 6 (2005); see also M. Gell-Mann and C. Tsallis eds, *Nonextensive Entropy: Interdisciplinary Applications*, Oxford University Press, New York (2004).
- [4] C. Tsallis, M. Gell-Mann and Y. Sato, Proc. Natl. Acad. Sc. USA 102, 15377 (2005); C. Tsallis, Milan Jour. of Math. 73, 145 (2005); L.G. Moyano, C. Tsallis and M. Gell-Mann, Europhys. Lett. (2006), in press, cond-mat/0509229.
- [5] C. Tsallis, Phys. Rev. E  $\mathbf{69}$ , 038101 (2004) [cond-mat/0304696]; M. Nauenberg, Phys. Rev. E  $\mathbf{69}$ , 038102 (2004); C. Tsallis, Physica D  $\mathbf{193}$ , 3 (2004).
- [6] A. Rapisarda, A. Pluchino, p. 202 in ref. [3]; A. Pluchino, A. Rapisarda, Physica A (2006) in press cond-mat/0511570; A. Pluchino, V. Latora, A. Rapisarda, Physica D **193**, 315 (2004); V. Latora, A. Rapisarda, C. Tsallis, Phys Rev. E 64, 056134 (2001).
- [7] A. Pluchino, V. Latora, A. Rapisarda, Phys. Rev. E **69**, 056113 (2004); cond-mat/0506665 and cond-mat/0509031.
- [8] F. Tamarit and C. Anteneodo, p. 194 in ref. [3]; F.D. Nobre and C. Tsallis, Phys. Rev. E 68, 036115 (2003); M. Sagakami and A. Taruya, Cont. Mech. and Therm. 16,279 (2004) and Mon. Not. R. Astron. Soc. 364, 990 (2005).
- [9] H. Morita, K. Kaneko, Phys. Rev. Lett. 96, 050602 (2006).
- [10] C. Anteneodo, C. Tsallis Phys. Rev. Lett. 80, 5313 (1998); A. Campa, A. Giansanti, D. Moroni, J. Phys. Math. Gen. 36, 6897 (2003).
- [11] P-H. Chavanis, Eur. Phys. J. (2006) in press cond-mat/0509726 and references therein.