ULB-TH/05-27

Geometrical Thermodynamic Field Theory

Giorgio Sonnino* and Jarah Evslin[†]

EURATOM - Belgian State Fusion Association and

International Solvay Institutes

Department of Theoretical Physics and Mathematics

Free University of Brussels (U.L.B.), Blvd du Triomphe

Campus de la Plaine, C.P. 231, Building NO

Brussels, B-1050, Belgium

Abstract

A manifestly covariant, coordinate independent reformulation of the Thermodynamic Field Theory (TFT) is presented. The TFT is a covariant field theory that describes the evolution of a thermodynamic system, extending the near-equilibrium theory established by Prigogine in 1954. We introduce the Minimum Dissipation Principle, which is conjectured to apply to any system relaxing towards a steady-state. We also derive the thermodynamic field equations, which in the case of $\alpha - \alpha$ and $\beta - \beta$ processes have already appeared in the literature. In more general cases the equations are notably simpler than those previously encountered and they are conjectured to hold beyond the weak-field regime. Finally we derive the equations that determine the steady-states as well as the critical values of the control parameters beyond which a steady-state becomes unstable.

June 30, 2021

^{*}gsonnino@ulb.ac.be

[†]jevslin@ulb.ac.be

1 Introduction

In 1999 the *Thermodynamic Field Theory* (TFT) was proposed, which describes the behavior of thermodynamic systems beyond the linear regime [1]-[5]. This approach starts from the concept of *entropy production*, which can always be expressed as a bilinear form of *some* variables characterizing the departure from equilibrium. If these variables are the thermodynamic forces, it is possible to formulate a covariant TFT and to find the *thermodynamic field equations* whose solutions give the generalized relation between the thermodynamic forces and the conjugate flows.

In refs [1]-[5], the thermodynamic field equations have been obtained from three postulates:

- 1) The shortest path principle
- 2) The closedness of the thermodynamic field strength
- **3)** The principle of least action

where the thermodynamic field strength is determined from the skew-symmetric part of the tensor relating the thermodynamic forces with the conjugate flows. The second postulate reflects the observation that, in all cases examined so far, the sources of the thermodynamic fields i.e., the internal fluctuations, time-dependent boundary conditions and external perturbations, only affect the balance equation of the dual form of f. This experimental observation allows us to identify the system of thermodynamic field equations as an analogue of Maxwell's equations. In the present note we will see that the second postulate is analogue to the absence of magnetic sources in a higher-dimensional version of Maxwell's equations.

The validity of this theory in the weak-field approximation has been tested in many examples of $\alpha - \alpha$ or $\beta - \beta$ processes such as, for example, the thermoelectric effect and the unimolecular triangular chemical reaction (see refs [1] and [2]). In the first case, the TFT provides new predictions when the material is subjected to strong electric field and, in the second, the thermodynamic field equations reproduce the well-known De Donder law.

Transport processes in magnetically confined plasmas, which are of $\alpha - \alpha$ type, have also been analyzed using the TFT. In particular, in ref. [6] the thermodynamic field equations were solved in the weak-field approximation of the classical and the

Pfirsch-Schlüter regimes. We found that the TFT does not correct the expressions for the ionic heat fluxes predicted by the neoclassical theory. On the other hand, the fluxes of matter and electronic energy (heat flow) are enhanced in the nonlinear classical and Pfirsch-Schlüter regimes. This phenomenon would have been amplified had we used the strong (exact) thermodynamic field equations. These results are in line with the experimental observations.

The thermodynamic field equations, in the weak-field approximation, have also been applied to several $\alpha - \beta$ processes. For example the Field-Körös-Noyes model, in which the thermodynamic forces and flows are related by an asymmetric tensor, was analyzed in ref [7]. Even in this case the numerical solutions of the model are in agreement with the theoretical predictions of the TFT. Recently, the Hall effect [8], [9] and, more generally, the galvanomagnetic and thermomagnetic effects have been analyzed in nonlinear regimes. In each of these papers it was shown that the TFT successfully describes the known physics in the nonlinear regimes and also predicts new interesting effects such as the nonlinear Hall effect. The theoretical predictions of the nonlinear Hall effect have been confirmed experimentally. In ref [9] it was shown that, for materials with low thermoelectric power coefficients and in the temperature range of available experimental data, the theoretical predictions of TFT agree with experiments.

When the thermodynamic system approaches equilibrium, the solution of the field equations reduces to the Onsager-Casimir tensor and satisfies the Prigogine theorem of minimum entropy production [10]. Far from equilibrium Glansdorff and Prigogine have demonstrated that, for time independent boundary conditions, a thermodynamic system relaxes to a steady-state satisfying the Universal Criterion of Evolution [11]. Glansdorff and Prigogine obtained this result in 1954 using a purely thermodynamical approach. As indicated in refs [1]-[5], making use of the second principle of thermodynamics, the Shortest Path Principle should ensure the validity of the Universal Criterion of Evolution.

In the present manuscript, using the coordinate independent language of Riemannian geometry, we reformulate the TFT in a generally covariant way. This allows us to extend several weak-field results to the strong-field regime. After expressing the concepts of thermodynamic forces, conjugate flows and entropy production in this new language in sec 2, we go on to study the relaxation of a thermodynamic system in section 3. In particular we prove that if the shortest path principle is valid, then

the Universal Criterion of Evolution, written in a covariant form, is automatically satisfied. We also demonstrate that the term expressing the Universal Criterion of Evolution, written in a covariant form, has a *local minimum* at the geodesic line. Physically this means that a thermodynamic system evolves towards a steady-state with the least possible dissipation. We shall refer to this proposal, together with its corollary the shortest path principle, as the *Minimum Dissipation Principle*.

In section 4 we obtain the thermodynamic field equations. For $\alpha - \alpha$ or $\beta - \beta$ processes we obtain the same equations found in refs [1]-[5], but for general processes, we propose a new set of the field equations, which are conjectured to be also valid beyond the weak-field regime. These equations are notably simpler than the ones found in ref.[4]. In section 5, we establish the equations that determine the steady-states and the critical values of the control parameters of a generic thermodynamic system at which a steady state becomes marginally stable. These equations are also conjectured to hold in the strong-field regime and therefore generalize the weak-field equations reported in refs [3]-[4]. Examples of calculations of the geometric quantities used in the new formulation of TFT can be found in the appendices.

2 Forces, Flows and Entropy Production

The central object of study in Thermodynamic Field Theory (TFT) is the thermodynamic space, which is a smooth, path-connected, geodesically complete, real manifold M equipped with a Riemannian metric g. A Riemannian metric is a positive-definite quadratic form on a manifold's tangent spaces T_xM , which varies smoothly from a point x of the manifold M to another. The metric g yields a smoothly-varying inner product

$$\langle A, B \rangle = A^{\mu} g_{\mu\nu} B^{\nu}, \qquad A, B \in T_x M$$
 (2.1)

where following the Einstein summation convention repeated indices are summed, as they will be in the remainder of this note. The inner product allows one to define the lengths of curves, angles and volumes.

M contains one special point named x=0, which corresponds to the unique equilibrium state. Note however that in practice the thermodynamical forces, which provide local coordinates on patches of M, will be not be independent but rather will satisfy a set of constraints that determines a submanifold N of M. When the submanifold N of physical configurations does not include equilibrium the system

relaxes do a different steady-state.

Given two points $x \in M$ and $y \in M$, we may construct a shortest path $\gamma(t)$ such that

$$\gamma: [t_i, t_f] \to M, \quad \gamma(t_i) = x, \quad \gamma(t_f) = y$$
 (2.2)

The path γ is automatically a geodesic. Clearly for some manifolds and choices of x and y there will be multiple paths γ that minimize the length, thus it is crucial that observable quantities be independent of the choice of path. The metric $g_{\mu\nu}$ allows to define the invariant affine parameter τ :

$$d\tau^2 \equiv g_{\mu\nu} dx^{\mu} dx^{\nu} \tag{2.3}$$

and our manifold M is "metrized" defining the length of an arc L by the formula

$$L = \int_{\tau_1}^{\tau_2} (g_{\mu\nu} \dot{x}^{\mu} \dot{x}^{\nu})^{1/2} d\tau \tag{2.4}$$

The positive definiteness of the matrix $g_{\mu\nu}$ ensures that $d\tau^2 \geq 0$ and allows us to choose an affine parameter τ that increases monotonically as the thermodynamic system evolves to a stable state.

We introduce now the exponential map which is a map from the tangent space T_xM of a Riemannian manifold M at the point x to another point $\exp_x(V)$ in M (see for example ref. [12]). The point $\exp_x(V)$ is the point $\gamma_V(t_f)$ on the geodesic γ that starts at $\gamma_V(t_i) = x$, where it is tangent to $\dot{\gamma}_V(t_i) = V$. t_f is determined by the condition that the arc length along γ from x to $\exp_x(V) = \gamma_V(t_f)$ is the norm $\langle V, V \rangle^{1/2}$ of the vector V. This map is usually denoted by

$$\exp_x(V) = \gamma_V(t_f) \tag{2.5}$$

We next introduce the generalized thermodynamic force vector $U_x \in T_x M$ at each point $x \in M$, which is defined along a shortest path $\bar{\gamma}_{U_x}$ by the relation

$$\exp_x(U) = \bar{\gamma}_{U_x}(t_f) = 0 \tag{2.6}$$

Therefore, the exponential map Eq. (2.6) associates to every tangent vector $U_x \in T_x M$ the equilibrium point $\tilde{\gamma}_{U_x}(t_f) = 0$ in M. A thermodynamic system at x reaches the equilibrium state after traveling along the shortest path $\bar{\gamma}_{U_x}$ with velocity proportional to U_x at the point x. The norm $|U_x|$ indicates the geodesic distance from the point x to the equilibrium point measured along the shortest path $\bar{\gamma}_{U_x}$. The components U_x^{μ}

of U_x are interpreted as the generalized thermodynamic forces. The exponential map is a one-to-one correspondence between a ball in T_xM to a neighborhood of $x \in M$ and so Eq. (2.6) defines a correspondence between the generalized thermodynamic forces and the points of manifold M. From now on we shall omit the suffix x being implicitly understood the dependence of vector U on point x. In Appendix A and Appendix B we will provide two examples of calculations of the map $\exp_x(U)$. In the first example the thermodynamic space M is flat space and so matrix tensor coincides with the Onsager matrix, while in the second M is the two-sphere.

The thermodynamic flows J_{μ} , which are conjugate to the thermodynamic forces U^{μ} , are defined by

$$J_{\mu} = \lambda_{\mu\nu} U^{\nu} \tag{2.7}$$

where $\lambda_{\mu\nu}$ is an asymmetric tensor and, for brevity, we have suppressed the dependence of J, g and U on the point x and geodesic $\bar{\gamma}$. Any 2-tensor may be decomposed into a symmetric and antisymmetric piece. The symmetric piece of λ is identified with the metric tensor g and we shall name the skew-symmetric piece $f_{\mu\nu}$. Eq. (2.7) can then be re-written as

$$J_{\mu} = \lambda_{\mu\nu} U^{\nu} = (g_{\mu\nu} + f_{\mu\nu})U^{\nu} \tag{2.8}$$

 J_{μ} is a one form $J \in T^*M$ which, like U, only vanishes at the equilibrium point x = 0 although there may be points at which both U and J depend on a choice of $\bar{\gamma}$. Eq. (2.8) allows one to generalize the inner product $\langle A, B \rangle$ of two vectors A and B in T_xM from Eq. (2.1) to

$$\langle A, B \rangle \equiv (g_{\mu\nu} + f_{\mu\nu})A^{\mu}B^{\nu} \tag{2.9}$$

Finally we will define the entropy production σ at the point x to be

$$\sigma = \langle U, U \rangle = (g_{\mu\nu} + f_{\mu\nu})U^{\mu}U^{\nu} = g_{\mu\nu}U^{\mu}U^{\nu} \ge 0$$
 (2.10)

where the inequality corresponds to the second law of thermodynamics. As the metric tensor $g_{\mu\nu}$ is a positive definite matrix, this inequality is always satisfied. The entropy production σ is an observable quantity and so the consistency of the formulation demands that it be independent of the arbitrary choice of $\bar{\gamma}$. In fact σ is just the square of the length of the shortest geodesic from x to 0, and so it is independent of the choice among geodesics of equal length and thus is globally-defined as was required.

In this note we will restrict our attention to homogeneous systems, therefore points $x \in M$ will correspond to configurations of the entire system. In particular the entropy flux through the boundaries, will be constant. This implies that any variation in the entropy production σ is equal to the variation of the time derivative of the total entropy. The generalization to inhomogeneous systems is straightforward and follows the strategy employed in Ref. [5].

3 Relaxation to a Steady-State

In the TFT description of a thermodynamic system a homogeneous configuration corresponds to a point x in the thermodynamic space M. The corresponding thermodynamic forces, which are completely determined by the configuration x, are assembled into the vector U defined in Eq. (2.6). If the point x is not a steady-state (see section 5) then the system will evolve. In this section we will consider the process known as relaxation in which the initial velocity vanishes and the system relaxes to a steady-state y.

The TFT description of relaxation rests upon two conjectures

- a) The manifold (M, g) is a Riemannian manifold with metric g;
- b) The Minimum Dissipation Principle.

The Minimum Dissipation Principle will be formulated in Subsec. (3.3), where we will see that it implies that during the process of relaxation the configuration traces out a geodesic γ in thermodynamic space. In Refs. [1]-[4] the geodesic property was referred to as the *shortest path principle*. It allows one to write

$$x = \gamma(0), \qquad y = \gamma(t_f) = \exp_x(V) \tag{3.1}$$

where V is a tangent vector to γ at x whose norm is the geodesic distance from x to y. The system begins at x at time 0 and reaches the steady-state y at the time t_f . The process of relaxation begins as soon as the system is released, that is, immediately after time zero. In support of this conjecture, we will now argue that V automatically satisfies a manifestly covariant form of the Minimum Entropy Production, the Glansdorff-Prigogine Universal Criterion of Evolution and the Minimum Dissipation Principle.

3.1 The Minimum Entropy Production Theorem

In 1945-1947, I. Prigogine proved the *Minimum Entropy Production Theorem* [10], which concerns the relaxation of thermodynamic systems near equilibrium. This theorem states that:

Minimum Entropy Production Theorem: Regardless of the type of processes considered, a thermodynamic system, near equilibrium, relaxes towards a steady-state y in such a way that the inequality

$$\frac{d}{dt}(V_{\mu}V^{\mu}) = \frac{d}{dt}\langle V, V \rangle \le 0 \tag{3.2}$$

is satisfied during throughout the evolution and is only saturated at y. The Minimum Entropy Production Theorem has been proven by I. Prigogine, using a purely thermodynamical approach. We shall now demonstrate that this theorem is automatic in the framework of TFT.

Consider a geodesic $\gamma(\tau)$ parametrized by $\tau \in [\tau_i, \tau_f]$ equipped with a tangent vector field W normalized to have unit length

$$W = \frac{\partial}{\partial \tau} \gamma(\tau) = \gamma^* (\frac{\partial}{\partial \tau}), \qquad \langle W, W \rangle = 1$$
 (3.3)

Notice that τ is not necessarily the time t, as the velocity is not necessarily unity throughout the evolution. The vector field W evaluated at any point x on the curve γ is a vector $W(x) \in T_x \gamma \subset T_x M$. We consider parameterizations of γ such that $W(x) \neq 0$. We then have the following identity

$$\mathcal{L}_{W(\gamma(\tau))}\langle V, V \rangle = \frac{d}{d\tau}(V_{\mu}V^{\mu}) \tag{3.4}$$

where $\mathcal{L}_{W(\gamma(t))}$ denotes the Lie derivative with respect to the vector field W and, since we are near equilibrium, indices are raised and lowered using the Onsager matrix which is a metric for Euclidean space.

There always exists a neighborhood $U \subset M$ of y, which physically corresponds to the near-equilibrium approximation when y is near equilibrium, such that we may uniquely identify the coordinates x^{μ} of a point x with the vectors V(x) in a subset of a tangent fiber. One example of such a U is a ball whose radius is the injectivity radius, which in topologically trivial situations is roughly the inverse square root of the curvature, but any U such that each x and y in U are connected by a unique geodesic will suffice. To show that there is a correspondence between points and

vectors we need to show that given x we can determine V(x) and vice versa. We have already seen that we may uniquely define V(x) from the unique geodesic in U connecting x and y.

To choose a name for the coordinate x given a choice of V(x) we must first confront the fact that V(x) is an element of the tangent space at x, T_xM , and so to compare different V(x)'s we must somehow place them in the same space. This may be done canonically in U using the fact that in U there is a unique geodesic γ_x from x to the steady-state y. Thus we may define the vector $A_y(x) \in T_yM$ by parallel transporting V(x) from x to y along γ . We may now use any choice of coordinates for y to write x on the same coordinate chart

$$y - x = A_y(x) = V(x) \tag{3.5}$$

where we have used parallel transport to define a coordinate system for T_xM which allows us to identify V(x) and A. Notice that each V(x) arises from a unique x because V(x) is parallel to γ and the geodesic equation implies that parallel vectors remain parallel under parallel transport, and so A is also parallel to γ . Thus the uniqueness of the geodesic implies the uniqueness of A and so by Eq. (3.5) the uniqueness of x. The sign in Eq. (3.5) is a result of the fact that V(x) points from x to the point y. In these coordinates the metric is the identity at y, the Christoffel symbols vanish at y, the geodesic curves in U that pass through y are straight lines in U and the entropy production is given by

$$\sigma = |x|^2 + O(x^3) \tag{3.6}$$

Let us now evaluate the Lie derivative of $\langle V, V \rangle$ when the thermodynamic system relaxes from the point x to the stationary point $y = \gamma(t_f)$. According to TFT this relaxation follows the unique geodesic in U that connects them, which is the straight line

$$x = y + V(\tau), \qquad V(\tau) = V(x)f(\tau) \tag{3.7}$$

in the coordinates (3.5). We then find

$$\mathcal{L}_{W(\gamma(\tau))}\langle V(\tau), V(\tau)\rangle = 2\langle V(\tau), \frac{\partial}{\partial \tau} V(\tau)\rangle = 2f(\tau)f'(\tau)\langle V(x), V(x)\rangle \le 0$$
 (3.8)

where the last inequality is a consequence of the fact that $\langle V(x), V(x) \rangle$ is positive by the positive definiteness of the Onsager tensor g, while $f(\tau)$ is positive and $f'(\tau)$ is negative because $f(\tau)$ is the length of V which monotonically falls to zero during the relaxation. Only at the steady-state y is $f(\tau_f) = 0$ and so the inequality is saturated.

3.2 The Universal Criterion of Evolution

The minimum entropy production theorem is generally not satisfied far from equilibrium. However in 1964 P. Glansdorff and I. Prigogine demonstrated that a similar theorem continues to hold for any relaxation to a steady-state. We report the original version of the Glansdorff and Prigogine theorem [11]: When the thermodynamic forces and conjugate flows are related by a generic asymmetric tensor, regardless of the type of processes, for time independent boundary conditions, a thermodynamic system, even in strong non-equilibrium conditions, relaxes to a steady-state in such a way that the following Universal Criterion of Evolution is satisfied

$$(g_{\mu\nu} + f_{\mu\nu})V^{\nu}\frac{dV^{\mu}}{dt} = \langle V, \frac{dV}{dt} \rangle \le 0$$
 $\left(\langle V, \frac{dV}{dt} \rangle = 0 \text{ at the steady state}\right)$ (3.9)

Again Glansdorff and Prigogine have demonstrated this theorem using a purely thermodynamical approach. In this subsection we shall prove that the Shortest Path Principle implies that a covariant form of the Universal Criterion of Evolution is automatically satisfied for relaxation processes in TFT. Consider a geodesic $\gamma(t)$ parametrized by $\tau \in [\tau_i, \tau_f]$ equipped with a tangent vector field W

$$W = \frac{\partial}{\partial \tau} \gamma(\tau) = \gamma^* (\frac{\partial}{\partial \tau}), \qquad \langle W, W \rangle = 1$$
 (3.10)

Again we have parametrized γ so that the velocity vector W always has a unit norm. As the velocity of the relaxation is an observable and is not necessarily equal to one, we must stress that τ refers to a particular parametrization and not necessarily to time. However it is monotonically increasing with respect to time. As usual we also define a second vector field V on the curve γ such that at any point x the vector V(x) satisfies

$$\gamma(\tau_f) = \exp_x(V(x)) \tag{3.11}$$

We will consider curves γ which contain no closed loops, that is curves for which $\gamma(\tau) = \gamma(\tau')$ implies $\tau = \tau'$ and so the condition (3.11) uniquely defines the vector field V.

Now, our task is to evaluate the term in Eq. (3.9) when the thermodynamic system relaxes towards a steady-state along a geodesic. This term contains a time derivative of the vector V, which a priori is ill-defined as V at different times inhabits the tangent fibers over different points of M, which are not canonically isomorphic. The time derivative may be defined using the Lie derivative with respect to the vector W.

This leads to the Covariant Form of the Universal Criterion of Evolution

$$\langle V, \mathcal{L}_W V \rangle \le 0$$
 (with $\langle V, \mathcal{L}_W V \rangle = 0$ only at the steady state) (3.12)

From now on, the term $\langle V, \mathcal{L}_W V \rangle$ will be referred to as the Glansdorff-Prigogine quantity. Notice that $V(\gamma(\tau_f)) = 0$ and so in particular the Glansdorff-Prigogine quantity relaxes to zero

$$\langle V(\gamma(\tau_f)), \mathcal{L}_{W(\gamma(\tau_f))} V(\gamma(\tau_f)) \rangle = 0.$$
 (3.13)

The quantity defined by the Lie derivative differs from the quantity in the minimum entropy production theorem, in fact the difference between these two terms is similar to the difference between the minimum entropy production term and the non-covariant time derivative term. The decomposition for the noncovariant derivative is

$$\frac{d}{d\tau}\langle V, V \rangle = (g_{\mu\nu} + f_{\mu\nu})V^{\nu}\frac{dV^{\mu}}{d\tau} + V^{\mu}\frac{d}{d\tau}[(g_{\mu\nu} + f_{\mu\nu})V^{\nu}]$$

$$= \langle V, \frac{dV}{d\tau} \rangle + V^{\mu}\frac{d}{d\tau}[(g_{\mu\nu} + f_{\mu\nu})V^{\nu}] \qquad (3.14)$$

and for the Lie derivative it is

$$\frac{d}{d\tau}\langle V, V \rangle = \mathcal{L}_W \langle V, V \rangle = \langle V, \mathcal{L}_W V \rangle + V^{\mu} (\mathcal{L}_W(\mathbf{g} + \mathbf{f}) V)_{\mu}$$
(3.15)

We shall now prove the covariant form of the Universal Criterion of Evolution.

Theorem: For all $\tau < \tau_f$ we have the strict inequality $\langle V, \mathcal{L}_W V \rangle < 0$.

Proof: For every point x on the curve γ the vectors V(x) and W(x) are elements of the one-dimensional vector space $T_x\gamma$, therefore they are parallel and there exists a function $f(\gamma(\tau))$ such that

$$V(\gamma(\tau)) = f(\gamma(t))W(\gamma(\tau)). \tag{3.16}$$

We may now evaluate the Lie derivative

$$\mathcal{L}_{W(\gamma(\tau))}V(\gamma(\tau)) = [W(\gamma(\tau)), V(\gamma(\tau))] = [W(\gamma(\tau)), f(\gamma(\tau))W(\gamma(\tau))]$$

$$= [\gamma^*(\frac{\partial}{\partial \tau}), f(\gamma(\tau))\gamma^*(\frac{\partial}{\partial \tau})] = \frac{\partial f(\gamma(\tau))}{\partial \tau}W(\gamma(\tau)). \quad (3.17)$$

Combining Eqs. (3.16) and (3.17) we may evaluate $\langle V, \mathcal{L}_W V \rangle$ as a function of f

$$\langle V, \mathcal{L}_W V \rangle = \langle f W, \frac{\partial f}{\partial \tau} W \rangle = f \frac{\partial f}{\partial \tau} \langle W, W \rangle.$$
 (3.18)

The fact that g is Riemannian and so positive definite implies that $\langle W, W \rangle > 0$. We have seen that $V(\gamma(\tau_f)) = 0$ and we have imposed that $W \neq 0$ therefore $f(\tau_f) = 0$. In addition the fact that $W \neq 0$ implies that the distance from $\gamma(\tau)$ to $\gamma(\tau_f)$ has no critical points and so f has no critical points. Thus f is strictly monotonic in τ and at the maximal value $\tau = \tau_f$ it vanishes. As a result f and $\partial f/\partial \tau$ have opposite signs

$$f\frac{\partial f}{\partial \tau} < 0 \tag{3.19}$$

and so the right hand side of Eq. (3.18) is the product of a positive and a negative term. Therefore $\langle V, \mathcal{L}_W V \rangle$ is negative as desired.

At this point one may object that the Glansdorff-Prigogine quantity in Eq. (3.12) is not actually the quantity that appears in (3.9) because t and τ are not equal and so W is not actually the velocity of the system. This implies that the covariant form of the Universal Criterion of Evolution is not in general equivalent to the original form. However after a little algebra one can see that they are equivalent whenever the deceleration |a|, velocity v and affine length l of the system satisfy the following inequality

$$|a| < \frac{v^2}{l - \tau} \tag{3.20}$$

at every affine time τ . The acceleration is unconstrained, and so this is only a constraint on the dissipation of the system as it approaches the steady-state and slows. However near the steady-state the denominator of the right hand side of Eq. (3.20) shrinks to zero and so even near the end of the relaxation the constraint on the rate of dissipation appears to be quite weak. This constraint will be compared with experiment in a subsequent publication.

3.3 The Minimum Dissipation Principle

In refs [1]-[4] we have introduced the Shortest Path Principle. In Subsec 3.2, we have demonstrated that if the shortest path principle is valid then the Universal Criterion of Evolution, written in a covariant form, is automatically satisfied. In this subsection we will show that the expression $\langle V, \mathcal{L}_W V \rangle$ has a local minimum at the geodesic line. We refer to this theorem, together with the Shortest Path Principle, as the Minimum Dissipation Principle. The Minimum Dissipation Principle corresponds to the following physical conjecture:

For any type of physical process, even in strong non-equilibrium conditions, a thermodynamic system subjected to time-independent boundary conditions relaxes towards a steady-state following the path with the least possible dissipation.

Let us consider an arbitrary path $\gamma(t)$ parameterized by $t \in [0, t_f]$ equipped with a tangent vector field

$$W = \frac{\partial}{\partial t}\gamma(t) = \gamma^*(\frac{\partial}{\partial t})$$
 (3.21)

This time we fix the velocity of the trajectory, this is necessary in any minimization of dissipation because otherwise one could always trivially speed up the dissipation by accelerating the process. In particular we will parametrize $\gamma(t)$ by imposing that W is a unit vector

$$\langle W, W \rangle = 1 \tag{3.22}$$

At every time t there exists at least one geodesic $\alpha_t(\tau)$ such that

$$\alpha_t(0) = \gamma(t), \quad \alpha_t(\tau_f) = \gamma(t_f)$$
 (3.23)

There also exists a vector V_t in the tangent space $T_{\gamma(t)}\alpha_t \subset T_{\gamma(t)}M$ such that

$$\gamma(t_f) = \exp_{\gamma(t)}(V_t) \tag{3.24}$$

Eq. (3.24) defines the vector V_t at every point $\gamma(t)$ on the curve γ and so it defines a vector field $V(\gamma(t))$ on γ with vectors in the tangent space TM. Thus we may define the Lie derivative $\mathcal{L}_{W(\gamma(t))}V(\gamma(t))$ which will be another vector field on γ .

We will show that, using the above parametrization

$$\langle V, \mathcal{L}_W V \rangle = -|V|\cos(\theta)$$
 (3.25)

where θ is the angle between α and γ . This quantity is minimized when $\theta = 0$, corresponding to the case in which α and γ are parallel. Therefore the Glansdorff-Prigogine quantity is minimized everywhere iff α and γ coincide everywhere, in which case γ is a geodesic.

To prove this theorem we shall perform our calculations in Fermi coordinates [13]. In particular, we shall work in the Fermi coordinate system such that the first derivatives of the components of the metric along the geodesic α_t vanish

$$\frac{\partial}{\partial x^{\lambda}} g_{\mu\nu}(\alpha_t) = 0 \tag{3.26}$$




Figure 1: As one moves along the curve γ the minimal distance to $\gamma(t_f)$ shrinks at a speed of $cos(\theta)$. This speed, which is proportional to the negative quantity $\langle V, \mathcal{L}_W V \rangle$, is locally minimized when $\theta = 0$ corresponding to the case in which γ follows a geodesic.

Therefore in Fermi coordinates, a distance ϵ from the geodesic, the components of the metric will differ by a correction of order ϵ^2 :

$$g_{\mu\nu}(\alpha_{t+\epsilon}) = g_{\mu\nu}(\alpha_t) + \frac{\partial g_{\mu\nu}(\alpha_t)}{\partial x^{\lambda}} \epsilon^{\lambda} + \frac{1}{2} \frac{\partial^2 g_{\mu\nu}(\alpha_t)}{\partial x^{\lambda} \partial x^{\kappa}} \epsilon^{\lambda} \epsilon^{\kappa} + O(\epsilon^3)$$
 (3.27)

or

$$\Delta g_{\mu\nu} = g_{\mu\nu}(\alpha_{t+\epsilon}) - g_{\mu\nu}(\alpha_t) = \frac{\partial^2 g_{\mu\nu}(\alpha_t)}{\partial x^{\lambda} \partial x^{\kappa}} \epsilon^{\lambda} \epsilon^{\kappa} + O(\epsilon^3)$$
 (3.28)

In Fermi coordinates, we may interpret the Glansdorff-Prigogine quantity (3.25) using the fact that $\langle V, V \rangle$ is the length squared of the geodesic α_t and its Lie derivative

with respect to W is

$$\mathcal{L}_W\langle V, V \rangle = 2\langle V, \mathcal{L}_W V \rangle + V^{\mu} V^{\nu} (\mathcal{L}_W \mathbf{g})_{\mu\nu}$$
 (3.29)

Notice that the term $\mathcal{L}_W\langle V, V \rangle$ does not correspond to the Lie derivative of the entropy production, which as seen in the section 2, is defined to be $\mathcal{L}_W\langle U, U \rangle$.

We shall now show that the second term on the right-hand side is of order ϵ while the other two terms are of order one

$$V^{\mu}V^{\nu}(\mathcal{L}_W\mathbf{g})_{\mu\nu} \sim O(\epsilon), \qquad \mathcal{L}_W\langle V, V \rangle \sim 2\langle V, \mathcal{L}_W V \rangle \sim O(1)$$
 (3.30)

Up to now, we have defined the vector W only along the curve γ . However, Eq. (3.29) requires a definition of the derivative of W in directions that are not parallel to the curve. Such derivatives have not yet been defined, but they may be defined if we extend the definition of W to a neighborhood U of the point $\alpha_t(0)$. We define W at a point $\alpha_t(0) + \epsilon^{\lambda}$ in this neighborhood by parallel transporting $W(\alpha_t(0))$ along the unique geodesic $\delta x^{\lambda} = \epsilon^{\lambda}$ in U that connects $\alpha_t(0)$ and $\alpha_t(0) + \epsilon^{\lambda}$ [14] (see Fig. (2)):




Figure 2: Definition define the vector W in the neighborhood of the point $\gamma(t)$ by parallel transporting the vector $W(\gamma(t))$ along an edge $\delta x^{\lambda} = \epsilon^{\lambda}$.

$$W_{\epsilon}^{\mu}(x) = W_{0}^{\mu} + \Gamma_{\lambda\kappa}^{\mu}(x)\epsilon^{\lambda}W_{0}^{\kappa} + O(\epsilon^{2})$$
(3.31)

Eq. (3.31) allows us to generalize the definition of the quantity $\langle V, \mathcal{L}_W V \rangle$ to arbitrary curves. The last term of Eq. (3.29) is the derivative of the metric along the direction parallel to α_t . It is easy to check that this term is of order ϵ in our coordinate system:

$$V^{\mu}V^{\nu}(\mathcal{L}_{W}\mathbf{g})_{\mu\nu} = V^{\mu}V^{\nu}W^{\lambda}\frac{\partial g_{\mu\nu}}{\partial x^{\lambda}} + 2V_{\mu}V^{\nu}\frac{\partial W^{\mu}}{\partial x^{\nu}}$$
(3.32)

or

$$V^{\mu}V^{\nu}(\mathcal{L}_{W}\mathbf{g})_{\mu\nu} = 2V_{\mu}V^{\nu}W^{\lambda}\Gamma^{\mu}_{\nu\lambda} + 2V_{\mu}V^{\nu}\frac{\partial\Gamma^{\mu}_{\lambda\kappa}}{\partial x^{\nu}}\epsilon^{\lambda}W^{\kappa}_{0} + O(\epsilon^{2})$$
(3.33)

where we have used the identity

$$\frac{\partial g_{\mu\nu}}{\partial x^{\lambda}} = \Gamma^{\kappa}_{\lambda\mu} g_{\kappa\nu} + \Gamma^{\kappa}_{\lambda\nu} g_{\kappa\mu} \tag{3.34}$$

However in Fermi coordinates, a distance ϵ from the curve the components of the affine connection will differ by a correction of order ϵ :

$$\Gamma^{\lambda}_{\mu\nu}(\alpha_{t+\epsilon}) = \Gamma^{\lambda}_{\mu\nu}(\alpha_t) + \frac{\partial \Gamma^{\lambda}_{\mu\nu}(\alpha_t)}{\partial x^{\kappa}} \epsilon^{\kappa} + O(\epsilon^2) = \frac{\partial \Gamma^{\lambda}_{\mu\nu}(\alpha_t)}{\partial x^{\kappa}} \epsilon^{\kappa} + O(\epsilon^2)$$
 (3.35)

Observe that from Eq. (3.35), we also have that

$$\frac{\partial \Gamma^{\lambda}_{\mu\nu}(\alpha_t)}{\partial x^{\eta}} = \lim_{\epsilon^{\eta} \to 0} \frac{\Gamma^{\lambda}_{\mu\nu}(\alpha_{t+\epsilon}) - \Gamma^{\lambda}_{\mu\nu}(\alpha_t)}{\epsilon^{\eta}} \sim O(1)$$
 (3.36)

Therefore, the second term on the right-hand side of Eq. (3.28) is of order ϵ . As a check, we note that Eq. (3.36) reveals that also the Riemannian curvature tensor $R^{\mu}_{\nu\lambda\kappa}$ is of order O(1), as it must be since our choice of coordinates cannot affect the curvature.

Now, we shall prove that the derivative of the length squared of α_t along the curve γ is of order O(1). For this, we have to compare the lengths of the geodesics α_t and $\alpha_{t+\epsilon}$. This leads us to the notion of Jacobi fields (see, for example, ref. [15]). Let us consider a single infinite family of geodesics in the manifold M. Let τ be a parameter varying along each geodesic of the family, and let t a parameter constant along each geodesic of the family, but varying as we pass from a geodesic to another. A Jacobi field is a vector field on a geodesic that describes infinitesimal deformations that interpolate between different geodesics in a one-parameter family. For example, the family α_t yields the Jacobi field J_t

$$J_t^{\mu}(\tau) = \frac{\partial \alpha_t^{\mu}(\tau)}{\partial t} \tag{3.37}$$

on each geodesic $\alpha_t^{\mu}(\tau)$.

Consider now two nearby geodesics in the α family, α_t and $\alpha_{t+\epsilon}$. We call the points A on α_t and A' on $\alpha_{t+\epsilon}$ corresponding points if they have equal values of τ (see Fig. 3). Let $\lambda_t^{\mu} = \epsilon \sin \theta J_t^{\mu}$ be the infinitesimal Jacobi vector (see Fig. 1) such that [16]

$$q_{\mu\nu}\alpha_t^{\mu}\lambda_t^{\nu} = 0 \tag{3.38}$$

This equation tells us that the deviation λ_t^{μ} is perpendicular to α_t^{μ} . The vector λ_t^{μ} satisfies the Jacobi equation [16]

$$\frac{D}{D\tau^2} \lambda_t^{\mu}(\tau) + R_{\nu\lambda\kappa}^{\mu} \frac{\partial \alpha_t^{\nu}(\tau)}{\partial \tau} \lambda_t^{\lambda}(\tau) \frac{\partial \alpha_t^{\kappa}(\tau)}{\partial \tau} = 0$$
 (3.39)

where D denotes the covariant derivative. Thus both terms in the Jacobi equation (3.39) are linear in ϵ . This implies that, in the limit $\epsilon \to 0$, the second derivative of our Jacobi field goes to zero. The relative angle ϕ at the point A' (see Fig. 3), between α_t and $\alpha_{t+\epsilon}$, measured using parallel transport along the flow of the Jacobi field, can be easily evaluated taking into account the relation [16]

$$(\Delta \alpha_t^{\mu})_{A'}(Y_{\mu})_{A'} = \int \int Y_{\mu} R_{\nu\lambda\kappa}^{\mu} \alpha_t^{\nu} \frac{\partial \alpha_t^{\lambda}}{\partial \tau} \lambda_t^{\kappa} dt d\tau$$
 (3.40)

where Y_{μ} is a vector chosen arbitrarily at A'. Eq. (3.40) shows then that ϕ is of order ϵ .

Our task is to compute the limit of quantity

$$\lim_{\epsilon \to 0} \frac{\Delta \langle V, V \rangle}{\epsilon} = 2g_{\mu\nu} V^{\nu} \lim_{\epsilon \to 0} \frac{\Delta V^{\mu}}{\epsilon} + V^{\mu} V^{\nu} \lim_{\epsilon \to 0} \frac{\Delta g_{\mu\nu}}{\epsilon}$$
(3.41)

Using the relation (3.28), the last term on the right-hand side vanishes. As seen in Fig. 3, to leading order in ϵ , the difference between the vector V^{μ} at the geodesic α_t and its image on the curve $\alpha_{t+\epsilon}$ under $exp_{\alpha_t}(\epsilon \sin(\theta)J_t)$ is of order ϵ^2 . In conclusion, the limit of the difference between the squared length of the geodesic $\alpha_{t+\epsilon}$ and the portion of the geodesic α_t such that $\alpha_{t+\epsilon} = \exp_{\alpha(t)}(\epsilon \sin(\theta)J_t)$, vanishes.

However, the curve $\alpha_{t+\epsilon}$ is not quite the same as $exp_{\alpha_t}(\epsilon \sin(\theta)J_t)$, because we have parameterized $\alpha_{t+\epsilon}$ such that (see Fig. 1)

$$\alpha_{t+\epsilon}(\tau=0) = \gamma(t+\epsilon) \neq \exp_{\gamma(t)}(\epsilon \sin(\theta)J_t(0)).$$
 (3.42)

That is to say that the Jacobi field is not parallel to γ , so it does not take the beginning of the curve α_t to the beginning of the curve $\alpha_{t+\epsilon}$. Instead exp of the Jacobi field




Figure 3: When the relative angle ϕ at the point A' between α_t and $\alpha_{t+\epsilon}$ is of order ϵ , the difference between length A'B' and length AB is of order ϵ^2 . Indeed, we have $A'B' - AB \simeq \frac{1}{2}\phi CB' \sim O(\epsilon^2)$.

 $\epsilon \sin(\theta) J_t(0)$ takes the beginning of the curve α_t to a point $-\epsilon \cos(\theta)$ on the curve $\alpha_{t+\epsilon}$. Up to corrections of order ϵ^2 the curves α_t and $\alpha_{t+\epsilon}$ are the same length, and so the distance from $\alpha_{t+\epsilon}(0)$ to $\gamma(t_f)$ is that of α_t minus the correction $\epsilon \cos(\theta)$ that we have just calculated. The distance squared then differs by $|V|\epsilon \cos(\theta)$. By definition the Lie derivative is this quantity divided by ϵ , which yields the claim (3.25).

4 The Thermodynamic Field Equations

As seen in Eq. (2.10), the second principle of thermodynamics is satisfied if the metric g is positive definite. However the postulates that we have presented so far are insufficient to calculate g. In this section we will add a third postulate to TFT stating that the metric is the solution to a set of field equations which are in turn equal to the variations of an action functional. In short we include the postulate:

c) The Principle of Least Action.

In particular we will assert that the action is invariant with respect to general coordinate transformations and we will not include a cosmological constant term, as there has been no experimental evidence for the presence of such a term in TFT.

We will also place the following three restrictions on our physical systems

- 1) Internal fluctuations are neglected;
- 2) Boundary conditions are time-independent;
- **3)** The system is not subjected to external perturbations.

If these three conditions are simultaneously satisfied then the equations of motion will be *source-free*. The internal fluctuations can be analyzed using the Landau-Lifshitz theory and systems submitted to time-dependent boundary conditions have been examined by York in ref. [19]. Examples of systems subjected to external perturbations, those most commonly encountered in practice, can be found in ref. [20].

4.1 α - α and β - β Processes

In α - α and β - β processes the matrix $\lambda_{\mu\nu}$ that relates the thermodynamic forces and flows (2.7) is symmetric and so is equal to the metric g. We therefore expect the action to be constructed entirely from g. After the cosmological constant, the invariant constructed using the least number of derivatives of g is the Ricci scalar $R \equiv g^{\mu\nu}R_{\mu\nu}$ where $R_{\mu\nu}$ is the Ricci tensor [18]

$$R_{\mu\nu} \equiv \Gamma^{\lambda}_{\mu\lambda,\nu} - \Gamma^{\lambda}_{\mu\nu,\lambda} + \Gamma^{\eta}_{\mu\lambda}\Gamma^{\lambda}_{\eta\nu} - \Gamma^{\eta}_{\mu\nu}\Gamma^{\lambda}_{\eta\lambda}$$
 (4.1)

Of course one could add higher powers of R and its derivatives to the action and maintain general covariance, but so far Eq. (4.1) has been sufficient to reproduce the experimental data and numerical simulations. If such terms are indeed present in the action then, by dimensional analysis, they will be multiplied by characteristic scales of positive spatial dimension and they will be negligible at smaller distance scales. Thus it may well be that above some threshold length scale, corresponding to very strong fields, higher order terms in R will begin to play a role. For example in gravitational theories such terms may play a role deep inside of black holes and immediately after the big bang, but are irrelevant at the scales of all observed phenomena. By contrast in general relativity the nonlinearity of the R term considered here already provides a significant correction to the clocks of the GPS satellites and its effect on Mercury's orbit was observed a hundred years ago.

We therefore postulate that the action I takes the simple form

$$I = \int \sqrt{g} R_{\mu\nu} g^{\mu\nu} d\Omega \tag{4.2}$$

where $d\Omega$ is a volume element and the domain of integration is the entire manifold M. Here g denotes the determinant of the metric $g_{\mu\nu}$. The principle of least action implies that the variation of I with respect to the metric vanishes, which yields the familiar thermodynamic field equations:

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 0 \tag{4.3}$$

For $n \neq 2$ Einstein's equation (4.3) reduces to the Ricci flatness condition

$$R_{\mu\nu} = 0 \tag{4.4}$$

4.2 The General Case

Let us now consider the general case in which $\lambda_{\mu\nu}$ is not necessarily symmetric

$$J_{\mu} = \lambda_{\mu\nu} U^{\nu} = (g_{\mu\nu} + f_{\mu\nu})U^{\nu} \tag{4.5}$$

Any completely skew-symmetric tensor of type (0,2) defines a 2-form \tilde{f} . Given a skew-symmetric tensor f of type (2,0) with components $f^{\mu\nu}$, we can define an (n-2)-form $*\tilde{f}$ called the Hodge dual of f. In terms of components, the relations between $*\tilde{f}$ and f is

$$*\tilde{f}_{j_1 j_2 \dots J_{n-2}} \equiv \frac{1}{2!} g^{1/2} \varepsilon_{\mu \nu j_1 j_2 \dots j_{n-2}} f^{\mu \nu}$$
(4.6)

where $\varepsilon_{\mu\nu j_1j_2...j_{n-2}}$ is the Levi-Civita symbol:

$$\varepsilon_{i,j...n} = \varepsilon^{i,j...n} = \begin{cases} +1 & \text{if } ij \dots n \text{ is an even permutation of } 1, 2 \dots, n \\ -1 & \text{if } ij \dots n \text{ is an odd permutation of } 1, 2 \dots, n \\ 0 & \text{otherwise} \end{cases}$$
(4.7)

The inverse of Eq. (4.6) reads

$$f^{\mu\nu} = \frac{1}{(n-2)!} g^{1/2} \varepsilon^{j_1 j_2 \dots j_{n-2} \mu\nu} (*\tilde{f})_{j_1 j_2 \dots j_{n-2}}$$
(4.8)

To write the kinetic term for the f-field we will need to first define its antisymmetric 3-tensor field strength H

$$H_{\mu\nu\rho} = \partial_{\mu} f_{\nu\rho} + \partial_{\nu} f_{\rho\mu} + \partial_{\rho} f_{\mu\nu} \tag{4.9}$$

or simply

$$\tilde{H} = d\tilde{f} \tag{4.10}$$

where d is the exterior derivative. The square of the exterior derivative, d^2 contains the contraction of two ordinary derivatives with the ϵ tensor and so it vanishes by antisymmetry. Therefore H is closed

$$d\tilde{H} = d^2\tilde{f} = 0 \tag{4.11}$$

In principle the balance equation for \tilde{H} could also contain a source term $\tilde{\mathcal{J}}$

$$d\tilde{H} = \tilde{\mathcal{J}} \tag{4.12}$$

where \mathcal{J} is a closed 4-form, $d\tilde{\mathcal{J}} = 0$.

Given a third-rank skew-symmetric tensor H we can define an (n-3)-form $*\tilde{H}$, the Hodge dual of H, with components

$$*\tilde{H}_{j_1 j_2 \dots J_{n-3}} \equiv \frac{1}{3!} g^{1/2} \varepsilon_{\mu\nu\lambda j_1 j_2 \dots j_{n-3}} H^{\mu\nu\lambda}$$
 (4.13)

As we shall see shortly, $*\tilde{H}$ also satisfies a balance equation with a source term, which we shall denote $*\tilde{J}$. This source term may be nonzero only if the three restrictions on our physical system are relaxed. In other words internal fluctuations, external perturbations and time-dependent boundary conditions can serve as $*\tilde{J}$ sources for $*\tilde{H}$. However in all cases examined so far no potential \mathcal{J} sources have been found for \tilde{H} .

We are therefore led to the following postulate [3],[4]:

d) The 3-form \tilde{H} has no sources or, equivalently, the 3-form \tilde{H} is closed.

Notice that, from the mathematical point of view, to impose $\tilde{\mathcal{J}} = 0$ it is sufficient to require that the tensor field f does not possess any singularities and so Eq. (4.10) is well-defined everywhere.

We will now present an invariant action I for the thermodynamic system. As in the previous subsection, we will restrict our attention to terms that contain the least number of derivatives, as higher derivate terms will be suppressed by a characteristic length scale and so may only be relevant in the presence of very strong fields. Again the metric may enter via an Einstein-Hilbert term equal to the Ricci scalar R. The antisymmetric tensor f may enter via a 2-derivative term $\tilde{H} \wedge *\tilde{H}$ generalizing the Maxwell action and also via a 0-derivative mass term $\chi \tilde{f} \wedge *\tilde{f}$ where χ is a dimensionful parameter. Here \wedge is the wedge product of differential forms and * is the Hodge star defined in Eqs. (4.6) and (4.13). A cosmological constant term and more complicated functions of the mass term are not necessary to explain the data and so will not be included. In components the proposed Lagrangian density is then

$$\mathcal{L}_{kin} = \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} + \frac{\chi}{2} f_{\mu\nu} f^{\mu\nu}$$
 (4.14)

where indices are raised using g. The mass parameter χ is an inverse of the characteristic length squared of the thermodynamic system.

The source $J_{\mu\nu}$ can be included to the theory by adding an interaction term to the Lagrangian density

$$\mathcal{L}_{int} = S_{n-1} J_{\mu\nu} f^{\mu\nu} \tag{4.15}$$

where $S_{n-1} = 2\pi^{n/2}\Gamma(n/2)$ is the area of a unit (n-1)-sphere in n space dimensions $(\Gamma(n/2))$ is the Gamma function). Notice that if we do not add the source J, we could add a new term coupling the source J to the dual field $f^{(dual)}$ where $df^{(dual)} = *H$. However, f and $f^{(dual)}$ are not mutually local, since $f^{(dual)}$ at a point cannot be constructed from f at that point together with a finite number of derivatives and so no local theory can simultaneously contain both J and J. In line with our postulate, we impose J = 0 and so we only consider the source term in Eq. (4.15).

Combining (4.14), (4.15) and the Einstein-Hilbert action we arrive at the total action

$$I = \int \sqrt{g}\mathcal{L} = \int (\sqrt{g}R + \frac{1}{12}\tilde{H} \wedge \star \tilde{H} + \frac{\chi}{2}\tilde{f} \wedge \star \tilde{f} + S_{n-1}\tilde{J} \wedge \star \tilde{f})$$
(4.16)

The relative normalization of the first two terms in the action (4.16) may be modified by rescaling the metric, here we have chosen the convention known as the string frame in string theory [21].

*J will be treated as an external source, and so there are two variables whose variations yield the field equations. The variation with respect to the metric yields a modified Einstein's equation

$$R_{\mu\nu}-\frac{1}{2}g_{\mu\nu}R=-\frac{1}{4}(H_{\mu\lambda\rho}H_{\nu}^{\lambda\rho}-\frac{1}{6}g_{\mu\nu}H_{\lambda\kappa\rho}H^{\lambda\kappa\rho})-\chi\!\left(f_{\mu\kappa}f_{\nu}^{\kappa}-\frac{1}{4}g_{\mu\nu}f_{\lambda\kappa}f^{\lambda\kappa}\right)+\frac{S_{n-1}}{2}g_{\mu\nu}J_{\lambda\kappa}f^{\lambda\kappa}$$

$$(4.17)$$

while the variation with respect to the f-field yields

$$d * \tilde{H} + \chi * \tilde{f} = -S_{n-1} * \tilde{J} \tag{4.18}$$

We have seen that the antisymmetric tensor f modifies the field equations for the metric. One may wonder if it also affects the shortest path principle. In general a p-tensor potential can be covariantly coupled to a p-dimensional J-source and to a (d-p-2)-dimensional \mathcal{J} -source. For example in 4-dimensional Maxwell theory the 1-tensor potential couples to electric and magnetic particles, while in 5 dimensions the electric sources are still particles but the magnetic monopoles are strings. In our case p=2, so the J-source is a string and the \mathcal{J} -source is (d-4)-dimensional. In particular, the trajectories in the manifold of thermodynamic configurations are, being paths, 1-dimensional and so they cannot contain the J-source. Except in 5-dimensions they also cannot contain the \mathcal{J} -source. Thus, there is no covariant coupling of the trajectories to the f-field, as there would have been for an ordinary vector potential.

Concretely, this means that the action for a relaxing configuration is proportional to the proper length of its trajectory

$$S \propto \int d\tau \left(\frac{dX^{\mu}}{d\tau} \frac{dX_{\mu}}{d\tau}\right)^{1/2} \tag{4.19}$$

just as in the case with no f-field. The proper length of a path, by definition, is extremized by a geodesic. Thus the solutions of the classical equations of motion for the evolution of a state are still geodesics in thermodynamic space, even when we include the f field. This is not to say that the solutions with an f field are just the solutions to the system without the f-field, because the new Einstein equation (4.17) no longer admits Ricci-flat metrics, and so the Riemannian manifold M itself is changed by the inclusion of f.

5 Steady-States and Stability Criteria

In the thermodynamic theory of irreversible processes, a steady-state of order κ , where k is less than or equal to the dimension n of our thermodynamic space M, is a thermodynamic point x with coordinates

$$U^1, \ U^2 \dots U^{\kappa}$$
 kept constant $(\kappa \le n)$
$$J_a = 0 \qquad (a = \kappa + 1, \dots n) \qquad (5.1)$$

From now on, Latin indices will run from $\kappa + 1$ to n and Greek indices will run from 1 to n. Thus (5.1) may be written more compactly as

$$J_{\mu}dU^{\mu}\mid_{st.state}=0\tag{5.2}$$

where symbol d denotes the total differential on the submanifold N of solutions of the constraint equations.

Steady-states may be stable or unstable. According to the Layapunov stability theory the stability of a state is determined as follows [11]

if
$$\frac{d}{dt}\delta^2 S \mid_{U_0^{\mu}} < 0$$
 then U_0^{μ} is unstable for $t \ge t_0$
if $\frac{d}{dt}\delta^2 S \mid_{U_0^{\mu}} > 0$ then U_0^{μ} is asymptotically stable for $t \ge t_0$ (5.3)

where S denotes the total entropy of the thermodynamic system. In practice we can modify the value of $\delta^2 S$ by varying a series of control parameters λ . When these parameters reach a critical value λ_c , the sign of the inequality (5.3) will change, and a steady-state will lose its stability. A steady-state in a configuration with critical control parameters is referred to as a state with marginal stability:

$$\frac{d}{dt}\delta^2 S \mid_{U_0^{\mu}} (\lambda_c) = 0 \quad \text{for } t \ge t_0$$
 (5.4)

The total entropy S is related to the entropy production σ considered in this note via the entropy balance equation

$$\frac{d}{dt}S = \sigma + \Sigma_S \tag{5.5}$$

where Σ_S denotes the entropy flux through the boundaries. Eq. (5.5) may be varied twice to obtain

$$\frac{d}{dt}\delta^2 S = \delta^2 \sigma + \delta^2 \Sigma_S \tag{5.6}$$

where the variation of the Σ_S term vanishes because we consider homogeneous configurations in which Σ_S is constant. In ref. [11], the authors considered inhomogeneous thermodynamic systems subject to boundary conditions in which all fluxes vanish at infinity. We are instead considering finite homogeneous systems for which the entropy flux through the boundaries, Σ_S is constant. Notice that in the particular case $\Sigma_S = 0$ the only steady-state is equilibrium. In all other cases our systems do not satisfy their boundary conditions, as they are considering all of spacetime whereas we

consider only a region in which the system is homogeneous. As a result simplifications that occur in ref. [11] which rely upon their choice of boundary conditions do not apply here, but they will not be needed. Thus in our case the stability condition is

if
$$\delta^2 \sigma \mid_{U_0^{\mu}} < 0$$
 then U_0^{μ} is unstable for $t \ge t_0$
if $\delta^2 \sigma \mid_{U_0^{\mu}} > 0$ then U_0^{μ} is asymptotically stable for $t \ge t_0$ (5.7)

These conditions may be interpreted geometrically as follows. Recall that the point x is in an j dimensional submanifold $N \subset M$ of configurations that satisfy a set of physical constraints that relate the thermodynamic forces. Then x is a steady state of order κ if $j = n - \kappa$ and if the derivatives of σ with respect to all tangent vectors to N vanish. Diagonalizing the matrix of second derivatives we may also define the stable directions to be the positive eigenspace, the unstable directions to be the negative eigenspace, and the marginally stable directions to be the null eigenspace.

The steady-state condition (5.2) is not canonically defined as the U's at various points x live in different spaces. However using the geometric interpretation of steady-states as critical points of S one arrives at a unique covariant definition for steady-states, and thus a definition that we conjecture will hold beyond the weak-field regime. The steady states are critical points of σ which means that $d\sigma$ vanishes where d is the exterior derivative on the submanifold N. As σ is a scalar, the ordinary and covariant derivatives of σ are equal and so

$$0 = d\sigma = D\sigma = D(U^{\mu}U^{\nu}g_{\mu\nu}) = 2g_{\mu\nu}U^{\nu}DU^{\mu}$$
 (5.8)

where the f term vanishes because f is antisymmetric and UU is symmetric and Dg vanishes because the covariant derivative of the metric always vanishes. Thus the covariant form of the steady-state condition is just

$$g_{\mu\nu}U^{\nu}DU^{\mu} = 0. \tag{5.9}$$

The stability condition (5.7) is already covariant, as S is a scalar and so ordinary derivatives on S are canonically defined. But for completeness we evaluate the double variation

$$\delta^2 \sigma = \frac{\partial^2 \sigma}{\partial x^a \partial x^b} \delta x^a \delta x^b \tag{5.10}$$

where

$$\frac{\partial^{2} \sigma}{\partial x^{a} \partial x^{b}} = 2g_{\mu\nu} U^{\mu}_{;a} U^{\nu}_{;b} + 2f_{\mu\nu;a} U^{\nu}_{0} U^{\mu}_{;b} + 2(g_{\mu\nu} + f_{\mu\nu}) U^{\nu}_{0} \left(U^{\mu}_{;a;b} + \Gamma^{\kappa}_{ab} U^{\mu}_{;\kappa} + U^{\mu}_{;\kappa} \frac{\partial^{2} x^{\kappa}}{\partial x^{a} \partial x^{b}} \right)$$
(5.11)

We have again used the fact that g is covariantly constant.

There are marginally stable states when the determinant of the symmetric matrix $\frac{\partial^2 \sigma}{\partial x^a \partial x^b}$ vanishes

$$\left| (2g_{\mu\nu} + f_{\mu\nu}) U_0^{\nu} \left(U_{;a;b}^{\mu} + \Gamma_{ab}^{\kappa} U_{;\kappa}^{\mu} + U_{;\kappa}^{\mu} \frac{\partial^2 x^{\kappa}}{\partial x^a \partial x^b} \right) + 2g_{\mu\nu} U_{;a}^{\mu} U_{;b}^{\nu} + f_{\mu\nu;a} U_0^{\nu} U_{;b}^{\mu} + f_{\mu\nu;b} U_0^{\nu} U_{;a}^{\mu} \right|_{\lambda = \lambda_c}$$

$$(5.12)$$

We expect that, even beyond the weak-field approximation, Eqs (5.9) and (5.12) will respectively determine the steady-states U_0 and the critical values λ_c of control parameter λ .

6 Conclusions

We have presented a manifestly covariant and coordinate-independent formulation of the Thermodynamic Field Theory (TFT). This allows the theory to be extended, for example, to thermodynamic spaces M that cannot be covered by a single coordinate patch corresponding to theories for which different observable quantities are suitable in different regimes. Making use of the Shortest Path Principle, we have demonstrated the validity of the Universal Criterion of Evolution, expressed in a covariant form, and we have shown that this term has a local minimum at the geodesic line. We referred to this theorem, together with the Shortest Path Principle, as the Minimum Dissipation Principle. Physically, the Minimum Dissipation Principle affirms that, for time-independent boundary conditions, a thermodynamic system evolves towards a steady-state with the least possible dissipation.

Of course the test of any proposed formulation is its agreement with experiment. One physical system for which a large amount of data is already available, and a theoretical underpinning is lacking, is the relaxation of a magnetically confined plasma towards a steady-state in the nonlinear classical and Pfirsch-Schlüter regimes or in the nonlinear Banana and Plateau regimes. The metrics in these regimes can be found in [8].

We have also derived the thermodynamic field equations. For the $\alpha - \alpha$ or $\beta - \beta$ processes, these equations reduce to the ones already found in refs [1]-[4]. The valid-

ity of these equations, in the weak-field approximation, have been tested analyzing several thermodynamic systems. So far, we have found that numerical simulations and experiments are in agreement with the theoretical predictions of the TFT. When the skew-symmetric tensor $f_{\mu\nu}$ does not vanish, we have proposed a new set of field equations, which are conjectured to hold even beyond the weak-field regime. The strong field equations are, however, notably simpler than the ones obtained in ref. [4].

The equations that determine the steady-states and the critical values of the control parameters of a generic thermodynamic system have also been presented. These equations are also conjectured to hold beyond the weak-field regime and so they generalize the weak-field equations reported in refs [2]-[4]. We note that this extension of linear thermodynamics is quite different from the Prigogine-Glansdorff extension, which is based on hydro- (or plasma-) dynamical stability theory and bifurcation analysis [11]. The relation between the two approaches should be made explicit. This will be the subject of a further publication.

The current approach describes the trajectory followed by a relaxing system, but does not give the time dependence of the relaxation process. It may be possible to include the time direction in the thermodynamic space with a metric such as $g_{tt} = -\sigma$. The geodesic evolution in spacetime would then give a firm prediction for the time dependence of the spatial trajectory. In addition the steady-state condition and the Layapunov stability conditions may be derived from the constraint that particles follow spacetime geodesics. Of course in order for the system to stop at a steady state somehow dissipation will need to be added to the system, once this is done it may be possible to apply TFT to processes other than relaxation.

Appendix A Example: Flat Space

Let $M=\pi$ be a plane. We will compute the map $\exp:T\pi\mapsto\pi$. In this case the Riemannian manifold (M,g) coincides with (M,L) where L is the Onsager matrix. There is a well-defined exponential map from a ball \mathcal{U}_x in $T_x\pi$ to a neighborhood $\bar{\mathcal{U}}_x$ of $x\in\pi$:

$$\mathcal{U}_x \subset T_x \pi = R^2 \to \bar{\mathcal{U}}_x \subset \pi = R^2$$
 (A.1)

When the metric is flat, the geodesics $\gamma_V(\tau)$ are straight lines

$$x_f^{\mu} = \tau A^{\mu} + x^{\mu} \tag{A.2}$$

where A^{μ} is a constant vector and $A \in T_x \pi$, $\gamma_A(t_i) = x$ and $\gamma_A(t_f) = x_f$. Therefore the exp map on x is simply

$$\exp_x \tau A = \tau A + x \tag{A.3}$$

that is, the exponential map $\exp_p V$ (with $p \in \pi$ and $V \in T_p \pi$) simply performs a translation:

$$\exp_p V = p + V \tag{A.4}$$

Let us now consider the case $U \in \mathcal{U}_x \subset T_0M = R^2$ where $0 \in \pi$ denotes the origin, corresponding to the equilibrium state. In this case U points from x to the origin and, using the canonical identification of the plane π and its tangent fibers, which are planes of the same dimension, we find the relation x = -U (see Eq. (3.5)). If the coordinates of x are x = (a, b), the coordinates of U are then U = (-a, -b) and, from Eq. (A.4), we have

$$\exp_p U = p + U = (a, b) + (-a, -b) = 0 \tag{A.5}$$

This implies that the exponential map sends the tangent vector U at p to the equilibrium state, which is the definition of U in Eq. (2.6).

The entropy production of a system at the point x is just its length squared

$$\sigma = |U|^2 = |-x|^2 = |x|^2 \tag{A.6}$$

Appendix B Example: The Two-Sphere

Let M be the unit two-sphere S^2 with the usual rotation-invariant metric

$$ds^2 = d\theta^2 + \sin^2\theta d\phi^2. \tag{B.1}$$

We will use both spherical and Cartesian coordinate systems, which are related by

$$x = \sin \theta \cos \phi, \quad y = \sin \theta \sin \phi, \quad z = \cos \theta.$$
 (B.2)

The geodesics on the 2-sphere are the great circles, which are the equators corresponding to various choices of north pole. We will now fix the north pole to be $\theta = 0$, and so every geodesic γ will intersect the equator $\theta = \pi/2$ at two points $\phi = -\delta$ and $\phi = \pi - \delta$. Note that the equator itself is the geodesic that intersects itself an infinite number of times, to apply the following derivation to the equator itself it would be

necessarily to deform it by a small rotation and then take that rotation to zero at the end. However the final result will make sense for every geodesic, including the equator.

The map $exp(\vec{v})$ is easy to calculate in the case in which γ is the equator, it just increases ϕ by the norm v of the vector \vec{v} . To evaluate exp on an arbitrary point (θ_0, ϕ_0) with an arbitrary tangent vector \vec{v} , which yields an arbitrary geodesic γ , it will be simplest to rotate the coordinate system so that γ becomes the equator, then exp may be evaluated and the coordinate system may be rotated back to yield the final answer.

The rotation of the coordinate system will proceed in two steps. First we use δ to define a rotated set of coordinates

$$\tilde{x} = \sin \theta \cos(\phi + \delta), \quad \tilde{y} = \sin \theta \sin(\phi + \delta), \quad \tilde{z} = z = \cos \theta.$$
 (B.3)

Then we rotate about the \tilde{y} -axis until γ is the equator, which we parametrize with the azimuthal variable τ

$$\begin{pmatrix} \tilde{x} \\ \tilde{y} \\ \tilde{z} \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \cos \tau \\ \sin \tau \\ 0 \end{pmatrix}. \tag{B.4}$$

In coordinates the rotation (B.4) is

$$\sin \theta \cos(\phi + \delta) = \cos \tau$$
, $\sin \theta \sin(\phi + \delta) = \cos \alpha \sin \tau$, $\cos \theta = \sin \alpha \sin \tau$. (B.5)

The tangent vector \vec{v} will be decomposed in polar coordinates (v, β) where v is the norm of \vec{v} and β is the angle between \vec{v} and the azimuthal direction ϕ . In this section we will calculate $exp_{(\theta_0,\phi_0)}\vec{v}$. First we will find expressions satisfied by the unknown angles τ and α in terms of the given data θ_0 , ϕ_0, v, β . These will allow us to calculate the z component z' of the function exp, which is particularly simple as it is independent of δ . Then we will rotate the coordinates to construct the other components x' and y'.

 \vec{v} is a tangent vector to the geodesic γ and so the rotation (B.4) relates \vec{v} to the equator's tangent vector $\partial/\partial\tau$. For example the last expression in (B.5) yields

$$\frac{\partial}{\partial \tau} \cos \theta = \frac{\partial}{\partial \tau} \sin \alpha \sin \tau \tag{B.6}$$

where the derivative is evaluated at (θ_0, ϕ_0) . The left hand side may be evaluated using the chain rule, where we note that the derivative of θ with respect to τ along the geodesic is just the theta component $\sin \beta$ of the unit tangent vector \vec{v}/v . The right hand side is evaluated by noting that α is the angle by which the entire geodesic γ is rotated, and so it is independent of τ . Thus (B.6) reduces to

$$\sin \beta \, \sin \theta_0 = \sin \alpha \, \cos \tau. \tag{B.7}$$

The dependence on the unknown angle α may be removed using the last term in Eq. (B.4). We then divide both sides of the resulting equation by $\cos \theta_0$ to obtain

$$\cot \tau = \sin \beta \ \tan \theta_0. \tag{B.8}$$

Thus we have determined the unknown angle τ in terms of the initial point and tangent vector. While in principle we could also find α , it will suffice to recall the last term from (B.5)

$$\sin \alpha = \cos \theta_0 \, \csc \tau. \tag{B.9}$$

In our new coordinates the exp operation corresponds to a simple shift of the equator's longitudinal coordinate τ

$$\tau \mapsto \tau + v.$$
 (B.10)

Finally we need to reexpress this shift in terms of the old coordinates. In fact it will suffice to find the corresponding coordinate z'. Using the rotation (B.4) at the new coordinate $\tau + v$

$$\begin{pmatrix} x' \\ y' \\ z' \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} \cos(\tau + v) \\ \sin(\tau + v) \\ 0 \end{pmatrix}. \tag{B.11}$$

we find

$$z' = \sin \alpha \ \sin(\tau + v). \tag{B.12}$$

This may be expanded using (B.9) and the trigonometric identity for the sine of a sum to yield

$$z' = \cos \theta_0 \, \csc \tau (\sin \tau \, \cos v + \cos \tau \, \sin v) = \cos \theta_0 (\cos v + \cot \tau \, \sin v). \tag{B.13}$$

Finally we may use (B.8) to obtain

$$z' = \cos \theta_0 \cos v + \sin \beta \sin \theta_0 \sin v. \tag{B.14}$$

To find the corresponding formula for x' we may rotate the coordinates by $\pi/2$, yielding a new z coordinate, \tilde{z} , equal to the old x coordinate

$$\tilde{z} = \cos \tilde{\theta} = \sin \theta \cos \phi = x.$$
 (B.15)

This rotation is an isometry so v is unchanged, however the angle β is effected. The rotated value of β may be found similarly to (B.6)

$$\frac{\partial}{\partial \tau} \cos \tilde{\theta} = \sin \tilde{\beta} \sin \tilde{\theta} = \frac{\partial}{\partial \tau} (\sin \theta \cos \phi) = \sin \beta \cos \theta \cos \phi - \cos \beta \sin \phi. \quad (B.16)$$

In the last term of the last expression of (B.16) we have used

$$\cos \beta = g_{\phi\phi}^{1/2} \frac{\partial \phi}{\partial \tau}, \quad g_{\phi\phi} = \sin^2 \theta.$$
 (B.17)

We may now use (B.15) and (B.6) to rotate our equation (B.14) for the z component of the exp function to obtain the x component

$$x' = \sin \theta_0 \cos \phi_0 \cos v + (\sin \beta \cos \theta_0 \cos \phi_0 - \cos \beta \sin \phi_0) \sin v. \tag{B.18}$$

And at last we may rotate ϕ to obtain the y component

$$y' = \sin \theta_0 \sin \phi_0 \cos v + (\sin \beta \sin \theta_0 \cos \phi_0 + \cos \beta \cos \phi_0) \sin v.$$
 (B.19)

Putting together Eqs. (B.14,B.18,B.19) we obtain

$$exp_{(\theta_0,\phi_0)}(v,\beta) = (x',y',z').$$
 (B.20)

Taking the north pole to be equilibrium, the thermodynamic force at any point is calculated using the geodesic that runs straight north from any point to the north pole

$$U = (v = \theta, \beta = 0). \tag{B.21}$$

This geodesic is uniquely determined for all points except for the north pole $\theta = \pi$, for which there are an infinite number of northward geodesics and the coordinate β is ill-defined. The entropy production is then simply

$$\sigma = \langle U, U \rangle = \theta^2 \tag{B.22}$$

Note that at the south pole, where the geodesic is not unique, σ is independent of the geodesic chosen and so continues to be well defined.

Acknowledgments

We acknowledge our indebtedness to Prof. C.M. Becchi of the University of Genova - Physics Department and to Prof. E. Tirapegui of the University of Chile - Physics Department, for reading this manuscript and making helpful suggestions.

One of us (GS) would like to express his sincere gratitude to his hierarchy at the European Commission: Prof. A. Mitsos, Dr P. Fernandez Ruiz and Dr E. Rille. He is also very grateful to Dr U. Finzi and Dr M. Cosyns, of the European Commission, for their continuing encouragement.

GS would like to thank the members of the EURATOM Belgian State Fusion Association, in particular Dr D. Carati, Dr B. Knaepen and Dr B. Weyssow.

The work of JE is partially supported by IISN - Belgium (convention 4.4505.86), by the "Interuniversity Attraction Poles Programme – Belgian Science Policy" and by the European Commission RTN program HPRN-CT-00131, in which he is associated to K. U. Leuven.

References

- [1] Sonnino G. Il Nuovo Cimento, **115** B, (2000), 1057.
- [2] Sonnino G. 2001 Thermodynamic Field Theory (An Approach to Thermodynamics of Irreversible Processes), proceedings of the 9th International Workshop on Instabilities and Nonequilibrium Structures, Viña del Mar (Chile). Kluwer Academic Publishers, Nonlinear Phenomena and Complex Systems Editions, 9, (2004) 291.
- [3] Sonnino G. 2002 A Field Theory Approach to Thermodynamics of Irreversible Processes, (Thèse d'Habilitation à Diriger des Recherches H.D.R.) Institut Non Linèaire de Nice (France).
- [4] Sonnino G. Il Nuovo Cimento, 118 B, (2003) 1175.
- [5] Sonnino G. Int. J. of Quantum Chemistry 98 (2004) 191.
- [6] Sonnino G. 2005 A Field Theory Approach to Transport Processes in Confined Plasmas: the Nonlinear Classical and Pfirsch-Schlüter Regimes and Transport

- Equations in the Nonlinear Banana and Plateau Regimes (submitted to publication in the review I.A.E.A. Nuclear Fusion).
- [7] Peeters Ph. and Sonnino G. Il Nuovo Cimento 115 B (2000) 1083.
- [8] Sonnino G. Nuovo Cimento 118 B (2003) 1155.
- [9] Sonnino G. and Peeters Ph. Chaos 14 (2004) 910.
- [10] Prigogine I. 1954 Thermodynamics of Irreversible processes, (John Wiley & Sons).
- [11] Glansdorff P. and Prigogine I. 1971 Thermodynamics of Structures, Stability and Fluctuations, (John Wiley & Sons).
- [12] Katok A. and Hasselblatt B. 1995 Introduction to the Modern Theory of Dynamical Systems, (Cambridge University Press).
- [13] Fermi F. Atti R. Accad. Lincei, Cl. Sc. Fis. Mat. Nat. 31 (1922) 21.
- [14] Weinberg S. 1972 Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity, (John Wiley & Sons).
- [15] Frankel T. 1997 The Geometry of Physics An Introduction, (Cambridge University Press).
- [16] Synge J.L. and Schild A. 1949 Tensor Calculus, (Dover Publications, Inc. New York).
- [17] Eisenhart L.P. 1997 Riemannian Geometry, (Eight printing, for Princeton Landmarks in Mathematics and Physics series Princeton University Press).
- [18] Lovelock D. and Rund H. 1989 Tensors, Differential Forms and Variational Principles, (Dover Publications, Inc., New York).
- [19] York J.W. Jr. Found. Phys., 16, (1986) 249.
- [20] Horsthemke H. and Lefever R. 1984 Noise-Induced Transitions. Theory and Applications in Physics, Chemistry and Biology, (Springer-Verlag).
- [21] Green M.B., Schwarz J.H. and Witten E. 1987 Superstring Theory. Volume 1 Introduction, (Cambridge University Press).